

Pyšek P, Richardson DM (2007) Traits associated with invasiveness in alien plants: Where do we stand? In: Nentwig W. (ed), Biological invasions, Ecological Studies 193, Springer-Verlag, Berlin & Heidelberg, pp 97–126

## References to Fig. 7.2

- Anttila CK, Daehler CC, Rank NE, Strong DR (1998) Greater male fitness of a rare invader (*Spartina alterniflora*, Poaceae) threatens a common native (*Spartina foliosa*) with hybridization. *Amer J Bot* 85:1597-1601
- Baker HG (1965) Characteristics and modes of origin of weeds. In: Baker HG, Stebbins GL (eds) *The genetics of colonizing species*, Academic Press, New York, pp 147-172
- Baruch Z, Bilbao B (1999) Effects of fire and defoliation on the life history of native and invader C4 grasses in a Neotropical savanna. *Oecologia* 119:510-520
- Baruch Z, Gómez JA (1996) Dynamics of energy and nutrient concentration and construction cost in a native and two alien C4 grasses from two neotropical savannas. *Plant and Soil* 181:175-184
- Baruch Z, Ludlow MM, Davis R (1985) Photosynthetic responses of native and introduced C4 grasses from Venezuelan savannas. *Oecologia* 67:288-293
- Bilbao B, Medina E (1990) Nitrogen-use efficiency for growth in a cultivated African grass and a native South American pasture grass. *J Biogeogr* 17:421-425
- Bímová K, Mandák B, Pyšek P (2003) Experimental study of vegetative regeneration in four invasive *Reynoutria* taxa. *Plant Ecol* 166:1-16
- Black RA, Richards JH, Manwaring JH (1994) Nutrient uptake from enriched soil microsites by three great basin perennials. *Ecology* 75:110-22
- Burns JH (2004) A comparison of invasive and non-invasive dayflowers (Comelinaceae) across experimental nutrient and water gradients. *Diversity Distrib* 10:387-397
- Caldwell MM, Richards JH, Johnson DA, Nowak RS, Dzurec RS (1981) Coping with herbivory: photosynthetic capacity and resource allocation in two semiarid *Agropyron* bunchgrasses. *Oecologia* 50:14-24
- Callaway JC, Josselyn MN (1992) The introduction and spread of smooth cordgrass (*Spartina alterniflora*) in South San Francisco Bay. *Estuaries* 15:218-225
- Deng X, Ye W-H, Feng H-L, Yang Q-H, Cao H-L, Xu K, Zhang Y (2004) Gas exchange characteristics of the invasive species *Mikania micrantha* and its indigenous congener *M. cordata* (Asteraceae) in South China. *Bot Bull Acad Sin* 45:213-220

- Durand ZL, Goldstein G (2001) Photosynthesis, photoinhibition, and nitrogen use efficiency in native and invasive tree ferns in Hawaii. *Oecologia* 126:345-354
- Forcella F, Wood JT, Dillon SP (1986) Characteristics distinguishing invasive weeds within *Echium* (Bugloss). *Weed Res* 26:351-364
- Gerlach JD, Rice KJ (2003) Testing life history correlates of invasiveness using congeneric plant species. *Ecol Appl* 13:167-179
- Goergen E, Daehler CC (2001a) Reproductive ecology of native Hawaiian grass (*Heteropogon contortus*; Poaceae) versus its invasive alien competitor (*Pennisetum setaceum*; Poaceae). *Int J Plant Sci* 162:317-326
- Goergen E, Daehler CC (2001b) Inflorescence damage by insects and fungi in native Pili grass (*Heteropogon contortus*) versus alien fountain grass (*Pennisetum setaceum*) in Hawaii. *Pacific Sci* 55:129-136
- Goergen E, Daehler CC (2002) Factors affecting seedling recruitment in an invasive grass (*Pennisetum setaceum*) and a native grass (*Heteropogon contortus*) in the Hawaiian Islands. *Plant Ecol* 161:147-156
- Grotkopp E, Rejmánek M, Rost TL (2002) Towards a casual explanation of plant invasiveness: Seedling growth and life history strategies of 29 pine (*Pinus*) species. *Amer Natur* 159:396-419
- Gruberová H, Bendová K, Prach K (2001) Seed ecology of alien *Bidens frondosa* in comparison with native species of the genus. In: Brundu G, Brock J, Camarda C, Child L, Wade M (eds), *Plant invasions: Species ecology and ecosystem management*, Backhuys Publishers, Leiden, pp 99-104
- Harris GA (1967) Some competitive relationships between *Agropyron spicatum* and *Bromus tectorum*. *Ecol Monogr* 37:89-111
- Holmes TH, Rice KJ (1996) Patterns of growth and soil-water utilization in some exotic annuals and native perennial bunchgrasses of California. *Ann Bot* 78:233-243
- Honig MA, Cowling RM, Richardson DM (1992) The invasive potential of Australian banksias in South African fynbos: A comparison of the reproductive potential of *Banksia ericifolia* and *Leucadendron laureolum*. *Austr J Ecol* 17:305-314
- Kloeppel BD, Abrams MD (1995) Ecophysiological attributes of the native *Acer saccharum* and the exotic *Acer platanoides* in urban oak forests in Pennsylvania, USA. *Tree Physiol* 15:739-746
- Kolb A, Alpert P (2003) Effects of nitrogen and salinity on growth and competition between a native grass and an invasive congener. *Biol Invas* 5:229-238

- Lambrecht-McDowell SC, Radosevich SR (2005) Population demographics and trade-offs to reproduction of an invasive and noninvasive species of *Rubus*. *Biol Invas* 7:281-295
- Lambrinos JG (2001) The expansion history of a sexual and asexual species of *Cortaderia* in California, USA. *J Ecol* 89:88-98
- Lambrinos JG (2002) The variable invasive success of *Cortaderia* species in a complex landscape. *Ecology* 83:518-529
- Larson KC (2000) Circumnutation behavior of an exotic honeysuckle vine and its native congener: Influence on clonal mobility. *Amer J Bot* 87:533-538
- Mandák B (2003) Germination requirements of invasive and non-invasive *Atriplex* species: a comparative study. *Flora* 198:45-54
- Matsuo K (1999) Comparison of ecological distributions and life history characteristics between invasive and closely related native *Plantago* species in Japan. In: Yano E, Matsuo K, Shiyomi M, Andow DA (eds) *Biological invasions of ecosystems by pests and beneficial organisms*, National Institute of Agro-Environmental Sciences Series 3, Tsukuba, Japan, pp 15–26
- McDowell SCL (2002) Photosynthetic characteristics of invasive and noninvasive species of *Rubus* (Rosaceae). *Amer J Botany* 89:1431-1438
- Mihulka S, Pyšek P, Martínková J (2003) Invasiveness of *Oenothera* congeners in Europe related to their seed characteristics. In: Child LE et al. (eds), *Plant invasions: Ecological threats and management solutions*, Backhuys Publishers, Leiden, pp 213–225
- Morris LL, Walck JL, Hidayati SN (2002) Growth and reproduction of the invasive *Ligustrum sinense* and native *Forestiera ligustrina* (Oleaceae): implications for the invasion and persistence of a nonnative shrub. *Int J Plant Sci* 163:1001-1010
- Muth NZ, Pigliucci M (2006) Traits of invasives reconsidered: phenotypic comparisons of introduced invasive and introduced noninvasive plant species within two closely related clades. *Amer J Bot* 93:188-196
- Pammenter NW, Drennan PM, Smith VR (1986) Physiological and anatomical aspects of photosynthesis of two *Agrostis* species at a sub-Antarctic island. *New Phytol* 102:143-160
- Pantone DJ, Pavlik BM, Kelley RB (1995) The reproductive attributes of an endangered plant as compared to a weedy congener. *Biol Conserv* 71:305-311
- Perrins J, Fitter A, Williamson M (1993) Population biology and rates of invasion of three introduced *Impatiens* species in the British Isles. *J Biogeogr* 20:33-44
- Pyke DA (1986) Demographic responses of *Bromus tectorum* and seedlings of *Agropyron spicatum* to grazing by small mammals: Occurrence and severity of grazing. *J Ecol* 75:825-835

- Pyke DA (1987) Demographic responses of *Bromus tectorum* and seedlings of *Agropyron spicatum* to grazing by small mammals: The influence of grazing frequency and plant age. *J Ecol* 75:825-835
- Pyšek P, Brock JH, Bímová K, Mandák B, Jarošík V, Koukolíková I, Pergl J, Štěpánek J (2003a) Vegetative regeneration in invasive *Reynoutria* (Polygonaceae) taxa: the determinant of invasibility at the genotype level. *Amer J Bot* 90:1487-1495
- Radford IJ, Cousens RD (2000) Invasiveness and comparative life history traits of exotic and indigenous *Senecio* species in Australia. *Oecologia* 125:531-542
- Radho-Toly S, Majer JD, Yates C (2001) Impact of fire on leaf nutrients, arthropod fauna and herbivory of native and exotic eucalypts in Kings Park, Perth, Western Australia. *Austral Ecol* 26:500-506
- Rejmánek M, Richardson DM (1996) What attributes make some plant species more invasive? *Ecology* 77:1655-1661
- Richards JH (1984) Root growth response to defoliation in two *Agropyron* bunchgrasses: field observations with an improved root periscope. *Oecologia* 64:21-25
- Richardson DM, Van Wilgen BW, Mitchell DT (1987) Aspects of the reproductive ecology of four australian *Hakea* species (Proteaceae) in South Africa. *Oecologia* 71:345-354
- Sallabanks R (1993) Fruiting plant attractiveness to avian seed dispersers: native vs. invasive *Crataegus* in western Oregon. *Madrono* 40:108-116
- Sans FX, Serrano-Garcia H, Afán I (2004) Life-history traits of alien and native *Senecio* species in the Mediterranean region. *Acta Oecol* 26:167-178
- Sasek TW, Strain BR (1991) Effects of carbon dioxide enrichment on the growth and morphology of a native and an introduced honeysuckle vine. *Amer J Bot* 78:69-75
- Schierenbeck KA, Marshall JD (1993) Seasonal and diurnal patterns of photosynthetic gas exchange for *Lonicera sempervirens* and *L. japonica* (Caprifoliaceae). *Amer J Bot* 80:1292-1299
- Schierenbeck K, Mack RN, Sharitz RR (1994) Effects of herbivory on growth and biomass allocation in native and introduced species of *Lonicera*. *Ecology* 75:1661-1672
- Schweitzer JA, Larson KC (1999) Greater morphological plasticity of exotic honeysuckle species may make them better invaders than native species. *J Torrey Bot Soc* 126:15-23
- Smith MD, Knapp AK (2001) Physiological and morphological traits of exotic, invasive exotic, and native plant species in tallgrass prairie. *Int J Plant Sci* 162:785-792
- Van Clef M, Stiles EW (2001) Seed longevity in three pairs of native and non-native congeners: assessing invasive potential. *Northeastern Natur* 8:301-310

- Vila M, D'Antonio CM (1998a) Fruit choice and seed dispersal of invasive vs. noninvasive *Carpobrotus* (Aizoaceae) in coastal California. *Ecology* 79:1053-1060
- Vila M, D'Antonio C (1998b) Hybrid vigor for clonal growth in *Carpobrotus* (Aizoaceae) in coastal California. *Ecol Appl* 8:1196-1205
- Williams DG, Black RA (1994) Drought response of a native and introduced Hawaiian grass. *Oecologia* 97:512-519