

Resonance and Multiplicity in Periodic Boundary Value Problems with Singularity

Irena Rachůnková*, Milan Tvrdý† and Ivo Vrkoč

(Received August 13, 2001)

Summary. The paper deals with the boundary value problem

$$u'' + k u = g(u) + e(t), \quad u(0) = u(2\pi), \quad u'(0) = u'(2\pi),$$

where $k \in \mathbb{R}$, $g : (0, \infty) \mapsto \mathbb{R}$ is continuous, $e \in \mathbb{L}[0, 2\pi]$ and $\lim_{x \rightarrow 0^+} \int_x^1 g(s) ds = \infty$. In particular, the existence and multiplicity results are obtained by using the method of lower and upper functions which are constructed as solutions of related auxiliary linear problems.

Mathematics Subject Classification 2000: 34B15, 34C25

Keywords. Second order nonlinear ordinary differential equation, periodic problem, lower and upper functions.

1 . Introduction

In this paper we consider the periodic boundary value problems of the form

$$(1.1) \quad u'' + k u = g(u) + e(t), \quad u(0) = u(2\pi), \quad u'(0) = u'(2\pi),$$

where

$$(1.2) \quad g \in \mathbb{C}(0, \infty), \quad e \in \mathbb{L}[0, 2\pi], \quad k \in \mathbb{R}$$

and g has a strong singularity at 0, i.e.

$$(1.3) \quad \lim_{x \rightarrow 0^+} \int_x^1 g(s) ds = \infty.$$

This problem was studied by many authors, starting from Lazer and Solimini [6], where (1.1) with $k = 0$ and g positive was considered. Later, this work has been

*Supported by the grant No. 201/01/1451 of the Grant Agency of the Czech Republic and by the Council of Czech Government J14/98:153100011

†Supported by the grant No. 201/01/1199 of the Grant Agency of the Czech Republic

generalized or extended e.g. by del Pino, Manásevich and Montero [1], Fonda [2], Fonda, Manásevich and Zanolin[3], Mawhin [7], Ge and Mawhin [4], Omari and Ye [9], Rachůnková [10], Rachůnková and Tvrđý [12], Rachůnková, Tvrđý and Vrkoč [14], Yan and Zhang [16], Zhang [17] and others. In particular, the problems having nonlinearities with the asymptotic behaviour at $+\infty$ which corresponds to $k > 0$ and g bounded below in (1.1) were solved by some of the above authors. For example, the papers [2], [9], [10], [12], [14] and [17] dealt with problems characterized by a positive k which was less than the first positive Dirichlet eigenvalue μ_1 of $x'' + \mu x = 0$, while the cases corresponding to k lying between two adjacent higher eigenvalues were investigated in [1] or [16]. The existence results in the resonance case $k = \mu_1$ were reached in [14]. In [2] or [3] we can also find multiplicity results for subharmonics. Theorems about more solutions of (1.1) provided $k = 0$ are proved in [10].

Here we bring new results about the existence of one or two positive solutions of (1.1) for both nonresonance and resonance values of k . Moreover, our Theorem 4.4 generalizes for $n = 3$ Theorem 3.5 in [10] (even in the case $k = 0$), because the condition (3.4) in Theorem 4.4 is weaker than the corresponding condition (3.16) in [10, Theorem 3.5] which requires (for $i = 1$) $g(x) + \bar{e} > 0$ on $[a_1, b_1] \subset (0, \infty)$, where $b_1 - a_1 = \frac{\pi}{3} \|e - \bar{e}\|_1$.

Let J be a (possibly unbounded) subinterval of \mathbb{R} . We say that $f : [0, 2\pi] \times J \mapsto \mathbb{R}$ fulfils the Carathéodory conditions on $[0, 2\pi] \times \mathbb{R}$, if f has the following properties: (i) for each $x \in J$ the function $f(\cdot, x)$ is measurable on $[0, 2\pi]$; (ii) for almost every $t \in [0, 2\pi]$ the function $f(t, \cdot)$ is continuous on J ; (iii) for each compact set $K \subset J$ the function $m_K(t) = \sup_{x \in K} |f(t, x)|$ is Lebesgue integrable on $[0, 2\pi]$. The set of functions satisfying the Carathéodory conditions on $[0, 2\pi] \times J$ is denoted by $\text{Car}([0, 2\pi] \times J)$.

For a given (possibly unbounded) subinterval J of \mathbb{R} , $\mathbb{C}(J)$ denotes the set of functions continuous on J , $\mathbb{L}[0, 2\pi]$ stands for the set of functions (Lebesgue) integrable on $[0, 2\pi]$, $\mathbb{L}_2[0, 2\pi]$ is the set of functions square integrable on $[0, 2\pi]$, $\mathbb{L}_\infty[0, 2\pi]$ is the set of functions essentially bounded on $[0, 2\pi]$, $\mathbb{AC}[0, 2\pi]$ denotes the set of functions absolutely continuous on $[0, 2\pi]$, $\mathbb{AC}^1[0, 2\pi]$ is the set of functions $u \in \mathbb{AC}[0, 2\pi]$ with the first derivative absolutely continuous on $[0, 2\pi]$ and $\mathbb{BV}[0, 2\pi]$ denotes the set of functions of bounded variation on $[0, 2\pi]$. For $x \in \mathbb{L}_\infty[0, 2\pi]$, $y \in \mathbb{L}[0, 2\pi]$ and $z \in \mathbb{L}_2[0, 2\pi]$, we denote $\|x\|_\infty = \sup \text{ess}_{t \in [0, 2\pi]} |x(t)|$,

$$\bar{y} = \frac{1}{2\pi} \int_0^{2\pi} y(s) \, ds, \quad \|y\|_1 = \int_0^{2\pi} |y(t)| \, dt \quad \text{and} \quad \|z\|_2 = \left(\int_0^{2\pi} z^2(t) \, dt \right)^{\frac{1}{2}}.$$

Furthermore, $\mathbb{C}^1[0, 2\pi]$ is the space of functions from $\mathbb{C}[0, 2\pi]$ having a continuous first derivative on $[0, 2\pi]$ equipped with the norm $x \in \mathbb{C}^1[0, 2\pi] \mapsto \|x\|_\infty + \|x'\|_\infty$.

If $x \in \mathbb{BV}[0, 2\pi]$, $s \in (0, 2\pi]$ and $t \in [0, 2\pi)$, then the symbols $x(s-)$, $x(t+)$ and

$\Delta^+x(t)$ are defined respectively by

$$x(s-) = \lim_{\tau \rightarrow s-} x(\tau), \quad x(t+) = \lim_{\tau \rightarrow t+} x(\tau) \quad \text{and} \quad \Delta^+x(t) = x(t+) - x(t).$$

Furthermore, x^{ac} and x^{sing} stand for the absolutely continuous part of x and the singular part of x , respectively. We suppose $x^{\text{sing}}(0) = 0$. For $x \in \mathbb{L}[0, 2\pi]$, the symbols x^+ and x^- denote its nonnegative and nonpositive parts.

Besides (1.1) we will also consider a more general problem

$$(1.4) \quad x'' = f(t, x), \quad x(0) = x(2\pi), \quad x'(0) = x'(2\pi),$$

where $f \in \text{Car}([0, 2\pi] \times J)$ and $J \subset \mathbb{R}$.

1.1. Definition. By a *solution* of (1.4) we understand a function $x: [0, 2\pi] \mapsto \mathbb{R}$ such that $x' \in \text{AC}[0, 2\pi]$, $x(0) = x(2\pi)$, $x'(0) = x'(2\pi)$ and $x(t) \in J$ and $x''(t) = f(t, x(t))$ hold for a.e. $t \in [0, 2\pi]$.

1.2. Definition. Functions $(\sigma, \rho) \in \text{AC}[0, 2\pi] \times \text{BV}[0, 2\pi]$ are *lower functions* of (1.4) if $\sigma(t) \in J$ for a.e. $t \in [0, 2\pi]$, the singular part ρ^{sing} of ρ is nondecreasing on $[0, 2\pi]$, $\sigma'(t) = \rho(t)$ and $\rho'(t) \geq f(t, \sigma(t))$ for a.e. $t \in [0, 2\pi]$, $\sigma(0) = \sigma(2\pi)$ and $\rho(0+) \geq \rho(2\pi-)$.

Similarly, functions $(\sigma, \rho) \in \text{AC}[0, 2\pi] \times \text{BV}[0, 2\pi]$ are *upper functions* of (1.4) if $\sigma(t) \in J$ for a.e. $t \in [0, 2\pi]$, ρ^{sing} is nonincreasing on $[0, 2\pi]$, $\sigma'(t) = \rho(t)$ and $\rho'(t) \leq f(t, \sigma(t))$ for a.e. $t \in [0, 2\pi]$, $\sigma(0) = \sigma(2\pi)$ and $\rho(0+) \leq \rho(2\pi-)$.

1.3. Remark. If $J = \mathbb{R}$, then Definitions 1.1 and 1.2 reduce to those given for the regular case in [11].

If (1.2) is true and $J = (0, \infty)$, then each solution and each upper (lower) function must be positive a.e. on $[0, 2\pi]$.

Our proofs will be based on the following theorem which is contained in [11, Theorems 4.1, 4.2 and 4.3] and which concerns the nonsingular case with $J = \mathbb{R}$.

1.4. Theorem. Let (σ_1, ρ_1) and (σ_2, ρ_2) be respectively lower and upper functions of the problem (1.4), where $J = \mathbb{R}$. Furthermore, assume that there is $m \in \mathbb{L}[0, 2\pi]$ such that $f(t, x) \geq m(t)$ for a.e. $t \in [0, 2\pi]$ and all $x \in \mathbb{R}$ (or $f(t, x) \leq m(t)$ for a.e. $t \in [0, 2\pi]$ and all $x \in \mathbb{R}$). Then (1.4) has a solution x such that $\|x'\|_\infty \leq \|m\|_1$. Moreover, if

$$(1.5) \quad \sigma_1(t) \leq \sigma_2(t) \quad \text{for all } t \in [0, 2\pi],$$

then $\sigma_1(t) \leq x(t) \leq \sigma_2(t)$ is true for all $t \in [0, 2\pi]$ and if (1.5) does not hold, then there is $t_x \in [0, 2\pi]$ such that

$$\min\{\sigma_1(t_x), \sigma_2(t_x)\} \leq x(t_x) \leq \max\{\sigma_1(t_x), \sigma_2(t_x)\}.$$

In [12] and [13] we have presented conditions ensuring the existence and localization of lower and upper functions of (1.4). As an immediate consequence of these results we get propositions for the following special case of the problem (1.4):

$$(1.6) \quad x'' = h(x) + e(t), \quad x(0) = x(2\pi), \quad x'(0) = x'(2\pi),$$

where

$$(1.7) \quad h \in \mathbb{C}(J), \quad e \in \mathbb{L}[0, 2\pi] \quad \text{and } J \text{ is an open subinterval in } \mathbb{R}.$$

1.5. Proposition. *Suppose that (1.7) holds. Further, let $A \in J$ and let the inequality*

$$(1.8) \quad h(x) + \bar{e} \leq 0 \quad \text{for } x \in [A, B] \quad (h(x) + \bar{e} \geq 0 \quad \text{for } x \in [A, B])$$

be fulfilled, where $B \in J$ and

$$(1.9) \quad B - A \geq \frac{\pi}{3} \|e - \bar{e}\|_1.$$

Then there exist lower (upper) functions (σ, ρ) of (1.6) and

$$(1.10) \quad \sigma(t) \in [A, B] \quad \text{for all } t \in [0, 2\pi].$$

Proof. If $J = \mathbb{R}$, then the proof is given in [12, Propositions 2.4 and 2.5]. In the case that $J \neq \mathbb{R}$, we put

$$(1.11) \quad f(t, x) = e(t) + \begin{cases} h(A) & \text{if } x < A, \\ h(x) & \text{if } x \in [A, B], \\ h(B) & \text{if } x > B. \end{cases}$$

Since $f \in \text{Car}([0, 2\pi] \times \mathbb{R})$, we get by [12, Propositions 2.4 and 2.5] the existence of lower (upper) functions (σ, ρ) of (1.4) fulfilling (1.10). The conditions (1.10) and (1.11) guarantee that (σ, ρ) are lower (upper) functions of (1.6), as well. \square

1.6. Proposition. *Suppose that (1.7) holds. Further, let $A \in J$, $k \neq n^2$ for all $n \in \mathbb{N}$ and let the inequality*

$$(1.12) \quad \begin{aligned} h(x) + kx + \bar{e} &\leq k \frac{A+B}{2} \quad \text{for } x \in [A, B] \\ (h(x) + kx + \bar{e}) &\geq k \frac{A+B}{2} \quad \text{for } x \in [A, B], \end{aligned}$$

be fulfilled, where $B \in J$ and

$$(1.13) \quad B - A = 2 \Phi(k) \|e - \bar{e}\|_1$$

and

$$(1.14) \quad \Phi(k) = \begin{cases} \min\left\{\frac{\pi}{6}, \frac{\coth(\sqrt{|k|}\pi)}{4\sqrt{|k|}}\right\} & \text{if } k < 0, \\ \min\left\{\frac{\pi}{6(1-k)}, \frac{1}{4k} \sin(\sqrt{k}\pi)\right\} & \text{if } k \in (0, \frac{1}{4}], \\ \min\left\{\frac{\pi}{6(1-k)}, \frac{1}{2\sqrt{k} \sin(\sqrt{k}\pi)}\right\} & \text{if } k \in (\frac{1}{4}, 1), \\ \frac{1}{2\sqrt{k} |\sin(\sqrt{k}\pi)|} & \text{if } k > 1, k \neq n^2, n \in \mathbb{N}. \end{cases}$$

Then there exist lower (upper) functions (σ, ρ) of (1.6) satisfying (1.10).

Proof. If $J = \mathbb{R}$, then the proof follows from [13, Theorems 3.1-3.4], where we put $a = k \frac{A+B}{2}$. In the case $J \neq \mathbb{R}$, we can use the same arguments as in the proof of Proposition 1.5. \square

The next lemma will be helpful in what follows.

1.7. Lemma. *Let (1.7) be true and let x be an arbitrary solution of (1.6). Further, let $t_1 \in [0, 2\pi]$ be such that $x(t_1) = \max_{t \in [0, 2\pi]} x(t)$. Then the inequality*

$$\int_{x(t_0)}^A h(s) ds \leq \|e\|_1 \|x'\|_\infty + \int_A^{x(t_1)} |h(s)| ds$$

holds for all $t_0 \in [0, 2\pi]$ and all $A \in J$ such that $x(t_1) \geq A$.

Proof. In virtue of the periodicity of x , we have $x'(t_1) = 0$. Consequently, multiplying the equality $x''(t) = h(x(t)) + e(t)$ a.e. on $[0, 2\pi]$ by $x'(t)$ and integrating from t_0 to t_1 , we get

$$0 \geq -\frac{(x'(t_0))^2}{2} = \int_{t_0}^{t_1} x''(t) x'(t) dt = \int_{x(t_0)}^{x(t_1)} h(s) ds + \int_{t_0}^{t_1} e(t) x'(t) dt.$$

Hence,

$$\begin{aligned} \int_{x(t_0)}^A h(s) ds &= \int_{x(t_0)}^{x(t_1)} h(s) ds - \int_A^{x(t_1)} h(s) ds \\ &\leq -\int_{t_0}^{t_1} e(t) x'(t) dt - \int_A^{x(t_1)} h(s) ds \leq \|e\|_1 \|x'\|_\infty + \int_A^{x(t_1)} |h(s)| ds. \quad \square \end{aligned}$$

The proof of the next lemma is an easy modification of that of [11, Lemma 1.1].

1.8. Lemma. *Suppose that (1.7) is true. Furthermore, let $I \subset J$ and let $h_* \in \mathbb{R}$ be such that $h(x) \geq h_*$ for all $x \in I$. Then*

$$\|x'\|_\infty \leq \|e\|_1 + 2\pi |h_*|$$

holds for each solution x of (1.6) with the property $x(t) \in I$ for all $t \in [0, 2\pi]$. \square

2 . Existence theorem

The main result of this section is Theorem 2.1 which gives an existence principle for the problem (1.1) in terms of lower and upper functions. More effective results can be obtained if we replace the assumption of the existence of lower and upper functions by the corresponding conditions from Propositions 1.5 and 1.6.

2.1. Theorem. *Suppose that (1.2), (1.3),*

$$(2.1) \quad \liminf_{x \rightarrow \infty} \frac{g(x)}{x} > k - \frac{1}{4}$$

and

$$(2.2) \quad \liminf_{x \rightarrow 0^+} g(x) > -\infty$$

are satisfied. Further, let there be lower functions (σ_1, ρ_1) and upper functions (σ_2, ρ_2) of (1.1) such that

$$(2.3) \quad \sigma_2(t) > 0 \quad \text{on } [0, 2\pi].$$

Then the problem (1.1) has a positive solution.

Before proving Theorem 2.1 we will prove several auxiliary assertions. In particular, Lemmas 2.2 and 2.4 give a priori estimates for solutions of (1.6). The proof of Theorem 2.1 follows from Proposition 2.9.

2.2. Lemma. *Let $g \in \mathbb{C}(0, \infty)$, $k \in \mathbb{R}$ and suppose that (1.3) is true. Further, let $E, K \in [0, \infty)$, $A \in (0, \infty)$ and $R \in [A, \infty)$. Then there is $\varepsilon^* > 0$ such that $\min_{t \in [0, 2\pi]} x(t) > \varepsilon^*$ holds for each $h \in \mathbb{C}(\mathbb{R})$ satisfying*

$$(2.4) \quad h(x) = g(x) - kx \quad \text{on } [\varepsilon^*, R]$$

for each $e \in \mathbb{L}[0, 2\pi]$ with $\|e\|_1 \leq E$ and for each solution x of (1.6) fulfilling

$$(2.5) \quad \|x'\|_\infty \leq K \quad \text{and} \quad \max_{t \in [0, 2\pi]} x(t) \in [A, R].$$

Proof. Put

$$K^* = EK + \int_A^R |g(s) - ks| ds.$$

In view of (1.3), there is $\varepsilon^* > 0$ such that

$$(2.6) \quad \int_{\varepsilon^*}^A (g(s) - ks) ds > K^*.$$

Let $h \in \mathbb{C}(\mathbb{R})$ fulfil (2.4), let $e \in \mathbb{L}[0, 2\pi]$ be such that $\|e\|_1 \leq E$ and let x be a solution of (1.6) verifying (2.5). Suppose that $\min_{t \in [0, 2\pi]} x(t) \leq \varepsilon^*$ and denote by t_0 and t_1 the points in $[0, 2\pi]$ such that $t_0 < t_1$, $x(t_0) = \varepsilon^*$, $x(t) \geq \varepsilon^*$ on $[t_0, t_1]$ and $x(t_1) = \max_{t \in [0, 2\pi]} x(t)$. By virtue of (2.5) we have $x(t_1) \in [A, R]$. Thus, taking into account (2.4) and (2.6) and using Lemma 1.7, we get

$$\begin{aligned} K^* &< \int_{\varepsilon^*}^A (g(s) - k s) ds = \int_{x(t_0)}^A h(s) ds \\ &\leq EK + \int_A^R |h(s)| ds = EK + \int_A^R |g(s) - k s| ds = K^*, \end{aligned}$$

a contradiction. \square

In particular, we have:

2.3. Corollary. *Suppose (1.2) and (1.3). Then each solution of (1.1) is positive on $[0, 2\pi]$.*

2.4. Lemma. *Let $E, C \in [0, \infty)$, $\eta \in (0, \frac{1}{4})$ and $(0, \infty) \subset J \subset \mathbb{R}$. Then for any $B \in (0, \infty)$ there is $R \in (B, \infty)$ such that the estimate*

$$(2.7) \quad \max_{t \in [0, 2\pi]} x(t) \leq R$$

is valid for each $h \in \mathbb{C}(J)$ satisfying

$$h(x)x + \left(\frac{1}{4} - \eta\right)x^2 \geq -C|x| \quad \text{on } J$$

for each $e \in \mathbb{L}[0, 2\pi]$ with $\|e\|_1 \leq E$ and for each solution x of (1.6) fulfilling

$$(2.8) \quad x(t) \in J \quad \text{for } t \in [0, 2\pi] \quad \text{and} \quad \min_{t \in [0, 2\pi]} x(t) \leq B.$$

Proof. Suppose that no such R exists. Then for any $\ell \in \mathbb{N}$ we can find $h_\ell \in \mathbb{C}(J)$, $e_\ell \in \mathbb{L}[0, 2\pi]$ and a solution x_ℓ of

$$x'' = h_\ell(x) + e_\ell(t), \quad x(0) = x(2\pi), \quad x'(0) = x'(2\pi)$$

such that $\|e_\ell\|_1 \leq E$,

$$(2.9) \quad h_\ell(x)x + \left(\frac{1}{4} - \eta\right)x^2 \geq -C|x| \quad \text{for all } x \in J$$

and

$$(2.10) \quad \min_{t \in [0, 2\pi]} x_\ell(t) \leq B \quad \text{and} \quad \max_{t \in [0, 2\pi]} x_\ell(t) > \ell.$$

For $\ell \in \mathbb{N}$, denote by t_ℓ the point in $[0, 2\pi]$ for which $x_\ell(t_\ell) = B$. Furthermore, let us extend x_ℓ and e_ℓ to functions 2π -periodic on \mathbb{R} . We have

$$x_\ell''(t) = h_\ell(x_\ell(t)) + e_\ell(t) \quad \text{for a.e. } t \in \mathbb{R}.$$

Multiplying this equality by $x_\ell(t)$, integrating from t_ℓ to $t_\ell + 2\pi$ and making use of (2.9), we obtain

$$\begin{aligned} \|x_\ell'\|_2^2 &= - \int_{t_\ell}^{t_\ell+2\pi} h_\ell(x_\ell(t)) x_\ell(t) dt - \int_{t_\ell}^{t_\ell+2\pi} e_\ell(t) x_\ell(t) dt \\ &\leq \left(\frac{1}{4} - \eta\right) \|x_\ell\|_2^2 + C \|x_\ell\|_1 + E \|x_\ell\|_\infty. \end{aligned}$$

On the other hand, in virtue of (2.10) we have

$$(2.11) \quad \|x_\ell\|_\infty \leq |x_\ell(t_\ell)| + \int_{t_\ell}^{t_\ell+2\pi} |x_\ell'(t)| dt \leq B + \sqrt{2\pi} \|x_\ell'\|_2.$$

Thus,

$$(2.12) \quad \left(\|x_\ell'\|_2 - E \sqrt{\frac{\pi}{2}}\right)^2 \leq \left(\frac{1}{4} - \eta\right) \|x_\ell\|_2^2 + \sqrt{2\pi} C \|x_\ell\|_2 + E B + \frac{\pi}{2} E^2.$$

Inserting $x_\ell(t) = v_\ell(t) + B$ on \mathbb{R} into (2.12), we obtain

$$(2.13) \quad (\|v_\ell'\|_2 - c)^2 \leq \left(\frac{1}{4} - \eta\right) \|v_\ell\|_2^2 + a \|v_\ell\|_2 + b,$$

where $a, b, c \in \mathbb{R}$ do not depend on ℓ . Since $v_\ell(t_\ell) = v_\ell(t_\ell + 2\pi) = 0$, by Scheeffer's inequality [15, p. 207] (see also [8, II.2]), we have $\|v_\ell\|_2^2 \leq 4 \|v_\ell'\|_2^2$ and hence (2.13) reduces to

$$(2.14) \quad \left(\frac{\|v_\ell'\|_2 - c}{\|v_\ell'\|_2}\right)^2 \leq 1 - 4\eta + \frac{2a}{\|v_\ell'\|_2} + \frac{b}{\|v_\ell'\|_2^2}.$$

Now, (2.10) and (2.11) yield $\lim_{\ell \rightarrow \infty} \|v_\ell'\|_2 = \infty$ and, by virtue of (2.14), we have

$$1 = \lim_{\ell \rightarrow \infty} \left(\frac{\|v_\ell'\|_2 - c}{\|v_\ell'\|_2}\right)^2 \leq \lim_{\ell \rightarrow \infty} \left(1 - 4\eta + \frac{2a}{\|v_\ell'\|_2} + \frac{b}{\|v_\ell'\|_2^2}\right) = 1 - 4\eta,$$

a contradiction. □

Let $g \in \mathcal{C}(0, \infty)$ fulfil (1.3), (2.1) and (2.2). Denote

$$(2.15) \quad g_0(x) = g(x) - kx \quad \text{for } x \in (0, \infty).$$

Then we have $g_0 \in \mathbb{C}(0, \infty)$,

$$(2.16) \quad \lim_{x \rightarrow 0^+} \int_x^1 g_0(s) \, ds = \infty$$

and

$$(2.17) \quad \liminf_{x \rightarrow \infty} \frac{g_0(x)}{x} > -\frac{1}{4}.$$

Furthermore, (2.2) and (2.16) imply

$$(2.18) \quad \inf_{x \in (0, R]} g_0(x) \in \mathbb{R} \quad \text{for each } R \in (0, \infty)$$

and

$$(2.19) \quad \limsup_{x \rightarrow 0^+} g_0(x) = \infty.$$

Moreover, in view of (2.17) and (2.18), there exist $\eta \in (0, \frac{1}{4})$ and $C \in [0, \infty)$ such that

$$(2.20) \quad g_0(x) + \left(\frac{1}{4} - \eta\right)x \geq -C \quad \text{for all } x \in (0, \infty).$$

2.5. Lemma. *Assume (1.2), (1.3) and (2.2) and let (σ, ρ) be lower functions of (1.1). Then $\min_{t \in [0, 2\pi]} \sigma(t) > 0$.*

Proof. In view of (2.15), (2.16) and (2.18), there are $\delta \in (0, \infty)$ and $M \in [0, \infty)$ such that

$$(2.21) \quad \lim_{x \rightarrow 0^+} \int_x^{\delta'} g_0(s) \, ds = \infty \quad \text{for all } \delta' \in (0, \delta)$$

and

$$(2.22) \quad g_0(x) \geq -M \quad \text{for all } x \in (0, \delta).$$

Let an arbitrary $\varepsilon > 0$ be given. Since in view of Definition 1.2 we have $\sigma(t) > 0$ a.e. on $[0, 2\pi]$, we can choose $t_0 \in (0, \varepsilon]$ in such a way that $\sigma(t_0) > 0$. Put $t^* = \sup\{t \in [t_0, 2\pi] : \sigma(s) > 0 \text{ on } [t_0, t]\}$. Notice that $\sigma(t^*) = 0$ holds whenever $t^* < 2\pi$.

Assume $\sigma(t^*) = 0$. Then there is $t' \in (t_0, t^*)$ such that

$$(2.23) \quad \sigma(t) \in [0, \delta) \quad \text{for all } t \in [t', t^*].$$

As $\rho \in \mathbb{BV}[0, 2\pi]$, we have $r = \|\rho\|_\infty + 1 < \infty$ and, due to Definition 1.2, we get that

$$\rho'(t) (\rho(t) - r) \leq g_0(\sigma(t)) (\rho(t) - r) + e(t) (\rho(t) - r)$$

is satisfied for a.e. $t \in [0, 2\pi]$. Let $t_n \in (t', t^*)$ be an increasing sequence such that $\lim_{n \rightarrow \infty} t_n = t^*$. Then

$$(2.24) \quad \lim_{n \rightarrow \infty} \sigma(t_n) = \sigma(t^*) = 0$$

and

$$\begin{aligned} & \int_{t'}^{t_n} \rho'(t) (\rho(t) - r) dt \\ & \leq \int_{t'}^{t_n} g_0(\sigma(t)) (\rho(t) - r) dt + \int_{t'}^{t_n} e(t) (\rho(t) - r) dt \\ & = - \int_{\sigma(t_n)}^{\sigma(t')} g_0(s) ds - r \int_{t'}^{t_n} g_0(\sigma(t)) dt + \int_{t'}^{t_n} e(t) (\rho(t) - r) dt. \end{aligned}$$

In virtue of (2.22) and (2.23), for any $n \in \mathbb{N}$ we have

$$- \int_{t'}^{t_n} g_0(\sigma(t)) dt \leq 2\pi M \quad \text{and} \quad \left| \int_{t'}^{t_n} e(t) (\rho(t) - r) dt \right| \leq 2r \|e\|_1.$$

Moreover, by (2.21) and (2.24),

$$\lim_{n \rightarrow \infty} \int_{\sigma(t_n)}^{\sigma(t')} g_0(s) ds = \infty.$$

This implies that

$$\lim_{n \rightarrow \infty} \int_{t'}^{t_n} \rho'(t) (\rho(t) - r) dt = -\infty.$$

On the other hand,

$$\left| \int_{t'}^{t_n} \rho'(t) (\rho(t) - r) dt \right| \leq 2r \int_0^{2\pi} |\rho'(t)| dt \leq 2r \|\rho\|_{\mathbb{B}\mathbb{V}} < \infty,$$

a contradiction. Thus, $t^* = 2\pi$ and $\sigma(t^*) = \sigma(2\pi) > 0$. In particular, we have shown that $\sigma(t)$ is positive on any interval $(\varepsilon, 2\pi]$, $\varepsilon > 0$, and, as we also have $\sigma(0) = \sigma(2\pi) > 0$ in view of the periodicity condition, this completes the proof. \square

2.6. Remark. If g satisfies the assumptions of Lemma 2.5 and, in addition, $g(x) \geq k$ on $(0, \infty)$, $e(t) = (k-1) \sin t$ on $[0, 2\pi]$, $\sigma(t) = 1 + \sin t$ and $\rho(t) = \cos t$ on $[0, 2\pi]$, then (σ, ρ) are upper functions of (1.1) and $\min_{t \in [0, 2\pi]} \sigma(t) = 0$. This shows that for upper functions the analogue of Lemma 2.5 does not hold.

2.7. Definition. Let $g \in \mathbb{C}(0, \infty)$ and $k \in \mathbb{R}$. Then, for given $E, A \in (0, \infty)$ and $B \in [A, \infty)$, we denote by $\mathcal{E}(E, A, B)$ the set of functions $e \in \mathbb{L}[0, 2\pi]$ with $\|e\|_1 \leq E$ and such that (1.1) has lower functions (σ_1, ρ_1) and upper functions (σ_2, ρ_2) fulfilling

$$(2.25) \quad A \leq \sigma_i(t) \leq B \quad \text{on } [0, 2\pi] \quad \text{for } i = 1, 2.$$

2.8. Remark. Let $g \in \mathbb{C}(0, \infty)$ satisfy (1.3), (2.1) and (2.2) and let g_0 be given by (2.15). Then g_0 fulfils (2.16)-(2.20) and we can choose a sequence $\{\varepsilon_n\}_{n=1}^\infty \subset (0, 1)$ with the properties

$$(2.26) \quad \begin{cases} \varepsilon_{n+1} < \varepsilon_n & \text{and } g_0(\varepsilon_n) > 0 \text{ for all } n \in \mathbb{N}, \\ \lim_{n \rightarrow \infty} \varepsilon_n = 0 & \text{and } \lim_{n \rightarrow \infty} g_0(\varepsilon_n) = \infty \end{cases}$$

and choose $C \geq 0$, $\eta \in (0, \frac{1}{4})$ and the functions $g_{n,m} \in \mathbb{C}(\mathbb{R})$, $n, m \in \mathbb{N}$, in such a way that the relations

$$(2.27) \quad g_{n,m}(x) = g_0(x) \text{ for } x \in [\varepsilon_n, m],$$

$$(2.28) \quad g_{n,m}(x)x + \left(\frac{1}{4} - \eta\right)x^2 \geq -C|x| \text{ for all } x \in \mathbb{R}$$

and

$$(2.29) \quad g_{*,m} := \inf_{\substack{x \in \mathbb{R} \\ \ell \in \mathbb{N}}} g_{\ell,m}(x) \in \mathbb{R}$$

are valid for all $n, m \in \mathbb{N}$. Indeed, for given $n, m \in \mathbb{N}$, we can put e.g.

$$g_{n,m}(x) = \begin{cases} 0 & \text{if } x \leq 0, \\ g_0(\varepsilon_n) \frac{x}{\varepsilon_n} & \text{if } x \in [0, \varepsilon_n], \\ g_0(x) & \text{if } x \in [\varepsilon_n, m], \\ g_0(m) & \text{if } x \geq m. \end{cases}$$

In what follows, having a sequence $\{\varepsilon_n\}_{n=1}^\infty$ which satisfies (2.27) and (2.28) and functions $g_{n,m}$, $n, m \in \mathbb{N}$, satisfying (2.27)-(2.29), we will often work with the auxiliary regular boundary value problems

$$(2.30) \quad x'' = g_{n,m}(x) + e(t), \quad x(0) = x(2\pi), \quad x'(0) = x'(2\pi).$$

2.9. Proposition. *Suppose that $g \in \mathbb{C}(0, \infty)$, $k \in \mathbb{R}$, $E, A \in (0, \infty)$, $B \in [A, \infty)$ and let (1.3), (2.1), (2.2) be satisfied. Then there are $R \in (B, \infty)$, $\varepsilon^* \in (0, \infty)$ and $K \in [0, \infty)$ such that for each $e \in \mathcal{E}(E, A, B)$ the problem (1.1) has a solution u satisfying*

$$(2.31) \quad \varepsilon^* \leq u(t) \leq R \text{ on } [0, 2\pi] \quad \text{and} \quad \|u'\|_\infty \leq K.$$

Moreover, if (σ_1, ρ_1) and (σ_2, ρ_2) are respectively lower and upper functions of (1.1) with the property (2.25), then

$$(2.32) \quad A \leq \min\{\sigma_1(t_u), \sigma_2(t_u)\} \leq u(t_u) \leq \max\{\sigma_1(t_u), \sigma_2(t_u)\} \leq B \\ \text{for some } t_u \in [0, 2\pi].$$

Proof. Let $e \in \mathcal{E}(E, A, B)$ be given and let ε_n and $g_{n,m}$, $n, m \in \mathbb{N}$, be chosen in such a way that (2.26)-(2.29) are true and

$$(2.33) \quad \varepsilon_n < A \quad \text{for all } n \in \mathbb{N}.$$

By Lemma 2.4, there is $R \in \mathbb{N} \cap (B, \infty)$ such that the estimate (2.7) is valid for all $n, m \in \mathbb{N}$, for each $e \in \mathbb{L}[0, 2\pi]$ with $\|e\|_1 \leq E$ and for each solution x of (2.30) fulfilling (2.8) with $J = \mathbb{R}$. For $n \in \mathbb{N}$, consider the problems

$$(2.34) \quad x'' = g_{n,R}(x) + e(t), \quad x(0) = x(2\pi), \quad x'(0) = x'(2\pi).$$

In view of (2.29), we have $g_{n,R}(x) \geq g_{*,R} \in \mathbb{R}$ for all $x \in \mathbb{R}$, i.e.

$$g_{n,R}(x) + e(t) \geq g_{*,R} + e(t) \quad \text{for a.e. } t \in [0, 2\pi] \quad \text{and all } x \in \mathbb{R} \quad \text{and } n \in \mathbb{N}.$$

Let $e \in \mathcal{E}(E, A, B)$ and let (σ_1, ρ_1) and (σ_2, ρ_2) be respectively lower and upper functions of (1.1) fulfilling (2.25). In view of (2.33), (σ_1, ρ_1) and (σ_2, ρ_2) are lower and upper functions of (2.34) for all $n \in \mathbb{N}$, respectively. Thus, by Theorem 1.4 and Lemma 1.8, the problem (2.34) has for any $n \in \mathbb{N}$ a solution x_n such that $\|x'_n\|_\infty \leq E + 2\pi |g_{*,R}| = K$ and

$$A \leq \min\{\sigma_1(t_n), \sigma_2(t_n)\} \leq x_n(t_n) \leq \max\{\sigma_1(t_n), \sigma_2(t_n)\} \leq B$$

for some $t_n \in [0, 2\pi]$. In particular, we have

$$\max_{t \in [0, 2\pi]} x_n(t) \in [A, R] \quad \text{for all } n \in \mathbb{N}.$$

Now, let $\varepsilon^* > 0$ correspond to E, K, A and R by Lemma 2.2 and let $n_0 \in \mathbb{N}$ be such that $\varepsilon_{n_0} < \varepsilon^*$. Then, since $g_{n_0,R}(s) = g_0(s)$ on $[\varepsilon_{n_0}, R]$, Lemma 2.2 yields

$$\min_{t \in [0, 2\pi]} x_{n_0}(t) > \varepsilon^* > \varepsilon_{n_0}.$$

Therefore, $u = x_{n_0}$ is a solution to (1.1) with the properties (2.31) and (2.32). \square

Proof of Theorem 2.1. In virtue of (2.3) and of Lemma 2.5 we can find $A \in (0, \infty)$ and $B \in [A, \infty)$ such that $e \in \mathcal{E}(\|e\|_1, A, B)$, and the assertion of Theorem 2.1 follows from Proposition 2.9. \square

The next result shows that the assumption of the existence of upper functions for (1.1) in Theorem 2.1 can be replaced by (2.35).

2.10. Corollary. *Suppose that (1.2), (1.3), (2.1), (2.2) and*

$$(2.35) \quad \inf_{t \in [0, 2\pi]} \text{ess } e(t) > -\infty$$

are satisfied. Further, let (σ_1, ρ_1) be lower functions of (1.1). Then the problem (1.1) has a positive solution u such that

$$(2.36) \quad u(t_u) \leq \sigma_1(t_u) \quad \text{for some } t_u \in [0, 2\pi].$$

Proof. By (1.3) we have $\limsup_{x \rightarrow 0^+} g(x) = \infty$ and, according to (2.35) and Lemma 2.5, there is $\varepsilon^* \in (0, \min_{t \in [0, 2\pi]} \sigma_1(t))$ such that

$$g(\varepsilon^*) - k \varepsilon^* + e(t) \geq g(\varepsilon^*) - k \varepsilon^* + \inf_{t \in [0, 2\pi]} \text{ess } e(t) \geq 0 \quad \text{for a.e. } t \in [0, 2\pi].$$

Therefore the functions $(\sigma_2(t), \rho_2(t)) = (\varepsilon^*, 0)$ on $[0, 2\pi]$ are upper functions of (1.1) and Theorem 2.1 together with Proposition 2.9 yield the existence of a positive solution u which fulfils (2.36). \square

2.11. Remark. Assume (1.2), while (1.3) need not be satisfied, and let u be a solution of (1.1). Further, suppose that g has a singularity at 0, which means that g is unbounded at 0, i.e.

$$(2.37) \quad \limsup_{x \rightarrow 0^+} g(x) = \infty.$$

Definition 1.1 requires any solution u of (1.1) to be positive a.e. on $[0, 2\pi]$. In particular, u can touch the singularity point $x = 0$. Nevertheless, it can vanish only on the set of zero measure. Corollary 2.3 says that this is impossible provided the singularity is strong, i.e. if (1.3) is satisfied. Therefore, the problem (1.1) can possess nonnegative solutions with at least one zero only if $\lim_{x \rightarrow 0^+} \int_x^1 g(s) ds \in \mathbb{R}$. If this together with (2.37) hold, the singularity $x = 0$ of g is called *weak*.

3 . Existence criteria

The main result of this section is Theorem 3.3 which gives a more effective existence criterion without the a priori assumption of the existence of lower and upper functions. For its proof the following lemmas will be helpful.

3.1. Lemma. *Suppose (1.7). Furthermore, let $(d, A_0] \subset J$ and*

$$(3.1) \quad h(x) + \bar{e} > 0 \quad \text{for all } x \in (d, A_0].$$

Then $\max_{t \in [0, 2\pi]} x(t) > A_0$ holds for each solution x of (1.6) such that $x(t) \in J$ for all $t \in [0, 2\pi]$ and $\min_{t \in [0, 2\pi]} x(t) > d$.

Proof. Let x be a solution of (1.6) and let $\min_{t \in [0, 2\pi]} x(t) > d$ and $x(t) \in J$ for all $t \in [0, 2\pi]$. Integrating the equality

$$x''(t) = h(x(t)) + e(t) \quad \text{a.e. on } [0, 2\pi]$$

over $[0, 2\pi]$ and taking into account the periodicity of x , we get

$$(3.2) \quad \int_0^{2\pi} h(x(t)) dt + 2\pi \bar{e} = 0.$$

On the other hand, if $x(t) \leq A_0$ held for all $t \in [0, 2\pi]$, then using (3.1) we would have

$$\int_0^{2\pi} h(x(t)) \, dt + 2\pi \bar{e} > 0,$$

a contradiction to (3.2). \square

3.2. Lemma. *Let $A \in (0, \infty)$, $B \in [A, \infty)$, $E \in [0, \infty)$, $k \in \mathbb{R}$ and let $g \in \mathbb{C}(0, \infty)$ fulfil (1.3), (2.1) and (2.2). Then there are $R, \varepsilon^* \in (0, \infty)$ and $K \in [0, \infty)$ such that for each $e \in \mathbb{L}[0, 2\pi]$ with $\|e\|_1 \leq E$ and each solution u of (1.1) satisfying*

$$(3.3) \quad u(t_u) \in [A, B] \quad \text{for some } t_u \in [0, 2\pi],$$

the estimates (2.31) are true.

Proof. Let g_0 be given by (2.15). Then $g_0 \in \mathbb{C}(0, \infty)$ fulfils (2.20) with some $\eta \in (0, \frac{1}{4})$ and $C \in [0, \infty)$. By Lemma 2.4, there is $R \in (B, \infty)$ such that the estimate $u(t) \leq R$ on $[0, 2\pi]$ is true for each $e \in \mathbb{L}[0, 2\pi]$ with $\|e\|_1 \leq E$ and each solution u of (1.1) fulfilling (3.3). Furthermore, as $g_* := \inf_{x \in (0, R]} g_0(x) \in \mathbb{R}$ in view of (2.18), according to Corollary 2.3 and Lemma 1.8 we have $u(t) > 0$ on $[0, 2\pi]$ and $\|u'\|_\infty \leq E + 2\pi |g_*|$ for all such solutions. Thus, if we put $K = E + 2\pi |g_*|$, we can complete the proof by means of Lemma 2.2. \square

3.3. Theorem. *Suppose that (1.2), (1.3), (2.1) and*

$$(3.4) \quad \liminf_{x \rightarrow 0^+} g(x) > -\bar{e}$$

are satisfied. Further, let $A \in (0, \infty)$ be such that

$$(3.5) \quad g(x) - kx \leq -\bar{e} \quad \text{for all } x \in [A, B],$$

where B fulfils (1.9). Then the problem (1.1) has a positive solution u such that

$$(3.6) \quad u(t_u) \leq B \quad \text{for some } t_u \in [0, 2\pi].$$

Proof. i) First, assume that (3.5) is satisfied with the strict inequality, i.e.

$$(3.7) \quad g(x) - kx + \bar{e} < 0 \quad \text{for all } x \in [A, B].$$

For $n \in \mathbb{N}$ define

$$(3.8) \quad e_n(t) = \max\{e(t), -n\} \quad \text{a.e. on } [0, 2\pi]$$

and consider the problems

$$(3.9) \quad x'' + kx = g(x) + e_n(t), \quad x(0) = x(2\pi), \quad x'(0) = x'(2\pi).$$

There is $E \in (0, \infty)$ such that $\|e_n\|_1 \leq E$ for all $n \in \mathbb{N}$. Furthermore,

$$(3.10) \quad \inf_{t \in [0, 2\pi]} \text{ess } e_n(t) \geq -n \quad \text{for all } n \in \mathbb{N} \quad \text{and} \quad \lim_{n \rightarrow \infty} \overline{e_n} = \overline{e}.$$

In virtue of (3.4), we can choose $A_0 \in (0, \frac{A}{2})$ in such a way that $g(x) - kx + \overline{e_n} \geq g(x) - kx + \overline{e} > 0$ holds for all $x \in (0, A_0]$ and all $n \in \mathbb{N}$. By Lemma 3.1 this implies

$$(3.11) \quad \max_{t \in [0, 2\pi]} x(t) > A_0 \quad \text{for all } n \in \mathbb{N} \quad \text{and all positive solutions } x \text{ of (3.9).}$$

Let $\{\varepsilon_n\}_{n=1}^{\infty} \subset (0, A_0)$ be an arbitrary decreasing sequence which tends to 0 as $n \rightarrow \infty$ and satisfies the relation $g(\varepsilon_n) - k\varepsilon_n > n$ for any $n \in \mathbb{N}$. In particular, with respect to (3.10), the functions $(\sigma_{2,n}(t), \rho_{2,n}(t)) = (\varepsilon_n, 0)$ are upper functions for (3.9). Furthermore, in view of (3.7), (3.8) and (1.9) we can find $n_0 \in \mathbb{N}$ and $\nu \in (0, \frac{A}{2})$ such that $g(x) - kx + \overline{e_n} \leq 0$ and $\|e_n - \overline{e_n}\|_1 \leq \frac{3}{\pi}(B - A + \nu)$ hold for all $x \in [A - \nu, B]$ and all $n \geq n_0$. By Proposition 1.5, this means that, for each $n \geq n_0$, the problem (3.9) has lower functions $(\sigma_{1,n}, \rho_{1,n})$ such that $\sigma_{1,n}(t) \in [A - \nu, B]$ for all $t \in [0, 2\pi]$.

To summarize, we have $e_n \in \mathcal{E}(E, \varepsilon_n, B)$ for all $n \geq n_0$. Thus, due to (3.11), Proposition 2.9 implies that for each $n \geq n_0$ the problem (3.9) has a positive solution x_n such that

$$(3.12) \quad x_n(t_n) \in [A_0, B] \quad \text{for some } t_n \in [0, 2\pi].$$

Hence, we can use Lemma 3.2 with $A = A_0$ to get that there are $R, \varepsilon^* \in (0, \infty)$ and $K \in [0, \infty)$ such that the relations $\varepsilon^* \leq x_n(t) \leq R$ on $[0, 2\pi]$ and $\|x'_n\|_{\infty} \leq K$ hold for each $n \geq n_0$. This means that the set $\{g(x_n(t)) - kx_n(t) : t \in [0, 2\pi], n \geq n_0\}$ is bounded and it follows that the sequence $\{x_n\}_{n=n_0}^{\infty}$ is equibounded and equicontinuous in $\mathbb{C}^1[0, 2\pi]$ and so, by the Arzelà - Ascoli Theorem, we can assume without loss of generality that $\{x_n\}_{n=n_0}^{\infty}$ converges in $\mathbb{C}^1[0, 2\pi]$ to a function $u \in \mathbb{C}^1[0, 2\pi]$. Consequently, for each $t \in [0, 2\pi]$ we have

$$\lim_{n \rightarrow \infty} [x'_n(t) - x'_n(0) + k \int_0^t x_n(s) ds] = \lim_{n \rightarrow \infty} \int_0^t (g(x_n(s)) + e_n(s)) ds,$$

wherefrom, using the Lebesgue Dominated Convergence Theorem, we get

$$u'(t) - u'(0) + k \int_0^t u(s) ds = \int_0^t (g(u(s)) + e(s)) ds,$$

which means that $u \in \mathbb{A}\mathbb{C}^1[0, 2\pi]$ and u is a solution to (1.1). Moreover, due to (3.12) we have (3.6).

ii) It remains to get rid of the assumption (3.7). For $n \in \mathbb{N}$, let us define

$$p_n(t) = e(t) - \frac{1}{n} \quad \text{a.e. on } [0, 2\pi].$$

For each $n \in \mathbb{N}$ and $x \in [A, B]$ we have $g(x) - kx + \overline{p}_n < 0$. Further, in view of (3.4) there are $n_0 \in \mathbb{N}$ and $A_1 \in (0, \frac{A}{2})$ such that

$$g(x) - kx + \overline{p}_n > 0 \quad \text{for all } x \in (0, A_1) \quad \text{and} \quad n \geq n_0.$$

Thus, by the first part of the proof, we get a sequence $\{x_n\}_{n=n_0}^\infty$ of solutions of the problems

$$(3.13) \quad x'' + kx = g(x) + p_n(t), \quad x(0) = x(2\pi), \quad x'(0) = x'(2\pi).$$

Since $\|p_n\|_1 \leq E + 1$ for $n \geq n_0$, we get that the solutions x_n of (3.13) satisfy $\varepsilon^* \leq x_n(t) \leq R$ on $[0, 2\pi]$ and $\|x_n'\|_\infty \leq K$, where the constants R, K and ε^* are now determined for A_1, A, B and $E + 1$ instead of A_0, A, B, E . Therefore, we can use the limiting process as in the first part of this proof and get the desired solution $u(t) = \lim_{n \rightarrow \infty} x_n(t)$ to (1.1) with the property (3.6). \square

The proof of the next theorem can be done as the previous one with the only difference that instead of Proposition 1.5 we will use Proposition 1.6.

3.4. Theorem. *Suppose that $k \neq n^2$ for all $n \in \mathbb{N}$ and replace condition (3.5) in Theorem 3.3 by*

$$(3.14) \quad g(x) + \bar{e} \leq k \frac{A+B}{2} \quad \text{for all } x \in [A, B],$$

where $A > 0$ and B fulfil (1.13) and (1.14). Then the problem (1.1) has a positive solution. \square

4 . Multiplicity results

In this section we present sufficient conditions for the existence of at least two positive solutions of (1.1). First, we will give some necessary auxiliary assertions.

4.1. Lemma. *Assume (1.2) and let $A \in (0, \infty)$ and $B \in (A, \infty)$ be such that (3.7) and (1.9) are satisfied. Then there are $\gamma : [0, 2\pi] \times [0, 2\pi] \mapsto \mathbb{R}$ continuous on $[0, 2\pi] \times [0, 2\pi]$ such that if we put*

$$(4.1) \quad \sigma_1(t) = A + \frac{\pi}{6} \|e - \bar{e}\|_1 + \int_0^{2\pi} \gamma(t, s) (e(s) - \bar{e}) ds \quad \text{for } t \in [0, 2\pi],$$

then $\sigma_1 \in \mathbb{AC}[0, 2\pi]$, $\sigma_1(t) \in [A, B]$ for all $t \in [0, 2\pi]$ and (σ_1, σ_1') are lower functions of (1.1).

Furthermore, there is $\nu_0 > 0$ such that for each $\nu \in [-\nu_0, \nu_0]$ the functions $(\sigma_1 + \nu, \sigma_1')$ are lower functions of (1.1).

Proof. Due to (3.7), we can find $\nu_0 > 0$ in such a way that

$$(4.2) \quad g(x) - kx + \bar{e} \leq 0 \quad \text{for } x \in [A - \nu_0, B + \nu_0].$$

Let $\gamma_0(t, s)$ be the Green function of the problem $x'' = 0$, $x(0) = x(2\pi) = 0$ and let

$$\sigma_0(t) = \int_0^{2\pi} \gamma(t, s) (e(s) - \bar{e}) ds \quad \text{for } t \in [0, 2\pi],$$

where

$$(4.3) \quad \gamma(t, s) = \gamma_0(t, s) - \frac{1}{2\pi} \int_0^{2\pi} \gamma_0(\tau, s) d\tau \quad \text{for } t, s \in [0, 2\pi].$$

It is easy to verify that $\sigma_0 \in \mathbb{AC}^1[0, 2\pi]$,

$$(4.4) \quad \sigma_0''(t) = e(t) - \bar{e} \quad \text{a.e. on } [0, 2\pi], \quad \sigma_0(0) = \sigma_0(2\pi), \quad \sigma_0'(0) = \sigma_0'(2\pi).$$

Moreover, $\bar{\sigma}_0 = 0$ and therefore by the proof of [12, Proposition 2.4] we have

$$(4.5) \quad \|\sigma_0\|_\infty \leq \frac{\pi}{6} \|e - \bar{e}\|_1.$$

In particular, we have

$$(4.6) \quad \sigma_1(t) = \sigma_0(t) + A + \frac{\pi}{6} \|e - \bar{e}\|_1 \quad \text{on } [0, 2\pi].$$

Now, choose an arbitrary $\nu \in [-\nu_0, \nu_0]$ and put

$$(4.7) \quad \sigma(t) = \sigma_1(t) + \nu \quad \text{for } t \in [0, 2\pi].$$

Obviously, σ and $\sigma_1 \in \mathbb{AC}^1[0, 2\pi]$ fulfil (4.4) if they are inserted there instead of σ_0 . Furthermore, (4.5)-(4.7) imply that

$$\sigma_1(t) \in [A, B] \quad \text{and} \quad \sigma(t) \in [A - \nu_0, B + \nu_0] \quad \text{for all } t \in [0, 2\pi].$$

Finally, in view of (4.2) we have $k\sigma(t) - \bar{e} \geq g(\sigma(t))$ on $[0, 2\pi]$ and, consequently,

$$\sigma''(t) + k\sigma(t) = e(t) - \bar{e} + k\sigma(t) \geq e(t) + g(\sigma(t)) \quad \text{for a.e. } t \in [0, 2\pi],$$

i.e. $(\sigma_1 + \nu, \sigma_1')$ are lower functions of (1.1) for each $\nu \in [-\nu_0, \nu_0]$. \square

4.2. Proposition. *Suppose (1.2) and let (σ_1, ρ_1) and (σ_2, ρ_2) be respectively lower and upper functions of (1.1) such that $\sigma_1(t) \leq \sigma_2(t)$ on $[0, 2\pi]$. Then there is a solution u of (1.1) such that*

$$(4.8) \quad \sigma_1(t) \leq u(t) \leq \sigma_2(t) \quad \text{on } [0, 2\pi].$$

Proof. Choose an arbitrary $f \in \text{Car}([0, 2\pi] \times \mathbb{R})$ in such a way that $f(t, x) = g(x) - kx + e(t)$ for a.e. $t \in [0, 2\pi]$ and all $x \in [\sigma_1(t), \sigma_2(t)]$. Then Theorem 1.4 ensures the existence of a solution u of (1.4) with $J = \mathbb{R}$ satisfying the estimates (4.8), which means that u is a solution to (1.1), as well. \square

4.3. Theorem. *Suppose that (1.2), (1.3), (2.2) and (2.35) hold and let $A \in (0, \infty)$ and $B \in (A, \infty)$ be such that (3.7) and (1.9) are true. Further, assume that there are upper functions (σ_2, ρ_2) of (1.1) such that $\sigma_2(t) \geq B$ for all $t \in [0, 2\pi]$. Then the problem (1.1) has at least two positive solutions.*

Proof. First, notice that by Lemma 4.1 and Proposition 4.2 the problem (1.1) has lower functions (σ_1, σ'_1) and a solution u for which (4.8) is true. Moreover, we have $\sigma_1(t) \in [A, B]$ on $[0, 2\pi]$ and there is $\nu \in (0, A)$ such that $(\sigma_1 - \nu, \sigma'_1)$ are also lower functions of (1.1).

Consider the function g_0 from (2.15). By (2.19) and (2.35) there is $A_0 \in (0, A - \nu)$ such that

$$(4.9) \quad g_0(A_0) + e(t) \geq g_0(A_0) + \inf_{t \in [0, 2\pi]} \text{ess } e(t) \geq 0 \quad \text{for a.e. } t \in [0, 2\pi].$$

This means that $(A_0, 0)$ are upper functions of (1.1). Furthermore, for a.e. $t \in [0, 2\pi]$, put $R = \|\sigma_2\|_\infty$ and

$$(4.10) \quad m(t) = e(t) + \min\{0, \inf_{s \in (0, R]} g_0(s)\}.$$

Then, in view of (2.18), $m \in \mathbb{L}[0, 2\pi]$. Denote

$$K = \|m\|_1 \quad \text{and} \quad K^* = K \|e\|_1 + \int_{A_0}^R |g_0(s)| \, ds.$$

Due to (2.16), we can choose $\varepsilon^* \in (0, A_0)$ in such a way that $g_0(\varepsilon^*) > 0$ and

$$(4.11) \quad \int_{\varepsilon^*}^{A_0} g_0(s) \, ds > K^*.$$

Now, for a.e. $t \in [0, 2\pi]$, define

$$(4.12) \quad f(t, x) = e(t) + \begin{cases} \tilde{g}_0(x) & \text{if } x \leq \sigma_2(t), \\ \tilde{g}_0(\sigma_2(t)) + \frac{x - \sigma_2(t)}{x - \sigma_2(t) + 1} & \text{if } x > \sigma_2(t), \end{cases}$$

where

$$(4.13) \quad \tilde{g}_0(x) = \begin{cases} 0 & \text{if } x < 0, \\ g_0(\varepsilon^*) \frac{x}{\varepsilon^*} & \text{if } x \in [0, \varepsilon^*), \\ g_0(x) & \text{if } x \geq \varepsilon^*. \end{cases}$$

Consider the problem (1.4) with $J = \mathbb{R}$. We have $f \in \text{Car}([0, 2\pi] \times \mathbb{R})$. Further, since $\varepsilon^* < A_0$ and (4.9) are valid, the couple $(A_0, 0)$ defines upper functions of (1.4). Similarly, since $\sigma_1(t) - \nu < \sigma_2(t)$ on $[0, 2\pi]$, the functions $(\sigma_1 - \nu, \sigma_1')$ are lower functions for (1.4). Finally, (4.10) and (4.12) imply

$$f(t, x) \geq m(t) \quad \text{for all } x \in \mathbb{R} \quad \text{and a.e. } t \in [0, 2\pi].$$

Therefore, we can apply Theorem 1.4 to obtain a solution v of (1.4) such that

$$(4.14) \quad A_0 \leq v(t_v) \leq \sigma_1(t_v) - \nu \quad \text{for some } t_v \in [0, 2\pi].$$

Relations (4.8) and (4.14) ensure that u and v are different. It remains to prove that v is a solution to (1.1). To this aim we need to show that the inequalities

$$(4.15) \quad \varepsilon^* \leq v(t) \leq \sigma_2(t) \quad \text{on } [0, 2\pi]$$

are valid. First, let us put $z(t) = v(t) - \sigma_2(t)$ for $t \in [0, 2\pi]$ and suppose that $\max_{t \in [0, 2\pi]} z(t) = z(\tau_1) > 0$. Due to the periodic conditions we can assume that $\tau_1 \in [0, 2\pi)$ and using the procedure from the proof of [11, Lemma 2.3] we can show that there is $\delta > 0$ such that $z(\tau + \delta) > z(\tau_1)$, a contradiction. This means that

$$(4.16) \quad v(t) \leq \sigma_2(t) \quad \text{on } [0, 2\pi]$$

and, in particular,

$$(4.17) \quad \max_{t \in [0, 2\pi]} v(t) = v(t_1) \in [A_0, R].$$

Now, assume that $\min_{t \in [0, 2\pi]} v(t) = v(t_0) < \varepsilon^*$. Due to (4.12) and (4.16), we have

$$v''(t) = \tilde{g}_0(v(t)) + e(t) \quad \text{a.e. on } [0, 2\pi].$$

Therefore, using (4.11), (4.13), (4.17) and Lemmas 1.7 and 1.8 we obtain

$$K^* < \int_{\varepsilon^*}^{A_0} g_0(s) \, ds \leq \int_{v(t_0)}^{A_0} \tilde{g}_0(s) \, ds \leq \|e\|_1 K + \int_{A_0}^R |g_0(s)| \, ds = K^*,$$

a contradiction. This proves that $v(t_0) \geq \varepsilon^*$, wherefrom, by virtue of (4.16), the relations (4.15) follow. \square

4.4. Theorem. *Theorem 4.3 remains valid if (3.4) is assumed instead of (2.2) and (2.35).*

Proof. Due to Lemma 4.1, the functions (σ_1, σ'_1) with σ_1 given by (4.1) are lower functions of (1.1). Therefore, by Proposition 4.2, the problem (1.1) has a solution u satisfying (4.8).

Let g_0 and e_n , $n \in \mathbb{N}$, be given by (2.15) and (3.8), respectively. Recall that the sequence $\{e_n\}_{n=1}^\infty$ is nonincreasing for a.e. $t \in [0, 2\pi]$, the relations (3.10) are true and

$$(4.18) \quad \lim_{n \rightarrow \infty} \|e_n - e\|_1 = 0.$$

Consequently, there is $E \in [0, \infty)$ such that $\|e_n\|_1 \leq E$ for all $n \in \mathbb{N}$. Due to (3.4) and (3.10) there is $A_0 \in (0, \frac{A}{2})$ such that

$$(4.19) \quad g_0(x) + \bar{e}_n \geq g_0(x) + \bar{e} > 0 \quad \text{for all } x \in (0, A_0] \quad \text{and } n \in \mathbb{N}.$$

Choose a sequence $\{\varepsilon_n\}_{n=1}^\infty \subset (0, A_0)$ in such a way that (2.26) and

$$(4.20) \quad g_0(\varepsilon_n) \geq n \quad \text{for all } n \in \mathbb{N}$$

are satisfied. Now, for $n \in \mathbb{N}$ and a.e. $t \in [0, 2\pi]$, define

$$(4.21) \quad f_n(t, x) = e_n(t) + \begin{cases} \tilde{g}_n(x) & \text{if } x < \sigma_2(t), \\ g_0(\sigma_2(t)) + \frac{x - \sigma_2(t)}{x - \sigma_2(t) + 1} & \text{if } x \geq \sigma_2(t) \end{cases}$$

and

$$(4.22) \quad \tilde{g}_n(x) = \begin{cases} |\bar{e}| + 1 & \text{if } x < 0, \\ g_0(\varepsilon_n) \frac{x}{\varepsilon_n} + (|\bar{e}| + 1) \frac{\varepsilon_n - x}{\varepsilon_n} & \text{if } x \in [0, \varepsilon_n), \\ g_0(x) & \text{if } x \geq \varepsilon_n \end{cases}$$

and consider the problems

$$(4.23) \quad x'' = f_n(t, x), \quad x(0) = x(2\pi), \quad x'(0) = x'(2\pi).$$

We will show that for all n sufficiently large the problem (4.23) verifies the assumptions of Theorem 1.4. Indeed, put $R = \|\sigma_2\|_\infty$. Then

$$g_* = \min\{|\bar{e}| + 1, \inf_{x \in (0, R]} g_0(x)\} \in \mathbb{R}$$

and $f_n(t, x) \geq e_n(t) + g_*$ for all $n \in \mathbb{N}$, $x \in \mathbb{R}$ and a.e. $t \in [0, 2\pi]$. Furthermore, from (3.10) and (4.20) we get that $f_n(t, \varepsilon_n) \geq g_0(\varepsilon_n) - n \geq 0$ holds for a.e. $t \in [0, 2\pi]$ and

each $n \in \mathbb{N}$. This implies that $(\varepsilon_n, 0)$ are upper functions of (4.23). Finally, in view of (3.7), (3.10) and (4.18) there are $n_0 \in \mathbb{N}$ and $\nu \in (0, \frac{A}{2})$ such that the relations

$$g_0(x) + \bar{e}_n < 0 \quad \text{for all } x \in [A - \nu, B] \quad \text{and} \quad \frac{\pi}{3} \|e_n - \bar{e}_n\|_1 < \frac{\pi}{3} \|e - \bar{e}\|_1 + \frac{\nu}{2}$$

hold for all $n \geq n_0$. Moreover, by (1.9) we have

$$(B - \frac{\nu}{2}) - (A - \nu) = B - A + \frac{\nu}{2} > \frac{\pi}{3} \|e_n - \bar{e}_n\|_1 \quad \text{for } n \geq n_0.$$

Thus, according to Lemma 4.1, for each $n \geq n_0$ the functions $(\sigma_{1,n}, \sigma'_{1,n})$, where

$$(4.24) \quad \sigma_{1,n}(t) = A + \frac{\pi}{6} \|e_n - \bar{e}_n\|_1 + \int_0^{2\pi} \gamma(t, s) (e_n(s) - \bar{e}_n) ds - \nu, \quad t \in [0, 2\pi],$$

are lower functions of the problem

$$x'' = g_0(x) + e_n(t), \quad x(0) = x(2\pi), \quad x'(0) = x'(2\pi)$$

and

$$(4.25) \quad \sigma_{1,n}(t) \in [A - \nu, B - \frac{\nu}{2}] \quad \text{for all } t \in [0, 2\pi].$$

Since we have

$$\varepsilon_n < \frac{A}{2} < A - \nu \leq B \leq \sigma_2(t) \quad \text{for all } t \in [0, 2\pi] \quad \text{and} \quad n \geq n_0,$$

it is easy to see that the functions $(\sigma_{1,n}, \sigma'_{1,n})$ are also lower functions of (4.23).

Thus, we can use Theorem 1.4 and Lemma 1.8 to show that for each $n \geq n_0$ the problem (4.23) has a solution x_n such that

$$(4.26) \quad \|x'_n\|_\infty \leq K = E + 2\pi |g_*|$$

and

$$(4.27) \quad \varepsilon_n \leq x_n(t_n) \leq \sigma_{1,n}(t_n) \quad \text{for some } t_n \in [0, 2\pi].$$

Now, using the arguments from the proof of Theorem 4.3 (see the proof of (4.16)), we can show that

$$(4.28) \quad x_n(t) \leq \sigma_2(t) \quad \text{on } [0, 2\pi].$$

With regard to (4.21) this means that for each $n \in \mathbb{N}$ the function x_n is a solution of

$$x'' = \tilde{g}_n(x) + e_n(t), \quad x(0) = x(2\pi), \quad x'(0) = x'(2\pi).$$

Using (3.10), (4.19) and (4.22) we can verify that there is $n_1 \geq n_0$ such that

$$\widetilde{g}_n(x) + \overline{e}_n > 0 \quad \text{for all } x \in (-\infty, A_0] \quad \text{and all } n \geq n_1.$$

Therefore, by Lemma 3.1 we have

$$\max_{t \in [0, 2\pi]} x_n(t) > A_0 \quad \text{for all } n \geq n_1$$

and Lemma 2.2 yields that there is $\varepsilon^* > 0$ such that

$$(4.29) \quad x_n(t) \geq \varepsilon^* \quad \text{for all } t \in [0, 2\pi] \quad \text{and } n \geq n_1.$$

In view of (4.26), (4.28) and (4.29), the sequence $\{x_n\}_{n=n_1}^\infty$ is equibounded and equicontinuous in $\mathbb{C}^1[0, 2\pi]$ and thus we can assume without loss of generality that it converges in $\mathbb{C}^1[0, 2\pi]$ to some function v . With regard to (4.22), (4.28) and (4.29), we have

$$\varepsilon^* \leq v(t) \leq \sigma_2(t) \quad \text{on } [0, 2\pi].$$

Therefore v is a solution to (1.1).

It remains to show that v differs from u . First, notice that since $\|\gamma\|_\infty = \sup_{t,s \in [0, 2\pi]} |\gamma(t, s)| < \infty$, the relations (4.1), (4.18) and (4.24) yield

$$(4.30) \quad \lim_{n \rightarrow \infty} \|\sigma_1 - \nu - \sigma_{1,n}\|_\infty \leq \lim_{n \rightarrow \infty} (\|\gamma\|_\infty + \frac{\pi}{6}) \|e - e_n\|_1 = 0.$$

Furthermore, we can choose a subsequence $\{t_{n_\ell}\}_{\ell=1}^\infty$ in $\{t_n\}_{n=1}^\infty$ in such a way that $\lim_{\ell \rightarrow \infty} t_{n_\ell} = t^* \in [0, 2\pi]$. Therefore, in view of (4.30), (4.8) and (4.27) we have

$$v(t^*) = \lim_{\ell \rightarrow \infty} x_{n_\ell}(t_{n_\ell}) \leq \lim_{\ell \rightarrow \infty} \sigma_{1,n_\ell}(t_{n_\ell}) = \sigma_1(t^*) - \nu < \min_{t \in [0, 2\pi]} u(t),$$

which completes the proof of the theorem. \square

4.5. Remark. Notice that in contrast to Theorems 2.1 and 3.3, in Theorems 4.3 and 4.4 we do not need to assume (2.1).

In conclusion we will give two additional multiplicity results (cf. Theorem 4.7). Their proofs can be done as those of Theorems 4.3 and 4.4, only instead of Lemma 4.1 we have to use its modification given below, which is related to Proposition 1.6.

4.6. Lemma. *Assume (1.2). Let $k \neq n^2$ for all $n \in \mathbb{N}$ and let $A \in (0, \infty)$ and $B \in (A, \infty)$ be such that*

$$(4.31) \quad g(x) + \overline{e} < k \frac{A+B}{2} \quad \text{for } x \in [A, B],$$

(1.13) and (1.14) are true. Let

$$\sigma_1(t) = \frac{A+B}{2} + \int_0^{2\pi} \tilde{\gamma}(t,s)(e(s) - \bar{e}) ds \quad \text{for } t \in [0, 2\pi],$$

where $\tilde{\gamma}$ is the Green function of the problem $x'' + kx = 0$, $x(0) = x(2\pi)$, $x'(0) = x'(2\pi)$.

Then $\sigma_1 \in \mathbb{AC}^1[0, 2\pi]$, $\sigma_1(t) \in [A, B]$ for all $t \in [0, 2\pi]$ and there is $\nu_0 > 0$ such that for each $\nu \in [-\nu_0, \nu_0]$ the functions $(\sigma_1 + \nu, \sigma_1')$ are lower functions of (1.1).

Proof. Choose $\nu_0 > 0$ in such a way that

$$(4.32) \quad g(x) - k\nu + \bar{e} \leq k \frac{A+B}{2} \quad \text{for } x \in [A - \nu_0, B + \nu_0] \quad \text{and } \nu \in [-\nu_0, \nu_0].$$

By the proofs of [13, Theorems 3.1 and 3.2], the function

$$\sigma_0(t) = \int_0^{2\pi} \tilde{\gamma}(t,s)(e(s) - \bar{e}) ds, \quad t \in [0, 2\pi],$$

possesses the following properties: $\sigma_0 \in \mathbb{AC}^1[0, 2\pi]$, $\bar{\sigma}_0 = 0$,

$$\sigma_0''(t) + k\sigma_0(t) = e(t) - \bar{e} \quad \text{a.e. on } [0, 2\pi], \quad \sigma_0(0) = \sigma_0(2\pi), \quad \sigma_0'(0) = \sigma_0'(2\pi)$$

and $\|\sigma_0\|_\infty \leq \Phi(k) \|e - \bar{e}\|_1$. In view of (1.13) we have

$$\sigma_1(t) = A + \Phi(k) \|e - \bar{e}\|_1 + \sigma_0(t) \quad \text{on } [0, 2\pi].$$

Therefore $\sigma_1(t) \in [A, B]$ for all $t \in [0, 2\pi]$ and it follows from (4.32) that $(\sigma_1 + \nu, \sigma_1')$ are lower functions of (1.1) for each $\nu \in [-\nu_0, \nu_0]$. \square

4.7. Theorem. *Suppose that $k \neq n^2$ for all $n \in \mathbb{N}$ and replace conditions (3.7) and (1.9) in Theorem 4.3 (Theorem 4.4) with (4.31), (1.13) and (1.14). Then the problem (1.1) has at least two positive solutions.*

References

- [1] M. DEL PINO, R. MANÁSEVICH AND A. MONTERO. T -periodic solutions for some second order differential equations with singularities. *Proc. Royal Soc. Edinburgh* **120A**, 231-244 (1992).
- [2] A. FONDA. Periodic solutions of scalar second order differential equations with a singularity. *Mémoire de la Classe de Sciences, Acad. Roy. Belgique*, **8-IV**, 1-39 (1993).
- [3] A. FONDA, R. MANÁSEVICH AND F. ZANOLIN. Subharmonic solutions for some second-order differential equations with singularities. *SIAM J. Math. Anal.* **24**, 1294-1311 (1993).

- [4] W. GE AND J. MAWHIN. Positive solutions to boundary value problems for second-order ordinary differential equations with singular nonlinearities. *Result. Math.* **34**, 108-119 (1998).
- [5] P. HABETS AND L. SANCHEZ. Periodic solutions of some Liénard equations with singularities. *Proc. Amer. Math. Soc.* **109** (1990), 1035-1044.
- [6] A. C. LAZER AND S. SOLIMINI. On periodic solutions of nonlinear differential equations with singularities. *Proc. Amer. Math. Soc.* **99** (1987), 109-114.
- [7] J. MAWHIN. Topological degree and boundary value problems for nonlinear differential equations. M. Furi (ed.), *Topological methods for ordinary differential equations*. Berlin: Springer-Verlag, Lect. Notes Math. 1537 (1993), 73-142.
- [8] D. S. MITRINOVIĆ, J. E. PEČARIĆ AND A. M. FINK. *Inequalities Involving Functions and Their Integrals and Derivatives*. Kluwer, Dordrecht, 1991.
- [9] P. OMARI AND W. YE. Necessary and sufficient conditions for the existence of periodic solutions of second order ordinary differential equations with singular nonlinearities. *Differential Integral Equations* **8** (1995), 1843-1858.
- [10] I. RACHŮNKOVÁ. On the existence of more positive solutions of periodic BVP with singularity. *Appl. Anal.*, to appear.
- [11] I. RACHŮNKOVÁ AND M. TVRĎÝ. Nonlinear systems of differential inequalities and solvability of certain nonlinear second order boundary value problems. *J. Inequal. Appl.* **6** (2001), 199-226.
- [12] I. RACHŮNKOVÁ AND M. TVRĎÝ. Construction of lower and upper functions and their application to regular and singular boundary value problems. *Nonlinear Analysis, T.M.A.* **47** (2001), 3937-3948.
- [13] I. RACHŮNKOVÁ AND M. TVRĎÝ. Localization of nonsmooth lower and upper functions for periodic boundary value problems. *Math. Bohem.*, to appear.
- [14] I. RACHŮNKOVÁ, M. TVRĎÝ AND I. VRKOČ. Existence of nonnegative and nonpositive solutions for second order periodic boundary value problems. *J. Differential Equations*, **176** (2001), 445-469.
- [15] L. SCHEEFFER. Über die Bedeutung der Begriffe "Maximum und Minimum" in der Variationsrechnung. *Math. Ann.* **26** (1885), 197-208.
- [16] P. YAN AND M. ZHANG. Higher order nonresonance for differential equations with singularities. Preprint.
- [17] M. ZHANG. A relationship between the periodic and the Dirichlet BVP's of singular differential equations. *Proc. Royal Soc. Edinburgh* **128A** (1998), 1099-1114.

Irena Rachůnková, Department of Mathematics, Palacký University, 779 00 OLOMOUC, Tomkova 40, Czech Republic (e-mail: rachunko@risc.upol.cz)

Milan Tvrďý, Mathematical Institute, Academy of Sciences of the Czech Republic, 115 67 PRAHA 1, Žitná 25, Czech Republic (e-mail: tvrды@math.cas.cz)

Ivo Vrkoč, Mathematical Institute, Academy of Sciences of the Czech Republic, 115 67 PRAHA 1, Žitná 25, Czech Republic (e-mail: vrkoc@ns.math.cas.cz)