

References:

- 1) A.E.Shishkov.
Localized boundary blow-up regimes for general divergent parabolic equations of arbitrary order.
Proc. of the Steklov Institute of Math. **236** (2002).
- 2) A.Shishkov, A.Shchelkov.
Blow-up boundary regimes for general quasilinear parabolic equations in multidimensional domains.
Sbornik:Mathematics **190** (1999),190:3, pp. 447-479.
- 3) V.Galaktionov, A.Shishkov.
Saint-Venant principle in blow-up for higher order quasilinear PDE.
Proc. Roy.Soc.Edinburgh. **133A** (2003), pp. 1075-1119.
- 4) V.Galaktionov, A.Shishkov.
Structure of boundary blow-up for higher order quasilinear parabolic PDE.
Proc. Roy. Soc. London.A **460** (2004), pp. 3299-3325.
- 5) V.Galaktionov, A.Shishkov.
Self-similar boundary blow-up for higher order quasilinear PDE.
Proc. Roy. Soc. Edinburgh. **135A** (2005), pp.1195-1227.
- 6) V.A.Galaktionov, A.E.Shishkov.
Higher-order quasilinear PDE with singular initial data.
Preprint 05/01 of Bath Univ., 2005, 20 p.
(<http://www.maths.bath.ac.uk/MATHEMATICS/preprints.html>)
to appear in *Commun. Contemporary Math.*
- 7) A.Shishkov, L.Veron.
Diffusion versus absorption in semilinear parabolic problems.
(to appear in *Compt. Rend. Acad. Sci. Paris.*)