

Associations between physical activity and the androgenic/estrogenic status of men

M.Slowinska-Lisowska¹, Jozkow P¹, Medras M^{1,2},

¹Dept. of Sports Medicine, University School of Physical Education, Wrocław, Poland

²Dept. of Endocrinology, Diabetology and Isotope Treatment, Wrocław Medical University, Wrocław, Poland

Corresponding author:

Malgorzata Slowinska-Lisowska

Department of Sports Medicine

University School of Physical Education

ul. Paderewskiego 35

51-617 Wrocław, Poland

e-mail: slowik29@ gmail.com

Short title: **Physical activity and androgenic/estrogenic status of men**

Summary

Concentrations of numerous hormones decrease with age. Some authors imply that a syndrome of partial endocrine deficiency may occur in the aging men. Among many lifestyle factors that influence hormonal status is physical activity. Especially interesting are relations between physical activity and the androgenic/estrogenic status of men.

The aim of this study was to evaluate age-related changes of serum androgens, estradiol and SHBG in men presenting different levels of physical activity.

Hormonal parameters were measured in a cohort of 387 healthy Caucasian men from one administrative region of Poland. They aged from 24 to 72 years. Their level of physical activity was determined with use of the International Physical Activity Questionnaire (IPAQ).

We have found that contrary to SHBG concentration, total testosterone, free testosterone, bioavailable testosterone, calculated free testosterone and estradiol were negatively associated with age in the investigated subjects.

Apart from estradiol, physical activity did not influence concentrations of the studied parameters. In younger (24-48 y), physically active males estradiol was significantly higher than in subjects characterized by a low level of physical activity. The situation was opposite in older males (48-72 y.). In this age group low level of physical activity was associated with lower concentration of estradiol. Undertaking physical effort increased the decline of estradiol level with age.

Key words

Hormones, age, motor activity; lifestyle

Introduction

Concentration of numerous hormones decrease with age. This phenomenon has an individual dynamics and it depends on genetic, psychical and social factors (stress, depressive states). It is also influenced by lifestyle: tobacco smoking, alcohol drinking and physical activity (Field *et al.* 1994, Svartberg and Jorde 2007, Shiels *et al.* 2009, Wolin *et al.* 2007, Muller *et al.* 2003).

The influence of repeated physical efforts on the androgenic status of aging men is not fully explored. Testosterone deficiency is supposed to be associated with vanishing of the muscle mass, increase of the visceral fat, insulin resistance, blood hypertension, dyslipidemia as well as osteopenia (Denti *et al.* 2000, Vanderschueren 2000).

Estrogens warrant proper function of the reproductive, skeletal and nervous systems in men. Although it is assumed that male aging is accompanied by a gradual increase of estrogens, not all authors confirm this observation. Data on associations between estrogens and physical activity is scarce (Drafta *et al.* 1982, Shiels *et al.* 2009).

We wanted to evaluate the impact of physical activity on age-related trends in hormonal status of men living in the Lower Silesia Region.

Materials and Methods

Our investigation is part of a larger project named Health of Adults in Lower Silesia (HALS). The region of Lower Silesia is located in southern Poland and covers an area of 19 947 sq km. It has 2 880 000 inhabitants (100% Caucasian). The project of the study was approved by the Bioethics Committee at the University School of Physical Education, Wroclaw, Poland.

We sent invitations to participate in the investigation to 900 adult men living in Lower Silesia. They were randomly selected by the Local Data Bank (Regional Statistical Office).

As an operator the personal identification number was used. From the target group, 372 subjects (43%) submitted their written consents and were enrolled for the study.

The subjects aged from 24 to 72 years. They underwent a standard medical interview and a full physical examination. About 43% of subjects had university degree. According to the medical interviews 21% had arterial hypertension, 19% chronic low back pain, 5% benign prostate hyperplasia and 4% diabetes. The anthropological characteristics of the studied men is shown in Tables 1 and 2. Subjects distinguished upon the level of physical activity did not differ as to age, height, weight and body mass index (BMI).

Physical activity of the studied subjects was determined with use of the Polish version of the International Physical Activity Questionnaire (IPAQ, short form ,last 7-day recall). IPAQ is commonly used as a comparable and standardized self-report measure of habitual physical activity of populations from different sociocultural contexts. It has been validated in healthy, adults from 12 countries (14). The questionnaire was also compared against doubly labeled water what implicated that it may underestimate energy expenditure at higher levels of physical activity (15). In the continuous scale the amount of physical exercise was counted as multiples of resting metabolic rate by minutes of performance during a week (METs-minute/week). In our study group we adapted the criteria presented by the IPAQ scientific committee [<http://www.ipaq.ki.se/>]. According to above classification persons who moved at least 12 500 steps a day or the equivalent in moderate or vigorous activities were highly active. They had at least an hour more moderate-intensity activity over and above the basal level of activity or half an hour of vigorous-intensity activity over and above basal levels daily (basal is 5 000 steps/day). Subjects who did not fulfill above criteria were assigned to the subgroup of low physical activity.

Statistical methods

Models evaluating concentrations of hormones in groups of different physical activity levels together with anthropological parameters and age were built using least squares multiple linear ridge stepwise regression. The statistical significance of models was tested by an F-test from a one-way ANOVA. The statistical significance of variables was tested using Student t-test. Non-significant variables in the models were eliminated by forward selection. All calculations were performed using statistical package *Statistica* v.6 (Statsoft Inc.)

Hormone Measurements

Radioimmunoassay kits were used for measurements of total testosterone (TT), free testosterone and estradiol (Diagnostic Products Corporation, USA). The intra- and interassay coefficients of variation (CV) were 5.5% and 5.9 % for total testosterone, 3.2% and 8.5% for free testosterone, 3.1 % and 4.5 % for estradiol. SHBG was measured using immunoradiometric assay- IRMA kits (Immunotech, The Czech Republic). The intra- and interassay CV were 3.8 and 7.0 %.

The concentrations of bioavailable testosterone and calculated free testosterone were assessed with a calculator developed at the Hormonology Department, University Hospital of Ghent, Belgium (details on the calculation are available on the web site: <http://www.issam.ch/freetesto.htm>).

Results

The multiple regression models for hormonal levels with statistically significant regresses are shown in Table 3.

Diagrams of spread and the regression straight line of the concentrations of the investigated hormones with age are shown in Fig. 1- 6.

We have found that concentrations of total testosterone, free testosterone, bioavailable testosterone, calculated free testosterone, SHBG and estradiol were significantly influenced by age. Physical activity did not modify concentrations of androgens in the investigated men.

The mean estradiol level in men aged 24 – 48 was higher in the subgroup declaring high level of physical activity. Contrary to above, in men >48 y. estradiol was lower in those who were physically active.

It's worth noticing that estradiol decreased by approximately 5.2 nmol/l per 10 years in highly active men. In subjects who declared low level of physical activity this decrease was insignificant (0.2 nmol/l per 10 years).

Discussion

Previous studies of the Polish population suggested rather high incidence of sedentary lifestyle and a relatively low level of occupational physical activity (Kaleta *et al.* 2007). Unlike above, in our investigation about 50% of men declared high level of daily physical activity. Similar outcomes were reported by authors who showed that the amount of physical activity was higher when IPAQ questionnaires were filled by subjects themselves rather than by experienced interviewers (Biernat *et al.* 2008). We have to mention a suggestion that IPAQ may overestimate the level of physical activity (Rzewnicki *et al.* 2003). On the other hand, it is probable that the investigated men were more health-conscious than average.

Our results are consistent with observations pointing to a decreased production of androgens and increased serum concentration of sex hormone binding globulin in aging men (Gray *et al.* 1991, Kaufman and Vermeulen 2005, Van den Beld *et al.* 1999).

Influence of age on estradiol concentration is not so clear. Authors of several studies found that the level of total estradiol does not change with age in men (Ferrini and Barret-Connor 1998, Muller *et al.* 2003). Others observed a decrease of estradiol only in the old age (Van den Beld *et al.* 2000). However in a cross-sectional study of men aged 25-85, bioavailable E₂ decreased significantly with age (Khosla *et al.* 1998). Associations between E₂ levels and ER alpha polymorphisms were suggestive for an estrogen-related processes, possibly related to changes in the neuroendocrine regulation of testosterone production in elderly men (Lapauw *et al.* 2008).

In our investigation estradiol was negatively associated with age and it's concentration was influenced by physical activity. We have noticed that in men between 24–48 y. estradiol concentration was higher in physically active rather than inactive subjects. At 48 y. the mean

estradiol concentration was similar in both groups. In men >48 y. physically active subjects had lower concentration of estradiol.

The age-dependent decline of estradiol was apparent in physically active men. The decrease of estradiol in inactive subjects was statistically insignificant. One may suspect that exercise exacerbates estradiol reduction with age.

It was noticed previously that vigorous physical activity undertaken more than four times per week increased total estradiol concentration more than less frequent vigorous exercise. According to Shiels et al.2009. such results question the opinion that those who engage in vigorous physical activity tend to have lower body fat, and thus should have lower estradiol production by adipocytes given the same testosterone level.

We have no explanation for a steeper estradiol decline with age in active subjects and a higher concentration of estradiol in young, physically active men. We underline the fact that there were no differences in weight and BMI between subgroups distinguished upon physical activity level. This finding can be a result of differences of the aromatase activity, as the physiological balance between sex steroid hormones is largely controlled by the aromatase cytochrome P-450 (P450arom) (Conley and Hinshelwood 2001)

Studies performed in rats proved that single-time physical effort in male subjects evokes an increase in the aromatase cytochrome P-450 (P450arom) in the skeletal muscles, reversely to the female subjects. After an intensive effort, in tissues obtained from the male subjects the estradiol concentration increased, and in the female subjects – the testosterone concentration increased (Aizawa *et al.* 2008)

Associations between physical activity and concentrations of androgens are not fully elucidated. In our sample physical activity did not modify age-dependent changes of: total testosterone, free testosterone, bioavailable testosterone, calculated free testosterone and SHBG in men aged 24-72 y .

Some authors reported positive associations between physical activity and androgen levels. An exercise-induced testosterone increase was found to be smaller in the elderly (Swerdloff and Wang 1993). Allen et al found that vigorous exercise lasting three or more hours per week is positively associated with testosterone and SHBG, but not free testosterone (Allen *et al.* 2002). Muller suggested that an increase in total testosterone, bioavailable testosterone and SHBG appears in the highly active group (Muller *et al.* 2003).

On the other hand several studies did not reveal any associations between physical activity and androgens (Ponholzer *et al.*, 2005, Svartberg *et al.* 2003, Ukkola *et al.* 2001, Shiels *et al.* 2009). We also reported a lack of association between androgen status and physical activity in men aged 45-58. There were no statistically significant differences of total testosterone, free testosterone, dehydroepiandrosterone sulphate and estradiol concentrations between physically active (1.0-1.5 hours of exercise twice a week, for about ten years prior examination) and physically inactive middle-aged men (Medras *et al.* 2005).

It has been also suggested that low testosterone reduces the level of physical activity (Van den Beld *et al.* 2000). An individually defined low level of androgens may be the cause of decreased physical activity.

Conclusions

Total testosterone, free testosterone, bioavailable testosterone, calculated free testosterone and estradiol decrease with age, reversely to the concentration of SHBG.

The level of physical activity does not influence concentrations of androgens or SHBG. In younger, physically active males (24-48 y.) serum estradiol is higher than in less active ones. To the opposite, older males (48-72 y.) presenting higher level of physical activity have lower estradiol levels. Physical effort increases estradiol decline with age.

Acknowledgements

This work was supported by the Ministry of Science and Higher Education of Poland (grant code: 2PO5D 0730).

References

- AIZAWA K, IEMITSU M, OTSUKI T, MAEDA S, MIYAUCHI T, MESAHI N: Sex differences in steroidogenesis in skeletal muscle following a single bout of exercise in rats. *J Appl Physiol.* **104(1)**, 67-74, 2008.
- ALLEN NE, APPLEBY PN, DAVEY GK, KEY TJ. Lifestyle and nutritional determinants of bioavailable androgens and related hormones in British men. *Cancer Causes Control.* **13(4)**, 353-63, 2002.
- CONLEY A, HINSHELWOOD M :Mammalian aromatases. *Reproduction.* **121(5)**, 685-95, 2001.
- DENTI L, PASOLINI G, SANFELICI L, BENEDETTI R, CECCHETTI A, CEDA GP, ABLONDI F, VALENTI G. : Aging-related decline of gonadal function in healthy men: correlation with body composition and lipoproteins. *Am Geriatr Soc.* **48(1)**, 51-58, 2000
- DRAFTA D, SCHINDLER AE, STROE E, NEACȘU E : Age-related changes of plasma steroids in normal adult males. *J Steroid Biochem.* **17(6)**, 683-7, 1982
- FERRINI RL, BARRET-CONNOR E : Sex hormones and age: a cross-sectional study of testosterone and estradiol and their bioavailable fractions in community dwelling men. *American Journal of Epidemiology*, **147**,750-754, 1998.
- FIELD AE, COLDITZ GA, WILLETT WC, LONGCOPE C, MC KINLAY JB: The relation of smoking, age, relative weight, and dietary intake to serumadrenal steroids,

sex hormones, and sex hormone-binding globulin in middle-aged men. *J Clin Endocrinol Metab* **79(5)**, 1310-1316, 1994

GRAY A, FELDMAN HA, MCKINLAY JB, LONGCOPE C : Age, disease, and changing sex hormone levels in middle-aged men: results of the Massachusetts Male Aging Study. *J Clin Endocrinol Metab.* **73(5)**, 1016-25, 1991

KALETA D, POLAŃSKA K, JEGIER A, BIERNAT E, STUPNICKI R, LEBIEDZIŃSKI B, JANCZEWSKA L: Assesment of physical activity by applying IPAQ questionnaire . *Physical Education and Sport:* **52(2)**, 83-89 2008.

KAUFMAN JM, VERMEULEN A: The decline of androgen levels in elderly men and its clinical and therapeutic implications. *Endocr Rev.* **26(6)**, 833-76, 2005

KHOSLA S, MELTON LJ, ATKINSON EJ, O'FALLON WM, KLEE GG, RIGGS BL :Relationship of serum sex steroid levels and bone turnover markers with bone mineral density in men and women: a key role for bioavailable estrogen. *J Clin Endocrinol Metab.* **83(7)**, 2266-2274, 1998.

LAPAUW B, GOEMAERE S, ZMIERCZAK H, VAN POTTELBERGH I, MAHMOUD A, TAES Y, DEBACQUER D, VANSTEELANDT S, KAUFMAN JM :The decline of serum testosterone levels in community-dwelling men over 70 years of age: descriptive data and predictors of longitudinal changes. *European Journal of Endocrinology.* **159(4)**, 459-68, 2008.

MEDRAS M, SLOWINSKA-LISOWSKA M, JOZKOW P: Impact of recreational physical activity on bone mineral density in middle-aged men. *Aging Male.* **8(3-4)**, 162-165, 2005.

MULLER M., TONKELAAR I, THIJSEN J, GROBBEE D, SCHOUW YT :Endogenous sex hormones in men aged 40-80 year. *European Journal of Endocrinology* **149**, 583-589, 2003.

PONHOLZER A, PLAS E, SCHATZL G, STRUHAL G, BRÖSSNER C, MOCK K, RAUCHENWALD M, MADERSBACHER S SVARTBERG J, MIDTBY M, BØNAA KH, SUNDSFJORD J, JOAKIMSEN RM, JORDE R: Relationship between testosterone serum levels and lifestyle in aging men. *Aging Male* **8(3-4)**, 190-193, 2005.

RZEWNICKI R, VANDEN AUWEELE Y, DE BOURDEAUDHUIJ I: Addressing overreporting on the International Physical Activity Questionnaire(IPAQ) telephone survey with a population sample. *J Public Health Nutr.* **6(3)**, 299-305, 2003

SHIELS MS, ROHRMANN S, MENKE A, SELVIN E, CRESPO CJ, RIFAI N, DOBS A, FEINLEIB M, GUALLAR E, PLATZ EA : Association of cigarette smoking, alcohol consumption, and physical activity with sex steroid hormone levels in US men. *Cancer Causes Control.* 2009, [Epub ahead of print]

SVARTBERG J, JORDE R: Endogenous testosterone levels and smoking in men. The fifth Tromsø study *Int J Androl.* **30(3)**, 137-143, 2006

SVARTBERG J, MIDTBY M, BØNAA KH, SUNDSFJORD J, JOAKIMSEN RM, JORDE R :The associations of age, lifestyle factors and chronic disease with testosterone in men: the Tromsø Study. *Eur J Endocrinol.* **149(2)**, 145-152, 2003.

SWERDLOFF RS, WANG C :Androgens and aging in men. *Exp Gerontol.* **28(4-5)**,435-46, 1993.

UKKOLA O, GAGNON J, RANKINEN T, THOMPSON PA, HONG Y, LEON AS, RAO DC, SKINNER JS,WILMORE JH, BOUCHARD C.. Age, body mass index, race and other determinants of steroid hormone variability:the HERITAGE Family Study.*Eur J Endocrinol.* **145(1)**, 1-9, 2001

VAN DEN BELD A, HUHTANIEMI IT, PETTERSSON KS, POLS HA, GROBBEE DE, DE JONG FH, LAMBERTS SW: Luteinizing hormone and different genetic

variants, as indicators of frailty in healthy elderly men. *J Clin Endocrinol Metab.* **84(4)**, 1334-1339, 1999

VAN DEN BELD AW, DE JONG FH, GROBBEE DE, POLS HA, LAMBERTS SW: Measures of bioavailable serum testosterone and estradiol and their relationships with muscle strength, bone density, and body composition in elderly men. *J Clin Endocrinol Metab.* **85(9)**, 3276-3282, 2000.

VAN DEN BELD AW, DE JONG FH, GROBBEE DE, POLS HA, LAMBERTS SW :Measures of bioavailable serum testosterone and estradiol and their relationships with muscle strength, bone density, and body composition in elderly men. . *J Clin Endocrinol Metab.* **85(9)**, 3276-3282, 2000

VANDERSCHUEREN D, VANDENPUT L: Androgens and osteoporosis. *Andrologia.* **32(3)**, 125-130, 2000

WOLIN KY, COLANGELO LA, LIU K, STERNFELD B, GAPSTUR SM : Associations of androgens with physical activity and fitness in young black and white men: the CARDIA Male Hormone Study. *Prev Med.* **44(5)**, 426-31, 2007

Table 1. Basic data (mean \pm SD) of the study group

Parameter	mean \pm SD	High activity (n=191)	low activity (n=196)
Age [years]	48,3 \pm 14,6	47,1 \pm 14,3	49,5 \pm 14,6
Weight [kg]	84,9 \pm 13,4	85,3 \pm 13,5	84,7 \pm 13,4
Height [cm]	175,0 \pm 6,9	176,0 \pm 6,5	175,6 \pm 6,8
Body mass index [kg/(m) ²]	27,5 \pm 4,1	27,3 \pm 4,1	27,6 \pm 4,2

Table 2 Basic data (mean \pm SD) of the study group

Age [years]	24-30 N=91	31-40 N=68	41-50 N=96	51-60 N=98	61-72 N=34
Weight [kg]	81,8 \pm 12,5	85,7 \pm 13,1	87,0 \pm 13,8	85,8 \pm 15,1	81,9 \pm 11,7
Height [cm]	179,2 \pm 6,2	177,7 \pm 5,7	175,9 \pm 6,3	172,0 \pm 7,0	172,5 \pm 6,9
Body mass index [kg/(m) ²]	25,4 \pm 3,4	27,1 \pm 3,8	28,1 \pm 4,0	28,9 \pm 4,8	27,5 \pm 3,3




Fig. 1 Concentration of Estradiol by age for men with low (L) and high (H) physical activity




Fig. 2 Concentration of Total Testosterone by age for men with low (L) and high (H) physical activity




Fig. 3 Concentration of Bioavailable Testosterone by age for men with low (L) and high (H) physical activity




Fig. 4 Concentration of SHBG by age for men with low (L) and high (H) physical activity




Fig. 5 Concentration of Free Testosterone by age for men with low (L) and high (H) physical activity




Fig. 6 Concentration of Calculated Free Testosterone by age for men with low (L) and high (H) physical activity