

On the existence of a σ -closed dense subset*

Jindřich Zapletal^{†‡}
Academy of Sciences, Czech Republic
University of Florida

December 20, 2009

Abstract

It is consistent with the axioms of set theory that there are two co-dense partial orders, one of them σ -closed and the other one without a σ -closed dense subset.

1 Introduction

One of the oldest properties of partial orders occurring in forcing arguments is σ -closedness. A partial ordering P, \leq is σ -closed if every countable decreasing sequence of elements of P has a lower bound. This property easily implies that forcing with P adds no new reals, preserves stationary subsets of ω_1 and so on. In this note, partially answering a question of Bohuslav Balcar, I will prove that having a σ -closed dense subset is not a forcing property of partial orders—it is not invariant under the co-density equivalence. The story is somewhat parallel to the Axiom A case. While Axiom A is a property of posets that was used with great success in the early years of forcing and still occurs in many textbooks, it is not really a forcing property of posets in this sense. I will prove

Theorem 1.1. *It is consistent with ZFC set theory that there is a partial order $\langle P \cup Q, \leq \rangle$ such that both P and Q are dense parts in it, P is σ -closed, while Q has no σ -closed dense subset.*

The method of proof closely follows the argument of [3]. The result is perhaps not entirely satisfactory in the sense that the existence of such partial orders may be a theorem of ZFC, and it is even not excluded that the σ -closed part P may be isomorphic to one of the standard σ -closed partial orders such as adding \aleph_2 many subsets of ω_1 with countable approximations. In the model for

*2000 AMS subject classification 03E40.

[†]Partially supported by NSF grant DMS 0801114 and Institutional Research Plan No. AV0Z10190503 and grant IAA100190902 of GA AV ČR.

[‡]Institute of Mathematics of the AS CR, Žitná 25, CZ - 115 67 Praha 1, Czech Republic, zapletal@math.cas.cz

the theorem, the continuum hypothesis holds and the posets have size \aleph_2 , which is minimal possible by the results of Foreman [1] and Vojtáš [4].

The notation of the paper follows the set theoretic standard of [2].

2 The proof

Work in the theory ZFC+CH. The partial orders P and Q are added by countable approximations. Define a partial order R to consist of quintuples $r = \langle P_r, Q_r, \leq_r, C_r, F_r \rangle$ such that

1. P_r, Q_r are disjoint countable subsets of ω_2 ;
2. \leq_r is a partial order on $P_r \cup Q_r$ such that both P_r and Q_r are dense in it, and moreover $\leq_r \cap \in = 0$;
3. C_r is a countable set of descending chains in the poset Q_r, \leq_r with no lower bound in \leq_r ;
4. $F_r : P_r \times C_r \rightarrow Q_r$ is a function such that $F_r(p, c) \in c$ is an element of the chain c such that every common lower bound $p' \leq p, F_r(p, c)$ in P_r is incompatible with some element of the chain c in \leq_r .

The ordering on R is defined by $r_1 \leq r_0$ if each coordinate of r_0 is a subset of the corresponding coordinate of r_1 and moreover, if $p, q \in P_{r_0} \cup Q_{r_0}$ are incompatible (resp. incomparable) in \leq_{r_0} then they are also incompatible (resp. incomparable) in \leq_{r_1} .

A bit of explanation is necessary here. Let $G \subset R$ be a generic filter and look into the model $V[G]$. The partial order $\langle P \cup Q, \leq \rangle$ from the main theorem is obtained from the generic filter G as the unions of the first three coordinates of the conditions in the generic filter. The last requirement in the second item is necessary to avoid the possibility that the density of P is \aleph_1 , which would be impossible by Foreman's result. The σ -closure of P will be guaranteed by a density argument. The descending chains in the set $C = \bigcup_{r \in G} C_r$ will have no lower bounds and will be plentiful enough so that Q will contain no σ -closed dense subset. The function F_r is a technical tool that guarantees that adding a lower bound to a countable decreasing chain in P does not necessitate adding a lower bound to one of the chains in C .

I will proceed with a series of more or less immediate lemmas.

Lemma 2.1. *The forcing R is σ -closed.*

Proof. If $r_n : n \in \omega$ is a descending chain of conditions in R then its coordinatewise union is still a condition in R and is the lower bound. \square

Lemma 2.2. *The forcing R has \aleph_2 -c.c.*

Proof. Suppose that $r_\alpha : \alpha \in \omega_2$ is a collection of conditions in R . I must produce $\alpha \neq \beta$ such that the conditions r_α and r_β are compatible in R . Choose a large enough cardinal θ and countable elementary submodels M_α of H_θ containing the collection of conditions as well as the ordinal α . By standard Δ -system and counting arguments, using the continuum hypothesis assumption, I will be able to find ordinals $\alpha \in \beta$ such that the corresponding models are isomorphic via a function $\pi : M_\alpha \rightarrow M_\beta$ which is the identity on their intersection (the *root*) and satisfies $\pi(\alpha) = \beta$. I can also require that all ordinals in ω_2 and the root are smaller than all ordinals in $\omega_2 \cap M_\alpha$ and not the root, which are in turn smaller than all the ordinals in $\omega_2 \cap M_\beta$ and not the root. I will prove that the conditions r_α and r_β have a lower bound. Write $r_\alpha = \langle P_\alpha, Q_\alpha, \leq_\alpha, C_\alpha, F_\alpha \rangle$ and similarly for β and note that $\pi(r_\alpha) = r_\beta$.

The common lower bound r is defined as the coordinatewise union on the first three coordinates of r_α and r_β . The function F_r must extend $F_\alpha \cup F_\beta$. It is necessary to define $F_r(p, c)$ where $p \in P_\alpha \setminus P_\beta$ and $c \in C_\beta \setminus C_\alpha$, or vice versa, where $p \in P_\beta \setminus P_\alpha$ and $c \in C_\alpha \setminus C_\beta$. The latter case is just a mirror image of the former case. In the former case, note that c must contain some condition $q \notin M_\alpha$ (otherwise $c = \pi^{-1}c \in C_\alpha$) and let $F_r(p, c)$ be one such condition in the chain c .

It is not difficult to verify that indeed $r \in R$. Consider for example the condition (4) in the case where $p \in P_\alpha$ and the root and $c \in C_\beta$ and not the root. Then, $F_r(p, c)$ is a condition in Q_β and not the root. All conditions $\leq_r p$ are in $P_\alpha \cup Q_\alpha$ and not the root, all conditions $\leq_r F_r(p, c)$ are in $P_\beta \cup Q_\beta$ and not the root, these two sets are disjoint, therefore $p, F_r(p, c)$ are \leq_r -incompatible and (4) holds.

In order to verify that $r \leq r_\alpha, r_\beta$, I need to show that the incompatibility relation on \leq_r extends that of \leq_α and \leq_β . For this, note that if a condition $p \in P_\alpha \cup Q_\alpha$ does not belong to the root, it has no elements of $P_\beta \cup Q_\beta$ below it. \square

Lemma 2.3. $R \Vdash \dot{P}$ is σ -closed.

Proof. Suppose that $r \in R$ forces $\dot{a} = \langle \dot{p}_n : n \in \omega \rangle$ is a descending chain of elements of \dot{P} . I must find a stronger condition forcing a lower bound to this chain. A reference to genericity will then complete the argument.

Use the σ -closedness of R to strengthen r if necessary to decide the names \dot{p}_n to be certain specific elements $p_n \in P_r$. Define a condition $r' \leq r$ by extending the poset $P_r \cup Q_r$ by adding an element $p \in P_{r'}$ such that for every $q \in P_r \cup Q_r$, $p \leq_{r'} q$ if and only if there is $n \in \omega$ with $p_n \leq_r q$; and adding a countable chain b below p which contains alternately elements of $P_{r'}$ and $Q_{r'}$. Define $C_{r'} = C_r$ and $F_{r'}$ to be a certain extension of F_r . I must define the values $F_{r'}(p, c)$ for every chain $c \in C_r$. The values $F_{r'}(q, c)$ for $q \in b$ will be defined in the same way.

For the definition, write $d = \{q \in c : \exists n \in \omega p_n \leq q\}$ and note that $d \neq c$. Either $F_r(p_0, c) \notin d$, or else, if $F_r(p_0, c) \in d$ as witnessed by p_n , then p_n is incompatible with some element of c by the properties of the function F_r , and

this element then must fall out of d . In any case, let $F_{r'}(p, c)$ be any element of $c \setminus d$. It is immediate that $F_{r'}(p, c)$ is incompatible with p in $\leq_{r'}$ and therefore the condition (4) is satisfied in this case.

It is now not difficult to check that $r' \in R$, $r' \leq r$ and $r' \Vdash \check{p}$ is a lower bound of \dot{a} as desired. \square

Lemma 2.4. $R \Vdash \dot{Q}$ does not have a dense σ -closed subset.

Proof. Suppose that $r \Vdash \dot{D} \subset \dot{Q}$ is dense. I will find a condition $r' \leq r$ such that there is a chain $d \in C_{r'}$ such that for every $q \in d$, $r' \Vdash \check{q} \in \dot{D}$. Such condition of course forces that \check{d} is a descending chain in \dot{D} with no lower bound.

For every ordinal $\alpha \in \omega_2$ find a condition $r_\alpha \leq r$ such that there is a condition $q_\alpha \in Q_{r_\alpha}$ which is as an ordinal larger than α and $r_\alpha \Vdash \check{q}_\alpha \in \dot{D}$. This is possible since the second requirement in (2), $R \Vdash \dot{Q} \cap \alpha$ is not dense in \dot{Q} . Thinning out if necessary, I may assume that $q_\alpha : \alpha \in \omega_2$ in fact form an increasing sequence as ordinals. Now, let θ be a large enough cardinal number and for every ordinal $\alpha \in \omega_2$ choose countable elementary submodels $M_\alpha \prec H_\theta$ containing \dot{D} as well as r_α, q_α . By a standard Δ -system and counting arguments using the continuum hypothesis assumptions, find ordinals $\alpha_n : n \in \omega$ such that the models $M_{\alpha_n} : n \in \omega$ form a Δ -system, they are pairwise isomorphic via functions $\pi_{mn} : M_{\alpha_m} \rightarrow M_{\alpha_n}$ which form a commuting system and are equal to the identity on the root of the Δ -system, $\pi_{mn}(r_{\alpha_m}) = r_{\alpha_n}$, $\pi_{mn}(q_{\alpha_m}) = q_{\alpha_n}$, and moreover, whenever $m \in n$ then all ordinals in $\omega_2 \cap M_{\alpha_m} \setminus M_{\alpha_n}$ are smaller than all ordinals in $\omega_2 \cap M_{\alpha_n} \setminus M_{\alpha_m}$, but greater than all ordinals in the root and ω_2 . I will produce a lower bound r' of the conditions $r_{\alpha_n} : n \in \omega$ such that $d = \{q_{\alpha_n} : n \in \omega\} \in C_{r'}$. This will complete the proof.

In fact, there is a canonical such condition r' . In order to facilitate the notation during the construction, write $r_{\alpha_n} = \langle P_n, Q_n, \leq_n, C_n, F_n \rangle$ and $q_{\alpha_n} = q_n \in Q_n$ for every number $n \in \omega$. We are going to have $P_{r'} = \bigcup_n P_n$, $Q_{r'} = \bigcup_n Q_n$. The ordering $\leq_{r'}$ is the inclusion-minimal one which extends all $\leq_n : n \in \omega$ and contains d as a chain. Since I want to make sure to get a condition \leq_{r_n} for all n , I must verify that the incompatibility relation of $\leq_{r'}$ extends the incompatibility relations of all $\leq_n : n \in \omega$. Well, suppose that $n \neq m \in \omega$ and $p, p' \in P_n \cup Q_n$ are conditions and $q \in P_m \cup Q_m$ is their lower bound in $\leq_{r'}$; I must find their lower bound in \leq_n . There are two cases. Either q belongs to the root, in which case it is enough to observe that $\leq_n = \leq_{r'}$ on the root and therefore q is the required lower bound in P_n as well. Or q does not belong to the root. In such a case, the minimality condition on $\leq_{r'}$ implies that either $q \leq_m p$, or $n \in m$ and $q_n \leq_n p$ and $q \leq_m q_m$ (and the same condition on p'). In any case, this means that $\pi_{mn}(q)$ is the required lower bound of p, p' in \leq_n . A similar break into cases also proves the following implications for every $p \in P_n \cup Q_n$ and $q \in P_m \cup Q_m$: if $p \geq_{r'} q$ then $p \geq_n \pi_{mn}(q)$, and if p and q are compatible in $\leq_{r'}$ then p and $\pi_{mn}(q)$ are compatible in \leq_n .

Let $C_{r'} = \{d\} \cup \bigcup_n C_n$. Note that d has no lower bound in $P_{r'} \cup Q_{r'}$ since it is cofinal in this set with the ordinal ordering. Finally, the function $F_{r'}$ will extend $\bigcup_n F_n$. Note that $\bigcup_n F_n$ is indeed a function: if $p \in P_n$ and $c \in C_n$ for some

$n \in \omega$ then either c is not in the root and then $\langle p, c \rangle$ is not in the domain of the functions $F_m : m \neq n$, and if p, c both belong to the root then so does $F_n(p, c)$ and for every $m \in \omega$, $F_n(p, c) = \pi_{nm}(F_n(p, c)) = (\pi_{nm}F_n)(\pi_{nm}p, \pi_{nm}c) = F_m(p, c)$. To verify that (4) holds, suppose that $p \in P_n, c \in C_n$, and $q \leq_{r'} p, F(p, c)$. I must show that q is not compatible with all elements of the chain c . Indeed, if $q \in P_m \cup Q_m$ were compatible with all elements of the chain c (which are all in $P_n \cup Q_n$), by the last sentence of the previous paragraph $\pi_{mn}q$ would be \leq_n compatible with all elements of c , contradicting the property (4) of the function F_n .

I must define the values $F_{r'}(p, c)$ where $p \in P_n$ and not in the root, and $c \in C_m$ not in the root, for some $n \neq m \in \omega$. Here, observe that all but finitely many elements of c fall out of the root of the Δ -system: the π embeddings move countable sequences pointwise and if they fixed all elements of c , they would all fix c and put c in the root. Then note that all but finitely many elements of c are not above q_m in \leq_m because q_m is not a lower bound of c in that ordering. The definition of $F_{r'}(p, c)$ divides into two possibilities, $m \in n$ and $n \in m$. If $m \in n$, let $F_{r'}(p, c) = q$ be an element of c which is not above q_m and not in the root. The minimality of the ordering $\leq_{r'}$ then implies that p and q are incompatible and therefore (4) is satisfied. If $n \in m$ then let $F_{r'}(p, c) = q$ be an element of c which is not in the root, not above q_m , and below $F_m(q_m, c)$. The verification of (4) is more complicated here. If $p \not\leq_n q_n$ then p is incompatible with q and therefore (4) holds. If $p \geq_n q_n$ then indeed there may be a lower bound p' of p and q . By the minimality of $\leq_{r'}$ it must be the case that $p' \in P_m \cup Q_m$ but not in the root, and $p' \leq q_m$. Then p' is incompatible with one of the elements of c by (4) applied to $F_m(q_m, c)$ and the minimality of $\leq_{r'}$.

Finally, I have to define the values of $F_{r'}(p, d)$ for $p \in P_{r'}$. Just let $F_{r'}(p, d) = q_0$. To see that (4) holds, let $p' \leq p$ be an arbitrary element of $P_{r'}$ below q_0 . p' does not belong to the root, and must belong to P_n for some $n \in \omega$. The minimality of $\leq_{r'}$ implies that $p' \leq_{r'} q_n$. However, $p' \neq q_n$ since $q_n \notin P_n$, and the minimality of $\leq_{r'}$ implies that p' is incompatible with q_{n+1} . (4) follows. \square

Together, the lemma shows that $V[G]$ has the same cardinals and reals as V , and P, Q are codense partial orders, one of them σ -closed and the other without a σ -closed dense subset, proving the theorem.

References

- [1] Matthew Foreman. Games played on Boolean algebras. *Journal of Symbolic Logic*, 48:714–723, 1983.
- [2] Thomas Jech. *Set Theory*. Academic Press, San Diego, 1978.
- [3] Thomas Jech and Saharon Shelah. On countably closed complete boolean algebras. *Journal of Symbolic Logic*, 61:1380–1386, 1996. math.LO/9502203.

- [4] Peter Vojtáš. Game properties of Boolean algebras. *Comment. Math. Univ. Carolinae*, 24:349–369, 1983.