

Decorated Linear Order Types and the Theory of Concatenation

Vedran Čačić

*Department of Mathematics
University of Zagreb
Bijenička 30, 10000 Zagreb
Croatia
veky@math.hr*

Pavel Pudlák

*Mathematical Institute
Academy of Sciences of the Czech Republic
Žitná 25, 115 67 Praha 1
Czech Republic
pudlak@math.cas.cz*

Greg Restall

*Department of Philosophy
The University of Melbourne
Parkville 3010
Australia
restall@unimelb.edu.au*

Alasdair Urquhart

*Department of Philosophy
University of Toronto
215 Huron Street, Toronto
Canada
urquhart@cs.toronto.edu*

Albert Visser

*Department of Philosophy
Utrecht University
Heidelberglaan 8, 3584 CS Utrecht
The Netherlands
albert.visser@phil.uu.nl*

September 14, 2007

Abstract

We study the interpretation of Grzegorzczuk's Theory of Concatenation TC in structures of decorated linear order types satisfying Grzegorzczuk's axioms. We show that TC is incomplete for this interpretation. What is more, the first order theory validated by this interpretation interprets arithmetical truth. We also show that every extension of TC has a model that is not isomorphic to a structure of decorated order types.

We provide a positive result, to wit a construction that builds structures of decorated order types from models of a suitable concatenation theory. This construction has the property that if there is a representation of a certain kind, then the construction provides a representation of that kind.

1 Introduction

In his paper [Grz05], Andrzej Grzegorzczuk introduces a theory of concatenation TC. The theory has a binary function symbol $*$ for concatenation and two constants \mathbf{a} and \mathbf{b} . The theory is axiomatized as follows.

$$\text{TC1. } \vdash (x * y) * z = x * (y * z)$$

$$\text{TC2. } \vdash x * y = u * v \rightarrow ((x = u \wedge y = v) \vee \exists w ((x * w = u \wedge y = w * v) \vee (x = u * w \wedge y * w = v)))$$

$$\text{TC3. } \vdash x * y \neq \mathbf{a}$$

$$\text{TC4. } \vdash x * y \neq \mathbf{b}$$

$$\text{TC5. } \vdash \mathbf{a} \neq \mathbf{b}$$

Axioms TC1 and TC2 are due to Tarski. Grzegorzczuk calls axiom TC2: the editor axiom. We will consider two weaker theories. The theory TC_0 has the signature with just concatenation, and is axiomatized by TC1,2. The theory TC_1 is axiomatized by TC1,2,3. We will also use TC_2 for TC.

The theories we are considering have various interesting interpretations. first they are, of course, theories of strings with concatenation. I.o.w., they are theories of free semigroups. Secondly they are theories of wider classes of structures, to wit structures of *decorated linear order types*, which will be defined below.

The theories TC_i are theories for concatenation without the empty string, i.e., without the unit element. Adding a unit ε one obtains another class of theories TC_i^ε , theories of free monoids, or theories of structures of decorated linear order types including the empty linear decorated order type. The basic list of axioms is as follows.

$$\text{TC}^\varepsilon 1. \vdash \varepsilon * x = x \wedge x * \varepsilon = x$$

$$\text{TC}^\varepsilon 2. \vdash (x * y) * z = x * (y * z)$$

$$\text{TC}^\varepsilon 3. \vdash x * y = u * v \rightarrow \exists w ((x * w = u \wedge y = w * v) \vee (x = u * w \wedge y * w = v))$$

$$\text{TC}^\varepsilon 4. \vdash \mathbf{a} \neq \varepsilon$$

$$\text{TC}^\varepsilon 5. \vdash x * y = \mathbf{a} \rightarrow (x = \varepsilon \vee y = \varepsilon)$$

$$\text{TC}^\varepsilon 6. \vdash \mathbf{b} \neq \varepsilon$$

$$\text{TC}^\varepsilon 7. \vdash x * y = \mathbf{b} \rightarrow (x = \varepsilon \vee y = \varepsilon)$$

$$\text{TC}^\varepsilon 8. \vdash \mathbf{a} \neq \mathbf{b}$$

We take TC_0^ε to be the theory axiomatized by $\text{TC}^\varepsilon 1, 2, 3$. We take TC_1^ε to be $\text{TC}_0^\varepsilon + \text{TC}^\varepsilon 4, 5$ and $\text{TC}^\varepsilon := \text{TC}_2^\varepsilon$ to be $\text{TC}_1^\varepsilon + \text{TC}^\varepsilon 6, 7, 8$.

One can show that TC is *bi-interpretable* with TC^ε , in which a unit ε is added via one dimensional interpretations without parameters.¹ The theory TC_1 is bi-interpretable with TC_1^ε via two-dimensional interpretations with parameters. The situation for TC_0 seems to be more subtle. See also [Vis07]. In Section 6, we will study an extension of TC_0^ε .

Andrzej Grzegorzcyk and Konrad Zdanowski have shown that TC is essentially undecidable. This result can be strengthened by showing that Robinson’s Arithmetic Q is mutually interpretable with TC. Note that TC_0 is undecidable —since it has an extension that parametrically interprets TC— but that TC_0 is not essentially undecidable: it is satisfied by a one-point model. Similarly TC_1 is undecidable, but it has as an extension the theory of finite strings of \mathbf{a} ’s, which is a notational variant of Presburger Arithmetic and, hence decidable.

We will call models of TC_0 *concatenation structures*, and we will call models of TC_i *concatenation i -structures*. The relation of isomorphism between concatenation structures will be denoted by \cong . We will be interested in concatenation structures, whose elements are decorated linear order types with the operation *concatenation of decorated order types*. Let a non-empty class A be given. An A -decorated linear ordering is a structure $\langle D, \leq, f \rangle$, where D is a non-empty domain, \leq is a linear ordering on D , and f is a function from D to A . A mapping ϕ is an *isomorphism* between A -decorated linear order types $\langle D, \leq, f \rangle$ and $\langle D', \leq', f' \rangle$ iff it is a bijection between D and D' such that, for all d, e in D , we have $d \leq e \Leftrightarrow \phi d \leq' \phi e$, and $fd = f'\phi d$. Our notion of isomorphism gives us a notion of A -decorated linear order type. We have an obvious notion of concatenation between A -decorated linear orderings which induces a corresponding notion of concatenation for A -decorated linear order types. We use α, β, \dots to range over such linear order types. Since, linear order types are classes we have to follow one of two strategies: either to employ Scott’s trick to associate a set object to any decorated linear order type or to simply refrain from dividing out isomorphism but to think about decorated linear orderings modulo isomorphism. We will employ the second strategy.

We will call a concatenation structure whose domain consists of (representatives of) A -decorated order types, for some A , and whose concatenation *is* concatenation of decorated order types: a *concrete concatenation structure*. It seems entirely reasonable to stipulate that e.g. the interpretation of \mathbf{a} in a concrete concatenation structure is a decorated linear order type of a one element order. However, for the sake of generality we will refrain from making this stipulation.

Grzegorzcyk conjectured that every concatenation 2-structure is isomorphic to a concrete concatenation structure. We prove that this conjecture is false. (i) Every extension of TC_1 has a model that is not isomorphic to a concatenation 1-structure and (ii) The set of principles of valid in all concrete concatenation 2-structures interprets arithmetical truth.

¹Albert Visser thinks he can improve this to: TC and TC^ε are definitionally equivalent.

The plan of the paper is as follows. We show, in Section 2, that we have, for all decorated order types α , β and γ , the following principle:

$$(\dagger) \quad \beta * \alpha * \gamma = \alpha \Rightarrow \beta * \alpha = \alpha * \gamma = \alpha.$$

This fact was already known. It is due to Lindenbaum, credited to him in Sierpiński's book [Sie58] on p. 248. It is also problem 6.13 of [KT06]. Our proof, however, is different.

It is easy to see that every group is a concatenation structure and that (\dagger) does not hold in the two element group. We show, in Section 5, that every concatenation structure can be extended to a concatenation structure with any number of atoms. It follows that there is a concatenation structure with at least two atoms in which (\dagger) fails. Hence, TC is incomplete for concrete concatenation structures. In Section 3, we provide a counterargument of a different flavour. We provide a tally interpretation that defines the natural numbers (with concatenation in the role of addition) in every concrete concatenation 2-structure. It follows that for every extension of TC₁ is satisfied by a concatenation 1-structure that is not isomorphic to any concrete concatenation 1-structure, to wit any model of that extension that contains a non-standard element. In Section 4, we strengthen the result of Section 3, by showing that in concrete concatenation 2-structures we can add multiplication to the natural numbers. It follows that the set of arithmetically true sentences is interpretable in the concretely valid consequences of TC₂.

Finally, in Section 6 we prove a positive result. We provide a mapping from arbitrary models of a variant of an extension of TC₀ to structures of decorated order types. As we have shown such a construction cannot always provide a representation. We show that, for a restricted class of representations, we have: if a model has a representation in the class, then the construction yields such a representation.

Acknowledgments

Some of the results of this note were obtained during the Excursion to mountain Ślęza of the inspiring Logic Colloquium 2007 in Wrocław and, in part, in the evening after the Excursion. We thank the organizers for providing this wonderful opportunity.

We thank Dana Scott for his comments, insights and questions.

Pavel Pudlák was supported by grants A1019401 and 1M002162080.

2 A Principle for Decorated Order Types

In this section we prove a universal principle that holds in all concatenation structures, which is not provable in TC. There is an earlier proof of this principle. See: [KT06], problem 6.13. Our proof, however, is different.

Theorem 2.1 *Let $\alpha_0, \alpha_1, \alpha_2$ be decorated order types. Suppose $\alpha_1 = \alpha_0 * \alpha_1 * \alpha_2$. Then $\alpha_1 = \alpha_0 * \alpha_1 = \alpha_1 * \alpha_2$.*

Proof

Suppose $\alpha_1 = \alpha_0 * \alpha_1 * \alpha_2$. Consider a decorated linear ordering $\mathcal{A} := \langle A, \leq, f \rangle$ of type α_1 . By our assumption, we may partition A into A_0, A_1, A_2 , such that:

$$\langle A, \leq, f \rangle = \langle A_0, \leq \upharpoonright A_0, f \upharpoonright A_0 \rangle * \langle A_1, \leq \upharpoonright A_1, f \upharpoonright A_1 \rangle * \langle \leq \upharpoonright A_2, f \upharpoonright A_2 \rangle,$$

where $\mathcal{A}_i := \langle A_i, \leq \upharpoonright A_i, f \upharpoonright A_i \rangle$ is an instance of α_i . Let $\phi : \mathcal{A} \rightarrow \mathcal{A}_1$ be an isomorphism.

Let $\phi^n \mathcal{A}_{(i)} := \langle \phi^n[A_{(i)}], \leq \upharpoonright \phi^n[A_{(i)}], f \upharpoonright \phi^n[A_{(i)}] \rangle$. We have: $\phi^n \mathcal{A}_i$ is of order type α_i and $\phi^n \mathcal{A}$ is of order type α_1 .

Clearly, $\phi \mathcal{A}_0$ is an initial substructure of $\phi \mathcal{A} = \mathcal{A}_1$. So, \mathcal{A}_0 and $\phi \mathcal{A}_0$ are disjoint and $\phi \mathcal{A}_0$ adjacent to the right of \mathcal{A}_0 . Similarly, for $\phi^n \mathcal{A}_0$ and $\phi^{n+1} \mathcal{A}_0$. Take $A_0^\omega := \bigcup_{i \in \omega} \phi^i A_0$. We find that $\mathcal{A}_0^\omega := \langle A_0^\omega, \leq \upharpoonright A_0^\omega, f \upharpoonright A_0^\omega \rangle$ is initial in \mathcal{A} and of decorated linear order type α_0^ω . So $\alpha_1 = \alpha_0^\omega * \rho$, for some ρ . It follows that $\alpha_0 * \alpha_1 = \alpha_0 * \alpha_0^\omega * \rho = \alpha_0^\omega * \rho = \alpha_1$. The other identity is similar. \square

So, every concrete concatenation structure validates that $\alpha_1 = \alpha_0 * \alpha_1 * \alpha_2$ implies $\alpha_1 = \alpha_0 * \alpha_1 = \alpha_1 * \alpha_2$. We postpone the proof that this principle is not provable in TC to Section 5.

3 Definability of the Natural Numbers

In this section, we show that the natural numbers can be defined in every concrete concatenation 1-structure. We define:

- $x \subseteq y : \leftrightarrow x = y \vee \exists u (u * x = y) \vee \exists v (x * v = y) \vee \exists u, v (u * x * v = y)$.
- $x \subseteq_{\text{ini}} y : \leftrightarrow x = y \vee \exists v (x * v = y)$.
- $x \subseteq_{\text{end}} y : \leftrightarrow x = y \vee \exists u (u * x = y)$.
- $(n : \tilde{\mathbb{N}}_{\mathbf{a}}) : \leftrightarrow \forall m \subseteq_{\text{ini}} n (m = \mathbf{a} \vee \exists k \subseteq_{\text{ini}} m (k \neq m \wedge m = k * \mathbf{a}))$.

The use of ‘:’ in $n : \tilde{\mathbb{N}}_{\mathbf{a}}$ is derived from the analogous use in type theory. We could read it as: n is of sort $\mathbb{N}_{\mathbf{a}}$. We write $m, n : \tilde{\mathbb{N}}_{\mathbf{a}}$ for: $(m : \tilde{\mathbb{N}}_{\mathbf{a}}) \wedge (n : \tilde{\mathbb{N}}_{\mathbf{a}})$. Etc.

We prove the main theorem of this section.

Theorem 3.1 *In any concrete concatenation structure, we have:*

$$\tilde{\mathbb{N}}_{\mathbf{a}} = \{\mathbf{a}^{n+1} \mid n \in \omega\}.$$

I.o.w, $\tilde{\mathbb{N}}_{\mathbf{a}}$ is precisely the class of natural numbers in tally representation (starting with 1). Note that $$ on this set is addition.*

Proof

Consider any concrete concatenation 1-structure \mathfrak{A} . It is easy to see that every \mathfrak{a}^{n+1} is in $\tilde{\mathbb{N}}_{\mathfrak{a}}$.

Clearly, every element of $\tilde{\mathbb{N}}_{\mathfrak{a}}$ is either \mathfrak{a} or it has a predecessor. The axioms of TC_1 guarantee that this predecessor is unique. This justifies the introduction of the partial predecessor function pd on $\tilde{\mathbb{N}}_{\mathfrak{a}}$. Let α be the order type corresponding to \mathfrak{a} . Let β_0 be any element of $\tilde{\mathbb{N}}_{\mathfrak{a}}$. If, for some n , $\text{pd}^n \beta_0$ is undefined, then β_0 is clearly of the form α^{k+1} , for k in ω .

We show that the other possibility cannot obtain. Suppose $\beta_n := \text{pd}^n \beta_0$ is always defined. Let \mathcal{A} be a decorated linear ordering of type α and let \mathcal{B}_i be a decorated linear ordering of type β_i . We assume that the domain A of \mathcal{A} is disjoint from the domains B_i of the \mathcal{B}_i . Thus, we may implement $\mathcal{B}_{i+1} * \mathcal{A}$ just by taking the union of the domains.

Let ϕ_i be isomorphisms from $\mathcal{B}_{i+1} * \mathcal{A}$ to \mathcal{B}_i . Let $\mathcal{A}_i := (\phi_0 \circ \dots \circ \phi_i)(\mathcal{A})$. Then, the \mathcal{A}_i are all of type α and, for some \mathcal{C} , we have $\mathcal{B}_0 \cong \mathcal{C} * \dots * \mathcal{A}_1 * \mathcal{A}_0$. Similarly $\mathcal{B}_1 \cong \mathcal{C} * \dots * \mathcal{A}_2 * \mathcal{A}_1$. Let $\tilde{\omega}$ be the opposite ordering of ω . It follows that $\beta_0 = \gamma * \alpha^{\tilde{\omega}} = \beta_1 = \text{pd}(\beta_0)$. Hence, β_0 is not in $\tilde{\mathbb{N}}_{\mathfrak{a}}$.² A contradiction. \square

We call a concatenation structure *standard* if $\tilde{\mathbb{N}}_{\mathfrak{a}}$ defines the tally natural numbers. Since, by the usual argument, any any extension of TC_1 has a model with non-standard numbers, we have the following corollary.

Corollary 3.2 *Every extension of TC_1 has a model that is not isomorphic to a concrete concatenation 1-structure. In a different formulation: for every concatenation 1-structure there is an elementarily equivalent concatenation 1-structure that is not isomorphic to a concrete concatenation 1-structure.*

Note that the non-negative tally numbers with addition form a concrete concatenation 1-structure. Thus, the concretely valid consequences of $\text{TC}_1 + \forall x (x : \tilde{\mathbb{N}}_{\mathfrak{a}})$, i.e., the principles valid in every concrete concatenation 1-structure satisfying $\forall x (x : \mathbb{N}_{\mathfrak{a}})$ are decidable.

4 Definability of Multiplication

If we have two atoms to work with, we can add multiplication to our tally numbers. This makes the set of concretely valid consequences of TC non-arithmetical. The main ingredient of the definition of multiplication is the theory of relations on tally numbers. In TC , we can develop such a theory. We represent the relation $\{\langle x_0, y_0 \rangle, \dots, \langle x_{n-1}, y_{n-1} \rangle\}$, by:

$$\mathbf{bb} * x_0 * \mathbf{b} * y_0 * \mathbf{bb} * x_1 * \dots * \mathbf{bb} * x_{n-1} * \mathbf{b} * y_{n-1} * \mathbf{bb}.$$

We define:

$$\bullet r : \text{REL} \leftrightarrow \mathbf{bb} \subseteq_{\text{end}} r,$$

²Note that we are not assuming that γ is in \mathfrak{A} .

- $\emptyset := \mathbf{bb}$,
- $x[r]y := \leftrightarrow x, y : \tilde{\mathbf{N}}_a \wedge \mathbf{bb} * x * \mathbf{b} * y * \mathbf{bb} \subseteq r$.
- $\mathbf{adj}(r, x, y) := r * x * \mathbf{b} * y * \mathbf{bb}$.

Clearly, we have: $\mathbf{TC} \vdash \forall u, v \neg u[\emptyset]v$. To verify that this coding works we need the adjunction principle.

Theorem 4.1 *We have:*

$$\mathbf{TC} \vdash (r : \mathbf{REL} \wedge x, y, u, v : \tilde{\mathbf{N}}_a) \rightarrow (u[\mathbf{adj}(r, x, y)]v \leftrightarrow (u[r]v \vee (u = x \wedge v = y))).$$

We can prove this result by laborious and unspicuous case splitting. However, it is more to do the job with the help of a lemma. Consider any model of \mathbf{TC}_0 . Fix an element w . We call a sequence (w_0, \dots, w_k) a *partition* of w if we have that $w_0 * \dots * w_k = w$. The partitions of w form a category with the following morphisms. $f : (u_0, \dots, u_n) \rightarrow (w_0, \dots, w_k)$ iff f is a surjective and weakly monotonic function from $n + 1$ to $k + 1$, such that, for any $i \leq k$, $w_i = u_s * \dots * u_\ell$, where $f(j) = i$ iff $s \leq j \leq \ell$. We write $(u_0, \dots, u_n) \leq (w_0, \dots, w_k)$ for: $\exists f : (u_0, \dots, u_n) \rightarrow (w_0, \dots, w_k)$. In this case we say that (u_0, \dots, u_n) is a *refinement* of (w_0, \dots, w_k) .

Lemma 4.1 *Consider any concatenation structure. Consider a w in the structure. Then, any two partitions of w have a common refinement.*

Proof

Fix any concatenation structure. We first prove that, for all w , all pairs of partitions (u_0, \dots, u_n) and (w_0, \dots, w_k) of w have a common refinement, by induction of $n + k$.

If either n or k is 0, this is trivial. Suppose (u_0, \dots, u_{n+1}) and (w_0, \dots, w_{k+1}) are partitions of w . By the editor axiom, either (a) $u_0 * \dots * u_n = w_0 * \dots * w_k$ and $u_{n+1} = w_{k+1}$, or there is a v such that (b) $u_0 * \dots * u_n * v = w_0 * \dots * w_k$ and $u_{n+1} = v * w_{k+1}$, or (c) $u_0 * \dots * u_n = w_0 * \dots * w_k * v$ and $v * u_{n+1} = w_{k+1}$. We only treat case (b), the other cases being easier or similar. By the induction hypothesis, there is a common refinement (x_0, \dots, x_m) of (u_0, \dots, u_n, v) and (w_0, \dots, w_n) . Let this be witnessed by f , resp. g . It is easily seen that $(x_0, \dots, x_m, w_{k+1})$ is the desired refinement with witnessing functions f' and g' , where $f' := f[m + 1 \mapsto n + 1]$, $g' := g[m + 1 \mapsto k + 1]$. Here $f[m + 1 \mapsto n + 1]$ in the result of extending f to assign $n + 1$ to $m + 1$. \square

We turn to the proof of Theorem 4.1. The verification proceeds more or less as one would do it for finite strings.

Proof

Consider any concatenation 2-structure. Suppose $\text{REL}(r)$. The right-to-left direction is easy, so we treat left-to-right. Suppose x, y, u and v are tally numbers. and $u[\text{adj}(r, x, y)]v$. There are two possibilities. Either $r = \mathbf{bb}$ or $r = r_0 * \mathbf{bb}$. We will treat the second case. Let $s := \text{adj}(r, x, y)$. One the following four partitions is a partition of s : (i) $(\mathbf{b}, \mathbf{b}, u, \mathbf{b}, v, \mathbf{b}, \mathbf{b})$, or (ii) $(w, \mathbf{b}, \mathbf{b}, u, \mathbf{b}, v, \mathbf{b}, \mathbf{b})$, or (iii) $(\mathbf{b}, \mathbf{b}, u, \mathbf{b}, v, \mathbf{b}, \mathbf{b}, z)$, or (iv) $(w, \mathbf{b}, \mathbf{b}, u, \mathbf{b}, v, \mathbf{b}, \mathbf{b}, z)$. We will treat cases (ii) and (iv).

Suppose $\sigma := (w, \mathbf{b}, \mathbf{b}, u, \mathbf{b}, v, \mathbf{b}, \mathbf{b})$ is a partition of s . We also have that $\tau := (r_0, \mathbf{b}, \mathbf{b}, x, \mathbf{b}, y, \mathbf{b}, \mathbf{b})$ is a partition of s . Let (t_0, \dots, t_k) be a common refinement of σ and τ , with witnessing functions f and g . The displayed \mathbf{b} 's in these partitions must have unique places among the t_i . We define m_σ to be the unique i such that $f(i) = m$, provided that $\sigma_m = \mathbf{b}$. Similarly, for m_τ . (To make this unambiguous, we assume that if $\sigma = \tau$, we take σ as the common refinement with f and g both the identity function.)

We evidently have $7_\sigma = 7_\tau = k$ and $6_\sigma = 6_\tau = k - 1$. Suppose $4_\sigma < 4_\tau$. It follows that $\mathbf{b} \subseteq v$. So, v would have an initial subsequence that ends in \mathbf{b} , which is impossible. So, $4_\sigma \not\leq 4_\tau$. Similarly, $4_\tau \not\leq 4_\sigma$. So $4_\sigma = 4_\tau$. It follows that $v = y$. Reasoning as in the case of 4_σ and 4_τ , we can show that $2_\sigma = 2_\tau$ and, hence $u = x$.

Suppose $\rho := (w, \mathbf{b}, \mathbf{b}, u, \mathbf{b}, v, \mathbf{b}, \mathbf{b}, z)$ is a partition of s . We also have that $\tau := (r_0, \mathbf{b}, \mathbf{b}, x, \mathbf{b}, y, \mathbf{b}, \mathbf{b})$ is a partition of s . Let (t_0, \dots, t_k) be a common refinement of ρ and τ , with witnessing functions f and g . We consider all cases, where $1_\tau < 6_\rho$. Suppose $6_\rho = 1_\tau + 1 = 2_\tau$. Note that $7_\rho = 6_\rho + 1$, so we find: $\mathbf{b} \subseteq x$, quod non, since x is in $\tilde{\mathbf{N}}_a$. Suppose $2_\tau < 6_\rho < 4_\tau$. In this case we have a \mathbf{b} as substring of x . Quod non. Suppose $6_\rho = 4_\tau$. Since $7_\rho = 6_\rho + 1$, we get a \mathbf{b} in y . Quod non. Suppose $4_\tau < 6_\rho < 6_\tau$. In this case, we get a \mathbf{b} in y . Quod impossibile. Suppose $6_\rho \geq 6_\tau = k - 1$. In this place there is no place left for z among the t_i . So, in all cases, we obtain a contradiction. So the only possibility is $6_\rho \leq 1_\tau$. Thus, it follows that $u[r]v$. \square

We can now use our relations to define multiplication of tally numbers in the usual way. See e.g. Section 2.2 of [Bur05]. In any concrete concatenation 2-structure, we can use induction to verify the defining properties of multiplication as defined. It follows that we can interpret all arithmetical truths in the set of concretely valid consequences of TC.

Corollary 4.2 *We can interpret true arithmetic in the set of all principles valid in concrete concatenation 2-structures.*

5 The Sum of Concatenation Structures

In this section we show that concatenation structures are closed under sums. This result will make it possible to verify the claim that the universal principle

of Section 2 is not provable in TC. The result has some independent interest, since it provides a good closure property of concatenation structures.

Consider two concatenation structures \mathfrak{A}_0 and \mathfrak{A}_1 . We write \star for concatenation in the \mathfrak{A}_i . We may assume, without loss of generality, that the domains of \mathfrak{A}_0 and \mathfrak{A}_1 are disjoint. We define the sum $\mathfrak{B} := \mathfrak{A}_0 \oplus \mathfrak{A}_1$ as follows.

- The domain of \mathfrak{B} consists of non-empty sequences $w_0 \cdots w_{n-1}$, where the w_j are alternating between elements of the domains of \mathfrak{A}_0 and \mathfrak{A}_1 . In other words, if w_j is in the domain of \mathfrak{A}_i , then w_{j+1} , if it exists, is in the domain of \mathfrak{A}_{1-i} .
- The concatenation $\sigma \star \tau$ of $\sigma := w_0 \cdots w_{n-1}$ and $\tau := v_0 \cdots v_{k-1}$ is $w_0 \cdots w_{n-1} v_0 \cdots v_{k-1}$, in case w_{n-1} and v_0 are in the domains of different structures \mathfrak{A}_i . The concatenation $\sigma \star \tau$ is $w_0 \cdots (w_{n-1} \star v_0) \cdots w_{k-1}$, in case w_{n-1} and v_0 are in the same domain.

In case $\sigma \star \tau$ is obtained via the first case, we say that σ and τ are *glued together*. If the second case obtains, we say that σ and τ are *clicked together*.

Theorem 5.1 *The structure $\mathfrak{B} = \mathfrak{A}_0 \oplus \mathfrak{A}_1$ is a concatenation structure.*

Proof

Associativity is easy. We check the editor property TC2. Suppose $\sigma_0 \star \sigma_1 = z_0 \cdots z_{m-1} = \tau_0 \star \tau_1$. We distinguish a number of cases.

Case 1. Suppose both of the pairs σ_0, σ_1 and τ_0, τ_1 are glued together. Then, for some $k, n > 0$, we have $\sigma_0 = z_0 \cdots z_{k-1}$, $\sigma_1 = z_k \cdots z_{m-1}$, $\tau_0 = z_0 \cdots z_{n-1}$, and $\tau_1 = z_n \cdots z_{m-1}$.

So, if $k = n$, we have $\sigma_0 = \tau_0$ and $\sigma_1 = \tau_1$.

If $k < n$, we have $\tau_0 = \sigma_0 \star (z_k \cdots z_{n-1})$ and $\sigma_1 = (z_k \cdots z_{n-1}) \star \tau_1$. The case that $n < k$ is similar.

Case 2. Suppose σ_0, σ_1 is glued together and that τ_0, τ_1 is clicked together. So, there are $k, n > 0$, u_0 , and u_1 such that $\sigma_0 = z_0 \cdots z_{k-2} u_0$, $\sigma_1 = u_1 z_k \cdots z_{m-1}$, $u_0 \star u_1 = z_{k-1}$, $\tau_0 = z_0 \cdots z_{n-1}$, and $\tau_1 = z_n \cdots z_{m-1}$.

Suppose $k \leq n$. Then, $\tau_0 = \sigma_0 \star (u_1 z_k \cdots z_{n-1})$ and $\sigma_1 = (u_1 z_k \cdots z_{n-1}) \star \tau_1$. Note that, in case $k = n$, the sequence $z_k \cdots z_{n-1}$ is empty. The case that $k \geq n$ is similar.

Case 3. This case, where σ_0, σ_1 is clicked together and τ_0, τ_1 is glued together, is similar to case 2.

Case 4. Suppose that σ_0, σ_1 and τ_0, τ_1 are both clicked together. So, there are $k, n > 0$, u_0, u_1, v_0, v_1 such that $\sigma_0 = z_0 \cdots z_{k-2} u_0$, $\sigma_1 = u_1 z_k \cdots z_{m-1}$, $u_0 \star u_1 = z_{k-1}$, $\tau_0 = z_0 \cdots z_{n-2} v_0$, $\tau_1 = v_1 z_n \cdots z_{m-1}$ and $v_0 \star v_1 = z_{n-1}$.

Suppose $k = n$. We have $u_0 \star u_1 = z_{k-1} = v_0 \star v_1$. So, we have either (a) $u_0 = v_0$ and $u_1 = v_1$, or, for some w , either (b) $u_0 \star w = v_0$ and $u_1 = w \star v_1$,

or (c) $u_0 = v_0 \star w$ and $w \star u_1 = v_1$. In case (b), we have: $\sigma_0 \star w = \tau_0$ and $\sigma_1 = w \star \tau_1$. We leave (a) and (c) to the reader.

Suppose $k < n$. We have:

$$\sigma_0 \star (u_1 z_k \cdots z_{n-2} v_0) = \tau_0 \text{ and } \sigma_1 = (u_1 z_k \cdots z_{n-2} v_0) \star \tau_1.$$

The case that $k > n$ is similar. \square

It is easy to see that \oplus is a sum or coproduct in the sense of category theory. The following theorem is immediate.

Theorem 5.2 *If a is an atom of \mathfrak{A}_i , then a is an atom of $\mathfrak{A}_0 \oplus \mathfrak{A}_1$.*

Finally, we have the following theorem.

Theorem 5.3 *Let A be any set and let $\mathfrak{B} := \langle B, \star \rangle$ be any concatenation structure. We assume that A and B are disjoint. Then, there is an extension of \mathfrak{B} with at least A as atoms.*

Proof

Let A^* be the free semi-group on A . We can take as the desired extension of \mathfrak{B} , the structure $A^* \oplus \mathfrak{B}$. \square

Remark 5.4 The whole development extends with only minor adaptations, when we replace axiom TC2 by:

$$\bullet \vdash x \star y = u \star v \rightarrow ((x = u \wedge y = v) \dot{\vee} (\exists! w (x \star w = u \wedge y = w \star v) \vee \exists! w (x = u \star w \wedge y \star w = v)))$$

Here $\dot{\vee}$ is *exclusive or*. \square

6 A Canonical Construction

Although we know that not every concatenation structure can be represented by decorated linear orderings, i.e., as a concrete concatenation structure, there may exist a canonical construction of a concrete concatenation structure which is a representation whenever there exists any concrete representation. In this section we shall propose such a construction, but we can only show that it is universal in a restricted subclass of all concrete representations.

It will be now more convenient to work with a theory for monoids, rather than for semigroups, as we did in the previous sections. We will work in the theory TC_0^ε plus the following axiom.

$$\text{TC}^\varepsilon 9. \vdash x \star y \star z = y \rightarrow (x = \varepsilon \wedge z = \varepsilon).$$

We do not postulate the existence of irreducible elements, as they do not play any role in what follows, but they surely can be present.

We shall call elements of a model \mathcal{M} of TC_1^ε : words. When possible, the concatenation symbol $*$ will be omitted.

Lemma 6.1 *In a model \mathcal{M} of TC_1^ε the binary relation $\exists u(xu = y)$ defines an ordering on the elements of \mathcal{M} .*

Definition 6.1 Let w be a word.

- A k -partition of w is a k -tuple (w_1, \dots, w_k) such that $w_1 \dots w_k = w$; we shall often abbreviate it by $w_1 \dots w_k$.
- An ordering relation is defined on 3-partitions of w by

$$u_1 u_2 u_3 \leq v_1 v_2 v_3 \equiv \exists x_1, x_3 (v_1 x_1 = u_1 \wedge x_3 v_3 = u_3).$$

□

The axioms ensure that for any two partitions there is a unique common refinement.

Definition 6.2 [Word Ultrafilters] Let w be a word and S a set of 3-partitions of w . We shall call S a word ultrafilter (wuf) on w if

1. $\varepsilon w \varepsilon \in S$
2. $x \varepsilon y \notin S$ for any x, y
3. if $U \in S$, V is a 3-partition of w and $U \leq V$, then $V \in S$
4. if $xyz \in S$ and $y = y_1 y_2$ then exactly one of the following two cases holds: $(x, y_1, y_2 z) \in S$ or $(x y_1, y_2, z) \in S$.

□

Let S be a wuf on w and $xyz \in S$. Then we define the natural restriction of S to y which is a wuf S_y on y defined by:

$$(r, s, t) \in S_y \iff (xr, s, tz) \in S.$$

We shall define an ordering on wuf's on a fixed w and an equivalence on wuf's on all words of M .

- Let S and T be wuf's on w , then we define

$$S < T \iff \exists u, v ((\varepsilon, u, v) \in S \wedge (u, v, \varepsilon) \in T).$$

- Let S and T be wuf's on possibly different words, then we define

$$S \sim T \iff \exists x, x', y, z, z' (xyz \in S \wedge x' y z' \in T \wedge S_y = T_y).$$

Definition 6.3 Let w be a word. The *canonical decorated ordering* associated with w is the ordering of all wuf's on w , where each wuf S is decorated by $[S]_{\sim}$, the equivalence class of \sim containing S . This decorated ordering will be denoted by $C(w)$. \square

Here are some basic properties of $C(w)$.

- The topological space determined by the ordering is compact and totally disconnected. In particular, it has the largest and the smallest elements.
- For every proper prefix x of w , there is a uniquely determined pair of wuf's which forms a gap (no wuf in between). Thus there is a natural embedding of the ordering of the prefixes into $C(w)$. More precisely, we have two mappings ϕ_w^- and ϕ_w^+ such that for a proper prefix x the pair $\phi_w^-(x), \phi_w^+(x)$ is the gap corresponding to x . If $x = \varepsilon$ (or $x = w$) then only $\phi_w^+(x)$ (or $\phi_w^-(x)$) is defined and it is the least (largest) element of $C(w)$. Furthermore the images of these mappings are dense sets in $C(w)$.
- Vice versa, every gap in $C(w)$ corresponds to a prefix (or equivalently to a 2-partition).
- If a is an atom (irreducible element in \mathcal{M}), then it determines a principal wuf. For a given atom a all such principal wuf's are equivalent.

Definition 6.4 ρ is a *regular* representation of \mathcal{M} by decorated orderings, if for every 2-partition $x_1x_2 = w$ of $w \in \mathcal{M}$, there exists a *unique* 2-partition $A_1A_2 = \rho(w)$ such that $A_1 \cong \rho(x_1)$ and $A_2 \cong \rho(x_2)$. \square

We do not know if every concatenation structure that has a concrete representation also has a concrete regular representation.

If ρ is regular, we have an analogous property for k -partitions for every k . For a k -partition (x_1, \dots, x_k) of w in \mathcal{M} , we shall write

$$\rho^k(x_1, \dots, x_k) = (A_1, \dots, A_k),$$

where (A_1, \dots, A_k) is the uniquely determined k -partition of $\rho(x_1 \dots x_k)$ such that $A_i \cong \rho(x_i)$, for $i = 1, \dots, k$.

Theorem 6.5 1. *If the canonical mapping C is a representation of \mathcal{M} , then it is a regular representation of \mathcal{M} .*

2. *If there exists a regular representation ρ of \mathcal{M} , then so is also C .*

Proof

Ad 1. Let $uv = w$ be in \mathcal{M} and suppose we have two different 2-partitions $AB = C(w)$, $A'B' = C(w)$, with $A \cong A' \cong C(u)$, $B \cong B' \cong C(v)$. Suppose that A is a proper initial segment of A' . Since AB corresponds to a 2-partition uv , there is a gap between A and B . Since A is a proper initial segment of A'

the gap is in A' . As every gap corresponds to a 2-partition of the preimage in C , there exists y and D such that $A' = AD$ and $D \cong C(y)$. Hence $u = uy$, which is possible only if $y = \varepsilon$. But then D is empty, which is a contradiction.

Ad 2. Our strategy is

- (i) to construct an order preserving mapping $h : \text{Supp}(\rho(w)) \rightarrow \text{Supp}(C(w))$, for every $w \in M$, and then
- (ii) to show that if $\iota : C(u) \rightarrow C(v)$ is an isomorphism, then for every $S \in \text{Supp}(C(u))$ the fibers of S and $\iota(S)$, as decorated orderings, are isomorphic, i.e.,

$$h^{-1}(S) \cong h^{-1}(\iota(S)),$$

or they are both empty.

Ad (i). Let $w \in M$, let $j \in \text{Supp}(\rho(w))$. We define

$$h(j) = \{(x, y, z) \mid \exists A, B, D (j \in B \text{ and } \rho^3(x, y, z) = (A, B, D))\}.$$

One can readily verify that $h(j)$ is a wuf, and that h is order preserving.

Ad (ii). Let $S \in \text{Supp}(C(u))$ and $T \in \text{Supp}(C(v))$ such that $T = \iota(S)$. Then S and T have the same decoration, which means that $S \sim T$. By definition, there exist 3-partitions $(x, y, z) \in S$ and $(x', y, z') \in T$ such that $S_y = T_y$. Let $\rho^3(x, y, z) = (A, B, D)$ and $\rho^3(x', y, z') = (A', B', D')$. Then $B \cong B'$, as ρ is a representation. Let us denote this isomorphism by κ .

Take an arbitrary 3-partition $y_1 y_2 y_3 = y$ and let

$$\rho^5(x, y_1, y_2, y_3, z) = (A, B_1, B_2, B_3, D),$$

$$\rho^5(x', y_1, y_2, y_3, z') = (A', B'_1, B'_2, B'_3, D').$$

Then $B_i \cong B'_i$, for $i = 1, 2, 3$. By the regularity of ρ , the segments B_1, B_2, B_3 in B and the segments B'_1, B'_2, B'_3 in B' are uniquely determined by their isomorphism types, whence:

$$\kappa(B_i) = B'_i, \text{ for } i = 1, 2, 3. \tag{1}$$

The fiber $\iota^{-1}(S)$ is defined as the intersection of all segments B_2 that belong to 5-partitions (x, y_1, y_2, y_3, z) such that $(y_1, y_2, y_3) \in S_y = T_y$. Similarly, the fiber $\iota^{-1}(T)$ is defined as the intersection of all segments B'_2 that belong to 5-partitions (x', y_1, y_2, y_3, z') such that $(y_1, y_2, y_3) \in S_y = T_y$. According to (1), for all such 3-partitions, $\kappa : (B_1, B_2, B_3) \cong (B'_1, B'_2, B'_3)$. Hence κ is also an isomorphism of $\iota^{-1}(S)$ onto $\iota^{-1}(T)$, or both fibers are empty. \square

References

- [Bur05] John Burgess. *Fixing Frege*. Princeton Monographs in Philosophy. Princeton University Press, Princeton, 2005.
- [Grz05] Andrzej Grzegorzcyk. Undecidability without arithmetization. *Studia Logica*, 79:163–230, 2005.
- [KT06] P. Komjáth and V. Totik. *Problems and Theorems in Classical Set Theory*. Springer-Verlag, New York, 2006.
- [Sie58] W. Sierpiński. *Cardinal and ordinal numbers*, volume 34 of *Monografie Matematyczne*. Polska Akademia Nauk, Warszawa, 1958.
- [Vis07] A. Visser. Growing commas –a study of sequentiality and concatenation. Logic Group Preprint Series 257, Department of Philosophy, Utrecht University, Heidelberglaan 8, 3584 CS Utrecht, <http://www.phil.uu.nl/preprints/lgps/>, 2007.