

Rules with parameters in modal logic

Emil Jeřábek

`jerabek@math.cas.cz`

`http://math.cas.cz/~jerabek/`

Institute of Mathematics of the Academy of Sciences, Prague

Overview

The plan for this talk:

- General remarks on unification and admissibility
- Known results on unification and admissibility in transitive modal logics
- Unification and admissibility with parameters in transitive modal logics
- Unification and admissibility with parameters in intuitionistic logic

Unification and admissibility in propositional logics

Propositional logics

Propositional logic L :

Language: formulas Form_L built freely from **atoms** (variables) $\{x_n : n \in \omega\}$ using a fixed set of **connectives** of finite arity

Consequence relation \vdash_L : finitary structural Tarski-style consequence operator

I.e.: a relation $\Gamma \vdash_L \varphi$ between finite sets of formulas and formulas such that

- $\varphi \vdash_L \varphi$
- $\Gamma \vdash_L \varphi$ implies $\Gamma, \Gamma' \vdash_L \varphi$
- $\Gamma \vdash_L \varphi$ and $\Gamma, \varphi \vdash_L \psi$ imply $\Gamma \vdash_L \psi$
- $\Gamma \vdash_L \varphi$ implies $\sigma(\Gamma) \vdash_L \sigma(\varphi)$ for every substitution σ

Algebraization

L is **finitely algebraizable** wrt a class K of algebras if there is a finite set $F(u, v)$ of formulas and a finite set $E(x)$ of equations such that

- $\Gamma \vdash_L \varphi \Leftrightarrow E(\Gamma) \vDash_K E(\varphi)$
- $\Theta \vDash_K t \approx s \Leftrightarrow F(\Theta) \vdash_L F(t, s)$
- $x \not\vdash_L F(E(x))$
- $u \approx v \not\vDash_K E(F(u, v))$

We may assume K is a quasivariety

In our case we will always have:

$E(x) = \{x \approx 1\}$, $F(u, v) = \{u \leftrightarrow v\}$, K is a variety

Equational unification

Let Θ be an equational theory (or a variety of algebras):

- Θ -unifier of a set Γ of equations:
a substitution σ s.t. $\models_{\Theta} \sigma(t) \approx \sigma(s)$ for all $t \approx s \in \Gamma$
- Γ is Θ -unifiable if it has a Θ -unifier
- $\sigma \equiv_{\Theta} \tau$ iff $\models_{\Theta} \sigma(u) \approx \tau(u)$ for every variable u
- $\sigma \preceq_{\Theta} \tau$ (τ is more general than σ) if $\exists \rho \sigma \equiv_{\Theta} \rho \circ \tau$
- Complete set of unifiers of Γ : a set X of unifiers of Γ such that every unifier of Γ is less general than some $\tau \in X$
- Θ has finitary unification type if every finite Γ has a finite complete set of unifiers

Unification in propositional logics

If L is a logic finitely algebraizable wrt a variety K , we can express K -unification in terms of L :

An L -unifier of a formula φ is σ such that $\vdash_L \sigma(\varphi)$

Then we have:

- L -unifier of $\varphi = K$ -unifier of $E(\varphi)$
- K -unifier of $t \approx s = L$ -unifier of $F(t, s)$
- $\sigma \equiv_L \tau$ iff $\vdash_L F(\sigma(x), \tau(x))$ for every x
(in our case: $\vdash_L \sigma(x) \leftrightarrow \tau(x)$)
- ...

Admissible rules

Single-conclusion rule: Γ / φ (Γ finite set of formulas)

Multiple-conclusion rule: Γ / Δ (Γ, Δ finite sets of formulas)

- Γ / Δ is **L -derivable** (or **valid**) if $\Gamma \vdash_L \delta$ for some $\delta \in \Delta$
- Γ / Δ is **L -admissible** (written as $\Gamma \sim_L \Delta$) if every L -unifier of Γ also unifies some $\delta \in \Delta$

$$E(\Gamma / \Delta) := \bigwedge_{\gamma \in \Gamma} E(\gamma) \rightarrow \bigvee_{\delta \in \Delta} E(\delta):$$

- Γ / Δ is derivable iff $E(\Gamma / \Delta)$ holds in **all** K -algebras
- Γ / Δ is admissible iff $E(\Gamma / \Delta)$ holds in **free** K -algebras

Note: Γ is unifiable iff $\Gamma \not\sim_L \emptyset$

Multiple-conclusion consequence relations

Single-conc. admissible rules form a consequence relation

Multiple-conc. admissible rules form a (finitary structural) **multiple-conclusion consequence relation**:

- $\varphi \vdash \varphi$
- $\Gamma \vdash \Delta$ implies $\Gamma, \Gamma' \vdash \Delta, \Delta'$
- $\Gamma \vdash \varphi, \Delta$ and $\Gamma, \varphi \vdash \Delta$ imply $\Gamma \vdash \Delta$
- $\Gamma \vdash \Delta$ implies $\sigma(\Gamma) \vdash \sigma(\Delta)$ for every substitution σ

A set B of rules is a **basis** of L -admissible rules if \vdash_L is the smallest m.-c. c. r. containing \vdash_L and B

Admissibly saturated approximation

Γ is **admissibly saturated** if $\Gamma \sim_L \Delta$ implies $\Gamma \vdash_L \Delta$ for any Δ

Assume for simplicity that L has a well-behaved conjunction.

Admissibly saturated approximation of Γ :

a finite set of formulas Π_Γ such that

- each $\pi \in \Pi_\Gamma$ is admissibly saturated
- $\Gamma \sim_L \Pi_\Gamma$
- $\pi \vdash_L \varphi$ for each $\pi \in \Pi_\Gamma$ and $\varphi \in \Gamma$

Application of admissible saturation

Assuming every Γ has an a.s. approximation Π_Γ :

- Reduction of \sim_L to \vdash_L :

$$\Gamma \sim_L \Delta \quad \text{iff} \quad \forall \pi \in \Pi_\Gamma \exists \psi \in \Delta \pi \vdash_L \psi$$

- If $\Gamma \mapsto \Pi_\Gamma$ is computable and \vdash_L is decidable, then \sim_L is decidable
- If Γ / Π_Γ is derivable in \vdash_L + a set of rules $B \subseteq \sim_L$, then B is a basis of admissible rules
- If each $\pi \in \Pi_\Gamma$ has an mgu σ_π , then $\{\sigma_\pi : \pi \in \Pi_\Gamma\}$ is a complete set of unifiers for Γ
 \Rightarrow finitary unification

Projective formulas

π is **projective** if it has a unifier σ such that $\pi \vdash_L x \leftrightarrow \sigma(x)$
(in general: $\pi \vdash_L F(x, \sigma(x))$) for every variable x

- Every projective formula is **admissibly saturated**
- σ is an **mgu** of π : if τ is a unifier of π , then $\tau \equiv_L \tau \circ \sigma$

Projective approximation := admissibly saturated approximation consisting of projective formulas

If projective approximations exist:
convenient tool for analysis of unification and admissibility

Parameters

In real life, propositional atoms model both “variables” and “constants”

We don't want to allow substitution for constants

⇒ Generalize the set-up to use **two kinds of atoms**:

- **variables** $\{x_n : n \in \omega\}$
- **parameters** $\{p_n : n \in \omega\}$
(aka metavariables, constants, coefficients)

Unification with parameters

Substitutions only modify variables, we require $\sigma(p_n) = p_n$

Adapt accordingly the definitions of other notions:

- Unifier, $\sigma \preceq_L \tau$, admissible rule, m.-c. consequence relation, basis, a.s. formula and approximation, projective formula

Exception: “Propositional logic” is always assumed to be closed under substitution for parameters

Transitive modal logics

Transitive modal logics

Normal modal logics with a single modality \Box , include the transitivity axiom $\Box x \rightarrow \Box\Box x$ (i.e., $L \supseteq \mathbf{K4}$)

Common examples: various combinations of

logic	axiom (on top of $\mathbf{K4}$)	finite rooted transitive frames
S4	$\Box x \rightarrow x$	reflexive
D4	$\Diamond \top$	final clusters reflexive
GL	$\Box(\Box x \rightarrow x) \rightarrow \Box x$	irreflexive
K4Grz	$\Box(\Box(x \rightarrow \Box x) \rightarrow x) \rightarrow \Box x$	no proper clusters
K4.1	$\Box \Diamond x \rightarrow \Diamond \Box x$	no proper final clusters
K4.2	$\Diamond \Box x \rightarrow \Box \Diamond x$	unique final cluster
K4.3	$\Box(\Box x \rightarrow y) \vee \Box(\Box y \rightarrow x)$	linear (chain of clusters)
K4B	$x \rightarrow \Box \Diamond x$	lone cluster
S5	$= \mathbf{S4} \oplus \mathbf{B}$	lone reflexive cluster

Some classes of transitive logics

Cofinal-subframe (csf) logics:

- complete wrt a class of frames closed under the removal of a subset of non-final points
- all combinations of logics from the table are csf

Extensible logics:

- If a frame F has a unique root r whose reflexivity is compatible with L , and $F \setminus \{r\} \models L$, then $F \models L$
- K4, S4, GL, K4Grz, S4Grz, D4, K4.1, ... (not K4.2, ...)

Linear extensible logics:

- K4.3, S4.3, GL.3, ...

Unification in transitive modal logics

A lot is known about admissibility without parameters:

- Admissibility is **decidable** in a large class of logics (Rybakov)
- Extensible logics have **projective approximations** (Ghilardi)
 - finitary unification type
 - complete sets of unifiers computable
- **Bases** of admissible rules for extensible logics (J.)
- **Computational complexity** of admissibility (J.)
 - Lower bounds for a quite general class of logics
 - Matching upper bounds for csf extensible logics
- ... and more ...

Projectivity in modal logics

Fix $L \supseteq \mathbf{K4}$ with the finite model property (fmp)

Extension property: if F is a finite L -model with a unique root r and $x \vDash \varphi$ for every $x \in F \setminus \{r\}$, then we can change valuation of variables in r to make $r \vDash \varphi$

Theorem [Ghilardi]: The following are equivalent:

- φ is projective
- φ has the extension property
- θ_φ is a unifier of φ

where θ_φ is an explicitly defined composition of substitutions of the form $\sigma(x) = \Box\varphi \wedge x$ or $\sigma(x) = \Box\varphi \rightarrow x$

Semantics of admissible rules

If L is an extensible logic with fmp, tfae:

- $\Gamma \vdash_L \Delta$
- Γ / Δ holds in every L -frame W s.t. $\forall X \subseteq W$ finite:
 - If $L \not\supseteq \text{S4}$, X has an **irreflexive tight predecessor** t :

$$t\uparrow = X\uparrow$$

- If $L \not\supseteq \text{GL}$, X has a **reflexive tight predecessor** t :

$$t\uparrow = \{t\} \cup X\uparrow$$

For linear extensible L , take only $|X| \leq 1$

Notation: $x\uparrow = \{y : x R y\}$, $x\downarrow = \{x\} \cup x\uparrow$, $X\downarrow = \bigcup_{x \in X} x\downarrow$

Bases of admissible rules

If L is an extensible logic, it has a basis of admissible rules consisting of

$$\frac{\Box y \rightarrow \Box x_1 \vee \cdots \vee \Box x_n}{\Box y \rightarrow x_1, \dots, \Box y \rightarrow x_n} \quad (n \in \omega)$$

if L admits an irreflexive point, and

$$\frac{\Box(y \leftrightarrow \Box y) \rightarrow \Box x_1 \vee \cdots \vee \Box x_n}{\Box y \rightarrow x_1, \dots, \Box y \rightarrow x_n} \quad (n \in \omega)$$

if L admits a reflexive point

For L linear extensible, take only $n = 0, 1$

Complexity of admissible rules

Lower bound:

Assume $L \supseteq \mathbf{K4}$ and every depth-3 tree is a skeleton of an L -frame with prescribed final clusters.

Then L -admissibility is **coNEXP-hard**.

Upper bounds: Admissibility in

- csf extensible logics is **coNEXP-complete**
- csf linearly extensible logics is **coNP-complete**

Unification with parameters in modal logic

Known results

Less is known about admissibility in transitive modal logics in the presence of parameters:

- Rybakov's results on **decidability** of admissibility also apply to admissibility with parameters
- Recently, he expanded the results to effectively construct complete sets of unifiers \Rightarrow **finite unification type**

Terminology: From now on, admissibility and unification always allow parameters

New results

In this talk, we will show:

- Ghilardi-style characterization of projective formulas
- Existence of projective approximations for cluster-extensible (clx) logics [defined on the next slide]
- Semantic description of admissibility in clx logics
- Explicit bases of admissible rules for clx logics
- Computational complexity:
 - Lower bounds on unification in wide classes of transitive logics
 - Matching upper bounds for admissibility in csf clx logics
- Translation of these results to intuitionistic logic

Cluster-extensible logics

Let L be a transitive modal logic with fmp, $n \in \omega$, and C a finite cluster.

A finite rooted frame F is of **type** $\langle n, C \rangle$ if its root cluster $\text{rcl}(F)$ is isomorphic to C and has n immediate successor clusters.

L is **$\langle n, C \rangle$ -extensible** if:

For every type- $\langle n, C \rangle$ frame F , if $F \setminus \text{rcl}(F)$ is an L -frame, then so is F .

L is **cluster-extensible (clx)**, if it is $\langle n, C \rangle$ -extensible whenever there exists a type- $\langle n, C \rangle$ L -frame.

Examples: All combinations of K4, S4, GL, D4, K4Grz, K4.1, K4.3, K4B, S5, \pm bounded branching

Nonexamples: K4.2, S4.2, ...

Projective formulas: the extension property

Fix $L \supseteq \mathbf{K4}$ with the fmp, and P and V finite sets of parameters and variables, resp.

- If F is a rooted model with valuation of $P \cup V$, its **variant** is any model F' which differs from F only by changing the value of some variables $x \in V$ in $\text{rcl}(F)$
- A set M of **finite rooted L -models** evaluating $P \cup V$ has the **model extension property**, if:
every L -model F whose all rooted generated proper submodels belong to M has a variant $F' \in M$
- A **formula** φ in atoms $P \cup V$ has the **model extension property** if $\text{Mod}_L(\varphi) := \{F : \forall x \in F (x \models \varphi)\}$ does

Projective formulas: Löwenheim substitutions

Let φ be a formula in atoms $P \cup V$

- For every $D = \{\beta_x : x \in V\}$, where each β_x is a **Boolean function** of the **parameters** P , define the substitution

$$\theta_D(x) = (\Box\varphi \wedge x) \vee (\neg\Box\varphi \wedge \beta_x)$$

- Let θ_φ be the composition of substitutions θ_D for all the $2^{2^{|P|}|V|}$ possible D 's, in arbitrary order

Projective formulas: a characterization

Theorem:

Let $L \supseteq \mathbf{K4}$ have the fmp, and φ be a formula in finitely many parameters P and variables V . Tfae:

- φ is projective
- φ has the model extension property
- θ_φ^N is a unifier of φ

where $N = (|B| + 1)(2^{|P|} + 1)$, $B = \{\psi : \Box\psi \subseteq \varphi\}$

Remark: If $P = \emptyset$, we have $N \leq 2^{|\varphi|}$.

Ghilardi's original proof gives N nonelementary (tower of exponentials of height $\text{md}(\varphi)$)

Projective approximations

Theorem:

If L is a **clx** logic, every formula φ has a **projective approximation** Π_φ .

Moreover, every $\pi \in \Pi_\varphi$ is a Boolean combination of subformulas of φ .

Corollary:

- $\{\theta_\pi^N : \pi \in \Pi_\varphi\}$ is a complete set of unifiers of φ
- Admissibility in L is decidable (if L r.e.?)
- If $n = |\varphi|$, then $|\Pi_\varphi| \leq 2^{2^n}$, and $|\pi| = O(n2^n) \forall \pi \in \Pi_\varphi$
- $|\theta_\pi^N|$ is doubly exponential in $|B| + |V|$, and triply exponential in $|P|$. This is likely improvable.

Size of projective approximations

The bounds $|\Pi_\varphi| = 2^{2^{O(n)}}$ and $|\pi| = 2^{O(n)}$ for $\pi \in \Pi_\varphi$ are asymptotically optimal, even if $P = \emptyset$:

- If L is $\langle 2, \bullet \rangle$ -extensible (e.g., K4, GL), consider

$$\varphi_n = \bigwedge_{i < n} (\Box x_i \vee \Box \neg x_i) \rightarrow \Box y \vee \Box \neg y$$

$$\Pi_{\varphi_n} = \left\{ \bigwedge_{i < n} (\Box x_i \vee \Box \neg x_i) \rightarrow (y \leftrightarrow \beta(\vec{x})) \mid \beta: \mathbf{2}^n \rightarrow \mathbf{2} \right\}$$

- Similar examples work for $\langle 2, \circ \rangle$ -extensible logics (S4)

Irreflexive extension rules

Let $n < \omega$, and P a finite set of parameters.

$\text{Ext}_{n,\bullet}^P$ is the set of rules

$$\frac{P^e \wedge \Box y \rightarrow \Box x_1 \vee \cdots \vee \Box x_n}{\Box y \rightarrow x_1, \dots, \Box y \rightarrow x_n}$$

for each assignment $e: P \rightarrow \mathbf{2}$

Notation:

$$\varphi^1 = \varphi, \varphi^0 = \neg\varphi, P^e = \bigwedge_{p \in P} p^{e(p)}, \mathbf{2}^P = \{e \mid e: P \rightarrow \mathbf{2}\}$$

Reflexive extension rules

Let C be a finite reflexive cluster

$\text{Ext}_{n,C}^P$ is the set of the following rules:

Pick $E: C \rightarrow 2^P$ and $e_0 \in E(C)$, and consider

$$\frac{P^{e_0} \wedge \Box \left(y \rightarrow \bigvee_{e \in E(C)} \Box (P^e \rightarrow y) \right) \wedge \bigwedge_{e \in E(C)} \Box \left(\Box (P^e \rightarrow \Box y) \rightarrow y \right)}{\Box y \rightarrow x_1, \dots, \Box y \rightarrow x_n} \rightarrow \Box x_1 \vee \dots \vee \Box x_n$$

Tight predecessors

P a finite set of parameters, C a finite cluster, $n < \omega$

- A **P - L -frame** is a (Kripke or general) L -frame W together with a fixed valuation of parameters $p \in P$
- If $X = \{w_1, \dots, w_n\} \subseteq W$ and $E: C \rightarrow 2^P$, a **tight E -predecessor (E -tp)** of X is $\{u_c : c \in C\} \subseteq W$ such that

$$u_c \models P^{E(c)}, \quad u_c \uparrow = X \uparrow \cup \{u_d : d \in c \uparrow\}$$

(Note: $c \uparrow = C$ if C is reflexive, $c \uparrow = \emptyset$ if irreflexive)

- W is **$\langle n, C \rangle$ -extensible** if every $\{w_1, \dots, w_n\} \subseteq W$ has an E -tp for every $E: C \rightarrow 2^P$
- If L is a clx logic, W is **L -extensible** if it is $\langle n, C \rangle$ -extensible whenever L is

Correspondence and completeness

Theorem: If P is a finite set of parameters and W is a descriptive or Kripke P -K4-frame, tfae:

- $W \models \text{Ext}_{n,C}^P$
- W is $\langle n, C \rangle$ -extensible

Corollary: For a logic $L \supseteq \text{K4}$, tfae:

- L is $\langle n, C \rangle$ -extensible
- $\text{Ext}_{n,C}^P$ is L -admissible for every P

Theorem: If L has fmp and is $\langle n, C \rangle$ -extensible for all $\langle n, C \rangle \in X$, then $L + \{\text{Ext}_{n,C}^P : \langle n, C \rangle \in X\}$ is complete wrt **locally finite** (= all rooted subframes finite) P - L -frames, $\langle n, C \rangle$ -extensible for each $\langle n, C \rangle \in X$

Semantics and bases of admissible rules

Theorem:

Let L be a clx logic, and Γ / Δ a rule in a finite set of parameters P . Then tfae:

- $\Gamma \vdash_L \Delta$
- Γ / Δ holds in every [locally finite] L -extensible P - L -frame
- Γ / Δ is derivable in \vdash_L extended by the rules $\text{Ext}_{n,C}^P$ such that L is $\langle n, C \rangle$ -extensible

Corollary: If L is a clx logic, it has a **basis** of admissible rules consisting of $\text{Ext}_{n,C}^P$ for all finite P and all $\langle n, C \rangle$ such that L is $\langle n, C \rangle$ -extensible

Complexity: wide logics

Theorem:

If $L \supseteq \mathbf{K4}$ has width ≥ 2 , then unification (and thus inadmissibility) in L is NEXP-hard.

Theorem:

If L is a csf clx logic of width ≥ 2 and bounded cluster size, then inadmissibility (and thus unification) in L is NEXP-complete.

Examples: GL, K4Grz, S4Grz, ...

Complexity: fat logics

Theorem:

If $L \supseteq \mathbf{K4}$ has unbounded cluster size, then unification in L is coNEXP -hard.

Theorem:

If L is a clx logic of width ≤ 1 and unbounded cluster size, then inadmissibility in L is coNEXP -complete.

Examples: $\mathbf{S5}$, $\mathbf{K4.3}$, $\mathbf{S4.3}$, ...

Complexity: wide and fat logics

L is “chubby” if for all $n > 0$ there is a finite rooted L -frame containing an n -element cluster C and an element incomparable with C

Recall: $\Sigma_2^{\text{EXP}} = \text{NEXP}^{\text{NP}}$

Theorem:

If $L \supseteq \mathbf{K4}$ is chubby, then unification in L is Σ_2^{EXP} -hard.

Theorem:

If L is a csf clx logic of width ≥ 2 and unbounded cluster size, then inadmissibility in L is Σ_2^{EXP} -complete.

Examples: $\mathbf{K4}$, $\mathbf{S4}$, $\mathbf{S4.1}$, ...

Complexity: slim logics

Theorem:

If $L \supseteq \mathbf{K4}$, then unification in L is PSPACE-hard, unless L is a tabular logic of width 1.

Theorem:

If L is a **clx** logic of width 1, bounded cluster size, and depth > 1 , then admissibility in L is PSPACE-complete.

Examples: GL.3, K4Grz.3, S4Grz.3, ...

Theorem:

If L is a tabular logic of width 1 and depth d , then unification and inadmissibility in L are Π_{2d}^P -complete.

Examples: S5 + Alt $_n$, K4 + $\Box\perp$, ...

Complexity: summary

We get the following classification for csf clx logics:

logic		$\not\vdash_L$	param'r-free		with param's		example
cluster size	branching		unif'n	$\not\vdash_L$	unif'n	$\not\vdash_L$	
$< \infty$	0	NP-complete			Π_2^P -c.		S5 + Alt_n
	1				PSPACE-c.		GL.3
∞	≤ 1				coNEXP-c.		S5, S4.3
$< \infty$	≥ 2	PSPACE-c.	?	NEXP-complete		GL, Grz	
∞	(∞)			Σ_2^{EXP} -c.		K4, S4	

Intuitionistic logic

Results for modal logics can be transferred to intermediate logics by means of the Blok–Esakia isomorphism

The following result by Rybakov can be generalized to admissibility with parameters:

Theorem:

If $L \supseteq \text{IPC}$ and σL is its **largest** modal companion, then

$$\Gamma \vdash_L \Delta \Leftrightarrow \mathsf{T}(\Gamma) \vdash_{\sigma L} \mathsf{T}(\Delta),$$

where T is the Gödel translation

[However, $\bigwedge_{p \in P} \Box(p \rightarrow \Box p) \rightarrow \mathsf{T}(\varphi)$ is often more convenient.]

Corollaries

Note: The only “clx” $L \supseteq \text{IPC}$ are IPC itself and the bounded branching logics T_n (incl. $\text{T}_1 = \text{LC}$, $\text{T}_0 = \text{CPC}$)

The translation yields:

- Char. of **projective formulas** in $L \supseteq \text{IPC}$ with fmp
- Existence of **projective approximations** and **semantic description** of \sim_L for IPC and T_n
- **Complexity** (lower bounds need an extra argument):
unification and inadmissibility is
 - NEXP-complete for IPC
 - PSPACE-complete for LC
 - Π_{2d}^{P} -complete for G_{d+1}
 - NEXP-hard for any other intermediate logic

Intuitionistic extension rules

Bases of admissible rules require a separate construction:

A basis for IPC and \mathbf{T}_n is given by the rules

$$\frac{\bigwedge P \wedge \left(\bigvee_{i=1}^n x_i \vee \bigvee Q \rightarrow y \right) \rightarrow \bigvee_{i=1}^n x_i \vee \bigvee Q}{\bigwedge P \wedge y \rightarrow x_1, \dots, \bigwedge P \wedge y \rightarrow x_n}$$

where P, Q are disjoint finite sets of parameters

Questions

- Is there a general **reduction** of admissibility to nonunifiability (with parameters)?
- What is the complexity of **parameter-free** unification for non-linear csf clx logics?
 - NP-hard and NEXP-easy
 - If L includes **D4** or **GL** or $\Box x \rightarrow x \vee \Box \perp$: NP-complete (the universal frame of rank 0 is very simple)
 - Otherwise?

Thank you for attention!

References

- S. Ghilardi, *Best solving modal equations*, Ann. Pure Appl. Log. 102 (2000), 183–198.
- E. Jeřábek, *Admissible rules of modal logics*, J. Log. Comp. 15 (2005), 411–431.
- _____, *Complexity of admissible rules*, Arch. Math. Log. 46 (2007), 73–92.
- V. Rybakov, *Admissibility of logical inference rules*, Elsevier, 1997.
- _____, *Unifiers in transitive modal logics for formulas with coefficients (meta-variables)*, Log. J. IGPL, accepted.