

A posteriori error estimates in the finite element method

A survey of techniques

Tomáš Vejchodský (vejchod@math.cas.cz)
<http://www.math.cas.cz/vejchod>

Institute of Mathematics, Academy of Sciences
Žitná 25, 115 67 Prague 1
Czech Republic



Outline



- ▶ Poisson equation
- ▶ Finite element method
- ▶ Motivation
 - ▶ Adaptive algorithm
 - ▶ Numerical example
- ▶ A priori error estimates
- ▶ A posteriori error estimates
 - ▶ Explicit residual
 - ▶ Implicit residual
 - ▶ Dirichlet type
 - ▶ Neumann type
 - ▶ Hierarchic (residual)
 - ▶ Error majorants
 - ▶ Postprocessing
 - ▶ Quantity of interest

Poisson problem

- ▶ Classical formulation: find $u \in C^2(\Omega) \cap C(\bar{\Omega})$:

$$-\Delta u = f \quad \text{in } \Omega \qquad u = 0 \quad \text{on } \partial\Omega$$

- ▶ Weak formulation: $V = H_0^1(\Omega)$

$$\begin{aligned} u \in V : \quad & \underbrace{\mathcal{B}(u, v)}_{\int_{\Omega} \nabla u \cdot \nabla v \, dx} = \underbrace{\mathcal{F}(v)}_{\int_{\Omega} fv \, dx} \quad \forall v \in V \end{aligned}$$


- ▶ Galerkin method $V_h \subset V \quad \dim V_h < \infty$

$$u_h \in V_h : \quad \mathcal{B}(u_h, v_h) = \mathcal{F}(v_h) \quad \forall v_h \in V_h \quad \Leftrightarrow \quad Ay = F$$

$$u_h(x) = \sum_{j=1}^N y_j \varphi_j(x) \quad \sum_{j=1}^N y_j \underbrace{\mathcal{B}(\varphi_j, \varphi_i)}_{A_{ij}} = \underbrace{\mathcal{F}(\varphi_i)}_{F_i}$$

Finite element method (FEM)

- ▶ FEM $V_h = \{v_h \in V : v_h|_K \in P^1(K) \ \forall K \in \mathcal{T}_h\}$
 $\varphi_1, \dots, \varphi_N \dots$ FEM basis functions $\varphi_i(x_j) = \delta_{ij}$



Motivation – adaptive algorithm


1. **Initialize:** Construct the initial mesh \mathcal{T}_h .
2. **Solve:** Find u_h on \mathcal{T}_h .
3. **Estimate error:** Compute η_K for all $K \in \mathcal{T}_h$.
4. **Stopping criterion:** If $\sum_{K \in \mathcal{T}_h} \eta_K^2 \leq \text{TOL}^2 \Rightarrow \text{STOP}$.
5. **Mark:** If $\eta_K \geq \Theta \max_{K \in \mathcal{T}_h} \eta_K \Rightarrow \text{mark } K$. $0 < \Theta < 1$
6. **Refine:** Refine marked elements and build the new mesh \mathcal{T}_h .
7. GO TO 2.

Motivation – example

$$-\Delta u = 0 \quad \text{in } \Omega$$

$$u = g_D \quad \text{on } \partial\Omega$$

$$u = r^{\frac{2}{3}} \sin \frac{2\theta - \pi}{3}$$




Motivation – example


$$-\Delta u = 0 \quad \text{in } \Omega$$

$$u = g_D \quad \text{on } \partial\Omega$$


$$u = r^{\frac{2}{3}} \sin \frac{2\theta - \pi}{3}$$




Adaptive algorithm



Adaptive algorithm




Adaptive algorithm




Adaptive algorithm


Marked elements 1




Adaptive algorithm



Adaptive algorithm




Adaptive algorithm




Adaptive algorithm


Marked elements 2




Adaptive algorithm



Adaptive algorithm




Adaptive algorithm




Adaptive algorithm

Marked elements 3




Adaptive algorithm


Refined mesh 3



Adaptive algorithm




Adaptive algorithm




Adaptive algorithm

Marked elements 4




Adaptive algorithm


Refined mesh 4



Adaptive algorithm



Adaptive algorithm



A priori error estimates

- ▶ Discretization error: $e = u - u_h$
- ▶ Energy norm: $\|u\|^2 = \mathcal{B}(u, u) = \|u\|_{H^1(\Omega)}^2$
- ▶ Céa's lemma: $\|e\| = \inf_{v_h \in V_h} \|u - v_h\| = \text{dist}(u, V_h)$
- ▶ Lagrange interpolation: $\pi_h^{\text{Lag}} : C(\bar{\Omega}) \mapsto V_h$
 $v \in H^2(\Omega) \cap C(\bar{\Omega}) \quad \Rightarrow \quad \|v - \pi_h^{\text{Lag}} v\|_{H^1(\Omega)} \leq C h |v|_{H^2(\Omega)}$
- ▶ $u \in H^2(\Omega) \cap C(\bar{\Omega}) \quad \Rightarrow \quad \|e\| \leq C h |u|_{H^2(\Omega)}$

A posteriori error estimates I

Definition

- ▶ $\|e\| \approx \eta$ (or $\|e\| \leq \eta$, or $\eta \leq \|e\|$)
- ▶ $\eta = \eta(u_h, f, \Omega, \mathcal{T}_h, \dots)$

Properties

- ▶ Local: $\eta^2 = \sum_{K \in \mathcal{T}_h} \eta_K^2$
- ▶ Guaranteed upper (lower) bound: $\|e\| \leq \eta$ ($\eta \leq \|e\|$)
- ▶ Asymptotic exactness: $\lim_{h \rightarrow 0} I_{\text{eff}} = 1$, $I_{\text{eff}} = \frac{\eta}{\|e\|}$
- ▶ Efficient and reliable: $C_1 \eta \leq \|e\| \leq C_2 \eta$
- ▶ Robust: C_1 and C_2 are independent from quantities like coefficients in the equation, mesh aspect ratio etc.

A posteriori error estimates II

Remarks:

- ▶ Locality
⇒ fast evaluation of η
- ▶ Guaranteed upper bound
⇒ adaptive algorithm guarantees $\|e\| \leq \text{TOL}$
- ▶ Efficiency and reliability
⇒ convergence of adaptive algorithm
- ▶ Assymptotic exactness
⇒ efficiency and reliability (for sufficiently small h)

Proof:

$$\forall \varepsilon > 0 \exists \delta > 0 : 1 - \varepsilon \leq \frac{\eta}{\|e\|} \leq 1 + \varepsilon$$

Residual estimates

$$u \in V : \quad \mathcal{B}(u, v) = \mathcal{F}(v) \quad \forall v \in V$$

$$u_h \in V_h : \quad \mathcal{B}(u_h, v_h) = \mathcal{F}(v_h) \quad \forall v_h \in V_h$$

- ▶ Residual: $\mathcal{R}(v) = \mathcal{F}(v) - \mathcal{B}(u_h, v) \quad \forall v \in V$
- ▶ Residual equation: $e \in V : \quad \mathcal{B}(e, v) = \mathcal{R}(v) \quad \forall v \in V$

$$\left[\mathcal{B}(u, v) - \mathcal{B}(u_h, v) = \mathcal{F}(v) - \mathcal{B}(u_h, v) \quad \forall v \in V \right]$$
- ▶ Galerkin orthogonality: $\mathcal{B}(e, v_h) = 0 \quad \forall v_h \in V_h$
- ▶ $\|e\| = \sup_{0 \neq v \in V} \frac{|\mathcal{B}(e, v)|}{\|v\|} = \sup_{0 \neq v \in V} \frac{|\mathcal{R}(v)|}{\|v\|} = \|\mathcal{R}\|_{V^*}$

Explicit residual estimates I

- Residual splitting: $\mathcal{R}(v) = \sum_{K \in \mathcal{T}_h} \int_K r v \, dx + \sum_{\ell} \int_{\ell} J_{\ell} v \, ds$
 $r = f + \Delta u_h \quad J_{\ell} = (\nabla u_h^+ - \nabla u_h^-) \cdot \nu_{\ell} \quad \ell \text{ are edges in } \mathcal{T}_h$

$$\begin{aligned}\mathcal{R}(v) &= \mathcal{F}(v) - \mathcal{B}(u_h, v) = \sum_{K \in \mathcal{T}_h} \left(\int_K f v \, dx - \int_K \nabla u_h \cdot \nabla v \, dx \right) \\ &= \sum_{K \in \mathcal{T}_h} \left(\int_K f v \, dx + \int_K \Delta u_h v \, dx - \int_{\partial K} \nabla u_h \cdot \nu_K v \, ds \right) \\ &= \sum_{K \in \mathcal{T}_h} \int_K r v \, dx + \sum_{\ell} \int_{\ell} (\nabla u_h^+ - \nabla u_h^-) \cdot \nu_{\ell} v \, ds\end{aligned}$$

Explicit residual estimates II

- Clément inter.: $\pi_h^{\text{Cl}} : V \mapsto V_h$ $\|v - \pi_h^{\text{Cl}} v\|_{0,K} \leq C_1 |K|^{1/2} \|\nabla v\|_{0,\omega_K}$
 $\|v - \pi_h^{\text{Cl}} v\|_{0,\ell} \leq C_2 |\ell|^{1/2} \|\nabla v\|_{0,\omega_\ell}$

$$\begin{aligned}
 \|e\|^2 &= \mathcal{B}(e, e) = \mathcal{R}(e) = \mathcal{R}(e - \pi_h^{\text{Cl}} e) \\
 &= \sum_{K \in \mathcal{T}_h} \int_K r(e - \pi_h^{\text{Cl}} e) \, dx + \sum_{\ell} \int_{\ell} J_{\ell}(e - \pi_h^{\text{Cl}} e) \, ds \\
 &\leq \sum_{K \in \mathcal{T}_h} C_1 \|r\|_{0,K} |K|^{1/2} \|\nabla e\|_{0,\omega_K} + \sum_{\ell} C_2 \|J_{\ell}\|_{0,\ell} |\ell|^{1/2} \|\nabla e\|_{0,\omega_\ell} \\
 &\leq C_3 \left(\sum_{K \in \mathcal{T}_h} |K| \|r\|_{0,K}^2 + \sum_{\ell} |\ell| \|J_{\ell}\|_{0,\ell}^2 \right) + \varepsilon \|e\|^2 \\
 \|e\|^2 &\leq C_4 \underbrace{\left(\sum_{K \in \mathcal{T}_h} |K| \|r\|_{0,K}^2 + \sum_{\ell} |\ell| \|J_{\ell}\|_{0,\ell}^2 \right)}_{(\eta^{\text{expl}})^2} \equiv C_4 (\eta^{\text{expl}})^2
 \end{aligned}$$

Implicit residual estimates – Dirichlet I



Construction:

- ▶ Local Dirichlet problems:

$$e_K^{\text{Dir}} \in H_0^1(K) : \quad \mathcal{B}_K(e_K^{\text{Dir}}, v) = \mathcal{R}_K(v) \quad \forall v \in H_0^1(K)$$

- ▶ Approximate local problems: $V_{0,h}(K) \subset H_0^1(K)$

$$e_{K,h}^{\text{Dir}} \in V_{0,h}(K) : \quad \mathcal{B}_K(e_{K,h}^{\text{Dir}}, v_h) = \mathcal{R}_K(v_h) \quad \forall v_h \in V_{0,h}(K)$$

- ▶ $\eta_K^{\text{Dir}} = \|e_{K,h}^{\text{Dir}}\|_K \quad (\eta^{\text{Dir}})^2 = \sum_{K \in \mathcal{T}_h} (\eta_K^{\text{Dir}})^2$

- ▶ Notation:
 - $\mathcal{B}_K(u, v) = \int_K \nabla u \cdot \nabla v \, dx$
 - $\mathcal{R}_K(v) = \int_K fv \, dx - \int_K \nabla u_h \cdot \nabla v \, dx$
 - $\|v\|_K^2 = \mathcal{B}_K(v, v)$

Implicit residual estimates – Dirichlet II

Guaranteed lower bound:

- ▶ $e^{\text{Dir}}|_K = e_K^{\text{Dir}}$ $\forall K \in \mathcal{T}_h$; $V_0 = \{v \in V : v|_K \in H_0^1(K)\} \subset V$
- ▶ $e_h^{\text{Dir}}|_K = e_{K,h}^{\text{Dir}}$ $\forall K \in \mathcal{T}_h$; $V_{0,h} = \{v \in V : v|_K \in V_{0,h}(K)\} \subset V_0$
- ▶ Theorem: $\|e_h^{\text{Dir}}\| \leq \|e^{\text{Dir}}\| \leq \|e\|$

Proof:

- ▶ $\|e^{\text{Dir}}\| \leq \|e\|:$

$$e^{\text{Dir}} \in V_0 \quad v \in V_0$$

$$\mathcal{B}(e - e^{\text{Dir}}, v) = \mathcal{R}(v) - \mathcal{R}(v) = 0$$

$$\Rightarrow \mathcal{B}(e, e^{\text{Dir}}) = \|e^{\text{Dir}}\|^2$$

$$\begin{aligned} \Rightarrow \|e - e^{\text{Dir}}\|^2 &= \|e\|^2 - 2\mathcal{B}(e, e^{\text{Dir}}) + \|e^{\text{Dir}}\|^2 \\ &= \|e\|^2 - \|e^{\text{Dir}}\|^2 \geq 0 \end{aligned}$$

- ▶ $\|e_h^{\text{Dir}}\| \leq \|e^{\text{Dir}}\|:$

$$\mathcal{B}(e^{\text{Dir}} - e_h^{\text{Dir}}, v_h) = \mathcal{R}(v_h) - \mathcal{R}(v_h) = 0 \quad \forall v_h \in V_{0,h}$$

Similarly.

Implicit residual estimates – Neumann I

Construction:

- Weak f.: $e_K^{\text{Neu}} \in H_E^1(K) = \{v \in H^1(K) : v = 0 \text{ on } \partial K \cap \partial \Omega\}$:

$$\mathcal{B}_K(e_K^{\text{Neu}}, v) = \int_K fv \, dx - \mathcal{B}_K(u_h, v) + \int_{\partial K} g_K v \, ds \quad \forall v \in H_E^1(K)$$

- Classical f.: $-\Delta(e_K^{\text{Neu}} + u_h) = f \quad \text{in } K$

$$\nabla(e_K^{\text{Neu}} + u_h) \cdot \nu_K = g_K \quad \text{on } \partial K \setminus \partial \Omega$$

$$e_K^{\text{Neu}} + u_h = 0 \quad \text{on } \partial K \cap \partial \Omega$$

- $g_K|_\ell \in P^1(\ell)$, $\ell \subset \partial K$, $K \in \mathcal{T}_h$, $g_K \approx \nabla u|_K \cdot \nu_K$ on ∂K
- Compatibility condition: $g_K|_\ell + g_{K^*}|_\ell = 0$ for $\ell = \partial K \cap \partial K^*$
- p -order equilibration condition ($p = 0, 1$):

$$\int_K f \varphi \, dx - \mathcal{B}_K(u_h, \varphi) + \int_{\partial K} g_K \varphi \, ds = 0 \quad \forall \varphi \in P^p(K)$$

- $\eta_K^{\text{Neu}} = \|e_K^{\text{Neu}}\|_K \quad (\eta^{\text{Neu}})^2 = \sum_{K \in \mathcal{T}_h} (\eta_K^{\text{Neu}})^2$

Implicit residual estimates – Neumann II

Guaranteed upper bound:

- Theorem: Compatibility cond. $\Rightarrow \|e\| \leq \eta^{\text{Neu}}$

Proof: $v \in V$

$$\mathcal{B}(e, v) = \mathcal{R}(v) = \sum_{K \in \mathcal{T}_h} \left(\int_K fv \, dx - \mathcal{B}_K(u_h, v) + \int_{\partial K} g_K v \, ds \right)$$

$$= \sum_{K \in \mathcal{T}_h} \mathcal{B}_K(e_K^{\text{Neu}}, v) \leq \sum_{K \in \mathcal{T}_h} \|e_K^{\text{Neu}}\|_K \|v\|_K \leq \left(\sum_{K \in \mathcal{T}_h} \|e_K^{\text{Neu}}\|_K^2 \right)^{\frac{1}{2}} \|v\|$$

- Remark: Approximate Neumann problems:

$$e_{K,h}^{\text{Neu}} \in V_h^{\text{Neu}} \subset H_E^1(K) \Rightarrow \sum_{K \in \mathcal{T}_h} \|e_{K,h}^{\text{Neu}}\|_K^2 \leq (\eta^{\text{Neu}})^2$$

- In general: $\|e\| \not\leq \eta^{\text{Neu}}$

Hierarchic (residual) estimates

- ▶ $\widehat{V}_h = V_h \oplus Y_h \quad Y_h \subset V \quad V_h \cap Y_h = \{0\}$
- ▶ $\widehat{u}_h \in \widehat{V}_h : \quad \mathcal{B}(\widehat{u}_h, \widehat{v}_h) = \mathcal{F}(\widehat{v}_h) \quad \forall \widehat{v}_h \in \widehat{V}_h$
- ▶ $\|e\| \approx \|\widehat{u}_h - u_h\| \equiv \|\widehat{e}_h\|$
- ▶ $\overline{e}_h \in Y_h : \quad \mathcal{B}(\overline{e}_h, y_h) = \mathcal{R}(y_h) \quad \forall y_h \in Y_h$
- ▶ $\|e\| \approx \|\overline{e}_h\| \equiv \eta^{\text{Hie}}$
- ▶ Saturation assumption:
 $\exists \beta < 1: \quad \|u - \widehat{u}_h\| \leq \beta \|u - u_h\|$
- ▶ Strenghtened Cauchy-Schwarz inequality:
 $\exists \gamma < 1: \quad |\mathcal{B}(v_h, y_h)| \leq \gamma \|v_h\| \|y_h\| \quad \forall v_h \in V_h, y_h \in Y_h$
- ▶ $\|\overline{e}_h\| \leq \|\widehat{e}_h\| \leq \|e\| \leq \frac{1}{(1 - \beta^2)^{\frac{1}{2}}} \|\widehat{e}_h\| \leq \frac{1}{(1 - \beta^2)^{\frac{1}{2}}(1 - \gamma^2)^{\frac{1}{2}}} \|\overline{e}_h\|$

Error majorants

- ▶ Friedrichs inequality: $\|v\|_{0,\Omega} \leq C_\Omega \|\nabla v\|_{0,\Omega} \quad \forall v \in V = H_0^1(\Omega)$
- ▶ $\|e\| \leq C_\Omega \|f + \operatorname{div} \mathbf{y}\|_{0,\Omega} + \|\mathbf{y} - \nabla u_h\|_{0,\Omega} \quad \forall \mathbf{y} \in \mathbf{H}(\operatorname{div}, \Omega)$

Proof:

$$v \in V \quad \mathbf{y} \in \mathbf{H}(\operatorname{div}, \Omega)$$

$$\begin{aligned} \mathcal{B}(e, v) &= \mathcal{R}(v) + \int_{\Omega} \mathbf{y} \cdot \nabla v \, dx + \int_{\Omega} \operatorname{div} \mathbf{y} v \, dx + \int_{\partial\Omega} \mathbf{y} \cdot \nu v \, ds \\ &= \int_{\Omega} (f + \operatorname{div} \mathbf{y}) v \, dx + \int_{\Omega} (\mathbf{y} - \nabla u_h) \cdot \nabla v \, dx \\ &\leq \|f + \operatorname{div} \mathbf{y}\|_{0,\Omega} \|v\|_{0,\Omega} + \|\mathbf{y} - \nabla u_h\|_{0,\Omega} \|\nabla v\|_{0,\Omega} \\ &\leq \left(C_\Omega \|f + \operatorname{div} \mathbf{y}\|_{0,\Omega} + \|\mathbf{y} - \nabla u_h\|_{0,\Omega} \right) \|\nabla v\|_{0,\Omega} \end{aligned}$$

Put $v = e$.

Postprocessing

- ▶ Recovered gradient: $\nabla u_h \mapsto \mathcal{G}(u_h)$
- ▶ $\|e\| \approx \eta^{\text{post}} = \|\mathcal{G}(u_h) - \nabla u_h\|_{0,\Omega}$
- ▶ Superconvergence: $\|\nabla u - \mathcal{G}(u_h)\|_{0,\Omega} \leq C_1 h^{1+\epsilon}$
- ▶ Assumption: $\|e\| \geq C_2 h$
- ▶ Theorem (asymptotic exactness): $\lim_{h \rightarrow 0} \frac{\eta^{\text{post}}}{\|e\|} = 1$

Proof:

$$\frac{\eta^{\text{post}}}{\|e\|} \leq \frac{\|\nabla u - \nabla u_h\|_0}{\|e\|} + \frac{\|\nabla u - \mathcal{G}(u_h)\|_0}{\|e\|} \leq 1 + \frac{C_1 h^{1+\epsilon}}{C_2 h} \rightarrow 1$$

$$\frac{\eta^{\text{post}}}{\|e\|} \geq \frac{\|\nabla u - \nabla u_h\|_0}{\|e\|} - \frac{\|\nabla u - \mathcal{G}(u_h)\|_0}{\|e\|} \geq 1 - \frac{C_1 h^{1+\epsilon}}{C_2 h} \rightarrow 1$$

Quantity of interest

- ▶ Quantity of interest: $\Phi \in V^*$
- ▶ Adjoint problem: $z \in V : \quad \mathcal{B}(v, z) = \Phi(v) \quad \forall v \in V$
- ▶ Approx. adjoint prob.: $z_h \in V_h : \quad \mathcal{B}(v_h, z_h) = \Phi(v_h) \quad \forall v_h \in V_h$
- ▶ Error representation formula:

$$\Phi(e) = \mathcal{B}(e, z) = \mathcal{R}(z) = \mathcal{R}(z - z_h) = \mathcal{B}(u - u_h, z - z_h)$$

$$\begin{aligned} |\Phi(e)| &\leq \|u - u_h\| \|z - z_h\| \\ &\leq \eta^{\text{pri}} \eta^{\text{adj}} \end{aligned}$$

A posteriori error estimators

- ▶ Explicit residual – fast, simple, reliable, (efficient)
- ▶ Implicit residual
 - ▶ Dirichlet type – guaranteed lower bound, (reliable)
 - ▶ Neumann type – upper bound (not guaranteed), (efficient)
- ▶ Hierarchic (residual) – efficient and reliable
- ▶ Error majorants – guaranteed upper bound, demanding
- ▶ Postprocessing – fast, simple, superconvergence \Rightarrow asympt. exact
- ▶ Quantity of interest – if energy norm is not the goal

Recommended books



-  M. Ainsworth, J. T. Oden, *A posteriori error estimation in finite element analysis*, Wiley, New York, 2000.
-  I. Babuška, T. Strouboulis, *The finite element method and its reliability*, Clarendon Press, Oxford University Press, New York, 2001.
-  P. Neittaanmäki, S. Repin, *Reliable methods for computer simulation, error control and a posteriori estimates*, Elsevier, Amsterdam, 2004.
-  S. Repin, *A posteriori estimates for partial differential equations*, de Gruyter, Berlin, 2008.
-  R. Verfürth, *A review of a posteriori error estimation and adaptive mesh-refinement techniques.*, Wiley-Teubner, Chichester/Stuttgart, 1996.