

Physiological Research Pre-Press Article

Involvement of actin microfilament in regulation of pacemaking activity increased by hypotonic stress in cultured ICCs of murine intestine

Zuo Yu Wang¹, Xu Huang¹, Dong Hai Liu¹, Hong Li Lu¹, Young Chul Kim², Wen Xie Xu^{1*}

1. Department of Physiology, Shanghai Jiaotong University School of Medicine Shanghai, 200240, China

2. Department of Physiology, Chungbuk National University College of Medicine, 12 Gaeshin-dong, Hungduk-gu, Cheongju, Chungbuk, 361-763, Korea

* Correspondence to Professor Wen Xie Xu

Department of Physiology, Shanghai Jiaotong University School of Medicine

800 Dongchuan Rd, 328 Wenxuan Medical Building, Shanghai 200240, China

Tel: +86-21-34205639 E-mail: wenxiexu@sjtu.edu.cn

Short title: actin microfilament and ICCs pacemaking activity

Abstract

Distension is a regular mechanical stimulus in gastrointestinal (GI) tract. This study was designed to investigate the effect of hypotonic stress on pacemaking activity and determine whether actin microfilament is involved in its mechanism in cultured murine intestinal interstitial cells of Cajal (ICCs) by using whole-cell patch-clamp and calcium imaging techniques. Hypotonic stress induced sustained inward holding current from the baseline to -650 ± 110 pA and significantly decreased amplitudes of pacemaker current. Hypotonic stress increased the intensity of basal fluorescence ratio (F/F₀) from baseline to 1.09 ± 0.03 and significantly increased Ca²⁺ oscillation amplitude. Cytochalasin-B (20 μM), a disruptor of actin microfilaments, significantly suppressed the amplitudes of pacemaker currents and calcium oscillations, respectively. Cytochalasin-B also blocked hypotonic stress-induced sustained inward holding current and hypotonic stress-induced increase of calcium oscillations. Phalloidin (20 μM), a stabilizer of actin microfilaments, significantly enhanced the amplitudes of pacemaker currents and calcium oscillations, respectively. Despite the presence of phalloidin, hypotonic stress was still able to induce an inward holding current and increased the basal fluorescence intensity. These results suggest that hypotonic stress induces sustained inward holding current via actin microfilaments and the process is mediated by alteration of intracellular basal calcium concentration and calcium oscillation in cultured intestinal ICCs.

Key words: ICCs; hypotonic stress; actin microfilament; pacemaking activity; calcium oscillation

1. Introduction

The intrinsic (i.e. non-neural, non-hormonal) regulation of gut motility has been referred to in the literature as "myogenic", but it is now recognized that the traditional use of this term includes, besides smooth muscle mechanisms, the behaviour of intestinal interstitial cells of Cajal (ICCs) (Sanders 2008). ICCs are electrically coupled to smooth muscle cell, and affect resting membrane potentials of the smooth muscle/ICCs syncytium and impose pacemaking activity, sensitivity to stretch and responsiveness to neurotransmitters (Sanders *et al.* 2006) on the behaviour of smooth muscle cell. Mechanosensitivity underlies several of the fundamental processes that are required for effective gastrointestinal (GI) function. The GI tract is required to "sense" force and translate it to a chemical or electrical signal (Kraichely and Farrugia 2007).

ICCs are responsible for spontaneous and rhythmic electrical activity in GI tract. Although the mechanosensitivity underlying several fundamental processes of GI smooth muscle has been studied considerably, little is known about the mechanosensitivity underlying the pacemaking activity of ICCs. Stretch-dependent responses in antrum were mediated by ICCs-IM because no response was observed in antral muscles of W/W^V mice which lack ICCs-IM (Burns *et al.* 1996). The transient receptor potential (TRP) superfamily of non-selective channels has been identified in GI smooth muscle and ICCs (Kim *et al.* 2006). The TRP channels, in general, have multiple gating mechanisms. Several TRP channels including TRPA1 (Corey *et al.* 2004) and TRPC1 (Maroto *et al.* 2005) have been demonstrated to have mechanosensitivity. Strege *et al.* (2003) observed a mechanosensitive sodium channel current in human intestinal ICCs. They suggested that the channel may play an important role in normal physiological control of human intestinal motor function by contributing to setting of membrane potential, the rate of rise of slow wave and mechanosensitive regulation of slow wave frequency. A previous study at the tissue level also demonstrated that increasing muscle length resulted in significant changes in the resting potentials and intrinsic frequency of antral pacemakers (Won *et al.*

2005). Recently, ICCs have also been suggested to act as a stretch receptor to detect circumferential expansion and swelling of GI tract wall and trigger the contraction of smooth muscle (Aranishi *et al.* 2009).

The actomyosin cytoskeleton is a large network of structural, motor and signaling proteins that coordinate a plethora of cellular functions including cell division, adhesion and migration. Actin microfilaments were necessary in regulation of cell response to physical forces (Higashida *et al.* 2013). The activities of various transport proteins as well as ion and water permeable channels have been shown to be dependent on the extent of actin polymerization (Constantin 2013, Prat *et al.* 1993). Our previous study also demonstrated that actin microfilament is involved in regulation of pacemaker currents in cultured intestinal ICCs (^①Wang *et al.* 2010). However, little is known about the direct effect of hypotonic stress on ICCs pacemaker current. Accordingly, the present study was aimed to clarify the effect of hypotonic stress on ICCs pacemaker current and to test whether actin cytoskeleton takes part in this mechanism in cultured murine intestinal ICCs by using patch clamp and Ca²⁺ fluorescence techniques.

2. Materials and Methods

2.1 Preparation of cells

Balb/C mice (7-13 days old) of either sex were obtained from the Experimental Animal Center of the Chinese Academy of Science. Animals were killed by cervical dislocation. The small intestine was dissected from 1 cm below the pyloric ring to the cecum, and luminal contents were washed out with Krebs-Ringer bicarbonate (KRB). Tissues were pinned to the base of a **Sylgard** dish and the mucosae removed. Small strips of intestinal muscle were incubated at 37 °C for 23 min in enzyme solution comprising: 1.3 mg/ml collagenase (type II, Worthington), 2 mg/ml bovine serum albumin (Sino-American Biotechnology Co., China), 2 mg/ml trypsin inhibitor (Amresco, OH, USA) and 0.27 mg/ml ATP (Sigma Aldrich, St. Louis, MO, US). The muscle strips were

then washed three times to remove the enzyme before being triturated through blunt pipettes. The resulting cell suspension was plated on murine collagen-coated (2.5 µg/ml) sterile glass cover slips in culture dishes. Cells were allowed to settle for 30 min before adding SMGM medium (Clonetics Corp, San Diego, CA, USA) supplemented with 2% antibiotics/antimycotics (Gibco Invitrogen Corp, Grand Island, USA) and murine stem cell factor (SCF, 5 ng/ml, Sigma Aldrich, St. Louis, MO, US). Cells were then incubated at 37 °C in a 5% CO₂ incubator. The above methods were performed as described in ref. Zhang and the cultured cells have been confirmed to be ICCs (Zhang *et al.* 2006).

All experimental protocols were approved by the local animal care committee and conformed with the Guide for the Care and Use of Laboratory Animals published by the Science and Technology Commission of P.R.C (STCC Publication No.2, revised 1988).

2.2 Ca²⁺ fluorescent imaging

Cultured ICCs (48-72 h) were loaded with the Ca²⁺-sensitive indicator Fluo-3AM (8 µM) and Pluronic F-127 (0.03%) for 2 h at 37 °C and washed for a further 30 min in Physiological Salt Solution (PSS) for de-esterification. A microscope (Olympus BX51, Japan) equipped with a Hamamatsu 1394 Orca-ER CCD camera (Hamamatsu, Japan) was used to monitor and record digital images. Simple PCI 6 software (Compix Inc., USA) was used for data analysis. Cells were illuminated at 488 nm and emission light of 515–565 nm was detected. Fluorescence intensity was expressed as a fluorescence ratio (F/F₀), elucidating Ca²⁺ fluorescence intensity from its arbitrary resting fluorescence intensity (F₀). All recordings were made at room temperature with digital images recorded every second.

2.3 Patch clamp experiments

The whole cell patch clamp technique was used to record inward currents (voltage clamp) and membrane potential (current clamp) from cultured ICCs (48-72 h). Glass pipettes with a resistance of 3-5 MΩ were used to

form giga seals. Electric signals were amplified with an EPC-10 patch clamp amplifier (HEKA Instrument, Germany) and digitized with a PCI-16 A/D converter (HEKA Instrument, Germany). Data were analyzed using Origin 6.0 and Sigmaplot 2.0. All recordings were made at 30 °C.

2.4 Solutions and drugs

The KRB solution comprised (mM): KCl 4.7, NaCl 117, CaCl₂ 2.6, Glucose 11, MgSO₄ 1.2, KH₂PO₄ 1.2 and NaHCO₃ 25; and was bubbled with 95% O₂ / 5% CO₂ (final pH 7.4). PSS comprised (mM): KCl 4.5, NaCl 135, CaCl₂ 2, Glucose 10, MgCl₂ 1, 4-(2-Hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) 10 (final pH adjusted to 7.4 with Tris). Isotonic bath solution (310 mOsm) contained (mM): KCl 4.5, NaCl 95, CaCl₂ 2, Glucose 5, MgCl₂ 1, HEPES 10, sucrose 90; the pH was adjusted to 7.4 with Tris. Hypotonic solution (220 mOsm) had no sucrose but the other ingredients were the same as for the isotonic solution. The pipette solution comprised (mM): KCl 140, MgCl₂ 5, K₂ATP 2.7, Na₂GTP 0.1, creatine phosphate (disodium salt) 2.5, HEPES 5 and ethylene glycol tetraacetic acid (EGTA) 0.1 (final pH adjusted 7.2 with Tris).

Cytochalasin-B and phalloidin were dissolved in dimethyl sulfoxide (DMSO) to provide stock solutions stored at -20 °C until use. EGTA was dissolved in distilled water. Fluo-3AM and Pluronic F-127 were purchased from AnaSpec Inc. All other drugs/agents were purchased from Sigma (Aldrich, St. Louis, MO, US).

2.5 Statistical analysis and presentation

Data are expressed as mean ± SEM. Differences between data sets were evaluated using paired t-tests. *P* values <0.05 were considered significant. The 'n' values reported correspond to the number of cells analyzed in each experiment.

3. Results

3.1 Effect of hypotonic stress on ICCs pacemaker current and $[Ca^{2+}]_i$

Under whole cell voltage clamp mode the spontaneous and rhythmic inward currents were recorded in cultured intestinal ICCs at -60mV of holding potential. Hypotonic stress activated sustained inward holding current and decreased amplitude of pacemaker current (Fig.1Aa). The peak inward holding current was increased from the baseline to **-650±110 pA** (Fig.1Ab, n=6, $P<0.05$) and the amplitude of pacemaker current was decreased from **220±50 pA** to **140±50 pA** (Fig.1Ac, n=6, $P<0.05$). However, hypotonic stress did not affect frequencies of pacemaker current and the frequencies were 11.38 ± 1.31 cycles min⁻¹ in control and 11.63 ± 1.15 cycles min⁻¹ in hypotonic condition, respectively (n=6, $P>0.05$). The above results have already published in *J Biomech.* (Wang *et al.* 2010)

It has been demonstrated that calcium and its oscillation are very important in generation of pacemaker currents (Lowie *et al.* 2011). So we observed the effect of hypotonic stress on calcium oscillation of ICCs. Cultured ICCs (48-72 h) preloaded with fluo-3, a high affinity Ca²⁺-sensitive indicator, exhibited spontaneous periodic calcium oscillation. When cells were exposed to hypotonic solution (220 mOsm), mean amplitude of spontaneous Ca²⁺ oscillation increased from 0.08 ± 0.01 to 0.19 ± 0.03 ($\Delta F/F_0$) and basal fluorescence intensity increased from baseline to 1.09 ± 0.03 (F/F₀) (Fig.1B, n=6, $P<0.05$). However hypotonic stress did not significantly change the frequencies of Ca²⁺ transients (5.50 ± 0.55 cycles min⁻¹ in control and 6.30 ± 0.91 cycles min⁻¹ in hypotonic condition, respectively, n=6, $P>0.05$).

3.2 Effect of cytochalasin-B on hypotonic stress -induced increase of pacemaker current and [Ca²⁺]_i

Actin microfilaments were necessary in regulation of cell response to physical forces (Higashida *et al.* 2013). Our previous study also demonstrated actin microfilament involves in regulation of pacemaker currents in cultured intestinal ICCs (Wang *et al.* 2010). Cytochalasin B or D and phalloidin are the most common chemical compound to depolymerize and stabilize actin microfilaments (Mazzochi *et al.* 2006, Gravante *et al.* 2004, Faussonne-Pellegrini and Thuneberg 1999). So in this study we used cytochalasin-B and phalloidin as disruptor

and stabilizer of actin microfilaments.

Under whole cell voltage clamp mode the spontaneous rhythmic inward pacemaker currents were recorded at -60 mV of holding potential. Addition of 20 μ M cytochalasin-B to the external perfusion solution significantly suppressed pacemaker currents (Fig.2Aa). The amplitude and frequency of pacemaker currents were reduced from **490 \pm 160 pA** and 11.73 \pm 0.79 cycles/min to **230 \pm 90 pA** and 10.29 \pm 0.76 cycles/min following cytochalasin-B treatment, respectively (Fig.2B, n=6, P <0.05). Pretreatment of 20 μ M cytochalasin-B completely blocked the inward holding current induced by hypotonic stress (Fig.2Ab, n=6). The amplitude of pacemaker currents was reduced from **230 \pm 90 pA** to **110 \pm 40 pA** following cytochalasin-B plus hypotonic stress treatment (Fig.2Ca, n=6, P <0.05). The frequency of pacemaker currents was changed from 10.29 \pm 0.76 cycles/min to 9.65 \pm 0.85 cycles/min following cytochalasin-B plus hypotonic stress treatment, (Fig.2Cb, n=6, P >0.05).

Cytoskeleton regulate Ca^{2+} influx and release from intracellular calcium stores (Bose and Thomas, 2009). To investigate the mechanisms of hypotonic stress on pacemaker currents, we tested the effect of cytochalasin-B on hypotonic stress -induced increase of periodic $[\text{Ca}^{2+}]_i$ oscillation. 20 μ M cytochalasin-B significantly suppressed calcium oscillations amplitude ($\Delta F/F_0$) from 0.30 \pm 0.06 to 0.09 \pm 0.02 (Fig.3A, n=6, P <0.05). Pretreatment of cytochalasin-B completely blocked hypotonic stress-induced increase of the basal fluorescence intensity and the amplitude of calcium oscillations. The basal fluorescence intensity and amplitude ($\Delta F/F_0$) of calcium oscillations were 1.05 \pm 0.03 and 0.09 \pm 0.02 in cytochalasin-B group and 1.04 \pm 0.03 and 0.08 \pm 0.03 in cytochalasin-B plus hypotonic stress group (Fig.3B, n=6, P >0.05).

3.3 Effect of phalloidin on hypotonic stress-induced increase of pacemaker current and $[\text{Ca}^{2+}]_i$

The spontaneous rhythmic inward pacemaker currents were recorded in same as the previous condition. 20 μ M phalloidin significantly enhanced pacemaker currents (Fig.4Aa). The amplitude and frequency of pacemaker currents were increased from **750 \pm 280 pA** and 13.93 \pm 1.00 cycles/min to **1230 \pm 610 pA** and 14.68 \pm 1.00

cycles/min following phalloidin treatment, respectively (Fig.4B, n=6, $P<0.05$). Moreover, pretreatment of phalloidin did not blocked Hypotonic stress -induced inward holding current (Fig.4Ab, n=6). The amplitude of pacemaker currents was reduced from 1230 ± 610 pA to 200 ± 60 pA following phalloidin plus hypotonic stress treatment (Fig.4Ca, n=6, $P<0.05$). The frequency of pacemaker currents was changed from 14.68 ± 1.00 cycles/min to 14.00 ± 1.00 cycles/min following phalloidin plus hypotonic stress treatment, (Fig.4Cb, n=6, $P>0.05$).

20 μ M phalloidin significantly potentiated calcium oscillations amplitude ($\Delta F/F_0$) from 0.56 ± 0.13 to 0.72 ± 0.08 (Fig.5A, Ba, n=6, $P<0.05$). Hypotonic stress-induced increase in the amplitude ($\Delta F/F_0$) of calcium oscillations was blocked by pretreatment with phalloidin. The calcium oscillations amplitude ($\Delta F/F_0$) were 0.72 ± 0.08 in phalloidin group and 0.68 ± 0.12 in phalloidin plus hypotonic stress group (Fig.5A, Ba, n=6, $P>0.05$ vs hypotonic stress plus phalloidin group). However, pretreatment of phalloidin did not blocked hypotonic stress-induced increase in basal fluorescence intensity and the basal $\Delta F/F_0$ was enhanced from baseline to 1.32 ± 0.07 by hypotonic stress after pretreatment of phalloidin (Fig.5A, Bb, n=6, $P<0.05$).

4. Discussion

Distension is one of the major stimuli to induce motor activity in GI smooth muscle. It is speculated that parts of the cellular apparatus including ionic conductance(s) regulate membrane potential and excitability during distension of the bowel wall and this may be an important aspect of the 'myogenic response' to stretch (Won *et al.* 2013). Also the myogenic mechanism for the stretch-induced increase in contractility of smooth muscle in GI tract is not clear and it has been reported that intact cytoskeleton is related to mechanosensitivity (Kraichely and Farrugia 2007). Our previous study indicated that actin microfilament is involved in generation of pacemaker current in cultured intestinal ICCs and the mechanism is related to IP3-induced calcium release

(IICR). (Wang *et al.* 2010). The present study demonstrates that hypotonic stress activated sustained inward holding current and actin microfilaments were implicated in the process. Our results suggest that hypotonic stress may be important stimuli to regulate the pacemaking activity in intestinal ICCs via actin microfilaments.

Several mechanosensitive channels have been reported to be related to the myogenic mechanism, such as L-type Ca^{2+} channel (Kraichely *et al.* 2009), stretch-dependent K^+ (SDK) channels (Won *et al.* 2013) and stretch-activated non-selective cation channels (Wang *et al.* 2003). ICCs generate spontaneous electrical slow wave activity, conduct a rhythmic impulse to smooth muscle cells and mediate the propagation of the slow waves (Ward and Sanders 2001). This may indicate that ICCs may play a functional role in regulation of stretch-induced rhythmic motor activity. Ca^{2+} -dependent plasma membrane ion channels are responsible for mediating ICCs pacemaker potential (Lowie *et al.* 2011). In this study, we have found that hypotonic stress elicited a large inward holding current and reduced the amplitude of the pacemaker current generated by ICCs (Fig.1A). These findings suggest that distension or stretch stimulus may regulate motor activity of GI tract via changing pacemaking activity of ICCs. This is coincidence with the report that many kinds of mechanosensitivity channels are existed in ICCs of GI tract and participate in regulating cell response to force with a change in their open probability (Kraichely and Farrugia 2007). These results may support the previous observation showing a mechanosensitive role for interstitial cells of Cajal in smooth muscle tissues (Won *et al.* 2005).

The actomyosin cytoskeleton is a large network of structural, motor and signaling proteins that coordinate a plethora of cellular functions including cell division, adhesion and migration. Actin microfilaments were necessary in regulation of cell response to physical forces (Higashida *et al.* 2013). The activities of various transport proteins as well as ion and water permeable channels have been shown to be dependent on the extent of actin polymerization (Constantin 2013, Prat *et al.* 1993). Some mechanosensitive channels have been reported to

be related to cytoskeleton, like Stretch-dependent potassium channels (Koh and Sanders 2001) and muscarinic current (Wang *et al.* 2003) in GI smooth muscle and Nav1.5 and voltage-gated Ca^{2+} channel (Kraichely and Farrugia 2007) in ICCs of GI tract. In this study, depolymerized actin microfilament inhibited pacemaker current of ICCs and blocked hypotonic stress-induced inward holding current (Fig.2). In contrast, more polymerized actin microfilament increased pacemaker current of ICCs and did not block the inward holding current induced by hypotonic stress (Fig.4). These results indicate that actin microfilaments participate in the generation of pacemaking activity of ICCs and also play a key role in the action of hypotonic stress to intensify electrical activities of ICCs.

The oscillations of intracellular calcium ($[\text{Ca}^{2+}]_i$) observed in ICCs can periodically activate plasma membrane low Ca^{2+} -dependent ion channels, thereby generating pacemaker potentials (Nose *et al.* 2000, van Helden *et al.* 2000). Our previous studies have indicated that hypotonic stress potentiates muscarinic currents and depolarizes membrane potential by triggering ryanodine sensitive $[\text{Ca}^{2+}]_i$ store in gastric myocytes of guinea-pig (Li *et al.* 2002, Yu *et al.* 2002). To investigate the mechanism by which hypotonic stress increases the pacemaking activity, we further examined the effect of hypotonic stress on calcium oscillations by using the calcium imaging technique. Our data showed that hypotonic stress significantly enhances intracellular basal calcium concentrations and potentiates spontaneous Ca^{2+} oscillation amplitudes (Fig.1B). This result suggests that the increase in $[\text{Ca}^{2+}]_i$ induced by hypotonic stress maybe one of the key mechanism underlying the effect of hypotonic stress on pacemaker current. Our previous study also observed that 2-APB, an IP_3 receptor blocker, blocked both pacemaker current and Ca^{2+} transients but did not alter the effect of hypotonic stress on pacemaker current and Ca^{2+} transients. In contrast, ryanodine inhibited Ca^{2+} transients but not pacemaker current, and completely blocked hypotonic stress -induced inward holding current and hypotonic stress -induced increase of basal $[\text{Ca}^{2+}]_i$ (Wang *et al.* 2010). Thus, we presume that hypotonic stress -induced holding inward current may

be mediated by enhancement of $[Ca^{2+}]_i$ from ryanodine-sensitive store.

The cytoskeleton plays an important role in regulating Ca^{2+} influx or releasing (Bose and Thomas 2009). Young *et al.* (1997) found that cultured myocytes from the circular muscle layer of the rabbit distal colon responded to brief focal mechanical deformation of the plasma membrane with a transient increase in intracellular calcium concentration. They suggested that a communication between the plasma membrane and an internal Ca^{2+} store may be mediated by direct mechanical link via actin filaments. This is also supported by our results that depolymerized actin microfilament inhibited calcium oscillation amplitudes of ICCs and blocked hypotonic stress-induced increase of basal calcium concentrations (Fig.3). In contrast, the stabilizer of actin microfilament increased calcium oscillations amplitude of ICCs and did not inhibit the increase of basal calcium concentration (Fig.5). The results suggest that actin microfilament regulates the effect of hypotonic stress on pacemaker current via changing intracellular calcium concentration.

5. Conclusion

Hypotonic stress can regulate gastrointestinal smooth motility by potentiating ICCs pacemaking activity. Actin microfilaments are involved in the regulation of pacemaker currents by hypotonic stress via alteration of $[Ca^{2+}]_i$. Polymerized types of actin is vital for ICCs pacemaker activity response to stretch stimulation.

Conflict of interests statement

The authors declare that there are no conflicts of interest associated with this work.

Acknowledgments

This study was funded by Grant No.10672103; 31271236 provided by National Natural Science Foundation of China.

Reference

- ARANISHI H, KUNISAWA Y, KOMURO T: Characterization of interstitial cells of Cajal in the subserosal layer of the guinea-pig colon. *Cell Tissue Res* **335**: 323-329, 2009.
- BOSE DD, THOMAS DW: The actin cytoskeleton differentially regulates NG115-401L cell ryanodine receptor and inositol 1,4,5-trisphosphate receptor induced calcium signaling pathways. *Biochem. Biophys. Res. Commun* **379**: 594-599, 2009.
- BURNS AJ, LOMAX AE, TORIHASHI S, SANDERS KM, WARD SM: Interstitial cells of Cajal mediate inhibitory neurotransmission in the stomach. *Proc Natl Acad Sci USA* **93**: 12008-12013, 1996.
- Corey DP, Garc ía-A ñoveros J, Holt JR, Kwan KY, Lin SY, Vollrath MA, Amalfitano A, Cheung EL, Derfler BH, Duggan A, G ómez GS, Gray PA, Hoffman MP, Rehm HL, Tamasauskas D, Zhang DS: TRPA1 is a candidate for the mechanosensitive transduction channel of vertebrate hair cells. *Nature* **432**: 723-730, 2004.
- CONSTANTIN B: Dystrophin complex functions as a scaffold for signalling proteins. *Biochim Biophys Acta* **1838**: 635-642, 2014.
- FAUSSONE-PELLEGRINI MS, THUNEBERG L: Guide to the identification of interstitial cells of Cajal. *Microsc Res Tech* **47**: 248-266, 1999.
- GRAVANTE B, BARBUTI A, MILANESI R, ZAPPI I, VISCOMI C, DIFRANCESCO D: Interaction of the pacemaker channel HCN1 with filamin A. *J Biol Chem* **279**: 43847-43853, 2004.
- HIGASHIDA C, KIUCHI T, AKIBA Y, MIZUNO H, MARUOKA M, NARUMIYA S, MIZUNO K, WATANABE N: F- and G-actin homeostasis regulates mechanosensitive actin nucleation by formins. *Nat Cell Biol* **15**: 395-405, 2013.
- KIM BJ, SO I, KIM KW: The relationship of TRP channels to the pacemaker activity of interstitial cells of Cajal in the gastrointestinal tract. *J Smooth Muscle Res* **42**: 1-7, 2006.

- KRAICHELY RE, FARRUGIA G: Mechanosensitive ion channels in interstitial cells of Cajal and smooth muscle of the gastrointestinal tract. *Neurogastroenterol Motil* **19**: 245-252, 2007.
- KRAICHELY RE, STREGE PR, SARR MG, KENDRICK ML, FARRUGIA G: Lysophosphatidyl choline modulates mechanosensitive L-type Ca²⁺ current in circular smooth muscle cells from human jejunum. *Am J Physiol Gastrointest Liver Physiol* **296**: G833- G839, 2009.
- LOWIE BJ, WANG XY, WHITE EJ, HUIZINGA JD: On the origin of rhythmic calcium transients in the ICC-MP of the mouse small intestine. *Am J Physiol Gastrointest Liver Physiol* **301**: G835-G845, 2011.
- KOH SD, SANDERS KM: Stretch-dependent potassium channels in murine colonic smooth muscle cells. *J Physiol* **533**: 155-163, 2001.
- LI L, JIN NG, PIAO L, HONG MY, JIN ZY, LI Y, XU WX: Hyposmotic membrane stretch potentiated muscarinic receptor agonist-induced depolarization of membrane potential in guinea-pig gastric myocytes. *World J Gastroenterol* **8**: 724-727, 2002.
- MAZZOCHI C, BENOS DJ, SMITH PR: Interaction of epithelial ion channels with the actin-based cytoskeleton. *Am J Physiol Renal Physiol* **291**: F1113-F1122, 2006.
- MAROTO R, RASO A, WOOD TG, KUROSKY A, MARTINAC B, HAMILL OP: TRPC1 forms the stretch-activated cation channel in vertebrate cells. *Nat Cell Biol* **7**: 179-185, 2005.
- NOSE K, SUZUKI H, KANNAN H: Voltage dependency of the frequency of slow waves in antrum smooth muscle of the guinea-pig stomach. *Jpn J Physiol* **50**: 625-633, 2000.
- PRAT AG, BERTORELLO AM, AUSIELLO DA, CANTIELLO HF: Activation of epithelial Na⁺ channels by protein kinase A requires actin filaments. *Am J Physiol* **265**: C224- C233, 1993.
- SANDERS KM: Regulation of smooth muscle excitation and contraction. *Neurogastroenterol Motil* **20 Suppl 1**: 39-53, 2008.

SANDERS KM, KOH SD, WARD SM: Interstitial cells of Cajal as pacemakers in the gastrointestinal tract.

Annu Rev Physiol **68**: 307-343, 2006.

STREGE PR, OU Y, SHA L, RICH A, GIBBONS SJ, SZURSZEWski JH, SARR MG, FARRUGIA G:

Sodium current in human intestinal interstitial cells of Cajal. *Am J Physiol Gastrointest Liver Physiol* **285**: 1111-1121, 2003.

VAN HELDEN DF, IMTIAZ MS, NURGALIYEVA K, VON DER WEID P, DOSEN PJ: Role of calcium stores and membrane voltage in the generation of slow wave action potentials in guinea-pig gastric pylorus.

J Physiol **524 Pt 1**: 245-265, 2000.

WANG ZY, YU YC, CUI YF, LI L, GUO HS, LI ZL, XU WX: Role of actin microfilament in hyposmotic membrane stretch-induced increase in muscarinic current of guinea-pig gastric myocytes. *Sheng Li Xue Bao*

55: 177-182, 2003.

^①WANG ZY, HAN YF, HUANG X, LU HL, GUO X, KIM YC, XU WX: Actin microfilament involved in regulation of pacemaking activity in cultured interstitial cells of Cajal from murine intestine. *J Membr Biol*

234: 217-225, 2010.

^②WANG ZY, HAN YF, HUANG X, ZHAO P, LU HL, KIM YC, XU WX: Pacemaking activity is regulated by membrane stretch via the CICR pathway in cultured interstitial cells of Cajal from murine intestine. *J. Biomech* **43**: 2214-2220, 2010.

Biomech **43**: 2214-2220, 2010.

WARD SM, SANDERS KM: Physiology and pathophysiology of the interstitial cells of Cajal; from the bench to bedside I. Functional development and plasticity of interstitial cells of Cajal networks. *Am J Physiol Gastrointest Liver Physiol* **281**: G602-G611, 2001.

Gastrointest Liver Physiol **281**: G602-G611, 2001.

WON KJ, SANDERS KM, WARD SM: Interstitial cells of Cajal mediate mechanosensitive responses in the stomach. *Proc Natl Acad Sci USA* **102**: 14913-14918, 2005.

WON KJ, SANDERS KM, WARD SM: Stretch-dependent sensitization of post-junctional neural effectors in colonic muscles. *Neurogastroenterol Motil* **25**: e101-e113, 2013.

YOUNG SH, ENNES HS, MAYER EA: Mechanotransduction in colonic smooth muscle cells. *J Membr Biol* **160**: 141-150, 1997.

YU YC, GUO HS, PIAO L, LI L, LI ZL, XU WX: Intracellular calcium was involved in muscarinic currents increased by hypoosmotic membrane stretch in gastric myocytes of guinea pig. *Acta Pharmacol Sin* **23**: 1007-1012, 2002.

ZHANG Y, PIAO LH, HUANG X, HAN YF, ZHAO P, GAO L, KIM Y, XU WX: Pacemaker currents modulated by C-type natriuretic peptide in interstitial cells of cajal from murine small intestine. *J Physiol Biochem* **62**: 281-291, 2006.

Figure Legends

Fig.1. Effect of hypotonic stress on spontaneous rhythmic pacemaker current (**A**) and Ca^{2+} oscillation (**B**) in cultured murine small intestine ICCs. **A**) raw traces of pacemaker current under whole cell voltage clamp mode (holding potential=-60mV) under hypotonic stress conditions (**a**); summarizes the effect of hypotonic stress-induced inward holding current (**b**) and hypotonic stress on amplitude of pacemaker current (**c**). **B**) images (**a**) and raw traces (**b**) of Ca^{2+} oscillation fluorescence intensity (F/F0) under control and hypotonic stress conditions; summarizes the effect of hypotonic stress on ICCs Ca^{2+} oscillation amplitude ($\Delta\text{F}/\text{F0}$) (**c**) and basal fluorescence intensity (F/F0) (**d**). * $P < 0.05$ vs control, n=6.

Fig.2. Role of cytochalasin-B in regulating pacemaker current and hypotonic stress-induced increase effect in cultured murine small intestine ICCs. **A**) raw traces of cytochalasin B-induced inhibition effect on pacemaker currents (a) and hypotonic stress-induced inward holding current (b). **B**) summarize the effect of cytochalasin-B on pacemaker currents amplitude (a) and frequency (b). * $P < 0.05$ vs control, n=6. **C**) summarize the effect of cytochalasin-B plus hypotonic stress conditions on pacemaker currents amplitude (a) and frequency (b). * $P < 0.05$ vs cytochalasin-B, n=6.

Fig.3. Role of cytochalasin-B in regulating Ca^{2+} oscillation and hypotonic stress-induced increase effect in cultured murine small intestine ICCs. **A**) raw traces of Ca^{2+} oscillation fluorescence intensity (F/F0) of ICCs under control, cytochalasin and cytochalasin plus hypotonic stress conditions respectively; **B**) summarizes the effect of cytochalasin-B on Ca^{2+} oscillation amplitude ($\Delta\text{F}/\text{F0}$) (a) and basal fluorescence intensity (F/F0) (b) under normal and hypotonic stress condition. * $P < 0.05$ vs control, n=6.

Fig.4. Role of phalloidin in regulating pacemaker current and hypotonic stress-induced increase effect in cultured murine small intestine ICCs. **A)** raw traces of phalloidin-induced potentiation effect on pacemaker currents (**a**) and hypotonic stress-induced inward holding current (**b**). **B)** summarize the effect of phalloidin on pacemaker currents amplitude (**a**) and frequency (**b**). * $P < 0.05$ vs control, $n=6$. **C)** summarize the effect of phalloidin plus hypotonic stress conditions on pacemaker currents amplitude (**a**) and frequency (**b**). * $P < 0.05$ vs phalloidin, $n=6$.

Fig.5. Role of phalloidin in regulating Ca^{2+} oscillation and hypotonic stress-induced increase effect in cultured murine small intestine ICCs. **A)** raw traces of Ca^{2+} oscillation fluorescence intensity (F/F_0) of ICCs under control, phalloidin and phalloidin plus hypotonic stress conditions respectively; **B)** summarizes the effect of phalloidin on Ca^{2+} oscillation amplitude ($\Delta F/F_0$) (**a**) and basal fluorescence intensity (F/F_0) (**b**) under normal and hypotonic stress condition. * $P < 0.05$ vs control, $n=6$.

Fig.1




Fig.2




Fig.3




Fig.4




Fig.5

