



2012

ISSN 2226-0773

МЕЖДУНАРОДНЫЙ АЛЬМАНАХ  
INTERNATIONAL ALMANAC

<http://www.humanityspace.com>



Том 1 No 2 Volume 1 No 2

ГУМАНИТАРНОЕ ПРОСТРАНСТВО  
HUMANITY SPACE

Том 1 No 2  
Volume 1 No 2  
2012



MOSCOW

МОСКВА  
MOSCOW

**Contribution to the knowledge of longhorn beetles from Dhofar region in sultanate of Oman (Coleoptera: Cerambycidae)**

**Richard Ambrus<sup>1</sup>, Walter Grosser<sup>2</sup>**

<sup>1</sup> Trnkovo nám. 1112/1, CZ-152 00 Prague 5, Czech Republic  
e-mail: ambrus@centrum.cz

<sup>2</sup> U Střelnice 17, CZ-746 01 Opava, Czech Republic; e-mail: walterg@atlas.cz

**Key words:** Coleoptera, Cerambycidae, faunistics, new records, Dhofar, Sultanate of Oman, Palaearctic region.

**Abstract:** Totally 16 Cerambycidae species are reported for Dhofar in Sultanate of Oman. Ten species are reported for the first time for Oman: *Lygrus longicornis* (Pic, 1895); *Kabatekiella dalihodi* Holzschuh, 2008; *Yementallyrama nasheri* Adlbauer, 2007; *Idactus coquereli* (Fairmaire, 1890); *Apomecyna lameerei* (Pic, 1895); *Enaretta caudata* Fähræus, 1873; *Eunidia breuningae* Villiers, 1951; *Eunidia kristenseni* Aurivillius, 1911; *Eunidia nebulosa nebulosa* Erichson, 1843 and *Coptops aedificator* Fabricius, 1792. The genus *Jordanoleiopus* Lepesme & Breuning, 1955 is new recorder for the fauna of Palaearctic Region.

## INTRODUCTION

The cerambycid fauna of Sultanate Oman is very little known, since most of the species hitherto recorded from the Arabian Peninsula, have been collected in Saudi Arabia and Yemen.

The paper presents results of the Czech entomological expedition to Sultanate of Oman in September/October 2011. During this expedition were visited, with a special attention to interesting biotopes in Dhofar region, especially Jabal Samhan and Jabal al Qamar. The expedition included four Czech entomologists, Richard Ambrus, Walter Grosser, Pavel Kučera and Jiří Voříšek.

The beetles were collected individually using generally known methods such as beating, sweeping, individual collecting on wood, flower and leaves. The most effective method for collecting longhorn beetles were used UV light.

The total number of longhorn beetles recorded by Czech expedition from Sultanate of Oman is 16 species, but this is probably only a small number of species actually occurring there.

## Richard Ambrus, Walter Grosser

All the species of longhorn beetles were determined by Dr. Karl Adlbauer (Graz, Austria), Petr Kabátek (Prague, Czech Republic) and Pierpaolo Rapuzzi (Prepotto, Italy).

Abbreviations:

JV - Jiří Voříšek, Jirkov, CZ

PK - Pavel Kučera, Liberec, CZ

RA - Richard Ambrus, Prague, CZ

WG - Walter Grosser, Opava, CZ

## RESULTS

### Family CERAMBYCIDAE

#### Subfamily Prioninae

#### Tribe Cantharocnemi

### 1) *Cantharocnemis spondyloides* Audinet-Serville, 1832

**Material.** Oman, Dhofar, Jabal al Qamar, 20 km NW Dhalqut, 16°42'39.31"N 53°9'12.60"E, 21. 9. 2011, 1 adult attracted by light and remnants of 1 dead adult on ground (PK, RA); Oman, Dhofar, Jabal al Qamar, 5 km NE Dhalqut, 16°43'22.48"N 53°16'27.26"E, 22. 9. 2011, 1 adult attracted by light (WG); Oman, Dhofar, Jabal al Qamar, 15 km NW Rakhyut, 16°46'7.54"N 53°20'13.92"E, 24. 9. 2011, 1 adult attracted by light (WG); Oman, Dhofar, Jabal Samhan, 15 km NW Jufa, 17°11'10.14"N 54°56'34.26"E, 27. 9. 2011, 1 adult attracted by light and remnants of 1 dead adult on ground (PK, WG); Oman, Dhofar, Jabal Samhan, 15 km SE Tawi Atayr, 17°3'57.31"N 54°36'17.24"E, 28. 9. 2011, 1 adult attracted by light (PK).

**Distribution.** Afrotropical Region; Palaearctic Region: North Africa (Egypt, Marocco- Western Sahara), Asia (Oman, Saudi Arabia, Yemen).

**Richard Ambrus, Walter Grosser**

**Subfamily Cerambycinae  
Tribe Cerambycini**

**2) *Neoplocaederus denticornis* (Fabricius, 1801)**

**Material.** Oman, Dhofar, Jabal al Qamar, 20 km W Al Mughsayl, 16°50'41.89"N 53°41'10.14"E, 19. 9. 2011, 4 adults attracted by light (PK, RA); Oman, Dhofar, Jabal al Qamar, 10 km W Dhalqut, 16°42'9.90"N 53°11'40.56"E, 20. 9. 2011, 4 adults attracted by light (PK, RA); Oman, Dhofar, Jabal al Qamar, 20 km NW Dhalqut, 16°42'39.31"N 53°9'12.60"E, 21. 9. 2011, 15 adults attracted by light (PK, RA, WG); Oman, Dhofar, Jabal al Qamar, 5 km NE Dhalqut, 16°43'22.48"N 53°16'27.26"E, 22. 9. 2011, 12 adults attracted by light (JV, PK, RA); Oman, Dhofar, Jabal al Qamar, Rakhyut, 16°45'12.53"N 53°23'54.38"E, 23. 9. 2011, 7 adults attracted by light (PK, WG); Oman, Dhofar, Jabal al Qamar, 15 km NW Rakhyut, 16°46'7.54"N 53°20'13.92"E, 24. 9. 2011, 7 adults attracted by light (PK, WG); Oman, Dhofar, Jabal al Qamar, 15 km W Al Mughsayl, 16°51'44.56"N 53°43'18.38"E, 26. 9. 2011, 9 adults attracted by light (PK, WG); Oman, Dhofar, Jabal Samhan, 15 km NW Jufa, 17°11'10.14"N 54°56'34.26"E, 27. 9. 2011, 33 adults attracted by light (JV, PK, RA, WG); Oman, Dhofar, Jabal Samhan, Tawi Atayr env., 17°5'16.94"N 54°37'23.27"E, 1. 10. 2011, 6 adults attracted by light (JV, PK).

**Distribution.** Palaearctic Region: Asia (Oman, Saudi Arabia, Yemen).

**Tribe Lygrini**

**3) *Lygrus longicornis* (Pic, 1895)**

**Material.** Oman, Dhofar, Jabal al Qamar, 10 km W Dhalqut, 16°42'9.90"N 53°11'40.56"E, 20. 9. 2011, 2 adults attracted by light (PK, RA); Oman, Dhofar, Jabal al Qamar, 20 km NW Dhalqut, 16°42'39.31"N 53°9'12.60"E, 21. 9. 2011, 7 adults attracted by light (PK, RA, WG); Oman, Dhofar, Jabal al Qamar, 15 km NW Rakhyut,

**Richard Ambrus, Walter Grosser**

16°46'7.54"N 53°20'13.92"E, 24. 9. 2011, 2 adults attracted by light (PK, WG).

This species is new for the fauna of Oman.

**Distribution.** Palaearctic Region: Asia (Oman, Saudi Arabia, Yemen).

**Tribe Oemini**

**4) *Kabatekiella dalihodi* Holzschuh, 2008**

**Material.** Oman, Dhofar, Jabal al Qamar, 20 km NW Dhalqut, 16°42'39.31"N 53°9'12.60"E, 21. 9. 2011, 8 adults attracted by light (PK, RA); Oman, Dhofar, Jabal al Qamar, 5 km NE Dhalqut, 16°43'22.48"N 53°16'27.26"E, 22. 9. 2011, 12 adults attracted by light (PK, RA, WG); Oman, Dhofar, Jabal al Qamar, Rakhyut, 16°45'12.53"N 53°23'54.38"E, 23. 9. 2011, 9 adults attracted by light (PK, WG); Oman, Dhofar, Jabal al Qamar, 15 km NW Rakhyut, 16°46'7.54"N 53°20'13.92"E, 24. 9. 2011, 68 adults attracted by light (PK, WG); Oman, Dhofar, Jabal Samhan, Tawi Atayr env., 17°5'16.94"N 54°37'23.27"E, 1. 10. 2011, 12 adults attracted by light (JV).

This species is new for the fauna of Oman.

**Distribution.** Palaearctic Region: Asia (Oman, Yemen).

**5) *Yementallyrama nasheri* Adlbauer, 2007**

**Material.** Oman, Dhofar, Jabal al Qamar, 20 km W Al Mughsayl, 16°50'41.89"N 53°41'10.14"E, 19. 9. 2011, 3 adults attracted by light (PK, RA); Oman, Dhofar, Jabal al Qamar, 10 km W Dhalqut, 16°42'9.90"N 53°11'40.56"E, 20. 9. 2011, 6 adults attracted by light (PK, RA); Oman, Dhofar, Jabal al Qamar, 20 km NW Dhalqut, 16°42'39.31"N 53°9'12.60"E, 21. 9. 2011, 29 adults attracted by light (PK, RA, WG); Oman, Dhofar, Jabal al Qamar, 5 km NE Dhalqut, 16°43'22.48"N 53°16'27.26"E, 22. 9. 2011, 9 adults

**Richard Ambrus, Walter Grosser**

attracted by light (JV, PK); Oman, Dhofar, Jabal al Qamar, Rakhyut, 16°45'12.53"N 53°23'54.38"E, 23. 9. 2011, 4 adults attracted by light (JV, PK); Oman, Dhofar, Jabal al Qamar, 15 km NW Rakhyut, 16°46'7.54"N 53°20'13.92"E, 24. 9. 2011, 6 adults attracted by light (WG); Oman, Dhofar, Jabal Samhan, 15 km NW Jufa, 17°11'10.14"N 54°56'34.26"E, 27. 9. 2011, 9 adults attracted by light (PK, WG).

This species is new for the fauna of Oman.

**Distribution.** Palaearctic Region: Asia (Oman, Yemen).

**Subfamily Lamiinae**  
**Tribe Acanthocinini**

**6) *Jordanoleiopus* sp.**

**Material.** Oman, Dhofar, Jabal al Qamar, 10 km W Dhalqut, 16°42'9.90"N 53°11'40.56"E, 20. 9. 2011, 9 adults attracted by light (PK, RA, WG); Oman, Dhofar, Jabal al Qamar, 20 km NW Dhalqut, 16°42'39.31"N 53°9'12.60"E, 21. 9. 2011, 3 adults attracted by light (PK, RA); Oman, Dhofar, Jabal al Qamar, 5 km NE Dhalqut, 16°43'22.48"N 53°16'27.26"E, 22. 9. 2011, 165 adults attracted by light (JV, PK, RA, WG); Oman, Dhofar, Jabal al Qamar, 15 km NW Rakhyut, 16°46'7.54"N 53°20'13.92"E, 24. 9. 2011, 26 adults attracted by light (PK, WG); Oman, Dhofar, Jabal al Qamar, 15 km W Al Mughsayl, 16°51'44.56"N 53°43'18.38"E, 26. 9. 2011, 14 adults attracted by light (PK, WG); Oman, Dhofar, Jabal Samhan, Tawi Atayr env., 1. 10. 2011, 17°5'16.94"N 54°37'23.27"E, 5 adults attracted by light (JV, PK).

We were not able to determine this species, but according to Dr. Karl Adlbauer it belongs close to *Jordanoleiopus monoxenus* (Kolbe, 1894) from Afrotropical Region (Tanzania).

The genus *Jordanoleiopus* Lepesme & Breuning, 1955 is new for the fauna of Palaearctic Region.

**Richard Ambrus, Walter Gresser**

**Distribution.** Afrotropical Region; Palaearctic Region: Asia (Oman).

**Tribe Ancytonotini**

**7) *Idactus coquereli* (Fairmaire, 1890)**

**Material.** Oman, Dhofar, Jabal al Qamar, 20 km W Al Mughsayl, 16°50'41.89"N 53°41'10.14"E, 19. 9. 2011, 3 adults attracted by light (PK, RA); Oman, Dhofar, Jabal al Qamar, 10 km W Dhalqut, 16°42'9.90"N 53°11'40.56"E, 20. 9. 2011, 7 adults attracted by light (JV, PK, RA); Oman, Dhofar, Jabal al Qamar, 20 km NW Dhalqut, 16°42'39.31"N 53°9'12.60"E, 21. 9. 2011, 9 adults attracted by light (PK, RA, WG); Oman, Dhofar, Jabal al Qamar, 5 km NE Dhalqut, 16°43'22.48"N 53°16'27.26"E, 22. 9. 2011, 6 adults attracted by light (PK, RA); Oman, Dhofar, Jabal al Qamar, 15 km NW Rakhyyut, 16°46'7.54"N 53°20'13.92"E, 24. 9. 2011, 8 adults attracted by light (PK, WG); Oman, Dhofar, Jabal Samhan, 15 km NW Jufa, 17°11'10.14"N 54°56'34.26"E, 27. 9. 2011, 7 adults attracted by light (JV, PK, WG).

This species is new for the fauna of Oman.

**Distribution:** Afrotropical Region, Palaearctic Region: Asia (Oman, Yemen).

**8) *Idactus iranicus* Breuning, 1975**

**Material.** Oman, Dhofar, Jabal Samhan, 15 km NW Jufa, 17°11'10.14"N 54°56'34.26"E, 27. 9. 2011, 4 adults attracted by light (JV, PK, WG).

**Distribution:** Palaearctic Region: Asia (Arab Emirate, Iran, Oman).

**Richard Ambrus, Walter Grosser**

**Tribe Apomecynini**

**9) *Apomecyna lameerei* (Pic, 1895)**

**Material.** Oman, Dhofar, Jabal al Qamar, 5 km NE Dhalqut, 16°43'22.48"N 53°16'27.26"E, 22. 9. 2011, 1 adult attracted by light (JV).

This species is new for the fauna of Oman.

**Distribution.** Afrotropical Region; Palaearctic Region: North Africa (Egypt, Morocco), Asia (Arab Emirate, Egypt-Sinai, Iran, Iraq, Israel, Oman, Pakistan, Saudi Arabia, Yemen).

**10) *Enaretta caudata* Fahraeus, 1873**

**Material.** Oman, Dhofar, Jabal al Qamar, 10 km W Dhalqut, 16°42'9.90"N 53°11'40.56"E, 20. 9. 2011, 4 adults attracted by light (PK, RA); Oman, Dhofar, Jabal al Qamar, 5 km NE Dhalqut, 16°43'22.48"N 53°16'27.26"E, 22. 9. 2011, 11 adults attracted by light (JV, PK, RA, WG); Oman, Dhofar, Jabal al Qamar, 15 km NW Rakhyut, 16°46'7.54"N 53°20'13.92"E, 24. 9. 2011, 4 adults attracted by light (PK, WG); Oman, Dhofar, Jabal al Qamar, 15 km W Al Mughsayl, 16°51'44.56"N 53°43'18.38"E, 26. 9. 2011, 3 adults attracted by light (PK, WG); Oman, Dhofar, Jabal Samhan, 15 km NW Jufa, 17°11'10.14"N 54°56'34.26"E, 27. 9. 2011, 5 adults attracted by light (PK, WG).

This species is new for the fauna of Oman.

**Distribution.** Afrotropical Region; Palaearctic Region: Asia: Oman, Yemen.

**11) *Eunidia breuningae* Villiers, 1951**

**Material.** Oman, Dhofar, Jabal al Qamar, 20 km W Al Mughsayl, 16°50'41.89"N 53°41'10.14"E, 19. 9. 2011, 5 adults attracted by



**Richard Ambrus, Walter Grosser**

light (PK, RA); Oman, Dhofar, Jabal al Qamar, 20 km NW Dhalqut, 16°42'39.31"N 53°9'12.60"E, 21. 9. 2011, 11 adults attracted by light (PK, RA, WG); Oman, Dhofar, Jabal al Qamar, 5 km NE Dhalqut, 16°43'22.48"N 53°16'27.26"E, 22. 9. 2011, 6 adults attracted by light (JV, PK, RA); Oman, Dhofar, Jabal Samhan, 15 km NW Jufa, 17°11'10.14"N 54°56'34.26"E, 27. 9. 2011, 5 adults attracted by light (PK, WG).

This species is new for the fauna of Oman.

**Distribution.** Afrotropical Region; Palaearctic Region: North Africa (Egypt), Asia (Oman, Saudi Arabia, Yemen).

**12) *Eunidia kristenseni* Aurivillius, 1911**

**Material.** Oman, Dhofar, Jabal al Qamar, 10 km W Dhalqut, 16°42'9.90"N 53°11'40.56"E, 20. 9. 2011, 1 adult attracted by light (RA).

This species is new for the fauna of Oman.

**Distribution:** Afrotropical Region, Palaearctic Region: Asia (Oman, Saudi Arabia, Yemen).

**13) *Eunidia nebulosa nebulosa* Erichson, 1843**

**Material.** Oman, Dhofar, Jabal al Qamar, 20 km W Al Mughsayl, 16°50'41.89"N 53°41'10.14"E, 19. 9. 2011, 3 adults attracted by light (PK, RA); Oman, Dhofar, Jabal al Qamar, 20 km NW Dhalqut, 16°42'39.31"N 53°9'12.60"E, 21. 9. 2011, 4 adults attracted by light (PK, RA); Oman, Dhofar, Jabal al Qamar, 5 km NE Dhalqut, 16°43'22.48"N 53°16'27.26"E, 22. 9. 2011, 7 adults attracted by light (JV, PK, RA); Oman, Dhofar, Jabal Samhan, 15 km NW Jufa, 17°11'10.14"N 54°56'34.26"E, 27. 9. 2011, 10 adults attracted by light (JV, PK, WG).

This species is new for the fauna of Oman.

**Richard Ambrus, Walter Grosser**

**Distribution.** Afrotropical Region; Palaearctic Region: Asia (Oman, Saudi Arabia, Yemen).

**Tribe Batocerini**

**14) *Batocera rufomaculata rufomaculata* (DeGeer, 1775)**

**Material.** Oman, Dhofar, Jabal al Qamar, 15 km NW Rakhyut, 16°46'7.54"N 53°20'13.92"E, 24. 9. 2011, 3 adults sitting on *Ficus* sp. (PK, WG); Oman, Dhofar, Jabal Samhan, 15 km SE Tawi Atayr, 28. 9. 2011, 17°3'57.31"N 54°36'17.24"E, 2 adults sitting on *Ficus* sp. (PK).

**Distribution.** Afrotropical Region; Neotropical Region; Oriental Region; Palaearctic Region:

Asia (Egypt-Sinai, China-Hongkong and Xizang, India-Uttar Pradesh, Israel, Lebanon, Nepal, Oman, Pakistan, Syria, Turkey, Yemen).

**Tribe Crossotina**

**15) *Crossotus arabicus* Gahan, 1896**

**Material.** Oman, Dhofar, Jabal al Qamar, 20 km NW Dhalqut, 16°42'39.31"N 53°9'12.60"E, 21. 9. 2011, 3 adults reared ex larva from death branches of *Acacia* sp., 1 dead adult was found in a dead branch of *Acacia* sp. and 3 adults attracted by light (PK, RA, WG).

**Distribution.** Palaearctic Region: Asia (Oman, Saudi Arabia, Yemen).

**Tribe Mesosini**

**16) *Coptops aedificator* Fabricius, 1792**

**Material.** Oman, Dhofar, Jabal al Qamar, 10 km W Dhalqut, 16°42'9.90"N 53°11'40.56"E, 20. 9. 2011, 7 adults attracted by light (PK, RA); Oman, Dhofar, Jabal al Qamar, 15 km NW Rakhyut, 16°46'7.54"N 53°20'13.92"E, 24. 9. 2011, 5 adults attracted by light

**Richard Ambrus, Walter Grosser**

and 2 adults reared ex larva from dead branches and trunks of *Euphorbia* sp. (PK, RA, WG).

This species is new for the fauna of Oman.

**Distribution.** Afrotropical Region; Oriental Region; Palaearctic Region: Asia (Oman, Saudi Arabia, Taiwan, Yemen).

**Acknowledgement.** We are very grateful to Dr. Karl Adlbauer (Graz, Austria), Petr Kabátek (Prague, Czech Republic) and Pierpaolo Rapuzzi (Prepetto, Italy) for determination and comments on the collected species.

**REFERENCES**

- Adlbauer K., 2005. Neue Bockkäfer von Yemen (Coleoptera, Cerambycidae). – Linzer biol. Beitr., 37(2): 1007-1011.
- Adlbauer K., 2010. Cerambycidae, Western Palaearctic taxa: Arabian Peninsula, pp. 84-334. In: Löbl I. & Smetana A. (eds.): Catalogue of Palaearctic Coleoptera. 6. Chrysomeloidea. Apollo Books, Stenstrup: 924 pp.
- Holzschuh C., 1979. Insects of Saudi Arabia. Coleoptera: fam. Cerambycidae. Fauna of Saudi Arabia, 1: 293-294.
- Holzschuh C., 1993. Cerambycidae (Coleoptera) of Saudi Arabia: Part II, Prioninae and Cerambycinae. Fauna of Saudi Arabia, Basel 13: 110-129.
- Holzschuh C. & Teocchi P., 1991. Cerambycidae of Saudi Arabia: part I. Lamiinae. Fauna of Saudi Arabia, 12: 295-311.
- Janikova M. Coleoptera. Beetles Checklist [of Sultanate of Oman]. Sultanate of Oman, Ministry of Heritage & Culture. 39 pp.
- Sama G., 2000. Descrizione di un nuovo *Crossotus* Serville, 1835 del Medio Oriente. Quad. Studi Nat. Romagna, 13 (suppl.): 107-112.
- Sama G., Rapuzzi P., 2006. Preliminary report on a recent survey of the Egyptian Cerambycidae, with description of three new species. Quad. Studi Nat. Romagna, 23: 179-194
- Sama G., 2007. Description of two new genera and one new species of Cerambycidae from the Arabian peninsula (Insecta Coleoptera Cerambycidae). Quad. Studi Nat. Romagna, 24: 97-105.
- Sama G., Sudre J., 2010. Description of *Crossotus kadleci* from Yemen (Coleoptera, Cerambycidae, Lamiinae). Munis Entomology & Zoology, Vol. 5, Suppl.: 953-956.
- Strumia F., Dapporto L., Dellacasa M., Scaramozzino P.L., 2007. Notes on some Insects associated to Frankincense Tree (*Boswellia sacra* Flückiger, 1867, Burseraceae) in Dhofar (Sultanate of Oman). Atti Soc. tosc. Sci. nat., Mem., Serie B, 114: 135-139.

*Получена / Received: 27.03.2012*

*Принята / Accepted: 30.03.2012*