

1 **Time and dose-dependent effects of *Viscum album quercus* on rabbit** 2 **spermatozoa motility and viability *in vitro***

3
4 Marko HALO Jr.¹, Peter MASSANYI^{1,2}, Agnieszka GREN², Agnieszka LASAK²,
5 Tomas SLANINA¹, Lubomir ONDRUSKA³, Renata MUCHACKA², Drahomir
6 GALBAVY⁴, Peter IVANIC⁵, Eric Rendon SCHNEIR⁶, Grzegorz FORMICKI²

7
8 ¹Department of Animal Physiology, Faculty of Biotechnology and Food Sciences,
9 Slovak University of Agriculture, Trieda Andreja Hlinku 2, 949 76 Nitra, Slovak
10 Republic

11 ²Department of Animal Physiology and Toxicology, Institute of Biology, Faculty of
12 Geography and Biology, Pedagogical University of Cracow, ul. Podchorazych 2, 30-
13 084 Cracow, Poland

14 ³Institute of Farm Animals, Animal Production Research Centre Nitra, Hlohovecka 2,
15 951 41 Luzianky, Slovak Republic

16 ⁴Avelane Clinic, Krcmeryho 837/2, 949 01 Nitra, Slovak Republic

17 ⁵Slovak Biological Services, Kremnicka 2, 974 05 Banska Bystrica, Slovak Republic

18 ⁶Faculty of Planning, The National University Agraria La Molina, Av. La Molina s/n.
19 Lima 12, Peru

20

21 *Corresponding author: Marko Halo, Department of Animal Physiology, Faculty of
22 Biotechnology and Food Sciences, Slovak University of Agriculture, Trieda Andreja
23 Hlinku 2, 949 76 Nitra, Slovak Republic, Tel. +421 908 441 517, E-mail address:

24 markohalo@yahoo.com

25

26 Short title: *Viscum album quercus* affects spermatozoa motility

27

28 **Summary**

29

30 The target of this study was to evaluate the effect of extract of the European mistletoe –
31 *Viscum album quercus* L. on spermatozoa motility and viability *in vitro*. The CASA
32 system was used to determine the spermatozoa motility parameters at different time
33 intervals (0; 1; 2 and 3 h) and spermatozoa viability was determined in five different
34 doses of *Viscum album quercus* L [10 (QA); 6.6 (QB); 3.3 (QC); 2.5 (QD) and 2 (QE)
35 mg/mL]. Results in experimental groups detected a significant deterioration on rabbit
36 spermatozoa after 1, 2 and 3 hours, compared to the control. The initial total
37 spermatozoa motility showed increased value for all doses of *Viscum album quercus* in
38 comparison to control. After *in vitro* culture a dose–dependent decrease (QA: reduction
39 of 69.7%; QB: reduction of 40.9%) was found. For the progressive spermatozoa most
40 significant decrease (86.8% for QA vs. 48.5% for QB) was detected compared to the
41 control after 3 hours of culture. Spermatozoa viability (MTT test) was decreased in all
42 experiment groups at the end of experiment, but the differences were not significant.
43 Significant alterations of membrane integrity were found in groups with the highest
44 *Viscum album quercus* concentration (QA, QB), but acrosome integrity showed no
45 significant changes. Results suggest negative dose– and time–dependent effect of
46 *Viscum album quercus* at higher doses on spermatozoa motility and viability parameters
47 *in vitro*.

48

49 **Key words:** *Viscum album*; spermatozoa; CASA; viability; *in vitro*

50

51 **Introduction**

52

53 Anticancer preparations made from plants have been an object of scientific interest for
54 many years. It is worth noting that as many as 25% of cytostatics used in the anticancer
55 chemotherapy are obtained from plants.

56 Extracts of the European mistletoe (*Viscum album* L.) have been widely used for
57 decades as alternative, complementary treatment (Kovacs *et al.* 2006, Felenda *et al.*
58 2019, Suveren *et al.* 2017). In clinical practice mistletoe therapy is often given
59 concomitantly to conventional chemotherapy. Mistletoe plants are generally growing on
60 different host trees, like apple, oak, or pine. Cytotoxic glycoproteins, the mistletoe
61 lectins, are active component of mistletoe extracts and can stimulate effector cells of the
62 innate and adaptive immune system (Stein *et al.* 2002, Braedel-Ruoff 2010, Gren and
63 Formicki 2013, Gren and Massanyi 2016).

64 Experiments also indicate a statistically significant increase in albumin fraction level
65 and lymphocyte count. Moreover, decrease of the total protein content, protein fractions
66 globulins alpha2, beta, gamma and neutrophil, monocyte count in mouse serum was
67 observed (Gren 2009).

68 The reproductive ability and the semen quality of animal species can be affected by
69 many environmental sources, as well as age, stress, hormonal status, nutrition and
70 toxins (Mangelsdorf *et al.* 2003, Lukac *et al.* 2011, Mousa-Balabel and Mohamed 2011,
71 Fallas-López *et al.* 2011, Tirpak *et al.* 2017, Saha *et al.* 2019). In the many years,
72 following the increased success rate of cancer treatments, great efforts have been made
73 to improve quality of life in survivors, including fertility preservation in young patients
74 (Masopotova *et al.* 2018). Because of their gonadotoxic effects, chemo- and
75 radiotherapy can temporarily or permanently compromise fertility (Di Bisceglie *et al.*
76 2013). Oncological treatments present severe gonadotoxic effects on both germ and
77 Leydig cells. Of note, in a significant percentage of patients (20 – 50%)
78 spermatogenesis is impaired even before cancer treatments, probably due to the
79 malignancy itself. The recovery of normal spermatogenesis after treatment may require
80 several years, and mainly depends on various factors – initial spermatozoa count, type
81 and dose of specific oncological treatments and patient age. Disturbance of homeostasis
82 of reproductive system in some diseases can be seen from the decline in physical and
83 chemical parameters of spermatozoa, such as pH, semen volume, concentration,
84 motility, and the percentage of spermatozoa viability. These data justify the increasing
85 efforts in identifying prevention and treatment strategies to preserve reproductive
86 functions in young men with malignancies (Colpi *et al.* 2004, Maltaris *et al.* 2006;
87 Vitku *et al.* 2015; Heráček *et al.* 2018). The mechanisms underlying the male infertility
88 of *Viscum album* extracts have not been investigated.

89 The objective of this *in vitro* study was to determine the effect of various concentrations
90 of *Viscum album quercus* during various time periods (0 – 3 h) on the selected
91 parameters of rabbit spermatozoa motility and viability.

92

93

94

95

96 **Materials and methods**

97

98 **Animals, semen samples and *in vitro* culture**

99

100 Male rabbits (n=10; New Zealand White) kept under standard conditions at the
101 Experimental Station of the Animal Production Research Centre Nitra, Slovak Republic
102 were selected on the basis of age normally associated with reproduction (12 – 14
103 months). Animals were housed in a partially air-conditioned rabbit house (Animal
104 Production Research Centre, Nitra) under a photoperiod 16L : 8D (minimum light
105 intensity of 80 lux). Animals were kept in individual cages and fed with a commercial
106 diet and were provided water *ad libitum*. An air temperature of $20 \pm 2^\circ\text{C}$ and relative
107 humidity of $70 \pm 5\%$ was maintained in the rabbit house. Conditions of their care,
108 manipulations and use corresponded to the instruction of EC no. 178/2002 and related
109 EC documents, and they were approved by local ethics committee.

110 Semen samples (n = 5) in five replicates were collected on a single day (early in the
111 morning) with the help of artificial vagina (Krockova *et al.* 2012, Parkanyi *et al.* 2015).
112 Immediately after collection the individual doses of semen exhibiting a white color
113 without presence of any gel and artificial particles, were mixed together to obtain
114 pooled sample. The spermatozoa concentration in semen was $0.40 - 0.63 \times 10^9$ per mL.
115 The obtained semen samples were diluted according to routine methods (Chrenek *et al.*
116 2007, Roychoudhury and Massanyi 2008).

117 Later the spermatozoa were incubated in thermostat ($37 \pm 0.5^\circ\text{C}$) with various
118 concentrations of *Viscum album quercus* (Iscador Qu 10 mg; Weleda, Verein für
119 Krebsforschung Institute Hiscia – Arlesheim, Switzerland) dissolved in physiological
120 solution. **Concentrations of the Iscador Qu were based on manufacturer's values and
121 later diluted with physiological solution to reach decreased experimental concentration
122 (6.6 – 2.0 mg).** The control (QK) group was cultured only with physiological solution.
123 The scheme of experiments is presented in Table 1.

124

125 **2.2. Computer-assisted semen analysis**

126

127 Spermatozoa motility was used as an indicator of cell quality. The motility analysis was
128 carried out using a CASA (Computer Assisted Semen Analysis) system – Sperm Vision
129 TM program (MiniTub, Tiefenbach, Germany) with the Olympus BX 51 microscope
130 (Olympus, Tokyo, Japan) at cultivation times 0, 1, 2 and 3 hours (Time 0 – 3). Each
131 sample was placed into the Makler Counting Chamber (depth 10 μm , Sefi-Medical
132 Instruments, Haifa, Israel (Massanyi *et al.* 2008). This study was performed in five
133 replicates at each concentration. At least 1000 spermatozoa were analyzed in each
134 sample (Lukac *et al.* 2013, Halo *et al.* 2018). Using the rabbit specific set up the
135 following parameters have been evaluated – total motility (MOT; %), progressive
136 motility (PRO; %), distance average path (DAP; μm), distance curved line (DCL; μm),
137 distance straight line (DSL; μm), average path velocity (VAP; $\mu\text{m/s}$), velocity curved
138 line (VCL; $\mu\text{m/s}$), velocity straight line (VSL; $\mu\text{m/s}$), straightness (STR), linearity
139 (LIN), wobble (WOB), amplitude of lateral head displacement (ALH; μm) and beat-
140 cross frequency (BCF; Hz) as described previously (Roychoudhury *et al.* 2010, Tvrdá *et al.*
141 2015 Adamkovicova *et al.* 2016).

142

143

144 **Viability analysis – MTT test**

145

146 Viability of rabbit spermatozoa cultured with *Viscum album quercus* was evaluated by
147 the metabolic activity (MTT) assay after 3 hours of culture. This colorimetric assay
148 measures the conversion of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium
149 bromide (MTT; Sigma-Aldrich, St. Louis, USA) to purple formazan particles by
150 mitochondrial succinate dehydrogenase of intact mitochondria of living cells. Formazan
151 was measured spectrophotometrically by a microplate ELISA reader (Multiskan FC,
152 ThermoFisher Scientific, Finland). The data are expressed in percentage of control.
153 Results from the analysis were collected during four repeated experiments for each
154 concentration (Slanina *et al.* 2016).

155

156 **Viability – membrane integrity – Eosin-nigrosin**

157

158 The spermatozoa viability was evaluated using eosin-nigrosin staining methods (Slanina
159 *et al.* 2018). From all the samples smears were prepared after 3 hours of culture.
160 Experimental samples and the control sample were diluted in the ratio 1 : 2 : 2 with 5%
161 eosin (Eosin Y) and 10% nigrosin (Nigrosin) solution (both Sigma-Aldrich, St. Louis,
162 USA). For each slide 300 cells were counted under a light microscope (1000×, Leica
163 DMIL LED; Leica Microsystems CMS GmbH, Germany) and classified as viable
164 (intact membrane) and dead (damaged membrane). The experiment was realized in six
165 replicates. The results of viability evaluation were expressed as the percentage of viable
166 and dead spermatozoa (in %).

167

168 **Acrosomal integrity**

169

170 The acrosomal status was assessed after 3 hours of culture following the fast green-rose
171 Bengal staining protocol designed by Pope *et al.* (1991). This single-step staining
172 method applies a mixture consisting of 1% fast green (Sigma-Aldrich, St. Louis, USA),
173 1% rose bengal (Sigma-Aldrich, St. Louis, USA) and 40% ethyl alcohol (Centralchem,
174 Bratislava, Slovak Republic) in 0.1 M citric acid – 0.2 M disodium phosphate buffer
175 (Sigma-Aldrich, St. Louis, USA). Twenty microliters of the sample were mixed with 20
176 µl of the staining solution and incubated for 70 s at room temperature. Ten microliters
177 of the mixture were smeared on a tempered glass slide and air-dried at 37°C. Acrosomal
178 integrity was evaluated using bright field microscopy at 1000x using oil immersion. At
179 least 200 cells per slide were evaluated for the presence or absence of acrosome, and
180 expressed as a percentage rate (Tvrda *et al.* 2017).

181

182 **Statistical analysis**

183

184 The control group (medium without *Viscum album quercus*) was compared to the
185 experimental groups. Statistical analysis was carried out using the GraphPad Prism
186 program (version 3.02 for Windows; GraphPad Software, La Jolla California USA).
187 Descriptive statistical characteristics (mean, standard deviation) were evaluated at first.
188 One-way ANOVA with Dunnett's post-test was used for statistical evaluations. The
189 level of significance was set at *** ($p < 0.001$), ** ($p < 0.01$) and * ($p < 0.05$). For
190 individual measurements average value (x), minimum (min) and maximum (max) value,
191 standard deviation (SD) and coefficient of variation (CV) were recorded.

192 **Results**

193

194 **Spermatozoa motility**

195

196 The initial spermatozoa motility (Time 0) showed increased value for all doses of
197 *Viscum album quercus* in comparison to the control group. Statistically significant
198 increase was observed ($p < 0.05$) in the sample QA ($77.23 \pm 8.64\%$), and also ($p <$
199 0.001) for QB ($81.17 \pm 8.08\%$) and QC ($80.87 \pm 8.83\%$). After 1 h of the culture the
200 average spermatozoa motility in control group was $72.17 \pm 11.94\%$. At the same time a
201 statistically significant ($p < 0.001$) decrease (23.8%) was observed in group QA (54.98
202 $\pm 15.53\%$). After 2 hours of *in vitro* cultivation significantly decreased ($p < 0.001$ and
203 0.01) total spermatozoa motility in comparison to the control group was observed in
204 groups QA and QB (doses 10 and 6.6 mg/mL). Also, after 3 hours of *in vitro* cultivation
205 a dose-dependent decrease (QA: $19.39 \pm 7.7\%$; reduction of 69.7%; QB: $37.84 \pm$
206 18.29% ; reduction of 40.9%) was found (Figure 1).

207

208 **Progressive spermatozoa motility**

209

210 At Time 0 the average progressive spermatozoa motility was higher in all experimental
211 groups with *Viscum album quercus* in comparison with the control ($44.67 \pm 13.03\%$).
212 These changes were statistically significant. After 1 hour of culture the progressive
213 spermatozoa motility was the highest in the control group QK ($60.97 \pm 13.98\%$) and a
214 statistically significant decrease was detected in groups QA and QB. This decrease was
215 compared to control 42.4% for QA and 24.2% for group QB. After 2 hours of *in vitro*
216 culture, the progressive spermatozoa motility was significantly reduced in groups QA
217 and QB with a decrease up to 53% in QA ($22.43 \pm 16.34\%$) and 29.5% in group QB
218 ($33.63 \pm 13.31\%$). Even more dramatic reduction (86.8% for QA vs. 48.5% for QB) was
219 detected compared to the control ($45.12 \pm 19.61\%$) after 3 hours of culture (Figure 2).

220

221 **Distance parameters**

222

223 Spermatozoa distance path parameters confirm the negative dose and time-dependent
224 effect of *Viscum album quercus*. Spermatozoa distance average path significantly
225 decreased at Time 1 in experimental groups QA, QB and QC. Later (Time 2) a
226 significant decrease was detected in almost all experimental groups (except the lowest
227 concentration) and at Time 3 only in groups with the highest *Viscum album quercus*
228 concentrations (Table 2).

229

230 Spermatozoa distance curvilinear line showed similar trends as spermatozoa distance
231 path with various significant decrease at all time periods in experimental groups
232 compared to control (Table 3).

233

234 Spermatozoa distance straight line significantly decreased at all time periods in groups
235 QA - QC. Only at Time 2 group QD showed significant decrease (Table 4).

236

237

238

239

240 **Velocity parameters**

241

242 Also the spermatozoa velocity parameters were affected by the addition of *Viscum*
243 *album quercus* (Iscador Qu) to the culture medium. The spermatozoa velocity curved
244 line at all time periods in control group was 97.58 - 131.47 $\mu\text{m/s}$. After a significant
245 increase at Time 0 (groups QB, QD) a significant decrease was found in all experient
246 groups. Later the most significant decrease was detected in group QA compared to
247 control (131.47 ± 16.38 vs. 93.25 ± 10.19 at Time 1; 108.93 ± 17.72 vs. 70.77 ± 19.69
248 at Time 2 and 97.58 ± 21.79 vs. 47.86 ± 21.66 at Time 3). Also in group QB a
249 significant decrease was detected after 1, 2 and 3 hours of culture (Figure 3).

250

251 The spermatozoa velocity average path clearly confirm the data for spermatozoa
252 velocity curved line with similar significant decrease (Table 5).

253

254 Also the spermatozoa velocity straight line was negative affected by the *Viscum album*
255 *quercus* addition. In group with the highest concentration this parameter decreased at
256 Time 1 to 55.64%, at Time 2 to 54.00% and at Time 3 to 47.77% compared to control
257 (Table 6).

258

259 **Other fine motility parameters**

260

261 The initial amplitude of lateral head displacement (ALH) was $3.78 \pm 0.63 \mu\text{m}$. At Time
262 0 a statistically significant increase was detected in all experiment groups. After 1 hour
263 of cultivation the ALH was in control group $4.70 \pm 0.44 \mu\text{m}$ and the values decreased
264 by 15.1% for group QA and 15.9% for QB. After 2 hours the ALH was the highest in
265 control groups ($4.16 \pm 0.44 \mu\text{m}$) and the most significant decrease was noted in group
266 QA ($3.24 \pm 0.88 \mu\text{m}$, $p < 0.001$) with a decrease of 22.1%. After 3 h of *in vitro*
267 cultivation, the lowest value was found in the sample QA ($2.72 \pm 1.16 \mu\text{m}$) compared to
268 control ($3.98 \pm 0.79 \mu\text{m}$) and was statistically significant (Figure 4).

269

270 The average value of the initial beat cross frequency (BCF) was $29.79 \pm 2.98 \text{ Hz}$. After
271 1 h of *in vitro* cultivation the BCF in QK group increased to $31.91 \pm 2.60 \text{ Hz}$. In all
272 experimental samples a statistically significant decrease was found (time-dependent).
273 However, after 2 hours the BCF in control groups (QK) reached $29.55 \pm 2.18 \text{ Hz}$. The
274 most significant decrease (25.2%) was found for group QA ($22.11 \pm 8.10 \text{ Hz}$). After 3
275 hours the BCF in the control sample was $28.46 \pm 2.75 \text{ Hz}$. In the group QA a
276 statistically significant decrease (34.6%; $18.62 \pm 8.49 \text{ Hz}$), and also in the group QB
277 (17.9%; $23.36 \pm 6.03 \text{ Hz}$), compared to the control group was detected (Figure 5).

278

279 The straightness of spermatozoa movement was in all time periods significantly
280 decreased only in group QA (Table 7).

281

282 Very similar trends were found for spermatozoa linearity (Table 8).

283

284 The spermatozoa wobble was significantly affected only in groups QA and QB (Time 1
285 and 2) with the highest difference detected at Time 1 (0.51 ± 0.03 in group QK vs. 0.44
286 ± 0.03 in group QA (Table 9).

287

288 Viability, membrane integrity, acrosomal integrity

289
290 After 3 hours of incubation, viability of rabbit spermatozoa showed decreased values in
291 all doses of *Viscum album quercus* in comparison to the control group, but the
292 difference were not significant (Figure 6).

293
294 Significant decrease of membrane integrity (intact) was found in groups with the
295 highest *Viscum album quercus* concentrations QB ($64.00 \pm 2.65\%$; $p < 0.05$) and QA
296 ($60.33 \pm 5.51\%$; $p < 0.01$) compared to control group (QK) ($79.33 \pm 1.53\%$). Also, in
297 groups QE, QD, QC lower values in comparison to control group were detected ($p <$
298 0.05 ; Figure 7).

299
300 The values of acrosomal integrity showed very similar tendency (94.00 – 97.33%) in all
301 experimental groups with no significant difference (Figure 8).

302 Discussion

303
304
305 Researchers are unanimous about the negative impact of various factors on male
306 reproduction and cell/spermatozoa quality (Burton 2013, Lukac *et al.* 2013, Petrovova
307 *et al.* 2014, Lukacova *et al.* 2015, Kolarova *et al.* 2017). The mistletoe (*Viscum album*
308 L.) extract has been shown to be an effective complementary drug in the treatment of
309 cancer patients after surgical removal of the primary tumor. It improved survival,
310 recovery from damage caused by irradiation or cytostatic therapy, and quality of life. In
311 animal tests, clear anti-carcinogenic effects of Iscador were demonstrated. Mainly
312 immune stimulation but also direct cytotoxic activity is believed to be responsible for
313 the anti-carcinogenic activity of Iscador. Tests examining the acute toxicity, genotoxic
314 effects as well as effects on reproduction showed no adverse effects of Iscador
315 preparations. Genotoxic effects and effects on reproduction, which cannot be evaluated
316 in clinical use, have been cleared up in animal tests. Iscador was shown to be clearly
317 non-genotoxic and free of relevant toxic effects on reproduction *in vivo* (Maldacker
318 2006, Gardin 2009).

319 On the other hand, it has been stated that anticancer drugs could cause harmful effects
320 on the spermatozoa quality and spermatogenic cell arrangement of male. Therefore, this
321 study was designed to evaluate the *in vitro* effects of *Viscum album quercus*
322 (IsCADOR) on rabbit spermatozoa in order to assess possible beneficial and/or toxic
323 effects of this compound. Rabbits were chosen as the experimental animal in this
324 research for their well-defined reproductive systems (Paal *et al.* 2014, Rafay and
325 Parkanyi 2016). **Some herbal extracts have been proven to have effects on male**
326 **infertility, for example, gossypol, papaya seed, neem oil, neem seaweed and**
327 **verbascoside (Mosher and Pratt 1991, Dehghan *et al.* 2005, Roychoudhury *et al.* 2009,**
328 **Okab *et al.* 2013, Vizzarri *et al.* 2019).**

329 Since decades, *Viscum album* preparations have been used in Europe in oncology. They
330 show multi-faceted anti-tumor *in-vitro* activities, which include inhibition of tumor cell
331 proliferation, induction of apoptosis, inhibition of angiogenesis, modulation of immune
332 competence and gene signature expression. Recently, it was demonstrated *in vitro* that
333 *Viscum album* exerts an anti-inflammatory effect, mostly directed to chronic
334 inflammation by selectively inhibiting cytokine-induced expression of cyclooxygenase-
335 22 (Seibert *et al.* 1989, Bussing 2006, Bussing *et al.* 2008, Hegde *et al.* 2011, Hajto *et*

336 *al.* 2011). Iscador as an anticancer drug used in the treatment of a variety of neoplastic
337 lesions. On the other hand, treatment with *Viscum album quercus* (Iscador) is
338 accompanied by different toxic effects on different body organs.

339 An *in vitro* study investigated the effect of a standardized mistletoe preparation on the
340 action of Trastuzumab, a drug used for the treatment of Her-2 positive breast cancer. A
341 dose dependent anti-proliferative effect of *Viscum album* extract (VAE) was observed
342 at concentrations ≥ 10 $\mu\text{g/mL}$ after 3 days of incubation. After 7 days a significant
343 growth inhibition of 60% with the clinically relevant concentration 1 $\mu\text{g/mL}$ was
344 detected and no proliferating cells were left at VAE concentrations of 10 and 100 $\mu\text{g/ml}$.
345 With 10 $\mu\text{g/ml}$ VAE after 7 days the proportion of early apoptotic cells raised from
346 9.0 % in the control to 17.4 % and that of late apoptotic/necrotic cells from 18.7 % in
347 control to 78.7 %, respectively (Weissenstein *et al.* 2016).

348 Effects of *Viscum album quercus* (VA Qu) extract in various doses were also
349 investigated in an *in vitro* model with tumor cells: three multiple myeloma (MM) cell
350 lines (OPM-2, RPMI-8226, U-266) and three B cell lymphoma cell lines (U-698,
351 DOHH-2, WSU-1). *Viscum album/Qu* extract markedly downregulated the membrane
352 expression of IL-6R and gp130 in RPMI-8226 (down to 29% and 32%) and the
353 expression of gp130 in WSU-1 (down to 22%). There was a marked reduction of viable
354 cells of RPMI-8226 (down to 28%) and WSU-1 (down to 8 %) at 100 $\mu\text{g}/10^6$ cells /ml.
355 There was a clear relationship between the inhibition of proliferation and viability. VA
356 Qu was more effective in cells having a high proliferation rate than in those with a low
357 proliferation rate (Kovacs *et al.* 2006).

358 Our observations indicate that Iscador in higher doses decreases the spermatozoa
359 motility, progressive spermatozoa motility and some other fine spermatozoa motility
360 parameters (velocity curved line, amplitude of lateral head displacement and beat cross
361 frequency). Thus, Iscador (in high doses *in vitro*) can have long-term effects on
362 testicular function and can be a potent gonadal toxic drug.

363 Spermatozoa motility is one of the most important effective factors in male fertility. The
364 mechanism by which Iscador affects the spermatozoa motility has not been clearly
365 elucidated.

366 A decline in fructose level due to alteration in carbohydrate metabolism after Iscador
367 Qu treatment can be suggested (Rigau *et al.* 2001, Gren *et al.* 2011). Reason for
368 decreased spermatozoa motility, can be also a decreased level of androgen carrier
369 proteins involved in spermatozoa motility. Mistletoe lectins have been identified as
370 main active components and exhibit cytotoxic effects. Therefore, probably lectins have
371 a negative effect on the spermatozoa motility found in this study. Also a relationship
372 between diminished spermatozoa quality and anticancer treatment may a result of a
373 series of cascade events that cause a fall in intracellular ATP levels (Luria *et al.* 2002,
374 Turner 2006, Storey 2008), release of different apoptogenic factors (as pro-caspases,
375 cytochrome C, and apoptosis inducing factor) (Casao *et al.* 2015) from mitochondria
376 into the cytosol through disruption of mitochondrial membrane, inactivation of some
377 biochemical pathways, enzyme dysfunction, disturbed axonemal protein
378 phosphorylation, increased membrane permeability and generation of spermicidal
379 products, which have adverse effects on the spermatozoa functions. Similarly, the
380 spermatozoa motility is also reduced by ergonovine (Tash and Means 1982, Gallagher
381 and Senger 1989), and the ergot alkaloids induce motility changes in bovine sperm cells
382 by interacting with alpha-adrenergic receptors (Wang *et al.* 2009).

383

384 **Conclusion**

385

386 The results of this study indicate that spermatozoa are a useful *in vitro* model for the
387 toxicological evaluation of chemicals providing quantitative as well as qualitative data.
388 In conclusion, the present study shows reduction of spermatozoa quality after *Viscum*
389 *album quercus* addition *in vitro*.

390

391 **Acknowledgments**

392

393 This work was supported by projects VEGA 1/0539/18, APVV-16-0289 and KEGA
394 010/SPU-4/2018.

395

396 **Conflict of Interest**

397

398 Authors have no conflict of interest to declare.

399

400 **References**

401

- 402 ADAMKOVICOVA M, TOMAN, R, MARTINIAKOVA M, OMELKA R,
403 BABOSOVA R, KRAJCOVICOVA V, GROSSKOPF B, MASSANYI P: Sperm
404 motility and morphology changes in rats exposed to cadmium and diazinon. *Reprod*
405 *Biol Endocrinol* **14**: 42, 2016.
406 BRAEDEL-RUOFF S: Immunomodulatory effects of *Viscum album* extracts on natural
407 killer cells: Review of clinical trials. *Forsch Komplementarmed* **17**: 63-73, 2010.
408 BURTON A: Study suggests long-term decline in French sperm quality. *Environ Health*
409 *Perspect* **121**: a46, 2013.
410 BUSSING A, BUCKNER U, ENSER-WEIS U: Modulation of chemotherapy-
411 associated immunosuppression by intravenous application of *Viscum album* L. extract,
412 (Iscador): A randomised phase II study. *Eur J Integr Med* **1**: 2-3, 2008.
413 BUSSING A: Immune modulation using mistletoe (*Viscum album* L.) extracts Iscador:
414 Influencing cell function through subcutaneous and intravenous application. *Arzneim*
415 *Forschung* **56**: 508–515, 2006.
416 CASAO A, MATA-CAMPUZANO M, ORDAS L, CEBRIAN-PEREZ JA, MUINO-
417 BLANCO T, MARTINEZ-PASTOR F: Cleaved PARP-1, an apoptotic marker, can be
418 detected in ram spermatozoa. *Reprod Domest Anim* **50**: 688-691, 2015.
419 CHRENEK P, TRANDZIK J, MASSANYI P, MAKAREVICH A, LUKAC N,
420 PESKOVICOVA D, PALEYANDA R: Effect of transgenesis on reproductive traits of
421 rabbit males. *Anim Reprod Sci* **99**: 127-134, 2007.
422 COLPI GM, CONTALB GF, NERVA F, SAGONE P, PIEDIFERRO G: Testicular
423 function following chemo-radiotherapy. *Eur J Obstet Gynecol Reprod Biol* **113**: 2-6,
424 2004.
425 DEGHAN MH, MARTIN T, DEGHANAN R: Antifertility effect of Iranian neem
426 seed alcoholic extract on epididymal sperm of mice. *Iran J Reprod Med* **3**: 83-89, 2005.
427 DI BISCEGLIE C, BERTAGNA A, COMPOSTO ER, LANFRANCO F, BALDI M,
428 MOTTA G, BARBERIS AM, NAPOLITANO E, CASTELLANO E, MANIERI C:
429 Effects of oncological treatments on semen quality in patients with testicular neoplasia
430 or lymphoproliferative disorders. *Asian J Androl* **15**: 425-429, 2013.

431 FALLAS-LOPEZ M, RODRIGUEZ-DE LARA R, BARCENA-GAMA R, SANCHEZ-
432 TORRES ESQUEDA MT, HERNANDEZ-SANCHEZ D, MARTINEZ-HERNANDEZ
433 PA, AGUILAR-ROMERO O: Rabbit sexual behavior, semen and sperm characteristics
434 when supplemented with sprouted wheat. *Anim Reprod Sci* **129**: 221-228, 2011.
435 FELENDÁ JE, TUREK C, STINTZING FC: Antiproliferative potential from aqueous
436 *Viscum album L.* preparations and their main constituents in comparison with ricin and
437 purothionin on human cancer cells. *J Ethnopharmacol*: in press, 2019.
438 GALLAGHER GR, SENGER PL: Effect of phenylephrine, ergonovine, oxytocin and
439 norepinephrine as an extender ingredient on viability of bovine spermatozoa. *J Anim Sci*
440 **67**: 1573-1576, 1989.
441 GARDIN NE: Immunological response to mistletoe (*Viscum album L.*) in cancer
442 patients: A four-case series. *Phytotherapy Res* **23**: 407-411, 2009.
443 GREN A: Effects of Iscador preparations on the reactivity of mouse immune system.
444 *Neuro Endocrinol Lett* **30**: 530-534, 2009.
445 GREN A, FORMICKI G, STAWARZ R, MARTINIAKOVA M, OMEŁKA R: Effect
446 of Iscador on selected parameters of the metabolic block in the animal type Diabetes
447 induced by Alloksan. *J Microbiol Biotech Food Sci* **1**: 215, 2011.
448 GREN A, FORMICKI G: Effects of iscador and vincristine and 5-fluorouracil on brain,
449 liver, and kidney element levels in alloxan-induced diabetic mice. *Biol Trace Elem Res*
450 **152**: 219-224, 2013.
451 GREN A, MASSANYI P: Antidiabetic and antioxidant potential of plant extracts. Nitra:
452 Slovak University of Agriculture, Nitra, pp. 125, 2016. ISBN 978-80-552-1524-2
453 HAJTO T, FODOR K, PERJESI P, NEMETH P: Difficulties and perspectives of
454 immunomodulatory therapy with mistletoe lectins and standardized mistletoe extracts in
455 evidence-based medicine. *Evid Based Complement Alternat Med* **2011**:1-6, 2011.
456 HALO M JR, TIRPAK F, DANO A, ZBYNOVSKA K, KOVACIK A, ONDRUSKA L,
457 GREN A, LUKAC N, MASSANYI P. Zinc affects rabbit spermatozoa in vitro: effects
458 on motility and viability. *J Microbiol Biotech Food Sci* **8**: 901-904, 2018.
459 HEGDE P, MADDUR MS, FRIBOULET A, BAVRY J, KAVERI SV: *Viscum album*
460 exerts anti-inflammatory effect by selectively inhibiting cytokine-induced expression of
461 cyclooxygenase-2. *PLoS One* **6**: e26312, 2011.
462 HERÁČEK J, SOBOTKA V, KOLÁTOROVÁ L, KOČÁREK J, HAMPL R: Serum
463 and intratesticular sex steroids in azoospermic men: how do they correlate? *Physiol Res*
464 **67**: S521-S524, 2018.
465 KOLAROVA H, KRIZOVA J, HULKOVA M, HANSIKOVA H, HULKOVA H,
466 SMID V, ZEMAN J, HONZIK T, TESAROVA M: Changes in transcription pattern
467 lead to a marked decrease in COX, CS and SQR activity after the developmental point
468 of the 22(nd) gestational week. *Physiol Res* **67**: 79-91, 2017.
469 KOVACS E, LINK S, TOFFOL-SCHMIDT U: Cytostatic and cytotoxic effects of
470 mistletoe (*Viscum album L.*) quercus extract Iscador. *Arzneimittelforschung* **56**: 467-
471 473, 2006.
472 KROCKOVA J, MASSANYI P, TOMAN R, DANKO J, ROYCHOUDHURY S: In
473 vivo and in vitro effect of bendiocarb on rabbit testicular structure
474 and spermatozoa motility. *J Environ Sci Health, Part A* **47**: 1301-1311, 2012.
475 LUKAC N, BARDOS L, STAWARZ R, ROYCHOUDHURY S, MAKAREVICH AV,
476 CHRENEK P, DANKO J, MASSANYI P: In vitro effect of nickel on bovine
477 spermatozoa motility and annexin V-labeled membrane changes. *J Appl Toxicol* **31**:
478 144-149, 2011.

479 LUKAC N, LUKACOVA J, PINTO B, KNAZICKA Z, TVRDA E, MASSANYI P:
480 The effect of nonylphenol on the motility and viability of bovine spermatozoa in vitro. *J*
481 *Environ Sci Health, Part A* **48**: 973-979, 2013.

482 LUKACOVA J, JAMBOR T, KNAZICKA Z, TVRDA E, KOLESAROVA A, LUKAC
483 N: Dose- and time-dependent effects of bisphenol A on bovine spermatozoa in vitro. *J*
484 *Environ Sci Health, Part A* **50**: 669-676, 2015.

485 LURIA A, RUBINSTEIN S, LAX Y, BREITBART H: Extracellular adenosine
486 triphosphate stimulates acrosomal exocytosis in bovine spermatozoa via P2
487 purinoceptor. *Biol Reprod* **66**: 429-437, 2002.

488 MALDACKER J: Preclinical investigations with mistletoe (*Viscum album* L.) extract
489 Iscador. *Arzneimittelforschung* **56**: 497-507, 2006.

490 MALTARIS T, KOELBL H, SEUFERT R, KIESEWETTER F, BECKMANN MW,
491 MUELLER A, DITTRICH R: Gonadal damage and options for fertility preservation in
492 female and male cancer survivors. *Asian J Androl* **8**: 515-533, 2006.

493 MANGELSDORF I, BUSCHMANN J, ORTHEN B: Some aspects relating to the
494 evaluation of the effects of chemicals on male fertility. *Regul Toxicol Pharmacol* **37**:
495 356-369, 2003.

496 MASOPUSTOVA A, JEHLICKA P, HUML M, VOTAVA T, TREFIL L, KRESLOVA
497 M, SYKORA J: Plethysmographic and biochemical markers in the diagnosis of
498 endothelial dysfunction in pediatric acute lymphoblastic leukemia survivors - new
499 applications. *Physiol Res* **67**: 903-909, 2018.

500 MASSANYI P, CHRENEK P, LUKAC N, MAKAREVICH AV, OSTRO A, ZIVCAK
501 J, BULLA J: Comparison of different evaluation chambers for analysis of rabbit
502 spermatozoa motility using CASA system. *Slovak J Anim Sci* **41**: 60-66, 2008.

503 MOUSA-BALABEL TM, MOHAMED R A: Effect of different photoperiods and
504 melatonin treatment on rabbit reproductive performance. *Vet Q* **31**: 165-171, 2011.

505 MOSHER WD, PRATT WF: Fecundity and infertility in the United States: Incidence
506 and trends. *J Fertil Steril* **56**: 192-193, 1991.

507 OKAB AB, SAMARA EM, ABDOUN KA, RAFAY J, ONDRUSKA L, PARKANYI
508 V, PIVKO J, AYOUB MA, AL-HAIDARY AA, ALJUMAAH RS, MASSANYI P,
509 LUKAC N: Effects of dietary seaweed (*Ulva lactuca*) supplementation on the
510 reproductive performance of buck and doe rabbits. *J Appl Anim Res* **41**: 347-355, 2013.

511 PAAL D, KROCKOVA J, ONDRUSKA L, SLANINA T, STREJCEK F, MASSANYI
512 P: Effect of semen collection frequency on the progress in the motility of rabbit
513 spermatozoa. *Slovak J Anim Sci* **47**: 61-67, 2014.

514 PARKANYI V, ONDRUSKA L, MASSANYI P, LUKAC N: Effects of the HPA lectin
515 on rabbit sperm motility and fertility. *Slovak J Anim Sci* **48**: 186, 2015.

516 PETROVOVA E, LUPTAKOVA L, KRESAKOVA L, MALOVESKA M,
517 VDOVIAKOVA K, ELIAS MZJ: Effects of pirimicarb on cell proliferation in
518 immature 3D cell cultures. *Folia Veterinaria* **58**: 169-174, 2014.

519 POPE CE, ZHANG YZ, DRESSER BL: A simple staining method for quantifying the
520 acrosomal status of cat spermatozoa. *Theriogenology* **35**: 257, 1991.

521 RAFAY J, PARKANYI V: The rabbit as a model and farm animal at the Research
522 Institute for Animal Production Nitra: A review. *Slovak J Anim Sci* **49**: 141-146, 2016.

523 RIGAU T, FARRE M, BALLESTER J, MOGAS T, PENA A, RODRIGUEZ-GIL JE:
524 Effects of glucose and fructose on motility patterns of dog spermatozoa from fresh
525 ejaculates. *Theriogenology* **56**: 801-815, 2001.

526 ROYCHOUDHURY S, MASSANYI P: In vitro copper inhibition of the rabbit
527 spermatozoa motility. *J Environ Sci Health, Part A* **43**: 651-656, 2008.

528 ROYCHOUDHURY S, MASSANYI P, SLAMECKA J, CHLEBEC I, TRANDZIK J,
529 BULLA J, OKAB AB, TAHA TA, SALEM MH, AYOUB MA: In vitro gossypol
530 induced spermatozoa motility alterations in rabbits. *J Environ Sci Health, Part B* **44**:
531 730-741, 2009.

532 ROYCHOUDHURY S, MASSANYI P, BULLA J, CHOUDHURY MD, STRAKA L,
533 LUKAC N, FORMICKI G, DANKOVA M, BARDOS L: In vitro copper toxicity on
534 rabbit spermatozoa motility, morphology and cell membrane integrity. *J Environ Sci*
535 *Health, Part A* **45**: 1482-1491, 2010.

536 SAHA R, ROYCHOUDHURY S, KAR K, VARGHESE AC, NANDI P, SHARMA
537 GD, FORMICKI G, SLAMA P, KOLESAROVA A: Coenzyme Q10 ameliorates
538 cadmium induced reproductive toxicity in male rats. *Physiol Research* **68**: 141-145,
539 2019.

540 SEIBERT H, KOLOSSA M, WASSERMANN O: Bovine spermatozoa as animal in
541 vitro model for studies on the cytotoxicity of chemicals: Effects of chlorophenols. *Cell*
542 *Biol Toxicol* **5**: 315-330, 1989.

543 SLANINA T, KOVACOVA R, LUKAC N, MASSANYI P: Impact of tilmicosin on the
544 rabbit spermatozoa motility and viability. *J Microbiol Biotech Food Sci* **5**: 53-56, 2016..

545 SLANINA T, MIŠKEJE M, TIRPAK F, BŁASZCZYK M, FORMICKI G,
546 MASSANYI P: Caffeine strongly improves motility parameters of turkey spermatozoa
547 with no effect on cell viability. *Acta Vet Hung* **66**: 137-150, 2018.

548 STEIN GM, BUSSING A, SCHIETZEL M: Stimulation of the maturation of dendritic
549 cells in vitro by a fermented mistletoe extract. *Anticancer Res* **22**: 4215-4219, 2002.

550 STOREY BT: Mammalian sperm metabolism: Oxygen and sugar, friend and foe. *Int J*
551 *Dev Biol* **52**: 427-437, 2008.

552 SUVEREN E, BAXTER GF, ISKIT AB, TURKER AU: Cardioprotective effects of
553 *Viscum album L. subsp. album* (European mistletoe) leaf extracts in myocardial
554 ischemia and reperfusion. *J Ethnopharmacol* **209**: 203-209, 2017.

555 TASH JS, MEANS AR: Regulation of protein phosphorylation and motility of sperm by
556 cyclic adenosine monophosphate and calcium. *Biol Reprod* **26**: 745-763, 1982.

557 TIRPAK F, SLANINA T, KOVACIK A, ONDRUSKA L, MASSANYI P JR, HALO M
558 JR, MASSANYI P: Low taurine concentrations positively affect rabbit spermatozoa
559 properties in later time intervals. *J Microbiol Biotech Food Sci* **7**: 128-131

560 TVRDA E, LUKAC N, LUKACOVA J, JAMBOR T, MASSANYI P: Dose- and time-
561 dependent in vitro effects of divalent and trivalent iron on the activity of bovine
562 spermatozoa. *Biol Trace Elem Res* **167**: 36-47, 2015.

563 TVRDA E, MACKOVICH A, GREIFOVA H, HASHIM F, LUKAC N: Antioxidant
564 effects of lycopene on bovine sperm survival and oxidative profile following
565 cryopreservation. *Vet Med* **62**: 429-436, 2017.

566 TURNER RM: Moving to the beat: a review of mammalian sperm motility
567 regulation. *Reprod Fertil Dev* **18**: 25-38, 2006.

568 VIZZARRI F, PALAZZO M, CASAMASSIMA D, ONDRUSKA L, MASSANYI M,
569 TIRPAK F, FORMICKI G, GREN A, MASSANYI P: *Lippia citriodora* (verbascoside)
570 extract supplementation: Effect on rabbit semen quality in vivo and in vitro. *Czech J*
571 *Anim Sci* **64**: 1-10, 2019.


572 WANG H, LOOPER ML, JOHNSON ZB, RORIE RW, ROSENKRANS CF:
 573 Involvement of signaling pathways in bovine sperm motility, and effect of ergot
 574 alkaloids. *In Vitro Cell Dev Biol* **45**: 483-489, 2009.
 575 WEISSENSTEIN U, KUNZ M, URECH K, REGUEIRO U, BAUMGARTNER S:
 576 Interaction of a standardized mistletoe (*Viscum album*) preparation with antitumor
 577 effects of Trastuzumab in vitro. *BMC Complement Altern Med* **16**: 271, 2016.
 578 VITKU J, SOSVOROVA L, CHLUPACOVA T, HAMPL R, HILL M, SOBOTKA V,
 579 HERACEK J, BICIKOVA M, STARKA L: Differences in bisphenol A and estrogen
 580 levels in the plasma and seminal plasma of men with different degrees of infertility.
 581 *Physiol Res* **64**: S303-S311, 2018.

582
 583
 584
 585

Table 1 – Concentrations of *Viscum album quercus* used in the study.

Group	Semen (μL)	Iscador Qu 10 mg (μL)	Physiological solution (μL)	Concentration of Iscador Qu in samples (mg/mL)
QK – control	100	0	300	0
QA	100	300	0	10
QB	100	200	100	6.6
QC	100	100	200	3.3
QD	100	100	300	2.5
QE	100	100	400	2

586




587

588 Figure 1 – The effect of *Viscum album quercus* on the total spermatozoa motility (in %).

589 QK – 0; QE – 2; QD – 2.5; QC – 3.3; QB – 6.6; QA – 10 mg/mL of Iscador Qu. The

590 level of significance was set at *** (p < 0.001), ** (p < 0.01) and * (p < 0.05).



591
592
593
594
595
596
597

Figure 2 – The effect of *Viscum album quercus* on the progressive spermatozoa motility (in %). QK – 0; QE – 2; QD – 2.5; QC – 3.3; QB – 6.6; QA – 10 mg/mL of Iscador Qu. The level of significance was set at *** (p < 0.001), ** (p < 0.01) and * (p < 0.05).

Table 2 – Spermatozoa distance average path (DAP; μm) in experimental groups and time periods.

Group	Mean	S.D.	C.V.	Minimum	Maximum
Time 0					
QK	22.90	4.43	19.36	16.02	32.75
QA	24.56	3.79	15.43	16.82	31.60
QB	25.69	3.92	15.25	18.35	32.61
QC	25.41	5.21	20.52	11.97	33.70
QD	26.14	5.76	22.02	14.05	38.53
QE	22.97	3.99	17.37	16.40	31.49
Time 1					
QK	28.91	3.69	12.75	21.19	36.69
QA	18.05***	2.22	12.30	14.14	22.83
QB	20.24***	2.25	11.14	16.41	26.01
QC	22.92***	2.38	10.38	18.70	27.44
QD	26.23	2.42	9.21	21.15	30.61
QE	26.49	3.44	12.99	21.54	32.76
Time 2					
QK	22.51	3.34	14.84	15.29	31.10
QA	13.27***	3.36	25.34	0.00	16.81
QB	17.26***	2.41	13.97	12.68	21.90
QC	17.45***	1.88	10.80	13.81	21.56
QD	19.02**	2.72	14.30	14.45	27.30
QE	20.55	2.80	13.60	13.64	26.45
Time 3					

QK	19.54	4.31	22.06	11.80	27.42
QA	9.40 ^{***}	3.99	42.45	0.00	16.90
QB	14.17 ^{***}	4.06	28.65	0.00	22.14
QC	16.70	4.42	26.48	8.95	22.17
QD	18.56	4.40	23.72	11.09	27.61
QE	18.44	3.18	17.25	12.06	22.85

598 Legend: S.D. – standard deviation; C.V. – coefficient of variation; ^{***} ($p < 0.001$), ^{**}
599 ($p < 0.01$) (experimental group vs. control). QK – 0; QE – 2; QD – 2.5; QC – 3.3; QB –
600 6.6; QA – 10 mg/mL of Iscador Qu.

601
602 Table 3 – Spermatozoa distance curvilinear line (DCL; μm) in experimental groups and
603 time periods.

Group	Mean	S.D.	C.V.	Minimum	Maximum
Time 0					
QK	47.84	7.80	16.30	35.09	67.59
QA	52.25	6.81	13.03	36.55	65.73
QB	52.76	6.52	12.35	38.31	65.55
QC	50.01	7.19	14.37	34.42	63.38
QD	56.03 ^{**}	14.35	25.61	35.85	92.81
QE	47.66	6.48	13.59	33.89	62.09
Time 1					
QK	56.01	7.39	13.19	44.27	77.16
QA	41.19 ^{***}	4.74	11.52	33.71	53.96
QB	43.79 ^{***}	4.70	10.74	36.12	53.41
QC	45.98 ^{***}	4.07	8.86	40.28	54.68
QD	52.54	5.33	10.14	40.52	61.28
QE	53.61	7.27	13.56	41.29	66.05
Time 2					
QK	46.87	7.61	16.24	33.81	68.31
QA	31.38 ^{***}	9.01	28.71	0.00	44.65
QB	40.40 ^{***}	4.93	12.21	31.29	50.00
QC	38.71 ^{***}	5.07	13.09	29.61	50.94
QD	41.62 [*]	5.56	13.35	28.80	59.57
QE	45.42	5.46	12.03	31.92	58.97
Time 3					
QK	42.31	9.20	21.76	27.39	60.82
QA	20.69 ^{***}	9.53	46.07	0.00	37.80
QB	34.22 ^{**}	10.26	29.97	0.00	52.86
QC	37.83	9.40	24.85	21.49	49.89
QD	43.07	8.81	20.45	25.50	58.38
QE	41.58	7.61	18.31	25.81	54.31


604 Legend: S.D. – standard deviation; C.V. – coefficient of variation; ^{***} ($p < 0.001$), ^{**}
605 ($p < 0.01$) and ^{*} ($p < 0.05$) (experimental group vs. control). QK – 0; QE – 2; QD – 2.5;
606 QC – 3.3; QB – 6.6; QA – 10 mg/mL of Iscador Qu.

607

608 Table 4 – Spermatozoa distance straight line (DSL; μm) in experimental groups and
 609 time periods.

Group	Mean	S.D.	C.V.	Minimum	Maximum
Time 0					
QK	17.63	3.87	21.97	10.02	26.30
QA	16.62	3.16	19.00	12.07	23.31
QB	18.15	3.57	15.25	11.38	24.51
QC	19.18	4.75	20.52	8.00	26.47
QD	17.84	3.59	20.11	9.68	24.21
QE	16.91	3.24	19.17	11.06	23.93
Time 1					
QK	22.28	3.37	15.13	16.09	29.73
QA	12.84 ^{***}	1.52	11.88	9.67	16.74
QB	15.33 ^{***}	1.81	11.78	12.13	19.33
QC	17.72 ^{**}	1.94	10.92	13.87	22.38
QD	19.85	1.74	8.75	15.93	23.06
QE	19.88	2.18	10.95	16.33	23.88
Time 2					
QK	17.38	2.66	15.29	12.61	24.09
QA	9.60 ^{***}	2.48	25.82	0.00	13.09
QB	12.73 ^{***}	2.04	16.05	8.87	17.83
QC	13.53 ^{***}	1.57	11.62	10.57	17.16
QD	14.82 ^{**}	2.58	17.40	11.06	22.86
QE	15.58	1.95	12.52	10.86	19.14
Time 3					
QK	14.86	3.12	20.97	8.69	20.65
QA	6.72 ^{***}	2.81	41.89	0.00	10.76
QB	10.08 ^{***}	2.75	27.26	0.00	15.48
QC	12.12 ^{**}	3.16	26.10	5.58	16.52
QD	13.31	3.37	25.33	7.81	22.37
QE	13.60	2.12	15.59	9.66	17.87

610 Legend: S.D. – standard deviation; C.V. – coefficient of variation; *** ($p < 0.001$), **
 611 ($p < 0.01$) (experimental group vs. control). QK – 0; QE – 2; QD – 2.5; QC – 3.3; QB –
 612 6.6; QA – 10 mg/mL of Iscador Qu.



613
614
615
616
617
618
619

Figure 3 – The effect of *Viscum album quercus* on the spermatozoa velocity curved line ($\mu\text{m/s}$). QK – 0; QE – 2; QD – 2.5; QC – 3.3; QB – 6.6; QA – 10 mg/mL of Iscador Qu. The level of significance was set at *** ($p < 0.001$), ** ($p < 0.01$) and * ($p < 0.05$).

Table 5 – Spermatozoa velocity average path (VAP; $\mu\text{m/s}$) in experimental groups and time periods.

Group	Mean	S.D.	C.V.	Minimum	Maximum
Time 0					
QK	52.47	10.24	19.51	33.87	73.56
QA	57.45	9.21	16.03	39.74	73.75
QB	60.27	9.37	15.55	43.24	79.03
QC	59.22	12.53	21.16	27.74	76.57
QD	60.01	12.74	21.22	30.63	82.13
QE	53.54	9.79	18.28	37.27	73.56
Time 1					
QK	68.12	8.16	11.99	50.96	86.26
QA	41.16***	5.22	12.68	31.48	51.78
QB	46.62***	5.25	11.26	36.14	59.89
QC	53.53***	6.09	11.37	43.38	64.99
QD	60.81***	5.87	9.65	48.00	73.53
QE	61.79**	8.34	13.50	50.35	78.56
Time 2					
QK	52.56	7.84	14.91	35.50	72.26
QA	30.32***	7.56	24.93	0.00	38.99
QB	38.92***	5.39	13.84	29.51	49.88
QC	40.41***	4.07	10.07	32.51	48.94
QD	43.58***	6.71	15.41	31.77	60.93
QE	47.11*	6.80	14.43	33.09	64.36
Time 3					


QK	45.30	10.31	22.75	27.50	64.75
QA	22.61***	10.05	44.43	0.00	40.14
QB	31.86***	9.39	29.48	0.00	49.84
QC	38.17	10.45	27.38	19.32	52.12
QD	42.18	10.34	24.51	25.95	61.73
QE	41.81	7.56	18.08	28.83	52.11

620 Legend: S.D. – standard deviation; C.V. – coefficient of variation; *** ($p < 0.001$), **
621 ($p < 0.01$) and * ($p < 0.05$) (experimental group vs. control). QK – 0; QE – 2; QD – 2.5;
622 QC – 3.3; QB – 6.6; QA – 10 mg/mL of Iscador Qu.
623

624 Table 6 – Spermatozoa velocity straight line (VSL; $\mu\text{m/s}$) in experimental groups and
625 time periods.


Group	Mean	S.D.	C.V.	Minimum	Maximum
Time 0					
QK	40.53	8.81	21.73	21.06	60.28
QA	39.07	7.71	19.72	27.11	54.48
QB	42.93	8.68	20.22	26.91	59.89
QC	44.94	11.41	25.39	18.85	60.35
QD	41.47	9.28	22.36	21.05	58.10
QE	39.72	7.99	20.12	25.20	56.26
Time 1					
QK	52.75	7.60	14.41	38.78	70.71
QA	29.35***	3.70	12.60	21.52	37.00
QB	35.44***	4.17	11.78	27.66	44.88
QC	41.50***	5.04	12.14	31.08	52.56
QD	46.26***	4.27	9.24	36.38	55.44
QE	46.64***	5.25	11.25	39.21	57.29
Time 2					
QK	40.72	6.28	15.41	29.37	56.76
QA	21.99***	5.56	25.60	0.00	30.47
QB	28.76***	4.53	15.76	20.79	40.55
QC	31.40***	3.55	11.31	24.63	39.57
QD	34.09***	6.39	18.75	23.67	51.32
QE	35.68**	4.51	12.63	26.71	44.68
Time 3					
QK	34.52	7.51	21.77	20.31	49.01
QA	16.49***	7.56	45.86	0.00	31.82
QB	22.65***	6.40	28.23	0.00	34.89
QC	27.81**	7.45	26.80	12.09	38.72
QD	30.22	7.86	23.01	18.52	50.15
QE	30.80	4.97	16.14	21.21	40.57

626 Legend: S.D. – standard deviation; C.V. – coefficient of variation; *** ($p < 0.001$), **
627 ($p < 0.01$) (experimental group vs. control). QK – 0; QE – 2; QD – 2.5; QC – 3.3; QB –
628 6.6; QA – 10 mg/mL of Iscador Qu.



629
630
631
632
633
634

Figure 4 – The effect of *Viscum album quercus* on the amplitude of lateral head displacement (μm). QK – 0; QE – 2; QD – 2.5; QC – 3.3; QB – 6.6; QA – 10 mg/mL of Iscador Qu. The level of significance was set at *** ($p < 0.001$), ** ($p < 0.01$) and * ($p < 0.05$).



635
636
637
638
639
640

Figure 5 – The effect of *Viscum album quercus* on the beat cross frequency (Hz). QK – 0; QE – 2; QD – 2.5; QC – 3.3; QB – 6.6; QA – 10 mg/mL of Iscador Qu. The level of significance was set at *** ($p < 0.001$), ** ($p < 0.01$) and * ($p < 0.05$).

Table 7 – Spermatozoa straightness (STR) in experimental groups and time periods.

Group	Mean	S.D.	C.V.	Minimum	Maximum
Time 0					
QK	0.77	0.06	7.88	0.61	0.89
QA	0.67***	0.06	9.28	0.57	0.77
QB	0.70**	0.06	9.17	0.59	0.80
QC	0.75	0.06	8.50	0.63	0.87
QD	0.69***	0.11	15.98	0.45	0.84
QE	0.73	0.04	5.16	0.67	0.82
Time 1					
QK	0.77	0.04	5.57	0.62	0.86
QA	0.71***	0.05	6.91	0.58	0.79
QB	0.76	0.04	5.45	0.69	0.86
QC	0.77	0.03	3.59	0.71	0.81
QD	0.76	0.05	6.26	0.64	0.85
QE	0.75	0.03	4.51	0.68	0.82
Time 2					
QK	0.77	0.04	5.42	0.62	0.85
QA	0.70***	0.14	19.46	0.00	0.89
QB	0.73	0.05	6.56	0.65	0.84
QC	0.77	0.02	3.10	0.72	0.82
QD	0.77	0.05	5.89	0.69	0.85
QE	0.75	0.04	5.50	0.66	0.84
Time 3					
QK	0.76	0.05	6.45	0.66	0.91
QA	0.64 ^C	0.26	40.91	0.00	31.82
QB	0.69	0.14	20.23	0.00	0.92
QC	0.73	0.05	6.66	0.61	0.80
QD	0.71	0.06	8.58	0.61	0.81
QE	0.74	0.04	5.89	0.66	0.81

641 Legend: S.D. – standard deviation; C.V. – coefficient of variation; *** ($p < 0.001$), **
642 ($p < 0.01$) (experimental group vs. control). QK – 0; QE – 2; QD – 2.5; QC – 3.3; QB –
643 6.6; QA – 10 mg/mL of Iscador Qu.

644
645 Table 8 – Spermatozoa linearity (LIN) in experimental groups and time periods.

Group	Mean	S.D.	C.V.	Minimum	Maximum
Time 0					
QK	0.37	0.05	13.31	0.23	0.45
QA	0.32**	0.03	10.81	0.26	0.37
QB	0.34	0.05	14.89	0.27	0.44
QC	0.38	0.07	17.39	0.24	0.48
QD	0.33	0.09	25.70	0.18	0.47
QE	0.35	0.04	12.00	0.27	0.42
Time 1					
QK	0.40	0.04	10.28	0.29	0.50
QA	0.31***	0.03	9.77	0.24	0.36
QB	0.35***	0.03	8.88	0.29	0.46

QC	0.38	0.03	7.33	0.30	0.44
QD	0.38	0.04	9.56	0.30	0.47
QE	0.37*	0.02	6.42	0.32	0.41
Time 2					
QK	0.37	0.03	8.10	0.27	0.42
QA	0.31***	0.12	38.07	0.00	0.86
QB	0.31***	0.03	8.75	0.26	0.38
QC	0.35	0.03	7.42	0.30	0.40
QD	0.35	0.04	10.76	0.28	0.42
QE	0.34	0.02	6.99	0.29	0.39
Time 3					
QK	0.35	0.03	9.49	0.30	0.44
QA	0.32	0.18	56.85	0.00	0.74
QB	0.29	0.11	38.09	0.00	0.81
QC	0.32	0.04	12.17	0.24	0.40
QD	0.30	0.04	12.96	0.25	0.39
QE	0.33	0.03	10.36	0.27	0.40

646 Legend: S.D. – standard deviation; C.V – coefficient of variation; *** ($p < 0.001$), ** (p
647 < 0.01) and * ($p < 0.05$) (experimental group vs. control). QK – 0; QE – 2; QD – 2.5;
648 QC – 3.3; QB – 6.6; QA – 10 mg/mL of Iscador Qu.

649


650

Table 9 – Spermatozoa wobble (WOB) in experimental groups and time periods.


Group	Mean	S.D.	C.V.	Minimum	Maximum
Time 0					
QK	0.48	0.04	8.73	0.38	0.55
QA	0.47	0.02	3.71	0.43	0.50
QB	0.48	0.03	7.21	0.44	0.55
QC	0.50	0.05	9.99	0.36	0.57
QD	0.47	0.06	11.90	0.39	0.56
QE	0.48	0.04	7.73	0.40	0.53
Time 1					
QK	0.51	0.03	6.17	0.44	0.60
QA	0.44***	0.03	6.29	0.38	0.49
QB	0.46***	0.02	4.89	0.42	0.53
QC	0.50	0.03	5.35	0.43	0.56
QD	0.50	0.02	4.84	0.46	0.57
QE	0.49*	0.02	4.81	0.43	0.53
Time 2					
QK	0.48	0.02	4.70	0.43	0.52
QA	0.42**	0.13	30.50	0.00	0.86
QB	0.42**	0.02	5.45	0.39	0.47
QC	0.45	0.03	6.08	0.30	0.40
QD	0.46	0.03	6.83	0.39	0.51
QE	0.45	0.02	4.99	0.40	0.50
Time 3					
QK	0.46	0.03	6.50	0.39	0.57

QA	0.42	0.20	46.81	0.00	0.80
QB	0.41	0.12	28.25	0.00	0.87
QC	0.44	0.03	7.70	0.34	0.50
QD	0.43	0.03	6.84	0.38	0.49
QE	0.44	0.03	6.16	0.38	0.52


651 Legend: S.D. – standard deviation; C.V. – coefficient of variation; *** ($p < 0.001$), **
652 ($p < 0.01$) and * ($p < 0.05$) (experimental group vs. control). QK – 0; QE – 2; QD – 2.5;
653 QC – 3.3; QB – 6.6; QA – 10 mg/mL of Iscador Qu.
654



655 Figure 6 – The effect of *Viscum album quercus* on the viability (%) of rabbit
656 spermatozoa after 3 hours of incubation. QK – 0; QE – 2; QD – 2.5; QC – 3.3; QB –
657 6.6; QA – 10 mg/mL of Iscador Qu.
658
659



660
 661 Figure 7 – The effect of *Viscum album quercus* on the membrane integrity (%) of rabbit
 662 spermatozoa assessed after 3 hours of incubation. QK – 0; QE – 2; QD – 2.5; QC – 3.3;
 663 QB – 6.6; QA – 10 mg/mL of Iscador Qu. The level of significance was set at ** ($p <$
 664 0.01) and * ($p < 0.05$).
 665



666
 667 Figure 8 – The effect of *Viscum album quercus* on the acrosomal integrity(%) of rabbit
 668 spermatozoa assessed after 3 hours of incubation. QK – 0; QE – 2; QD – 2.5; QC – 3.3;
 669 QB – 6.6; QA – 10 mg/mL of Iscador Qu.
 670
 671