

Physiological Research Pre-Press Article

Cerebrovascular dynamics during continuous motor task.

Martin Müller und Mareike Österreich,

Neurocenter, Neurovascular Laboratory, Lucerne Kantonsspital, Spitalstrasse, CH 6000 Lucerne,
Switzerland;

Running head: Motor task cerebrovascular dynamics

Correspondence

Martin Müller, MD

Neurocenter, Kantonsspital Luzern

Spitalstasse

CH-6000 Lucerne

Martin.mueller@luks.ch

Phone: +41 41 205 5113

Abstract

Purpose. We investigated the cerebral autoregulation (CA) dynamics parameter phase and gain change when exposed to a longlasting motor task. **Methods.** 25 healthy subjects (mean age \pm SE, 38 ± 2.6 years; 13 females) underwent simultaneous recordings of spontaneous fluctuations in blood pressure (BP), cerebral blood flow velocity (CBFV), and end-tidal CO₂ (ETCO₂) over 5 minutes of rest followed by 5 minute of left elbow flexion at a frequency of 1 Hz. Tansfer function gain and phase between BP and CBFV were assessed in the frequency ranges of very low frequencies (VLF; 0.02-0.07 Hz), low frequencies (LF; 0.07-0.15), and high frequencies (HF; >0.15). **Results.** CBFV increased on both sides rapidly to maintain an elevated steady state until movement stopped. Cerebrovascular resistance fell on the right side (rest 1.35 ± 0.06 , movement 1.28 ± 0.06 ; $p < 0.01$), LF gain decreased from baseline (right side 0.97 ± 0.07 %/mm Hg; left 1.01 ± 0.09) to movement epoch (right 0.73 ± 0.08 ; left 0.76 ± 0.06 ; $p \leq 0.01$). VLF phase decreased from baseline (right 1.03 ± 0.05 radians; left 1.10 ± 0.06) to the movement epoch (right 0.81 ± 0.07 ; left 0.82 ± 0.10 ; $p \leq 0.05$). **Conclusion.** CA regulates continuous motor efforts by changes in resistance, gain and phase.

Keywords: cerebrovascular dynamics, motor task, transcranial Doppler sonography, transfer function, physiology

Introduction

Cerebral autoregulation (CA) is the ability of the brain's vasculature to maintain cerebral blood flow constant over a wide blood pressure (BP) range (Kontos et al. 1978; Harper et al. 1984). When performing cognitive, motor or other tasks the blood supply to the activated brain areas is then further modulated regionally via vasoconstriction and –dilation which are based on the metabolic demands. CA is assessable by a) static methods which are characterized by measuring cerebral blood flow or its velocity [CBF(V)] under stable conditions before and after a vasodilatory/vasoconstrictory stimulus (Leenders et al. 1999; Paulson et al. 2010; Powers 1991) and b) by dynamic methods which analyze the interaction between CBF(V) and BP at spontaneous or induced slow oscillations present in CBF(V) and BP (Claassen et al. 2016; Meel-van den Abeelen et al. 2014; Müller and Österreich 2014; Zhang et al. 1998). When CA fails an additional metabolic mechanism helps to keep oxygen concentration in the needed range by increasing the oxygen extraction rate as increasing the amount of the oxygen concentration released from hemoglobin (Leenders et al. 1999). In pathological conditions such as acute stroke the coordinated overlapping of these mechanisms seems to be disturbed (Caldas et al. 2018; Maggio et al. 2013; Reinhard et al. 2012; Rivera-Lara et al. 2017) which might be relevant for an early start of rehabilitation (Coleman et al. 2017; Li et al. 2018, Chi et al 2018).

In recent years the dynamics of the cerebrovascular system was studied by means of mathematical models which usually describe the BP – CBFV system over a wide range of BP and CBFV changes with CBF regulation occurring mostly between CBFV changes of 5 to 50 seconds (Claassen et al. 2016; Mitsis et al. 2006; Müller and Österreich 2014; Serrador et al. 2005; Zhang et al. 1998.). To characterize the system better its response to physiological stimuli like hand gripping, arm or leg movements, or visual tasks (Caldas et al. 2018; Giller et al 2000; Maggio et al. 2013; Panerai et al. 2012a; Salinet et al. 2013) is analyzed. The duration of the used stimuli is usually short (e.g. a minute or less) and repeated several times to allow for an analysis after averaging over all runs, a setting usually addressing neurovascular coupling (NVC). The characteristic NVC response is a rapid, often bilateral CBF(V) increase which continues as long as the stimulus is present. As far as it is deducible from such short interventions, CA in terms of the transfer function (TF) estimates phase and gain is unchanged compared to the baseline prior to the stimulus. Whether these findings are transferrable to continuous tasks is assumed, but not established. The aim of our study was to prove this assumption using as stimulus elbow flexion and extension (at a frequency of 1 Hz) as this task has been shown to provide a feasible CBFV increase (Llwyd et al. 2017; Panerai et al. 2012a) that enables to study autoregulatory mechanisms.

Material and Methods

The study was approved by the ethics committee of Northwest- and Central Switzerland and followed the Declaration of Helsinki as well as good clinical practice standards. Written informed consent was obtained from all subjects [n=25; women 13, mean age \pm standard error of means (SE) 38 ± 2.6 years (range 24-62 years)].

Experimental setting: all investigations followed the recommendation of Claassen et al. (2016), were performed in the late morning with the subject in a supine position with the head elevated approximately 30° in a room with dimmed light. Last coffee or tea uptakes were taken a minimum of 4

hours before the investigations. The subjects were dominantly right-handed (20 of 25); all were non-smokers, healthy and without cerebrovascular risk factors. Participants were carefully instructed to flex and extend the elbow over the full movement range (= one movement cycle) at a pace of 1 Hz which was signaled by a computerized metronome's beep. Before mounting the probes several elbow movement cycles were tried at the metronome's pacing. We assessed the cerebral blood flow velocity (CBFV) in both middle cerebral arteries (MCA) using transcranial Doppler sonography (TCD) by placing the 2-MHz probes of the TCD device (MultidopX, DWL; Compumedics, Sippligen, Germany) at the temporal skull and fixed them with a head holder provided by the manufacturer. The MCAs were identified as CBFV directed towards the probe/skull in a depth of 50-55 mm. End-tidal CO₂ (ETCO₂) was measured via a small nostril tube with the capnography embedded in the TCD device. BP was estimated by finger plethysmography (Finometer Pro; Finapres Medical Systems, Amsterdam, The Netherlands) placed at the fingertip of the right index finger; special attention was paid to its calibration to the brachial artery pressure. The BP signal was imported into the TCD device, and all parameters were recorded simultaneously. After all probes were placed and the subject familiar with the surroundings and the experimental setting, we started with a baseline recording of 5 minutes; thereafter the elbow movement paradigm was performed over 5 minutes.

Data preparation: BP, CBFV, and ETCO₂ data were collected at 100 Hz. The data were analyzed using Matlab (2015b; Math Works Inc., Natick, MA, USA). The data were visually inspected for artefacts, and only artefact-free data periods of 5 min. were used. Each raw data time series was averaged over 1 second interval. The coherence and the TF estimates of phase and gain between the BP and the CBFV time series were extracted from their respective power auto-spectra or cross-spectra using Welch's averaged periodogram method, with a Hann window length of 100 s, window overlap of 50%, and a total data length of 300 seconds. In each subject, the coherence, the phase (in radian), and the gain (in cm/s/mmHg and in %/mmHg) were calculated over a frequency range of 0.02–0.40 Hz in the epochs of baseline and motor task. Based on recent recommendations (Claassen et al 2016) only those phase and gain values were considered for further use when coherence was >0.44. The results of phase, gain and coherence are reported as their respective average in the very low frequency range (VLF, 0.02-0.07 Hz), low frequency range (LF, 0.07-0.15 Hz), and high frequency range (HF, 0.15-0.4 Hz). Over each epoch we also calculated the cerebrovascular resistance as mean BP/ mean CBFV, and heart rate which was derived from the BP signal.

Statistical analysis

Data analysis was performed with the Matlab Statistical Toolbox. Using a Kolmogorov-Smirnov test, all data showed a normal distribution, and the data are reported as mean \pm SE (standard error of the mean). To compare means between the two epochs (baseline and motor task) we used one-way ANOVA including a correction for multiple comparisons when indicated. $P \leq 0.05$ was considered statistically significant.

Results

In 24 of the 25 subjects recordings were possible in both MCAs. In one person only the right MCA was detected. Artefact-free recordings were achieved from all vessels. Figure 1 shows the time

course of mean CBFV (right n=25; left n=24), mean ETCO₂ (n=25) and mean BP (n=25) over the total experiment. After the baseline epoch CBFV increases rapidly with elbow movement and remained at an elevated steady state over the total time of elbow movement; CBFV increase is slightly higher on the right side (contralateral to the arm movement; table1). After starting of arm movement ETCO₂ and BP decreased (ETCO₂) or increased (BP). We defined the time point at which ETCO₂ decreased or BP increased as the time point at which the decreasing or increasing curve was clearly below or above the fluctuation range of the baseline recording curve. ETCO₂ dropped 5 ± 0.1 seconds after movement starting from 44.8 to 41.1 mmHg to remain thereafter relatively stable at a mean ETCO₂ of 41.7 mm Hg. In contrast to CBFV and ETCO₂, BP remained at the baseline level for the first 20 ± 0.3 seconds after starting of the stimulus before rising notably.

Table 1 shows the significant differences between the variables of the baseline and the motor task epochs. BP, CBFV on both sides and heart rate increased while ETCO₂ decreased in the motor task epoch. CVR decreased in the right MCA ($p \leq 0.01$) but not in the left MCA. Coherence and gain (in cm/s/mmHg) did not show differences between the two epochs within one side nor between the two sides. Gain in %/mmHg decreased significantly at both sides in the LF range from baseline (right side 0.97 ± 0.07 ; left 1.01 ± 0.09) to movement epoch (right 0.73 ± 0.08 ; left 0.76 ± 0.06 , $P \leq 0.01$); it remained unchanged in the VLF and HF ranges. Phase (in radian) in the VLF range decreased significantly from baseline (right 1.03 ± 0.05 ; left 1.10 ± 0.06) to the elbow movement epoch (right 0.81 ± 0.07 ; left 0.82 ± 0.10 , $P \leq 0.05$); it did not change in the LF and HF ranges. Heart rate (in heart beat/s) was $1.11-1.15 \pm 0.019$ (maximum SE) during the rest period; it was $1.12-1.16 \pm 0.022$ between starting elbow movement and 12 seconds thereafter (not significant different); it was 1.38 ± 0.011 ($p < 0.005$) at 15 seconds after elbow movement start and remained at this level until the end of movement (1.38 ± 0.022 ; $p < 0.005$ compared to baseline)

Discussion

The time course of and the amount of the evoked CBFV achieved by our setting corresponds to other motor tasks evoked flow studies targeting NVC insofar as flow is evoked bilaterally, remains elevated as long as the stimulus is present, and an attenuation does not ensue; a BP increase occurs comparably to those observed in NVC studies (Caldas et al. 2018; Giller et al 2000; Llwyd et al. 2017; Maggio et al. 2013; Nikolic et al. 2015). In contrast to these reports on NVC, however, our continuous motor task showed changes in the LF range (% /mmHg) and in the VLF range (phase) which are consistent with a disturbed dynamical CA (dCA; Zhang et al.1998; Panerai et al. 1999; Panerai et al. 2001; Serrador et al. 2005). At least in the right MCA, the vessel supplying the cortical areas in which the movement is primarily generated, CVR decreased which can be interpreted as a vasodilation. The CVR decrease seems in agreement with current concepts of neurovascular interaction derived from functional magnetic resonance imaging and from near infrared studies (Huber et al. 2017; Petridou et al. 2017; Schmid et al. 2017): with the beginning of the task a vasodilation occurs which leads to an increase in cerebral blood volume and an increase in the concentration of oxygenated blood which is considered to represent an increase in CBF. When such a vasodilation is induced by CO₂, in dCA the vasodilatory response consists usually of a phase decrease while gain remains not significantly changed (Müller et al. 2016; Panerai et al. 1999; Zhang et al. 1998). Thus, the phase decrease findings in our population could be the result of vasodilation. However, when BP is pharmacologically increased rapidly, dCA's response has been reported to consist of a decrease in both phase and gain

accompanied by a CVR increase (Zhang et al. 2009). Our findings with phase and gain decreases would then hint more in the direction of a BP related response with CVR decrease indicating some additional regional regulation.

The other variable to influence dCA is ETCO₂ change. An ETCO₂ decrease as recorded in our subjects leads usually to a phase increase and to a gain decrease (Birch et al. 1995; Müller et al. 2003). This is contrary to what we have found. An exercise induced lactate acidosis drives a compensatory hyperventilation with the result that dCA's phase and gain were reported unchanged (Ogoh et al. 2007). Thus, ETCO₂ decrease due to hyperventilation might have nothing to do with the dCA changes.

The BP rise in our subjects could be due to the (most likely exercise induced) heart rate increase or be a result of the brain's metabolic demands. It is beyond our data to clarify whether the dCA changes are primarily due to the metabolic demands or due to the BP increase. Maybe a study comparable to ours which includes an observation of the metabolic changes in the brain, e.g. by using NIRS devices able of recording cytochrome-oxidase changes (Obrig et al. 2000) could be of help for this purpose. Such a study might be relevant especially when patients are included (see next paragraph).

Our phase and gain findings could be considered a disturbed dCA induced by performing repetitive arm movements. This movement induced disturbed CA state will interfere in many stroke patients with a preexisting CA condition which by itself is the combined result of an elevated BP, the effect of cerebral ischemia (Llwyd et al. 2018; Reinhard et al. 2012) and/or an unstable circulatory response induced by orthostatic stress (Mitis et al. 2006; Serrador et al. 2005). In clinical practice, in most stroke patient's systolic BP values until 180 mm Hg are tolerated. But high BP handicaps dCA in several conditions: when it is rapidly increased (Zhang et al. 2009), when BP is newly diagnosed before treatment is sufficient (Serrador et al. 2005; Zhang et al. 2007), and in patients with long lasting chronic high BP even when BP is sufficiently treated (Müller et al. 2018). Hence, one future question is whether all disturbances will add up, and whether a risk for the patient will result.

Our study has limitations.

With TF estimates we used strict linear modeling to assess the relationship between BP and CBFV. This kept nonlinear and nonstationary aspects of the system unconsidered (Castro et al. 2017; Giller and Mueller 2003; Placek et al. 2017). All nonlinear approaches showed that the coefficient of variance as a sign of the models quality or analog measures become smaller compared to the strict linear models indicating that these models were likely more precise. However, in terms phase, gain or impulse response as measure of comparison such models did not exhibit large differences from linear differential equations models (Kouchakpour et al. 2014; Marmarelis et al. 2014; Meel-van den Abeelen et al. 2014; Panerai 1999a; Smirl et al 2015; Panerai et al. 2001). Regarding non-stationarity present in the data, recordings over time periods of several minutes can average out non-stationary effects with the result that time-variant models and time-invariant models produce close results (Kouchakpour et al. 2010; Marmarelis et al 2014; Nikolic et al. 2015; Placek et al 2017).

We used CBFV instead of CBF. The velocity signal depends strongly on the vessel's diameter and on multiple other influences apart from BP, such as CO₂, autonomic influences, myogenic and cognitive

activity (Panerai et al. 2012; Panerai et al. 2012a; Sohn 1998; Spronck et al. 2012). Nevertheless, when a Doppler signal derived index of cerebral blood flow (flow index) was compared to CBFV, this flow index was shown to increase during the hand grip task but the amount of increase was less high compared to the velocity signal (Giller et al. 2000).

We suggest that ETCO₂ at rest reflects blood partial CO₂ content, a relationship that might be less close and more variable while exercising (Brys et al. 2003; Jørgensen et al. 1992; Ogoh et al 2007; Robbins et al. 1990). This could be relevant to explain the transfer function result differences between the two cycling studies and our investigation because CO₂ reacts slowly and amounts for a considerable fraction of CBFV variations, and hence spectral density, in the LF and VLF range (Kouchakpour et al. 2010; Mitsis et al. 2006; Panerai et al. 2012; Panerai et al. 2012a).

Conclusions: Our findings suggest that CA adapt to a continuous motor task by modulating CVR, gain and phase. However, the observed changes could also be considered a CA impairment. BP elevation might be a co-factor to be considered. Whether this is of relevance for motor recovery e.g. after stroke needs further investigation.

Acknowledgments: We gratefully thank Prof RS Smith, Automatic Control Laboratory, ETH Zürich, for his helpful comments on transfer function analysis.

Conflict of interest: The authors declare that they have no conflict of interest.

References

Birch AA, Dirnhuber MJ, Hartley-Davies R, Iannotti F, Neil-Dwyer G.. Assessment of autoregulation by means of periodic changes in blood pressure. *Stroke*. 26: 834-7, 1995

Caldas JR, Panerai RB, Salinet ASM, Seng-Shu E, Ferreira GSR, Camara L, Passos RH, Galas FRBG, Almeida JP, Nogueira RC, de Lima Oliveira M, Robinson TG, Hajjar LA. Dynamic cerebral autoregulation is impaired during submaximal isometric handgrip in patients with heart failure. *Am J Physiol Heart Circ Physiol* **315**: H254-H261, 2018

Castro P, Freitas J, Santos R, Panerai R, Azevedo E. Indexes of cerebral autoregulation do not reflect impairment in syncope: insights from head-up tilt test of vasovagal and autonomic failure subjects. *Eur J Appl Physiol* **117**: 1817–1831, 2017

Chi N-F, Hu H-H, Wang C-Y, Chan L, Peng C-K, Novak V, Hu C-J. Dynamic cerebral autoregulation is an independent functional outcome predictor of mild acute ischemic stroke. *Stroke* **49**: 2605-2611, 2018

Claassen JA, Meel-van den Abeelen AS, Simpson DM, Panerai RB.; international Cerebral Autoregulation Research Network (CARNet). Transfer function analysis of dynamic cerebral autoregulation: A white paper from the International Cerebral Autoregulation Research Network. *J Cereb Blood Flow Metab.* **36**: 665-80, 2016

Coleman ER, Moudgal R, Lang K, Hyacinth HI, Awosika OO, Kissela BM, Feng W. Early rehabilitation after stroke: a narrative review. *Curr Atheroscler Rep.* (November 7, 2017). doi 10.1007/s11883-017-0686-6, 2017

Giller CA, Giller AM, Cooper Cr, Hatab MR. Evaluation of the cerebral hemodynamic response to rhythmic handgrip. *J Appl Physiol* **88**: 2205-2213, 2000

Giller CA, Mueller M. Linearity and non-linearity in cerebral hemodynamics. *Med Eng Phys* **25**: 633-646, 2003

Harper SL, Bohlen HG, Rubin MJ. Arterial and microvascular contributions to cerebral cortical autoregulation. *Am J Physiol* 246 (Heart Circ. Physiol. 15): H17-H24, 1984

Huber, Uludag K, Möller HE: Non-BOLD contrast for laminar fMRI in humans: CBF, CBV, and CRMO2. *NeuroImage* ; <http://dx.doi.org/10.1016/j.neuroimage.2017.07.041>, 2017

Jørgensen LG, Perko M, Hanel B, Schroeder TV, Secher NH. Middle cerebral artery flow velocity and blood flow during exercise and muscle ischemia in humans. *J Appl Physiol* **72**: 1123–1132, 1992

Kontos HA, Wie EP, Navari RM, Levasseur JE, Rosenblum WI, Patterson JL. Responses of cerebral arteries and arterioles to acute hypotension and hypertension. *Am J Physiol* 234: H371-H383, 1978

Kouchakpour H, Allen R, Simpson DM. Nonlinear, multiple-input modeling of cerebral autoregulation using Volterra Kernel estimation. *Conf Proc IEEE Eng Med Biol Soc* 2010: 2375-8, 2010

Leenders KL, Perani D, Lammertsma AA, Heather JD, Buckingham P, Healy MJ, Gibbs JM, Wise RJ, Hatazawa J, Herold S. Cerebral blood flow, blood volume and oxygen utilization. Normal values and effect of age. *Brain* **113**: 27-47, 1999.

Li Z, Zhang X, Wang K, Wen J. Effects of early mobilization after acute stroke: a meta-analysis of randomized control trials. *J Stroke Cerebrovasc Dis.* **27**: 1326-1337, 2018

Llwyd O, Panerai RB, Robinson TG. Effects of dominant and non-dominant passive arm manoeuvres on the neurovascular coupling response. *Eur J Appl Physiol* **117**: 2191-2199, 2017

Llwyd O, Salinet ASM, Panerai RB, Lam MY, Saeed NP, Brodie F, Bor-Seng-Shu, Robinson TG, Nogueira RC. Cerebral haemodynamics following acute stroke: effects of stroke severity and stroke subtype. *Cerebrovascular disease extra* 2018; **8**: 80-89, 2018

Maggio P, Salinet ASM, Panerai RB, Robinson TG. Does hypercapnia-induced impairment of cerebral autoregulation affect neurovascular coupling? A functional TCD study. *J Appl Physiol* **115**: 491-497, 2013

Marmarelis VZ, Shin DC, Orme M, Zhang R: Time-varying modeling of cerebral hemodynamics. *IEEE Trans Biomed Eng* **61**: 694–704, 2014.

Meel-van den Abeelen ASS, Simpson D, Zhang R, Tarumi T, Rickards CA, Payne S, Mitsis GD, Kostoglou K, Marmarelis V, Shin D, Tzeng Y-C, Ainslie PN, Gommer E, Müller M, Dorado AC, Smielewski P, Yelicich B, Puppo C, Liu X, Czosnyka M, Wang C-Y, Novak V, Panerai RB, Claassen JAHR. Between-center variability in transfer function analysis: a widely used method for linear quantification of the dynamic pressure-flow relation: the CARnet bootstrap study. *Med Biol Eng* **36**: 620-7, 2014

Mitsis GD, Zhang R, Levine BD, Marmarelis VZ. Cerebral hemodynamics during orthostatic stress assessed by nonlinear modeling. *J Appl Physiol* **101**: 354–366, 2006.

Müller M, Bianchi O, Erülkü S, Stock C, Schwerdtfeger K; for the Homburg Traumatic Brain Injury Group. Changes in Linear Dynamics of Cerebrovascular System After Severe Traumatic Brain Injury. *Stroke* **34**: 1197-1202, 2003

Müller MW, Österreich M. A comparison of dynamic cerebral autoregulation across changes in cerebral blood flow velocity for 200 s. *Front Physiol.* (August 26, 2014). doi: 10.3389/fphys.2014.00327., 2014

Müller M, Österreich M, Müller A, Lygeros L (2016). Assessment of the brain's macro- and micro-circulatory blood flow responses to CO₂ via transfer function analysis. *Front Physiol* (May 9, 2016). doi: 10.3389/fphys.2016.00162, 2016

Müller M, Österreich M, von Hessling A, Smith RS. Incomplete recovery of cerebral blood flow dynamics in sufficiently treated high blood pressure. *J Hypertens.* (July 10, 2018). doi: 10.1097/HJH.0000000000001854., 2018

Nikolic D, Birch AA, Panerai RB, Simpson DM. Assessing cerebral blood flow control from variability in blood pressure and arterial CO₂ levels. *Conf Proc IEEE Eng Med Biol Soc* 2015: 1785-8, 2015

Obrig H, Neufang M, Wenzel R, Kohl M, Steinbrink J, Einhupl K, Villringer A. Spontaneous low frequency oscillations of cerebral hemodynamics and metabolism in human adults. *Neuroimage* 12: 623–639, 2000

Ogoh S, Dalsgaard MK, Secher NH, Raven PB. Dynamic blood pressure control and middle cerebral artery mean blood velocity variability at rest and during exercise in humans. *Acta Physiol* 191: 3-14, 2007

Nogueira RC, Bor-Seng-Shu E, Santos MT, Negrrao CE, Teixeira MJ, Panerai RB (2013). Dynamic cerebral autoregulation changes during sub-maximal handgrip maneuver. *Plos One* 8: e70821. <https://doi.org/10.1371/journal.pone.0070821>

Panerai RB, Deverson ST, Mahony P, Hayes P, Evans DH. Effect of CO₂ on dynamic cerebral autoregulation measurement. *Physiol. Meas.* 20: 265–275, 1999

Panerai RB, Dawson SL, Potter JF. Linear and nonlinear analysis of human dynamic cerebral autoregulation. *Am J Physiol Heart Circ Physiol* 277:H1089-H1099., 1999a

Panerai RB, Dawson SL, Eames PJ, Potter JF. Cerebral blood flow velocity response to induced and spontaneous sudden changes in arterial blood pressure. *Am J Physiol Heart Circ Physiol* 280: H2162-2174, 2001

Panerai RB, Eyre M, Potter F. Multivariate modeling of cognitive-motor stimulation on neurovascular coupling: transcranial Doppler used to characterize myogenic and metabolic influences. *Am J Physiol Regul Integr Comp Physiol* 303: R395–R407, 2012

Panerai RB, Salinet ASM, Robinson TG. Contribution of arterial blood pressure and PaCO₂ to the cerebrovascular responses to motor stimulation. *Am J Physiol Heart Circ Physiol* 302: H459–H466., 2012a

Paulson OB, Hasselbalch SG, Rostrup E, Knudsen GM, Pelligrino D. Cerebral blood flow response to functional activation. *J Cereb Blood Flow Metab.* 30: 2-14, 2010

Petridou N, Siero JCW (2017). Laminar fMRI: what can the time domain tell us. Non-BOLD contrast for laminar fMRI in humans: CBF, CBV, and CMRO₂. *NeuroImage* 2017; <http://dx.doi.org/10.1016/j.neuroimage.2017.07.041>, 2017

Placek MM, Wachel P, Iskander DR, Smielewski P, Uryga A, Mielczarek A, Szczepański TA, Kasprovicz M. Applying time-frequency analysis to assess cerebral autoregulation during hypercapnia. *PLoS ONE* 12: e0181851. <https://doi.org/10.1371/journal.pone.0181851>, 2017

Powers WJ. Cerebral hemodynamics in ischemic cerebrovascular disease. *Ann Neurol* 29: 231-240, 1991

Reinhard M, Rutsch S, Lambeck J, Wihler C, Czosnyka M, Weiller C, Hetzel A. Dynamic cerebral autoregulation associates with infarct size and outcome after ischemic stroke. *Acta Neurol Scand.* 125: 156-62, 2012

Rivera-Lara L, Zorrilla-Vaca A, Geocadin R, Zia W, Healy W, Thompson R, Smielewski P, Czosnyka M, Hogue CW. Predictors of outcome with cerebral autoregulation monitoring: a systematic review and meta-analysis. *Crit Care Med* 45: 695–704, 2017

- Robbins PA, Conway J, Cunningham DA, Khamnei S, Paterson DJ. A comparison of indirect methods for continuous estimation of arterial PCO₂ in men. *J Appl Physiol* **68**: 1727–1731, 1990.
- Salinet ASM, Robinson TG, Panerai RB. Active, passive and motor imagery paradigms: component analysis to assess neurovascular coupling. *J Appl Physiol* **114**: 1406-1412, 2013
- Schmid F, Barrett MJP, Jenny P, Weber B. Vascular density and distribution in neocortex. *NeuroImage* 2017; <http://dx.doi.org/10.1016/j.neuroimage.2017.06.046>, 2017
- Serrador JM, Sorond FA, Vyas M, Gagnon M, Iloputaife ID, Lipsitz LA. Cerebral pressure-flow relations in hypertensive elderly humans: transfer gain in different frequency domains. *J Appl Physiol* **98**: 151–159, 2005.
- Smirl JD, Hoffmann K, Tzeng Y-C, Hansen A, Ainslie PN. Methodological comparison of active- and passive-driven oscillations in blood pressure; implications for the assessment of cerebral pressure-flow relationships. *J Appl Physiol* **119**: 487–501, 2015.
- Sohn YH. Cerebral hemodynamic changes induced by sympathetic stimulation tests. *Yonsei Med J* **39**: 322–327, 1998.
- Spronck B, Martens EGHJ, Gommer ED, van de Vosse FN. A lumped parameter model of cerebral blood flow control combining cerebral autoregulation and neurovascular coupling. *Am J Physiol Heart Circ Physiol* **303**: H1143-H1153, 2012
- Zhang R, Zuckerman JH, Giller CA, Levine BD. Transfer function analysis of dynamic cerebral autoregulation in humans. *Am J Physiol*. **274**: H233-41, 1998.
- Zhang R, Witkowski S, Fu Q, Claassen JAHR, Levine BD. Cerebral hemodynamics after short- and long-term reduction in blood pressure in mild and moderate hypertension. *Hypertension* **49**: 1149–1155, 2007.
- Zhang R, Behbehani K, Levine BD. Dynamic pressure–flow relationship of the cerebral circulation during acute increase in arterial pressure. *J Physiol* **587**: 2567–2577, 2009

Table 1 Hemodynamic baseline variables and results of transfer function estimates at baseline and at elbow movement over 5 minutes at a frequency of 1 Hz.

	Baseline		Elbow movement	
Mean Blood Pressure (mm Hg)	83 ± 2.6		87 ± 3.4†	
Mean ETCO ₂ (mm Hg)	44.6 ± 0.60		41.7 ± 0.68†	
Mean Heart rate (beats/minute)	65 ± 1.8		74 ± 2.2†	
Mean CBFV right (cm/s)	65 ± 2.4		69 ± 2.4†	
Mean CBFV left (cm/s)	64 ± 2.2		70 ± 2.6†	
Mean CVR right (mm Hg/cm/s)	1.35 ± 0.06		1.28 ± 0.06†	
Mean CVR left (mm Hg/cm/s)	1.32 ± 0.06		1.31 ± 0.06	
Mean Coherence	Right	Left	Right	Left
- Very low frequency	0.46 ± 0.03	0.48 ± 0.03	0.46 ± 0.03	0.38 ± 0.04
- Low frequency	0.77 ± 0.02	0.75 ± 0.02	0.72 ± 0.03	0.67 ± 0.05
- High frequency	0.70 ± 0.02	0.68 ± 0.03	0.70 ± 0.03	0.65 ± 0.03
Mean Gain (in cm/s/mmHg)				
- Very low frequency	0.24 ± 0.05	0.23 ± 0.3	0.26 ± 0.04	0.19 ± 0.05
- Low frequency	0.62 ± 0.05	0.63 ± 0.05	0.54 ± 0.06	0.53 ± 0.06
- High frequency	0.70 ± 0.07	0.65 ± 0.07	0.65 ± 0.08	0.59 ± 0.08
Mean Gain (in %cm/s/mmHg)				
- Very low frequency	0.37 ± 0.07	0.29 ± 0.05	0.35 ± 0.06	0.26 ± 0.06
- Low frequency	0.97 ± 0.07	1.01 ± 0.09	0.73 ± 0.08†	0.76 ± 0.06†
- High frequency	1.02 ± 0.10	0.99 ± 0.09	0.93 ± 0.10	0.98 ± 0.10
Mean Phase (radians)				
- Very low Frequency	1.03 ± 0.05	1.10 ± 0.06	0.81 ± 0.07†	0.82 ± 0.10*
- Low frequency	0.70 ± 0.06	0.67 ± 0.05	0.77 ± 0.05	0.70 ± 0.05
- High frequency	0.28 ± 0.10	0.28 ± 0.10	0.37 ± 0.11	0.28 ± 0.11

Values are mean ± SE


ETCO₂, end-tidal carbon dioxide; CBFV, cerebral blood flow velocity; CVR, cerebrovascular resistance;

ANOVA: *p ≤ 0.05; †p ≤ 0.01

Figures

Figure 1

Captation: Time course (group averaged) of blood pressure (BP), end-tidal carbon dioxide (ETCO2), and cerebral blood flow velocity (CBFV) in both middle cerebral arteries (MCA).



Legend: For simplicity, bars indicate the maximum standard error of mean (SE) achieved in the whole epoch (baseline or elbow movement); the SE bar for CBFV represents the SE in both MCAs. The movement epoch is indicated by the dashed perpendicular lines.