

Physiological Research Pre-Press Article

Sex Differences in Blood Pressure of Aged Ren-2 Transgenic Rats

H. RAUCHOVÁ, S. HOJNÁ, M. KADLECOVÁ, I. VANĚČKOVÁ, J. ZICHA

Institute of Physiology of the Czech Academy of Sciences, Prague, Czech Republic

Running title: Blood Pressure and Sex Differences in Ren-2 TGR

Corresponding author

H. Rauchová, Institute of Physiology v.v.i., Czech Academy of Sciences, Vídeňská 1083,
14220 Praha 4, Czech Republic

Fax: ++420 241 062 488.

E-mail: Hana.Rauchova@fgu.cas.cz

Summary

Sex-related differences were observed not only in human but also in experimental hypertension. The aim of our study was to compare blood pressure (BP) of aged male and female heterozygous transgenic rats (TGR) harboring Ren-2 mouse gene, with their normotensive Hannover Sprague-Dawley (HanSD) controls. At the age of 9 months, systolic (SBP) and diastolic blood pressure (DBP) were measured by a direct puncture of carotid artery in rats awaking from isoflurane anesthesia. Thiobarbituric acid-reactive species (TBARS) formation was monitored as indicator of lipid peroxidation damage in heart, kidney and liver, whereas intracellular content of reduced glutathione was determined in the same organs as the main intracellular antioxidant. Furthermore, plasma triglycerides and total cholesterol as well as high-density lipoprotein (HDL) and low-density lipoprotein (LDL) fractions of cholesterol were measured. As compared to HanSD rats, we found significantly elevated BP only in male TGR (MAP: 123 ± 1 vs. 171 ± 5 , SBP: 150 ± 2 vs. 208 ± 7 , and DBP: 99 ± 3 vs. 140 ± 4 mm Hg), but not between TGR and HanSD females, which were both normotensive. We also did not find any significant differences in TBARS and reduced glutathione in the three above mentioned organs as well as in plasma cholesterol or its HDL and LDL fractions between transgene-negative HanSD and TGR animals of either sex. However, we found significant sex differences in TBARS, glutathione and plasma lipids in both rat strains. Our results confirmed that aged TGR exhibit a marked sexual BP dimorphism, which does not seem to be dependent on oxidative stress or abnormal cholesterol metabolism.

Key words

Hypertension • Thiobarbituric acid-reactive species (TBARS) • Reduced glutathione • Total plasma cholesterol • HDL and LDL cholesterol fractions

Introduction

Over the years plenty of data on blood pressure (BP) has been gathered in men and women. Generally, aging is usually associated with an increase in BP and hypertension becomes the most common chronic disease in the world. Men have higher incidence of hypertension compared with age-matched women until the onset of menopause. Prominent sex-related differences in hypertension were observed repeatedly not only in humans but also in experimental animals such as dog, rabbit, mouse or rat and even in birds (Blank *et al.* 2016, Lerman *et al.* 2019, Sandberg and Ji 2012). Animal models are inevitable for both the understanding hypertension pathogenesis and possible novel treatment options. Rats are often used as hypertension animal model. There are several rat models in which sex differences in BP were described – genetic hypertension in spontaneously hypertensive rats (Maris *et al.* 2005), NO-deficient hypertension induced by chronic L-NAME administration (Sainz *et al.* 2004) or salt hypertension in salt-sensitive Dahl rats (Zicha *et al.* 2012). There is an interesting model of hypertension in transgenic Ren-2 rats (TGR), in which mouse Ren-2 renin gene was inserted into the genome of normotensive Hannover Sprague-Dawley rats (HanSD). The overexpression of mouse renin gene induces the development of angiotensin II-dependent hypertension with endogenous activation of renin-angiotensin system (RAS), which plays a central role in blood pressure control and its long-term dysregulation causes sustained BP elevation. Therefore the pharmacological RAS targeting is widely used in the clinical hypertension treatment and permanently tested in different experimental trials (Laurent 2017).

For a better understanding on the role of RAS in the pathogenesis of hypertension we can use TGR as a well-defined monogenetic model of hypertension (Mullins *et al.* 1990). Homozygous TGR display severe malignant hypertension accompanied by organ damage and

they die at 7-13 weeks of age (Vernerová *et al.* 2009). On the other hand, heterozygous TGR develop a less severe hypertension, which allows long-term studies (Vaněčková *et al.* 2011). It was described that the established hypertension in male TGR is maintained or only a slight BP decrease occurs in the late adulthood (Lee *et al.* 1998, Springate *et al.* 1992, Vaněčková *et al.* 2011). In contrast, BP decline in female TGR can even yield normotensive values (Cargnelli *et al.* 1998, Lee *et al.* 1998, Vaněčková *et al.* 2011).

The aim of our study was to search for sex differences a) in blood pressure of 9-month-old male and female TGR, b) in the content of TBARS (oxidative stress marker) or reduced glutathione (intracellular antioxidant) in heart, kidney and liver, and c) in plasma lipids (triglycerides and cholesterol).

Material and Methods

Animals

Adult male and female heterozygous (mRen-2)²⁷ transgenic rats (TGR) aged 9 months were housed at 23 °C under a 12 h light/dark cycle periods, fed a standard rat chow Altromin and given to tap water *ad libitum*. Transgene-negative Hannover Sprague-Dawley rats (HanDS) served as controls. At the end of the experiment blood pressure was measured. After blood collection, kidney, heart and liver were excised and used for tissue analysis. Plasma samples (in the EDTA presence) were prepared and stored at -80 °C until further analysis.

All the procedures and experimental protocols were performed in accordance with guidelines and practice established by the *Ethical Committee of the Institute of Physiology*

CAS, conformed to the *European Convention on Animal Protection and Guidelines on Research Animal Use*.

Measurement of blood pressure

Mean arterial (MAP), systolic (SBP) and diastolic (DBP) blood pressure and heart rate were measured by a direct puncture of carotid artery under isoflurane anesthesia (2.5 % isoflurane) and in awaking animals (0.5 % isoflurane). To eliminate the influence of circadian blood pressure (BP) variation, the measurements were always done approximately at the same time of day (between 8:00 and 10:00 a.m.).

Determination of thiol concentration

The intracellular content of reduced glutathione (GSH) in heart, kidney and liver was determined immediately in fresh tissues according to the methods described earlier (Ellman 1959). Briefly, the tissue samples were homogenized in 3 % sulfosalicylic acid and 10 % homogenates were centrifuged for 10 min at 3000 g. A portion of the supernatant was mixed with 0.02 M 5, 5'-dithiobis-(2-nitrobenzoic acid) in 0.1 M phosphate buffer (pH 8). The absorbance of a colored product was read at 412 nm, the concentration of GSH was calculated from calibration curve prepared by serial dilution of 1 mM stock solution. The results were expressed as $\mu\text{mol GSH/g tissue}$.

Measurement of lipid peroxidation

Lipid peroxidation in the samples was monitored by measuring thiobarbituric acid-reactive substances (TBARS) formation (Ohkawa *et al.* 1979). The frozen-thawed 10 % homogenates were incubated with thiobarbituric and acetic acid at 95 °C for 45 min.

Absorbance was measured at 535 nm using Tecan Infinite M200 multimode microplate spectrofluorometer. The results were expressed as nmol of TBARS/mg of protein.

Biochemical parameters

Folin method was used for the determination of protein concentration using bovine serum albumin as standard (Lowry *et al.* 1951).

The concentrations of plasma triglycerides (TG), total cholesterol (TC), and high-density lipoprotein-cholesterol (HDL) were estimated using commercial kits (Pliva-Lachema Diagnostika, Brno, Czech Republic). Low-density lipoprotein-cholesterol (LDL) was estimated indirectly using the formula: $LDL = TC - (TG/5) - HDL$.

Results

Figure 1 shows systolic (SBP), mean arterial (MAP) and diastolic (DBP) blood pressures in 9-month-old male and female TGR and their HanSD controls which were recorded in awaking animals. In TGR females, the blood pressures were decreased to normotensive values seen in HanSD females. On the other hand, older TGR males had still a considerable hypertension, their SBP and DBP being substantially elevated compared to their HanSD controls.

Body weights were higher in male but not in female TGR as compared to sex-matched HanSD animals (Table 1). The absolute weights of hearts and kidneys were always greater in males than in females of both genotypes, but there were no significant differences in the relative weights of hearts and kidneys between TGR and their sex- and age-matched HanSD

controls. Similarly, 9-month-old male TGR had greater absolute liver weight than HanSD rats, but their relative liver weight was not increased.

To evaluate the degree of organ damage by lipid peroxidation we measured TBARS concentrations as an indirect marker of lipid peroxidation in the heart, kidney and liver (Table 2). We did not find any significant difference in TBARS concentrations in the particular tissues between HanSD and TGR rats, but we found significantly lower TBARS concentrations in all examined organs of male rats, irrespective of the genotype.

We observed the lowest concentration of thiol groups in the heart and the highest one in the liver (Table 3). Concentrations of thiol groups were not significantly different between TGR and HanSD animals. However, male rats of both genotypes had significantly higher heart and liver SH concentration but a significantly lower kidney SH concentration than corresponding females.

Male rats of both genotypes had significantly higher values of serum triglycerides (TG), total cholesterol (TC) and low-density lipoprotein-cholesterol (LDL) as well as the ratio between total cholesterol and its HDL fraction (TC/HDL, atherogenic index) than corresponding females (Table 4). However, there were no significant differences in the above parameters between TGR and HanSD controls.

Discussion

TGR represents a model of hypertension with a well-defined genetic background. We confirmed a marked sexual dimorphism in this model because our data obtained in awaking rats indicated BP elevation only in TGR males, whereas BP of TGR females was close to that of HanSD females. Sex-related differences in BP values were reported earlier in this strain.

Our SBP values correspond well with previous two long-term studies (lasting 10-12 months) that followed both male and female TGR in comparison with their sex- and age-matched HanSD controls (Lee *et al.* 1996, Vaněčková *et al.* 2011). In both mentioned studies maximal SBP values were higher and the phase of established hypertension was longer in males than females. Cargnelli *et al.* (1998) showed that the established period of hypertension was followed by a progressive decrease of SBP towards the normal values in 35-week-old heterozygous females TGR, while SBP values of HanSD females were unaffected by ageing. Springate *et al.* (1994) described diminished SBP in 8-month-old male TGR compared to 2- and 4-month-old rats of this strain. We extended our study by measuring of MAP and DBP in two different conditions, i.e. under deep isoflurane anesthesia (data not shown) and in animals awaking from this anesthesia. The differences in BP were more pronounced in awaking animals.

TGR with elevated endogenous angiotensin II were sporadically described as a suitable model of spontaneous liver fibrosis and portal hypertension (non-alcoholic fatty liver disease), which is associated with hepatic mitochondrial oxidative damage and impaired mitochondrial fatty acid oxidation (Klein *et al.* 2019, Wei *et al.* 2008, 2009). However, our group of aged TGR males showed neither the higher relative liver weight nor the increased level of TBARS in liver.

Oxidative or nitrosative stress, which represents the imbalance between production and elimination of reactive oxygen or nitrogen species, is often considered as an important factor contributing to the pathogenesis of different forms of hypertension (Guzik and Touyz 2017, Reckelhoff *et al.* 2019). To examine the role of oxidative damage, we measured TBARS levels in three different organs (liver, heart and kidney). Vokurková *et al.* (2015) disclosed higher concentration of TBARS in renal cortex in 3-month-old TGR males but this difference was not confirmed by the parallel measurements of conjugated dienes or oxidative

index. However, we did not find significant differences in renal, cardiac or liver levels of TBARS between TGR and HanSD rats of either sex. Kopkan *et al.* (2009) reported increased levels of malondialdehyde in the whole kidney and left heart ventricle of 3-month-old male TGR and significantly reduced levels of malondialdehyde after the chronic administration of O₂⁻ scavenger tempol or NADPH oxidase inhibitor apocynin but this malondialdehyde decrease was not associated with any SBP change. Interestingly, in spite of higher blood pressure in our TGR males we found significantly lower concentrations of TBARS than in females for which we cannot offer a plausible explanation.

The main intracellular antioxidant reduced glutathione (GSH) plays important roles in peroxide detoxification, recycling of vitamins C and E, cysteine storage and other biochemical reactions (Aquilano *et al.* 2014, Gould and Pazdro 2019). We found sex-different GSH concentrations in all examined tissues (heart, kidney, liver). Males had always higher GSH concentrations in heart and liver but lower values in the kidney. On the other hand, GSH concentrations were not influenced by rat genotype. Similar GSH concentrations were also obtained in 3-month-old TGR males (Vokurková *et al.* 2015).

All parameters of lipid metabolism (with the exception of HDL cholesterol) were significantly higher in older males than females. This is a rather surprising finding because there are several reports of higher plasma cholesterol levels in female than male Sprague Dawley or Lewis rats of various age (Lee *et al.* 2008, Borbélyová *et al.* 2017). On the other hand, there were no differences between HanSD and TGR animals of either sex. To our knowledge, there are no data on alterations of lipid metabolism in TGR. The only relevant study (Vettor *et al.* 1994) reported similar fasting levels of plasma triglycerides in TGR and HanSD rats aged 5 months.

In summary, the present study shows a sexual dimorphism in blood pressure of heterozygous TGR. Additionally, we did not find any corresponding difference in TBARS

concentrations and reduced glutathione levels between transgene-negative normotensive (HanSD) and hypertensive Ren-2 transgenic rats (TGR).

Conflict of Interest

There is no conflict of interest.

Acknowledgements

The study was supported by an international research grant GACR 19-08260J and by an institutional support (RV0:67985823). A part of research results was presented on FEPS meeting in Bologna, Italy (Rauchová *et al.* 2019).

References

- AQUILANO K, BALDELLI S, CIRIOLO MR: Glutathione: new roles in redox signaling for an old antioxidant. *Front Pharmacol* **5**: 196, 2014.
- BLENCK CL, HARVEY PA, RECKELHOFF JF, LEINWAND LA: The importance of biological sex and estrogen in rodent models of cardiovascular health and disease. *Circ Res* **118**: 1294-1312, 2016.
- BORBÉLYOVÁ V, DOMONKOS E, BÁBÍČKOVÁ J, TÓTHOVÁ Ľ, KAČMÁROVÁ M, ULIČNÁ O, OSTATNÍKOVÁ D, HODOSY J, CELEC P: Does long-term androgen deficiency lead to metabolic syndrome in middle-aged rats? *Exp Gerontol* **98**:38-46, 2017.

- CARGNELLI G, ROSSI GP, PESSINA AC, LUCIANI S, DEBETTO P, GANTEN D, PETERS J, BOVA S: Changes of blood pressure and aortic strip contractile responses to ET-1 of heterozygous female transgenic rats, TGR(mRen2)27. *Pharmacol Res* **37**: 207-211, 1998.
- ELLMAN GL: Tissue sulfhydryl groups. *Arch Biochem Biophys* **82**: 70-77, 1959.
- FORTEPIANI LA, RECKELHOFF JF: Role of oxidative stress in the sex differences in blood pressure in spontaneously hypertensive rats. *J Hypertens* **23**: 801-805, 2005.
- GOULD RL, PAZDRO R: Impact of supplementary amino acids, micronutrients, and overall diet on glutathione homeostasis. *Nutrients* **11**: 1056, 2019.
- GUZIK TJ, TOUYZ RM: Oxidative stress, inflammation, and vascular aging in hypertension. *Hypertension* **70**: 660-667, 2017.
- KASPER SO, CARTER CS, FERRARIO CM, GANTEN D, FERDER LF, SONNTAG WE, GALLAGHER PE, DIZ DI: Growth, metabolism, and blood pressure disturbances during aging in transgenic rats with altered brain renin-angiotensin systems. *Physiol Genomics* **23**: 311-317, 2005.
- KLEIN S, KLEINE CE, PIEPER A, GRANZOW M, GAUTSCH S, HIMMIT M, KAHRMANN K, SCHIERWAGEN R, USCHNER FE, MAGDALENO F, NAOUM ME, KRISTIANSEN G, WALTHER T, BADER M, SAUERBRUCH T, TREBICKA J: TGR(mREN2)27 rats develop non-alcoholic fatty liver disease-associated portal hypertension responsive to modulations of Janus-kinase 2 and Mas receptor. *Sci Rep* **9**: 11598, 2019.
- KOPKAN L, HUSKOVÁ Z, VAŇOURKOVÁ Z, THŮMOVÁ M, ŠKAROUPKOVÁ P, MALÝ J, KRAMER HJ, DVOŘÁK P, ČERVENKA L: Reduction of oxidative stress does not attenuate the development of angiotensin II-dependent hypertension in Ren-2 transgenic rats. *Vascul Pharmacol* **51**: 175-181, 2009.

- LAURENT S: Antihypertensive drugs. *Pharmacol Res* **124**: 116-125, 2017.
- LEE CE, KANG JS, KIM KI: Effects of gender, gonadectomy and sex hormones on growth and plasma cholesterol level in rats. *Ann Nutr Metab* **53**:1-5, 2008.
- LEE MA, BÖHM M, PAUL M, BADER M, GANTEN U, GANTEN D: Physiological characterization of the hypertensive transgenic rat TGR(mREN2)27. *Am J Physiol* **270**: E919-E929, 1996.
- LERMAN LO, KURTZ TW, TOUYZ RM, ELLISON DH, CHADE AR, CROWLEY SD, MATTSON DL, MULLINS JJ, OSBORN J, EIRIN A, RECKELHOFF JF, IADECOLA C, COFFMAN TM: Animal models of hypertension: A scientific statement from the American Heart Association. *Hypertension* **73**: e87-e120, 2019.
- LOWRY OH, ROSEBROUGH NJ, FARR AL, RANDALL RJ: Protein measurement with the Folin phenol reagent. *J Biol Chem* **193**: 265-275, 1951.
- MARIS ME, MELCHERT RB, JOSEPH J, KENNEDY RH: Gender differences in blood pressure and heart rate in spontaneously hypertensive and Wistar-Kyoto rats. *Clin Exp Pharmacol Physiol* **32**: 35-39, 2005.
- MULLINS JJ, PETERS J, GANTEN D: Fulminant hypertension in transgenic rats harbouring the mouse Ren-2 gene. *Nature* **344**: 541-544, 1990.
- NAUTIYAL M, KATAKAM PV, BUSIJA DW, GALLAGHER PE, TALLANT EA, CHAPPELL MC, DIZ DI: Differences in oxidative stress status and expression of MKP-1 in dorsal medulla of transgenic rats with altered brain renin-angiotensin system. *Am J Physiol Regul Integr Comp Physiol* **303**: R799-R806, 2012.
- OHKAWA H, OHISHI N, YAGI K: Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem* **95**: 351-358, 1979.

- RAUCHOVÁ H, HOJNÁ S, KADLECOVÁ M, VANĚČKOVÁ I, ZICHA J: Sex-related differences of blood pressure in older Ren-2 transgenic rats. *Acta Physiol* **227** (Suppl 718): 173, 2019.
- RECKELHOFF JF, ROMERO DG, YANES CARDOZO LL: Sex, oxidative stress, and hypertension: insights from animal models. *Physiology (Bethesda)* **34**: 178-188, 2019.
- SÁINZ J, OSUNA A, WANGENSTEEN R, DE DIOS LUNA J, RODRÍGUEZ-GÓMEZ I, DUARTE J, MORENO JM, VARGAS F: Role of sex, gonadectomy and sex hormones in the development of nitric oxide inhibition-induced hypertension. *Exp Physiol* **89**:155-162, 2004.
- SANDBERG K, JI H: Sex differences in primary hypertension. *Biol Sex Differ* **3**: 7, 2012.
- SPRINGATE JE, FELD LG, GANTEN D: Renal function in hypertensive rats transgenic for mouse renin gene. *Am J Physiol* **266**: F731-F737, 1994.
- VANĚČKOVÁ I, HUSKOVÁ Z, VAŇOURKOVÁ Z, ČERVENKA L: Castration has antihypertensive and organoprotective effects in male but not in female heterozygous Ren-2 rats. *Kidney Blood Press Res* **34**: 46-52, 2011.
- VERNEROVÁ Z, KUJAL P, KRAMER HJ, BÄCKER A, ČERVENKA L, VANĚČKOVÁ I: End-organ damage in hypertensive transgenic Ren-2 rats: influence of early and late endothelin receptor blockade. *Physiol Res* **58** (Suppl 2): S69-S78, 2009.
- VETTOR R, CUSIN I, GANTEN D, ROHNER-JEANRENAUD F, FERRANNINI E, JEANRENAUD B: Insulin resistance and hypertension: studies in transgenic hypertensive TGR(mREN-2)²⁷ rats. *Am J Physiol* **267**: R1503-R1509, 1994.
- VOKURKOVÁ M, RAUCHOVÁ H, ŘEZÁČOVÁ L, VANĚČKOVÁ I, ZICHA J: NADPH oxidase activity and reactive oxygen species production in brain and kidney of adult male hypertensive Ren-2 transgenic rats. *Physiol Res* **64**: 849-856, 2015.

- WEI Y, CLARK SE, MORRIS EM, THYFAULT JP, UPTERGROVE GM, WHALEY-CONNELL AT, FERRARIO CM, SOWERS JR, IBDAH JA: Angiotensin II-induced non-alcoholic fatty liver disease is mediated by oxidative stress in transgenic TG(mRen2)27(Ren2) rats. *J Hepatol* **49**: 417-428, 2008.
- WEI Y, CLARK SE, THYFAULT JP, UPTERGROVE GM, LI W, WHALEY-CONNELL AT, FERRARIO CM, SOWERS JR, IBDAH JA: Oxidative stress-mediated mitochondrial dysfunction contributes to angiotensin II-induced nonalcoholic fatty liver disease in transgenic Ren2 rats. *Am J Pathol* **174**: 1329-1337, 2009.
- ZICHA J, DOBEŠOVÁ Z, VOKURKOVÁ M, RAUCHOVÁ H, HOJNÁ S, KADLECOVÁ M, BEHULIAK M, VANĚČKOVÁ I, KUNEŠ J: Age-dependent salt hypertension in Dahl rats: fifty years of research. *Physiol Res* **61** (Suppl 1): S35-S87, 2012.

Table 1 General characteristics of female and male Hannover Sprague-Dawley (HanSD) and transgenic rats (TGR) aged 9 months.

Females	HanSD (n=8)	TGR (n=8)
<i>Body weight (BW; g)</i>	353 ± 12	382 ± 8
<i>Heart weight (HW; mg)</i>	1165 ± 24	1253 ± 24
<i>Relative heart weight (HW/BW; mg/g)</i>	3.33 ± 0.12	3.31 ± 0.13
<i>Kidney weight (KW; mg)</i>	2071 ± 46	2272 ± 46*
<i>Relative kidney weight (KW/BW; mg/g)</i>	5.93 ± 0.29	5.97 ± 0.31
Males	HanSD (n=6)	TGR (n=5)
<i>Body weight (BW; g)</i>	690 ± 1 [#]	818 ± 24* [#]
<i>Heart weight (HW; mg)</i>	1756 ± 21 [#]	2130 ± 55* [#]
<i>Relative heart weight (HW/BW; mg/g)</i>	2.55 ± 0.03 [#]	2.61 ± 0.09 [#]
<i>Left ventricular weight (LVW; mg)</i>	1376 ± 34	1720 ± 89*
<i>Relative left ventricular weight (LVW/BW; mg/g)</i>	2.00 ± 0.02	2.12 ± 0.06
<i>Kidney weight (KW; mg)</i>	4289 ± 227 [#]	4762 ± 212 [#]
<i>Relative kidney weight (KW/BW; mg/g)</i>	6.22 ± 0.34	5.83 ± 0.26
<i>Liver weight (LW; mg)</i>	23757 ± 1921	28376 ± 2163*
<i>Relative liver weight (LW/BW; mg/g)</i>	34.5 ± 2.9	34.7 ± 2.3

All values are mean ± SEM, n, number of animals is given in parentheses. Significantly different: * $P \leq 0.05$ vs. HanSD, [#] $P \leq 0.05$ vs. females.

Table 2 Thiobarbituric acid-reactive substances in heart, kidney and liver of female and male Hannover Sprague-Dawley (HanSD) and transgenic rats (TGR) aged 9 months.

Females	HanSD (n=8)	TGR (n=8)
<i>Heart</i> ($\mu\text{mol/mg protein}$)	15.93 \pm 0.90	15.98 \pm 1.01
<i>Kidney</i> ($\mu\text{mol/mg protein}$)	28.79 \pm 2.20	28.96 \pm 1.34
<i>Liver</i> ($\mu\text{mol/mg protein}$)	13.46 \pm 0.98	15.13 \pm 1.77
Males	HanSD (n=6)	TGR (n=5)
<i>Heart</i> ($\mu\text{mol/mg protein}$)	11.70 \pm 1.52 [#]	11.74 \pm 1.23 [#]
<i>Kidney</i> ($\mu\text{mol/mg protein}$)	18.55 \pm 0.11 [#]	20.36 \pm 1.21 [#]
<i>Liver</i> ($\mu\text{mol/mg protein}$)	8.80 \pm 0.45 [#]	8.52 \pm 2.19 [#]

All values are mean \pm SEM, n, number of animals is given in parentheses Significantly different: [#] $P \leq 0.05$ vs. females.

Table 3 Concentration of thiol groups in heart, kidney and liver of female and male Hannover Sprague-Dawley (HanSD) and transgenic rats (TGR) aged 9 months.

Females	HanSD (n=8)	TGR (n=8)
<i>Heart</i> ($\mu\text{mol/g}$)	1.73 \pm 0.08	1.54 \pm 0.05
<i>Kidney</i> ($\mu\text{mol/g}$)	4.10 \pm 0.17	4.24 \pm 0.17
<i>Liver</i> ($\mu\text{mol/g}$)	4.89 \pm 0.39	4.82 \pm 0.28
Males	HanSD (n=6)	TGR (n=5)
<i>Heart</i> ($\mu\text{mol/g}$)	2.13 \pm 0.08 [#]	2.35 \pm 0.05 [#]
<i>Kidney</i> ($\mu\text{mol/g}$)	3.19 \pm 0.18 [#]	3.36 \pm 0.15 [#]
<i>Liver</i> ($\mu\text{mol/g}$)	8.58 \pm 0.29 [#]	8.03 \pm 0.56 [#]

All values are mean \pm SEM, n, number of animals is given in parentheses. Significantly different: [#] $P \leq 0.05$ vs. females.

Table 4 Plasma lipids in female and male Hannover Sprague-Dawley (HanSD) and transgenic rats (TGR) aged 9 months.

Females	HanSD (n=8)	TGR (n=8)
<i>Triglycerides (mmol/l)</i>	0.26 ± 0.01	0.26 ± 0.03
<i>Total cholesterol (mmol/l)</i>	1.96 ± 0.14	1.84 ± 0.10
<i>HDL (mmol/l)</i>	1.49 ± 0.15	1.44 ± 0.08
<i>LDL (mmol/l)</i>	0.42 ± 0.09	0.35 ± 0.04
<i>TC/HDL ratio</i>	1.36 ± 0.11	1.28 ± 0.03
Males	HanSD (n=6)	TGR (n=5)
<i>Triglycerides (mmol/l)</i>	0.85 ± 0.06 [#]	0.79 ± 0.15 [#]
<i>Total cholesterol (mmol/l)</i>	2.37 ± 0.08 [#]	2.59 ± 0.18 [#]
<i>HDL (mmol/l)</i>	1.28 ± 0.03	1.37 ± 0.12
<i>LDL (mmol/l)</i>	0.91 ± 0.04 [#]	1.06 ± 0.13 [#]
<i>TC/HDL ratio</i>	1.84 ± 0.02 [#]	1.91 ± 0.12 [#]

All values are mean ± SEM, n, number of animals is given in parentheses. Significantly different: [#] $P \leq 0.05$ vs. females. HDL, high-density lipoprotein-cholesterol; LDL, low-density lipoprotein-cholesterol; TC, total cholesterol

Legend to figure:

Fig. 1. Mean (MAP), systolic (SBP) and diastolic (DBP) blood pressure in awaking female (F) and male (M) Hannover Sprague-Dawley (HanSD) and transgenic rats (TGR) aged 9 months.

All values are mean \pm SEM. Significantly different: * $P \leq 0.05$ vs. HanSD, # $P \leq 0.05$ vs. females.

