

Factors affecting the function of the mitochondrial membrane permeability transition pore and their role in evaluation of calcium retention capacity values

Zdeněk Drahota^{1,2}, René Endlicher^{1,3}, Otto Kučera¹, David Rychtřmoc¹, Zuzana Červinková¹

¹Department of Physiology, Faculty of Medicine in Hradec Kralove, Charles University, Czech Republic

²Institute of Physiology, Czech Academy of Sciences, Prague, Czech Republic

³Department of Anatomy, Faculty of Medicine in Hradec Kralove, Charles University, Czech Republic

Corresponding author:

Zuzana Červinková

Charles University

Faculty of Medicine in Hradec Králové

Department of Physiology

Šimkova 870

500 03 Hradec Králové

Czech Republic

E-mail: wolff@lfhk.cuni.cz

Phone: +420 495 816 432

ORCID:

Zdenek Drahota 0000-0003-2117-6809

Rene Endlicher 0000-0002-5781-4115

Otto Kucera 0000-0002-5251-5343

David Rychtřmoc 0000-0003-4675-1523

Zuzana Cervinkova 0000-0002-3908-290X

Short title:

Evaluation of calcium retention capacity

Summary

Values of the calcium retention capacity (CRC) of rat liver mitochondria are highly dependent on the experimental conditions used. When increasing amounts of added calcium chloride are used (1.25-10 nmol), the values of the CRC increase 3-fold. When calcium is added in 75 s intervals, the CRC values increase by 30 % compared with 150 s interval additions. CRC values are not dependent on the calcium/protein ratio in the measured sample in our experimental design. We also show that a more detailed evaluation of the fluorescence curves can provide new information about mitochondrial permeability transition pore opening after calcium is added.

Key words

Liver mitochondria · Calcium Retention Capacity · Mitochondrial permeability transition pore

Introduction

The mitochondrial permeability transition pore (MPTP) activated by calcium ions plays an important role in the development of necrotic and apoptotic processes (Smith *et al.* 2012). This pore was discovered sixty-seven years ago (Raaflaub 1953) as a nonspecific pore in the inner mitochondrial membrane that enables penetration of molecules with molecular weights less than 1.5 kD. Its opening results in dissipation of the mitochondrial membrane potential and inhibition of ATP generation. Many studies have confirmed the existence of MPTP and described its properties and functions, e.g., its activation by calcium, phosphate, fatty acids and prooxidants; its inhibition by Mg ions and ADP; and the role of cyclophilin D and cyclosporine in calcium-dependent pore opening (Halestrap *et al.* 1997, Carafoli *et al.* 2001, Halestrap 2009). However, the molecular structure of the pore has not yet been elucidated. Many mitochondrial proteins have been proposed as components of its structure (Halestrap 2009, Bernardi 2013, Alavian *et al.* 2014), but none have been confirmed by molecular genetic studies (He *et al.* 2017).

All basic information about MPTP function, regulation, tissue-specificity (Panov *et al.* 2007, Endlicher *et al.* 2009), age-specificity (Drahota *et al.* 2012b) and sex-specificity (Milerová *et al.* 2016) and its participation in the pathogenesis of many diseases was obtained through the assessments with spectrophotometric measurements of mitochondrial swelling induced by calcium ions. We have improved this graphical method using derivation of the swelling curves. After swelling curve derivation, two further parameters, in addition to the extent of swelling, can be acquired in a numeric form: the maximum swelling rate and the time at which the maximum swelling rate was reached after calcium addition (Drahota *et al.* 2012a).

More recently, a new approach based on a different principle has been used to evaluate the function and regulation of the MPTP in various mitochondrial preparations. The rationale

of the method is based on the determination of the calcium retention capacity (CRC) value using the membrane impermeable fluorophore Calcium Green-5N (Ichas *et al.* 1997, Fontaine *et al.* 1998a). Calcium retention capacity indicates the number of calcium ions that must accumulate in mitochondria to induce pore opening. By this procedure, all information obtained from swelling measurements was confirmed. In addition, new data about pore regulation and function in various tissues and various pathological processes were obtained. However, this method of measuring CRC values has not yet been well standardized; therefore, the data obtained by various laboratories cannot be easily compared. According to our measurements, the CRC values are highly dependent on the composition of the incubation medium (Endlicher *et al.* 2019). In this communication, we describe that the CRC values are also dependent on the experimental protocol used. The main factors that affect the detection of CRC values are the calcium amount used for titration, and time interval between calcium additions. We also demonstrate that additional information about the process of pore opening during titration with calcium may be obtained from the resultant fluorometric curves.

Methods

Chemicals

All chemicals, unless otherwise stated, were of analytical grade and obtained from Sigma-Aldrich (Darmstadt, Germany). Calcium Green-5N was obtained from Thermo Fisher Scientific Inc. (Waltham, MA, USA).

Animals

Male Wistar rats (215±15 g) were obtained from Velaz (Lysá nad Labem, Czech Republic). The rats were housed at 23±1°C and 55±10 % humidity, with air exchange 12-14

times/h and a 12 h light-dark cycle period. The animals had free access to a standard laboratory diet (ST-1, Velaz, Czech Republic) and tap water. All animals received care according to the guidelines set by the Animal-Welfare Body of Charles University, Czech Republic, and the EU Directive 2010/63/EU for animal experiments. Protocols complied with ARRIVE guidelines. The animals were sacrificed during deep general inhalation anaesthesia by exsanguination from the aortic bifurcation. The livers were removed, washed in the cold isolation medium described below, and cut into small pieces.

Isolation of mitochondria

Liver mitochondria were isolated as previously described (Bustamante *et al.* 1977). The washed and cut 3 g liver tissue was homogenized at 0°C by a Teflon-glass homogenizer in an isolation medium containing 220 mM D-mannitol, 70 mM sucrose, 2 mM HEPES, 0.2 mM EGTA, and 0.5 g of fatty acid-free bovine serum albumin (BSA) per litre at pH 7.2. The 10 % homogenate was centrifuged for 4 min at 830 g, and the resulting supernatant was centrifuged for 15 min at 5,200 g. The mitochondrial sediment was washed during two 10 min centrifugations (at 11,200 and at 13,000 g) in isolation medium lacking EGTA and resuspended in the same medium to a final volume of 3 ml. Isolated mitochondria were stored at 0°C. The calcium retention capacity was measured immediately after isolation.

Determination of mitochondrial proteins

The mitochondrial protein concentration was determined using the Bradford method with bovine serum albumin as a standard (Bradford 1976).

Measurement of calcium retention capacity

The mitochondrial retention capacity for calcium was evaluated using the membrane-impermeable fluorescent probe Calcium Green-5N on an AMINCO-Bowman Series 2

spectrofluorometer (Thermo Electron Corporation) at an excitation wavelength of 506 nm and emission wavelength of 592 nm. Measurements were performed at room temperature. A total of 1 ml of medium (125 mM sucrose, 65 mM KCl, and 10 mM HEPES at pH 7.2), 1 μ M Calcium Green-5N, 10 mM succinate, 0.5 μ M rotenone and mitochondria (adjusted to a protein concentration of 0.4 g protein/l) were added. Then, calcium chloride (CaCl_2) was added at the amounts described in the figure captions at an interval of 75 or 150 s. The probe reversibly binds to calcium ions. After each calcium addition, the fluorescence of the probe increased, and then it decreased when the added calcium had accumulated. When the accumulated calcium reached the critical intramitochondrial concentration required for pore opening, it was released, and the fluorescence rose dramatically (Ichas *et al.* 1997, Fontaine *et al.* 1998a).

Statistical analysis

The experiments were performed at least five times; representative results are shown. Values are depicted as the means \pm SD; $p < 0.05$ was set as the threshold for statistical significance. Statistical evaluation was performed using GraphPad Prism 6.01 software (La Jolla, CA, USA). The data were first tested for normality by means of the Kolmogorov-Smirnov test. None of the data followed a Gaussian distribution and thus were all analysed by nonparametric tests. For the evaluation of the effects of different amounts of added calcium during the same interval, we used the Kruskal-Wallis test followed by Dunn's multiple comparisons test. For the comparison of changes between different time intervals and for evaluation of changes in the calcium/protein ratio, an unpaired two-tailed Mann-Whitney test was used.

Results

In this communication, we claim that CRC values are dependent on the experimental conditions used, namely on the amount of calcium added in various portions required for the pore opening, and on the intervals between calcium additions.

When 1.25, 2.5, 5 and 10 nmol CaCl₂ were added in 150 s intervals, the CRC values increased at 2.5 nmol Ca by 147.8 %, at 5 nmol Ca by 226 %, and at 10 nmol Ca by 303.5 % compared with those from 1.25 nmol Ca additions (Fig. 1A, Fig. 2AB, Fig. 3AB).

When different amounts of calcium were added in 75 s intervals (Fig. 2AB, Fig. 3AB), CRC values were significantly increased to 130-140 % at all Ca doses (1.25-10 nmol) compared with those from doses added in 150 s intervals (Table 1).

When the protein concentration in the incubation medium differed, the values of CRC did not change (Fig. 4).

All these data show that the novel method of determining CRC value must be standardized for obtaining reproducible and generally comparable CRC data from different laboratories. This standardization is needed because the CRC values are modified not only by the composition of the incubation medium but also by the experimental protocol used.

We also present data demonstrating that additional information may be acquired from classic CRC curves. When the fluorescence curves from Fig. 1A are evaluated at greater detail, we can see that after the first calcium addition with all calcium amounts used (1.25 - 10 nmol), the increased fluorescence values obtained after calcium addition return quickly back to the basal values (Fig. 1B). This finding indicates that, after all the added calcium had been accumulated in the mitochondria, the pore remained closed (Fig. 1B). The same picture was obtained after the second calcium addition. However, after the third calcium addition (Fig. 1C), there was a decrease in fluorescence to basal values only after lower amounts of calcium were added (1.25, 2.5 and 5 nmol). After the highest amount of 10 nmol was added, there was only a small fluorescence decrease over a short period of time, followed by a period of large

increase, which indicates opening of the pore and release of the accumulated calcium from the intramitochondrial space.

Moreover, another interesting detail can be inferred from Fig. 1A. It is obvious that, after the 8th and 6th additions of the two lowest calcium amounts (1.25 and 2.5 nmol), the calcium had accumulated, and the pore remained closed. However, after a short period of time, the pore was completely opened. At the two highest calcium amounts (5 and 10 nmol), we also observed two possible rates of pore opening. This finding suggests that, after those particular additions, some duration of calcium action was required for complete pore opening by the calcium accumulated in mitochondria (Fig. 1A).

To confirm this time-dependent effect of accumulated calcium in further experiments, we present other data indicating that opening of the pore may have transpired over several phases (Fig. 5). For this experiment, we added increasing calcium amounts (5 - 200 nmol CaCl₂) only once, and we followed the fluorescence changes for a longer period of time. At the lowest doses of calcium added (5 and 10 nmol), we observed a small increase in fluorescence due to the added calcium and a decrease in the fluorescence due to accumulation in the mitochondria. Fluorescence values returned to baseline and remained without change, which indicated that the pore was closed. However, after a long time period (900 to 1200 s), the fluorescence increased, indicating complete pore opening. These data show that even very low concentrations of accumulated calcium considered ineffective during initial pore opening may be effective after a longer period of action. Our data document that the time period required for pore opening after calcium accumulation in mitochondria was shorter when increased calcium concentrations were added (Fig. 5). Curves obtained after the addition of 15, 25 and 50 nmol of calcium reveal two different pore opening rates. Curves with 100 and 200 nmol of calcium show only a small fluorescence decrease followed by an increase in fluorescence, indicating that the pore had fully opened. This figure also demonstrates that our

experiments are performed at the calcium concentration range when Calcium Green is not yet fully saturated by calcium.

As shown in Fig. 5, the MPTP opens by the lowest calcium amount if it can act over a longer period. Additionally, even with higher calcium amounts added, it is obvious that some time passes before the accumulated calcium activates complete pore opening. Therefore, we may conclude that a dose-dependent time period is required before calcium accumulation in the matrix activates complete pore opening. Our data also show that the pore opening may transpire over several phases that are dependent on the amount of calcium in the mitochondrial matrix and on the duration of its action.

Discussion

Data presented in this communication add support to our previous findings indicating that the measurement of CRC values had not been well defined because CRC values depend on more factors than the medium composition used (Endlicher *et al.* 2019). We have shown in this communication that CRC also depends on the calcium amounts used to trigger MPTP opening and on the intervals between calcium additions but not on the calcium/protein ratio in measured preparations. Therefore, for comparability of CRC values described in the literature, it is necessary to provide and consider the factors mentioned above and standardize reporting the experimental conditions in more detail. Obviously, many other factors remain that might affect CRC measurements. The main problem in studies of the MPTP is that this pore has a very important role in the pathogenesis of many diseases, and although we know much about its function and regulation, our knowledge about its molecular structure is still insufficient.

Therefore, in this communication, we focused on elucidating the factors that can modify the CRC values, and we also showed that more data about the mechanism of the MPTP opening may be acquired from fluorescence curves because the fluorescence method is

much more sensitive than the swelling method. We have shown that not only the medium composition but also the experimental protocol used may considerably affect the values of calcium retention capacity (Endlicher *et al.* 2019). Here, we describe two factors that should be taken into account when data from different laboratories are compared. The first factor described is the amount of calcium added to induce pore opening. Some authors prefer high peaks caused by the adding greater amounts of calcium ions, which open the pore more quickly (Fontaine *et al.* 1998b). Some prefer low peaks caused by lower amounts of calcium, which take more time to reach complete pore opening (Pardo *et al.* 2015). The next important factor that must be controlled for is the interval between sequential additions of calcium. We observed that changes in both above-mentioned parameters resulted in very different CRC values. Therefore, it is important to obtain consensus on the experimental protocol for the measurement of CRC values.

However, even more important is the fact that 67 years after MPTP discovery, we still do not know its molecular structure. Discussions regarding whether MPTP is a pore or just an unspecific hole in the mitochondrial membrane continue (Szabo and Zoratti 2014). The titles of many recent reviews point out words such as “fixing a hole” (Di Lisa and Bernardi 2006), “MPTP: a mystery solved?” (Bernardi 2013), “MPTP: Where the known meets unknown” (Juhaszova *et al.* 2008), and “Enjoy the trip: Calcium in Mitochondria Back and Forth” (De Stefani *et al.* 2016).

These reviews demonstrate almost 50 years of unsuccessful work to resolve the problem of the molecular structure of the MPTP. Certainly new strategies and new methods are required. However, before new ideas and methods are accessible, there is still a chance to address remaining problems related to pore function, regulation and specificity, as well as its involvement in multiple pathologies, e.g., cardiovascular diseases (Weiss *et al.* 2003), cancer (Biasutto *et al.* 2010, Bonora and Pinton 2014), neurodegenerative diseases (Rao *et al.* 2014,

Nicholls 2017), diabetes mellitus (Skrha *et al.* 2011), and liver steatosis (Einer *et al.* 2018), by the two methods available. These data will be important for both diagnosis and treatment.

Conflict of interest

The authors declare that there is no conflict of interests regarding the publication of this manuscript.

Acknowledgements

This work was supported by the Charles University Program PROGRES Q40/02 and the Ministry of Education, Youth and Sports Project InterCost LTC17044. Language of this manuscript was corrected by the American Journal Experts, certificate verification key: D826-8D49-62EB-D714-E756.

Text to figures

Fig. 1. CRC values depend on the amount of calcium added to induce pore opening.

Determination of the CRC of isolated rat liver mitochondria incubated in the medium, as described in the Methods, with different amounts of added calcium. A total of 1.25, 2.5, 5 and 10 nmol of calcium chloride were added to the mitochondrial suspension in 150 s intervals. Fluorescence data (a.u. arbitrary unit) were registered in 1 s intervals (A). Evaluation of changes in fluorescence data after the 1st calcium addition at high resolution (B). Evaluation of changes in fluorescence data after the 3rd calcium addition at high resolution (C).

Fig. 2. CRC values depend on the interval between sequential additions of calcium.

Determination of the CRC of isolated rat liver mitochondria with the addition of 1.25 nmol calcium chloride at 150 and 75 s intervals (A) and with the addition of 10 nmol of calcium chloride in 150 and 75 s intervals (B). Fluorescence data are in arbitrary unit (a.u.).

Fig. 3. Statistical evaluation of CRC changes. Statistical evaluation of changes in CRC upon adding different amounts of calcium chloride (1.25, 2.5, 5, and 10 nmol) in 150 or 75 s intervals (A). Statistical evaluation of CRC changes (%) (B).

Fig. 4. CRC values in different calcium/protein ratio. Determination of CRC using different calcium/mitochondrial protein ratios. Amount of calcium added to both mitochondrial preparations was equal (1.25 nmol); intervals between calcium additions were also the same (150 s). Fluorescence is expressed in arbitrary unit (a.u.).

Fig. 5. MPTP opening after adding a single dose of calcium. Determination of fluorescence changes (a.u., arbitrary units) in mitochondrial suspension after adding a single dose of an increasing amount of calcium chloride (5 - 200 nmol). Experimental conditions are as described in the Methods.

Table 1. CRC values determined by adding different amounts of calcium chloride in 150 or 75 s intervals and statistical evaluation of CRC changes.

Ca²⁺ amount per addition	CRC (150 s interval) (nmol/mg prot.)	CRC (75 s interval) (nmol/mg prot.)	CRC ratio 75/150 s intervals (%)
1.25 nmol Ca ²⁺	27.2±3.5	39.1±5.1	142.5±19.6 **
2.5 nmol Ca ²⁺	40.2±6.1	54.2±9.4	134±19.2 *
5 nmol Ca ²⁺	60.7±8.6	77.1±5.1	129.4±16.0 **
10 nmol Ca ²⁺	82.1±12.2	108.3±12.9	130.6±4.3 *



n = 6 - 7; *p<0.05; **p<0.01

References

- ALAVIAN KN, BEUTNER G, LAZROVE E, SACCHETTI S, PARK HA, LICZNEKSKI P, LI H, NABILI P, HOCKENSMITH K, GRAHAM M, PORTER GA, JR., JONAS EA: An uncoupling channel within the c-subunit ring of the F1FO ATP synthase is the mitochondrial permeability transition pore. *Proc Natl Acad Sci U S A* **111**: 10580-10585, 2014.
- BERNARDI P: The mitochondrial permeability transition pore: a mystery solved? *Front Physiol* **4**: 95, 2013.
- BIASUTTO L, AZZOLINI M, SZABO I, ZORATTI M: The mitochondrial permeability transition pore in AD 2016: An update. *Biochim Biophys Acta* **1863**: 2515-2530, 2016.
- BIASUTTO L, DONG LF, ZORATTI M, NEUZIL J: Mitochondrially targeted anti-cancer agents. *Mitochondrion* **10**: 670-681, 2010.
- BONORA M, PINTON P: The mitochondrial permeability transition pore and cancer: molecular mechanisms involved in cell death. *Front Oncol* **4**: 302, 2014.
- BRADFORD MM: A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**: 248-254, 1976.
- BUSTAMANTE E, SOPER JW, PEDERSEN PL: A high-yield preparative method for isolation of rat liver mitochondria. *Anal Biochem* **80**: 401-408, 1977.
- CARAFOLI E, SANTELLA L, BRANCA D, BRINI M: Generation, control, and processing of cellular calcium signals. *Crit Rev Biochem Mol Biol* **36**: 107-260, 2001.
- DE STEFANI D, RIZZUTO R, POZZAN T: Enjoy the Trip: Calcium in Mitochondria Back and Forth. *Annu Rev Biochem* **85**: 161-192, 2016.
- DI LISA F, BERNARDI P: Mitochondria and ischemia-reperfusion injury of the heart: fixing a hole. *Cardiovasc Res* **70**: 191-199, 2006.
- DRAHOTA Z, ENDLICHER R, STANKOVA P, RYCHTRMOC D, MILEROVA M, CERVINKOVA Z: Characterization of calcium, phosphate and peroxide interactions in activation of mitochondrial swelling using derivative of the swelling curves. *J Bioenerg Biomembr* **44**: 309-315, 2012a.
- DRAHOTA Z, MILEROVÁ M, ENDLICHER R, RYCHTRMOC D, ČERVINKOVÁ Z, OŠŤÁDAL B: Developmental changes of the sensitivity of cardiac and liver mitochondrial permeability transition pore to calcium load and oxidative stress. *Physiol Res* **61 Suppl 1**: S165-172, 2012b.
- EINER C, HOHENESTER S, WIMMER R, WOTTKE L, ARTMANN R, SCHULZ S, GOSMANN C, SIMMONS A, LEITZINGER C, EBERHAGEN C, BORCHARD S, SCHMITT S, HAUCK SM, VON TOERNE C, JASTROCH M, WALHEIM E, RUST C, GERBES AL, POPPER B, MAYR D, SCHNURR M, VOLLMAR AM, DENK G, ZISCHKA H: Mitochondrial adaptation in steatotic mice. *Mitochondrion* **40**: 1-12, 2018.
- ENDLICHER R, DRAHOTA Z, CERVINKOVA Z: Modification of calcium retention capacity of rat liver mitochondria by phosphate and tert-butyl hydroperoxide. *Physiol Res* **68**: 59-65, 2019.
- ENDLICHER R, KRIVAKOVA P, LOTKOVA H, MILEROVA M, DRAHOTA Z, CERVINKOVA Z: Tissue specific sensitivity of mitochondrial permeability transition pore to Ca²⁺ ions. *Acta Medica (Hradec Kralove)* **52**: 69-72, 2009.
- FONTAINE E, ERIKSSON O, ICHAS F, BERNARDI P: Regulation of the permeability transition pore in skeletal muscle mitochondria. Modulation By electron flow through the respiratory chain complex i. *J Biol Chem* **273**: 12662-12668, 1998a.
- FONTAINE E, ICHAS F, BERNARDI P: A ubiquinone-binding site regulates the mitochondrial permeability transition pore. *J Biol Chem* **273**: 25734-25740, 1998b.
- HALESTRAP AP: What is the mitochondrial permeability transition pore? *J Mol Cell Cardiol* **46**: 821-831, 2009.
- HALESTRAP AP, CONNERN CP, GRIFFITHS EJ, KERR PM: Cyclosporin A binding to mitochondrial cyclophilin inhibits the permeability transition pore and protects hearts from ischaemia/reperfusion injury. *Mol Cell Biochem* **174**: 167-172, 1997.

- HE J, FORD HC, CARROLL J, DING S, FEARNLEY IM, WALKER JE: Persistence of the mitochondrial permeability transition in the absence of subunit c of human ATP synthase. *Proc Natl Acad Sci U S A* **114**: 3409-3414, 2017.
- ICHAS F, JOUAVILLE LS, MAZAT JP: Mitochondria are excitable organelles capable of generating and conveying electrical and calcium signals. *Cell* **89**: 1145-1153, 1997.
- JUHASZOVA M, WANG S, ZOROV DB, NUSS HB, GLEICHMANN M, MATTSON MP, SOLLOTT SJ: The identity and regulation of the mitochondrial permeability transition pore: where the known meets the unknown. *Ann N Y Acad Sci* **1123**: 197-212, 2008.
- MILEROVÁ M, DRAHOTA Z, CHYTILOVÁ A, TAUCHMANNOVÁ K, HOUŠTĚK J, OŠŤÁDAL B: Sex difference in the sensitivity of cardiac mitochondrial permeability transition pore to calcium load. *Mol Cell Biochem* **412**: 147-154, 2016.
- NICHOLLS DG: Brain mitochondrial calcium transport: Origins of the set-point concept and its application to physiology and pathology. *Neurochem Int* **109**: 5-12, 2017.
- PANOV A, DIKALOV S, SHALBUYEVA N, HEMENDINGER R, GREENAMYRE JT, ROSENFELD J: Species- and tissue-specific relationships between mitochondrial permeability transition and generation of ROS in brain and liver mitochondria of rats and mice. *Am J Physiol Cell Physiol* **292**: C708-718, 2007.
- PARDO AC, RINALDI GJ, MOSCA SM: Mitochondrial calcium handling in normotensive and spontaneously hypertensive rats: correlation with systolic blood pressure levels. *Mitochondrion* **20**: 75-81, 2015.
- RAAFLAUB J: Die Schwellung isolierter Leberzellmitochondrien und ihre physikalisch-chemische Beeinflussbarkeit. *Helv Physiol Pharmacol Acta* **11**: 142-156, 1953.
- RAO VK, CARLSON EA, YAN SS: Mitochondrial permeability transition pore is a potential drug target for neurodegeneration. *Biochim Biophys Acta* **1842**: 1267-1272, 2014.
- SKRHA J, JR., GALL J, BUCHAL R, SEDLACKOVA E, PLATENIK J: Glucose and its metabolites have distinct effects on the calcium-induced mitochondrial permeability transition. *Folia Biol (Praha)* **57**: 96-103, 2011.
- SMITH RAJ, HARTLEY RC, COCHEME HM, MURPHY MP: Mitochondrial pharmacology. *Trends in Pharmacological Sciences* **33**: 341-352, 2012.
- SZABO I, ZORATTI M: Mitochondrial Channels: Ion Fluxes and More. *Physiological Reviews* **94**: 519-608, 2014.
- WEISS JN, KORGE P, HONDA HM, PING P: Role of the mitochondrial permeability transition in myocardial disease. *Circ Res* **93**: 292-301, 2003.

FIGURES







Fig. 4.



CRC (0.4 g/l) (nmol/mg prot.)	CRC (0.2 g/l) (nmol/mg prot.)	CRC ratio 0.2/0.4 g/l (%)
27.1±4.3	26.0±4.7	96.7±14.8

n = 6

