

Peroxisome proliferator-activated receptor γ in white and brown adipocyte regulation and differentiation

Hui Wu¹, Xiaohua Li², Chunyan Shen¹

¹Nutritional department, Seventh People's Hospital of Shanghai University of TCM, No 358, Datong Road, Gaoqiao Town, Pudong New Area, Shanghai 200137, China

²Endocrinology department, Seventh People's Hospital of Shanghai University of TCM, No 358, Datong Road, Gaoqiao Town, Pudong New Area, Shanghai 200137, China

Correspondence

Chunyan Shen, Nutritional department, Seventh People's Hospital of Shanghai University of TCM, No 358, Datong Road, Gaoqiao Town, Pudong New Area, Shanghai 200137, China

Tel.: +86-021-58676230

Email: scy_qy@126.com

Running title: PPAR γ in adipocyte regulation and differentiation

Summary

In as early as 1997, the World Health Organization officially recognized obesity as a chronic disease. The current epidemic of obesity and overweightedness has aroused great interest in the study of adipose tissue formation. The transcription factor peroxisome proliferator-activated receptor γ (PPAR γ) binds to the target gene promoter regulatory sequences, acting as a key factor in regulating the differentiation of preadipocytes in the adipose tissue, and plays an important role in regulating the adipocyte metabolism. A further understanding of the structure and expression characteristics of PPAR γ , in addition to its mechanisms of action in adipocyte differentiation, may be applied to control obesity and prevent obesity-related diseases. In this article, recent studies investigating the effect of regulating *PPAR* γ on adipocyte differentiation are reviewed. In particular, the structural characteristics, expression patterns, and molecular mechanisms of PPAR γ function in adipocyte differentiation are considered.

Keywords: PPAR γ , adipocytes, white adipose tissue, brown adipose tissue, differentiation regulation

1 INTRODUCTION

The current epidemic of obesity and overweight has caused a surge of interest in the study of adipose tissue formation. Recent studies have shown that adipocytes are not only the site of lipid storage, but also the source of many adipose-derived secretory factors that can affect and regulate the metabolism of adipose tissue, other tissues, and consequently the whole organism (Divella *et al.* 2016). Therefore, it is of great clinical significance to study the regulatory mechanisms of adipocyte proliferation and differentiation to control the formation of adipocytes and regulate immune function.

The proliferation and differentiation of adipocytes are complex processes controlled by multiple genes, in which peroxisome proliferator-activated receptor γ (PPAR γ) is the main regulator. Defects in PPARs have been linked to lipodystrophy, obesity, and insulin resistance as a result of impaired adipose tissue expansion and functionality. We believe a greater understanding of PPARs has a significant potential for improving health outcomes and interventions related to obesity, diabetes, and diet.

2 PPAR γ

2.1 PPAR γ subtypes

PPARs were first discovered in 1990, and they belong to the steroid, thyroid, and retinoic acid receptor superfamily. The name of this group of nuclear receptors was derived from their ability to stimulate the proliferation of peroxisomes in rats (Issemann and Green 1990). In humans, three nuclear isoforms were discovered (PPAR α , PPAR β/δ , and PPAR γ) that differ in tissue distribution and specific function. PPAR α is mainly expressed in the liver, kidney, heart, brown adipose tissue (BAT), and muscle, and is closely linked to the lipid metabolism (Kersten *et al.* 2000). PPAR β/δ is expressed in most cell types, most widely in the brain, adipose tissue and skin, and plays roles in fatty acid oxidation and energy balance (Peters *et al.* 2000). PPAR γ is the most widely studied subtype; it is expressed predominantly in adipose tissues, both white and brown (Vella *et al.* 2016). PPAR γ is a key transcriptional regulatory factor for adipocyte production in humans, directly regulating adipocyte differentiation and the expression of genes related to lipid metabolism (Zhang and Li 2010). PPAR γ is also expressed in monocytes, macrophages (including foam cells), cardiomyocytes, vascular smooth muscle cells, and endothelial cells, among others (Sato *et al.* 2005). A large number of *in vitro* and *in vivo* experiments have shown that PPAR γ exerts anti-inflammatory effects by inhibiting the transcription of pro-inflammatory genes. Accordingly, activation of PPAR γ signaling reduces the levels of a large number of inflammatory factors (Peng *et al.* 2015), thereby playing a protective role against tissue injury, including the lungs and brain (Victor *et al.* 2006).

The human PPAR γ gene is located at position 3p25 of chromosome 3 and contains nine exons that are transcribed to generate four isomers. To date, seven diverse mRNA transcripts of PPAR γ ($\gamma 1$ - $\gamma 7$) have been

identified, which are differentially generated based on initiation and alternative splicing of five exons in the terminal region and six exons in the open reading frame (Kvandova *et al.* 2016). PPAR γ 1 is expressed only in cardiac and skeletal muscle cells, vascular smooth muscle cells, and endothelial cells, while PPAR γ 2 and PPAR γ 4 isoforms are primarily expressed in adipocytes, macrophages, and the colon (Kvandova *et al.* 2016). PPAR γ 3 is expressed in adipocytes and the colon. The N-terminal of human PPAR γ 2 protein has 28 amino acid residues, 13 longer than the N-terminal of PPAR γ 1. Moreover, the ability of PPAR γ 2 to induce adipocyte formation is higher than that of PPAR γ 1 (Aprile *et al.* 2014).

2.2 PPAR γ activation by ligands

PPAR γ is a ligand-dependent transcription factor. The ligands of PPAR γ are mainly a variety of free fatty acids (FFA) and their derivatives, which can be divided into natural and synthetic ligands. Natural ligands mainly come from food and metabolic products of the body, including long-chain polyunsaturated FFA such as linoleic acid, linolenic acid, and arachidonic acid, as well as some arachidonic acid derivatives, and prostaglandins (PG), including 1, 5-deoxyprostaglandin J2, and prostaglandin A. The specificity of natural ligands is low; however, specificity is higher for synthetic ligands. Synthetic ligands include the insulin-sensitizing thiazolidinediones (TZD), such as troglitazone and pioglitazone, and non-steroidal anti-inflammatory drugs, such as indomethacin and niflumic acid. (Hallenborg *et al.* 2016).

Following complex formation with a ligand, PPARs become activated and bind to the retinoid X receptor (RXR) to form a PPAR-RXR transcriptional complex (Gearing *et al.* 1993). Similar to other nuclear receptor proteins, the PPARs and RXRs have a standard blueprint with four structural and functional elements (Figure 1) (Liu *et al.* 2016). 1) Non-ligand-dependent transcriptional activation domain (NLD); region A/B, in which the activation function-1 motif (AF-1) domain is the target for phosphorylation, and phosphorylation of AF-1 inhibits PPAR activity. 2) DNA-binding domain (DBD); region C, in which the highly conserved zinc-finger structure promotes the binding of the receptor to the DNA sequence in the promoter region of the target gene. 3) Transcriptional activity regulatory domain (HD); region D, the binding of the nuclear factor to the binding site of its cofactor affects the activity of PPAR. 4) Ligand-binding domain (LBD); region E /F, located at the c-terminal ligand-binding region forms a heterodimer, and AF-2 accelerates the binding with a cofactor, and ultimately assists in the transcription process.

2.3 The regulatory mechanism of PPAR γ

Zinc fingers in the DBD bind to peroxisome proliferator response elements on the target gene promoter to regulate transcription and inhibit or activate the expression of target genes. Peroxisome proliferator response elements exist in the promoters of several genes related to glucose and lipid metabolism, such as fatty acid synthase, pyruvate carboxylase, and glucose transporter 4 (El-Jack *et al.* 1999). The

phosphorylation state of PPAR γ is regulated by its AF-1 domain (Lee & Kim, 2015). PPAR γ interacts with several other signaling pathways, including Janus kinase-signal transducer and activator of transcription (JAK-STAT), nuclear factor κ -light-chain-enhancer of activated B cells (NF- κ B), and activator protein 1 (AP-1). PPAR γ competitively binds to the synergistic activation factors CBP and P300 to inhibit AP-1 and STAT activity, blocking the production of associated pro-inflammatory cytokines (interleukin-6 (IL-6), tumor necrosis factor α (TNF α)). PPAR γ can also directly bind to the P50/P56 subunit of NF- κ B, forming an inhibitory transcriptional complex that inhibits the expression of NF- κ B-dependent genes (Hong *et al.* 2003). The exact mechanisms of interactions between PPAR γ and other signaling pathways remain to be fully elucidated. PPAR γ is a nutrient sensor that regulates several homeostatic functions, such as glucose and lipid metabolism, by regulating the uptake, utilization, oxidation, and storage of FFA, in addition to its role in modulating inflammatory pathways.

3. Adipocyte differentiation

3.1 Origin of adipocytes

Mesenchymal stem cells (MSCs) are non-hematopoietic pluripotent stem cells with high self-renewal and multidifferentiation potential in the mesoderm. Under certain conditions, they differentiate into specific somatic cells of each hypoderm (Goudarzi *et al.* 2018).

Adipocyte differentiation can be divided into two stages. First, in the decisive stage, MSCs receive external signals and start to differentiate into precursor adipocytes. Specific molecular mechanisms are still unknown (Chu & Tao, 2017). Second, terminal differentiation (the differentiation of precursor adipocytes into mature adipocytes), includes three stages: The proliferative, differentiative, and mature stages. During the proliferative stage, fusion and contact inhibition of the precursor adipocytes are followed by the entrance into the cell cycle. CCAAT enhancer-binding protein (C/EBP) family β and δ subtypes are expressed, and cell differentiation is initiated. In the terminal differentiative stage, resting cells move into the growth stage, and transcription factors such as PPAR γ and C/EBP α are expressed at high levels, activating the expression of related genes. Cells then accumulate lipid droplets and become mature adipocytes. The mature adipocyte stage is marked by the cell being filled with a single large fat droplet, high expression of adipocyte marker genes, and secretion of cytokines that are involved in insulin sensitivity, energy balance regulation, and other processes (Tong & Hotamisligil, 2001).

3.2 Differences between three types of adipocytes

3.2.1 White adipocyte cells (WACs)

WACs are thought to originate in the lateral mesoderm from non-myogenic factor 5 (Myf5)-expressing precursors. WACs contain large and stable lipid droplets and few mitochondria, and are found throughout

the body, mainly under the skin and around internal organs. WACs store energy in the form of triglycerides and have endocrine functions. For example, WACs secrete FFA and adipocytokines, such as leptin, adiponectin, TNF α and IL-6, which act on distal tissues, including the brain, liver and muscle to regulate food intake, energy homeostasis and insulin sensitivity (Rosen *et al.* 2006). White adipose tissue (WAT) is therefore referred to as an active endocrine organ.

3.2.2 Brown adipocyte cells (BACs)

BACs and skeletal muscle cells originate from the same paraxial mesodermal cells that specifically express Myf5. Myf5 stimulates the gene transcription profile, promoting muscle differentiation, which partly explains the high metabolic activity of classic BACs (Seale *et al.* 2008).

BACs contain lipid droplets that are small, abundant, and dispersed throughout the cytoplasm. These cells contain a large number of mitochondria, consistent with their function of burning fat and generating heat. BAT is mainly distributed around the spine, collarbone, and adrenal glands. Extremely high mitochondrial content, dense vascular network and extensive nerve supply enable BAT to have important metabolic functions (Qian *et al.* 2015). Uncoupling protein 1 (UCP1) is a specific brown fat cell marker, which exists in the mitochondrial intima and regulates the thermogenesis of BAT by uncoupling the oxidative phosphorylation of ATP synthesis. Through this uncoupling, BAT protects against the cold and also burns excess fat and sugar, preventing the body from storing excess fat. Thus, both WACs and BACs play critical roles in maintaining whole-body energy homeostasis (Pellegrinelli *et al.* 2016).

3.2.3 Beige

In 1992, Cousin *et al.* found that if rodents are exposed to a cold environment for a prolonged time, BAC-like cells are found in their subcutaneous WAT. Like typical BACs, these adipocytes display many small lipid droplets and mitochondria scattered inside the cell and can generate heat. However, without external stimulation, BAC-like cells are not found in the normal subcutaneous WAT. This special type of adipose tissue is called the “beige adipose tissue” (Cousin *et al.* 1992).

The three types of adipocytes differ in gene expression patterns. Specifically, WACs express transcription factor 21 (TCF21), BACs express zinc-finger protein of the cerebellum 1 (ZIC1), and beige adipocytes express homeobox C9 (HOXC9), transmembrane protein 26 (TMEM26), cluster of differentiation 137 (CD137), and T-box transcription factor 1 (TBX1) (Walden *et al.* 2012).

There are various hypotheses on the origin of beige adipose tissue. One widely accepted theory proposes that classical BACs and skeletal muscle cells are derived from the Myf5-positive precursor lineage, and WACs and beige adipocytes are derived from Myf5-negative lineages. However, Myf5- and non-Myf5-expressing lineages are all found in WAT (Sanchez-Gurmaches *et al.* 2012). Expression of beige adipocyte-specific genes is also closely related to the location of the adipocytes within the tissue. Analysis of samples from human adult neck subcutaneous adipose tissue biopsies showed that, from the shallowest to the

deepest tissue, the shallowest fat contains only WACs, while cells in the middle may be beige, and BACs are located in the muscle sheath. Furthermore, the deeper they observed into the biopsy sample, the higher the expression of the BAC-specific genes they found (Sanchez-Gurmaches *et al.* 2012). This diversity in the origin of beige adipocytes suggests that beige, white, and brown adipocytes cannot be distinguished solely by Myf5 expression.

3.3 Key factors in Adipocytes differentiation

3.3.1 PGC-1 α

PPAR γ coactivator-1 α (PGC-1 α) belongs to the PGC1 family and is a major transcriptional coactivator of mitochondrial synthesis and oxidative metabolism genes in most cells, including BAC and skeletal muscle cells. It promotes the expression of UCP1 and stimulates mitochondrial biosynthesis (Puigserver *et al.* 1998). *In vitro* experiments have demonstrated that the overexpression of *PGC-1 α* in MSC blocks the differentiation of MSC into adipocytes. When PGC-1 α is overexpressed in MSC, the expression of mitochondrial biosynthesis, respiration, and thermogenesis genes in MSC increase, suggesting that PGC-1 α mediates the differentiation of MSC into BACs. (Liu *et al.* 2004).

3.3.2 C/EBPs

CCAAT/enhancer-binding protein (C/EBPs) has three subtypes: C/EBP α , C/EBP β , and C/EBP δ . C/EBP β coexpression with positive regulatory domain zinc-finger protein 16 (PRDM16) induces differentiation of fibroblasts into BACs. In the absence of C/EBP β , the differentiation of fibroblasts into BACs, induced by PRDM16, is blocked, UCP1 expression is decreased, and skeletal muscle-specific gene expression is enhanced (Kajimura *et al.* 2009). Animal experiments have shown that high expression of C/EBP β improves the sensitivity of cAMP, leading to an upregulation of PGC-1 α and UCP1, which in turn leads to an increase in the number of beige adipocytes (Karamitri *et al.* 2009). Follow-up studies showed that C/EBP β and C/EBP δ adjust the promoter element of their adjacent gene, and thereby transcriptionally activate C/EBP α . Upon activation, C/EBP α leads to PPAR γ expression. Furthermore, PPAR γ activity is dependent on histone acetylase 1 (HDAC1), as protease enzymatic degradation dissociates HDAC1 from a complex of C/EBP β associated with the C/EBP α , thus promoting the expression of C/EBP β and C/EBP α (Birsoy *et al.* 2008). Thus, C/EBPs and PPAR γ mutually promote, and activate the PGC-1 α signaling pathway through cAMP, promoting browning/beiging of adipose tissue.

3.3.3 PRDM16

PRDM16 is highly expressed in BAT and is not only the determinant gene of BAC directional differentiation but also the "on/off" gene for thermogenesis initiation and regulation of skeletal muscle and BAT differentiation. Therefore, PRDM16 determines the direction of differentiation for Myf5-expressing

precursor cells (Harms *et al.* 2014). High expression of PRDM16 induces the differentiation of precursor cells to BACs and the expression of BAC-specific genes (including UCP1 and PGC-1 α) by inhibiting the expression of MyoD and MyoG in myoblasts. (Iida *et al.* 2015).

The PLDLS motif exists in the suppression domain of PRDM16, which is an evolutionarily conserved sequence consisting of five consecutive amino acids (AA 804-808). The C-terminal binding protein (CtBP) binds to the PLDLS motif to form a PRDM16/CtBP complex, and binds to specific genes in WACs to inhibit their expression. PGC-1 α transposable CtBP binds to PRDM16 and forms a PRDM16/PGC-1 α transcriptional complex that activates BAC-specific genes, including PGC-1 α (Shingo Kajimura *et al.* 2008). PRDM16 promotes the expression of BAC-specific genes by recruiting related proteins (PPAR α / γ , C/EBP β and MED1) to the promoter or enhancer region of these genes. PRDM16 binding sites are found in the promoter regions of BAC-specific genes (Harms *et al.* 2015). The transcription complex formed by the combination of PRDM16 with C/EBP β induces the expression of PPAR γ and PGC-1 α , and PRDM16 subsequently binds to PPAR γ and PGC-1 α , promoting the expression of UCP1 (Kajimura *et al.* 2009). These studies confirm that PRDM16 is the main regulator of BAC formation, by activating BAC-related genes and inhibiting the expression of WAC-related genes (Kajimura *et al.* 2008).

3.3.4 PPAR γ

PPAR γ plays an irreplaceable role in regulating the differentiation of WAT and BAT. MSCs expressing PPAR γ 2 differentiate into WACs, and MSCs expressing PPAR γ 2 and C/EBP β , or PPAR γ 2-C/EBP β -PRDM16 differentiate into BACs. No single cytokine has been found to promote adipocyte differentiation when PPAR γ is knocked out (Lee and Ge 2014). PPAR γ knockout mice result in the loss of adipogenic differentiation, which manifests as adipose atrophy and insulin resistance (Barak Y *et al.* 1999). Many of the key adipocyte-producing signaling pathways eventually also function through PPAR γ and CEBPs. For example, the dephosphorylated Rb protein and Cyclin D3-Cdk6 complex enhance the transcription activity of PPAR γ through phosphorylation, while Cyclin D1 inhibits its transcriptional activity (Rosen and MacDougald 2006). Cytokines, such as TNF α and IL-1, phosphorylate the AF-1 domain of PPAR γ and inhibit adipocyte differentiation (Dapeng and Lixing 2010).

In their study, Ohno *et al.* found that complete PPAR γ receptor agonists, such as rosiglitazone, binds and stabilizes the PPAR γ LBD, promoting the accumulation of PRDM16 (Ohno H *et al.* 2012). The process of adipocyte differentiation involves a very complex transcriptional regulatory network, and many mechanisms are still unknown and need further exploration.

3.3.5 Bone morphogenesis proteins (BMPs)

BMPs are a class of multifunctional growth factors, all of which belong to the transforming growth factor β (TGF β) superfamily except for BMP1. Kang *et al.* found that five BMPs (BMP2, 4, 6, 7, and 9) effectively

induce MSCs to form adipocytes *in vivo* and *in vitro* (Kang *et al.* 2009). In addition, BMP4 and BMP7 also promote the transformation of WACs into BACs.

BMP7 increases the expression of transcriptional regulators PRDM16, PGC-1 α , PPAR γ , and C/EBPs. In animal experiments, embryos with the depletion of BMP7 show a BAC deficiency, and adipocytes stimulated by BMP7 show a strong browning ability *in vitro* (Schulz *et al.* 2011). Continuous secretion of BMP7 promotes the browning of subcutaneous WACs and reduces the degree of obesity in mice (Townsend *et al.* 2014). Also, BMP7 acts on human preadipocytes to promote the transcription of UCP1, and the expression of the beige adipocyte-specific genes *CD137* and *TMEM26* (Okla *et al.* 2015).

4 Browning of WAT

Because browning of WAT changes the energy balance from energy storage to energy expenditure, it has become a new target for the treatment of increasingly severe and prevalent obesity, as well as metabolic syndromes (Frontini *et al.* 2013). The transdifferentiation of white to brown adipocyte involves a transcriptional cascade. In this transcriptional network, PPAR γ , PRDM16, and PGC-1 α are the core regulatory transcription factors (Kinyui & Lei, 2013). Most adipocyte differentiation transforming factors act directly or indirectly through these transcription factors. Furthermore, most key adipocyte-converting signaling pathways eventually converge on them as well. The roles of each of these factors are described below.

4.1 Environmental factors

4.1.1 Cold

Cold temperatures stimulate not only BAC proliferation but also browning of WAT (Zhang *et al.* 2011). This is because the cold stimulates the transient receptor potential of peripheral sensory neurons, activates the sympathetic nervous system to release norepinephrine, and activates BAT through the PKA/p38 MAPK signaling pathway. These signaling pathways also upregulate UCP-1 through transcription factors PGC-1 α and C/EBP β . Studies have shown that cold stimulation or β 3 adrenaline receptor agonists promote subcutaneous adipose tissue cell reconstruction, especially the differentiation of beige adipose cells, and increase the number and proportion of beige adipocytes in the subcutaneous tissue, thereby increasing the ability of subcutaneous fat cells to produce heat (Boss *et al.* 2012).

4.1.2 Hypoxia

Under low oxygen concentrations, the expression of PGC-1 α increases, and the combination of PGC-1 α and PPAR γ effectively promotes the transformation of WAT into BAT, consistent with the browning of white adipocytes under normal oxygen concentrations (Bostrom *et al.* 2012). In a previous study, Lu *et al.* exposed 3T3-L1 adipocytes to transient hypoxia for 4 h/d, and found that transient hypoxia regulates

adipocyte heat production, mitochondrial generation, and glucose and lipid metabolism through AMPK and its related genes, such as PGC-1 α , PPAR γ , and GLUT1 (Lu *et al.* 2015).

4.1.3 Exercise

During exercise, the activation of the central nervous system promotes the release of catecholamines. β -adrenergic receptor agonists promote UCP1 activation in acute (lipid breakdown) and chronic (mitochondrial synthesis, WAT browning) exercise states, increasing the expression and activity of mitochondrial synthesis genes, promoting the browning of subcutaneous white adipocytes and augmenting the WAT adipokine profile (Bostrom *et al.* 2012). Exercise also activates browning of white adipocytes by activating PGC-1 α (Stanford *et al.* 2013); however, this mechanism of action remains poorly understood (Fitzgibbons *et al.* 2014).

4.2 Gene transcriptional regulation

4.2.1 PPAR γ agonists

After treatment with the PPAR γ agonist rosiglitazone, UCP1 and PGC-1 α significantly increase in WAT *in vivo*, which indicates that PPAR γ indirectly acts as a medium to promote browning of white adipose cells (Petrovic *et al.* 2010). However, PPAR γ agonist-induced browning of white adipocytes is dependent on PRDM16.

PPAR γ not only activates BAC gene expression but inhibits WAC gene expression during the browning of adipocytes (Vernoche *et al.* 2009). Moreover, PPAR γ stimulation stabilizes the expression of PRDM16 mRNA, and the induced PGC-1 α interacts with PRDM16 to promote BAT specific gene expression (Ohno *et al.* 2012). However, other studies have shown that PPAR γ agonists induce brown-like cell phenotypes in undifferentiated preadipocytes, such as MSCs or poorly differentiated preadipocytes, but do not elicit brown-like changes in mature WACs (Corrales *et al.* 2018).

4.2.2 PRDM16

BAT from PRDM16 knockout mice completely loses its thermogenic function, while *in vitro* culture of BACs from PRDM16 knockout mice shows an almost complete loss of BAC characteristics (Seale *et al.* 2007). PRDM16 expression in white preadipocytes promotes differentiation into BACs, including the inhibition of white adipocyte-specific gene expression and increased expression of brown adipocyte-specific and mitochondrial genes (Kajimura *et al.* 2015). PPAR γ might act on PRDM16 to induce adipocyte precursors to differentiate toward BACs and regulate the expression of BAC-related thermogenic genes (Harms and Seale 2013). Furthermore, PRDM16- knockout mice are prone to obesity and insulin resistance, induced by a high-fat diet (Cohen *et al.* 2014).

4.2.3 PGC-1 α

Selective PGC-1 α knockout from mice results in decreased heat production by subcutaneous WACs and decreased expression of mitochondria-related genes. This suggests that PGC-1 α might easily induce the browning of subcutaneous WAT. In human subcutaneous WACs, overexpression of PGC-1 α leads to the presence of BACs (Mazzucotelli *et al.* 2007). The expression of PGC-1 α in BAT is dependent on cAMP. C/EBP β combines with the cAMP response element at the proximal end of the promoter region of PGC-1 α to promote the expression of PGC-1 α and mediate the browning of white adipocytes (Karamanlidis *et al.* 2007).

4.3 Cytokines

4.3.1 Fibroblast growth factor 21 (FGF21)

FGF21 is an endocrine factor that plays a key role in energy balance, glucose metabolism and lipid metabolism. Numerous studies have shown that exercise can promote the secretion of FGF21 in the muscle tissue. FGF21 is primarily derived from the liver, but is also synthesized and released from WAT, and is, therefore, considered an adipokine (Kim *et al.* 2013). FGF21 is a key regulator of brown differentiation of white adipocytes. A study by Veniant *et al.* demonstrated that FGF21 induces the browning of WACs by inducing PGC-1 α expression in mouse adipocytes (Veniant *et al.* 2015). Human experiments have shown that BAT also secretes FGF21 and acts on itself to promote the browning of WACs (Hondares *et al.* 2014).

4.3.2 Vascular endothelial growth factor (VEGF)

VEGF is highly expressed in the adipose tissue and plays a role in angiogenesis and energy metabolism. Hypoxia is an inducer of VEGF expression. In addition, VEGF signaling pathway is essential for the induction of beige adipocyte genes, β 3 adrenergic agonists, and PPAR γ after exercise (Veikkola *et al.* 2000). VEGF-A-induced angiogenesis can achieve continuous and adequate oxygen and nutrient exchange during cellulite expansion, thereby improving adipose tissue function. Inhibition of VEGF-A-induced VEGFR2 activation in the early stages of weight gain, induced by a high-fat diet leads to increased systemic insulin resistance. VEGF-B is co-expressed with the nuclear-encoded mitochondrial gene, *OXPHOS*. PGC-1 regulates VEGF-B expression to coordinate FFA uptake and β -oxidation (Mehlem *et al.* 2016). VEGF-B knockout mice showed reduced lipid accumulation in the heart, muscle, and BAT, and increased lipid accumulation in WAT (Hagberg *et al.* 2010). This confirms the role of VEGF-B in the regulation of energy metabolism and differentiation of beige adipocytes.

Mice lacking VEGF-A tend to utilize fat, while those lacking VEGF-B tend to accumulate fat (Lee *et al.* 2014). Experiments in VEGF-B knockout mouse models have shown that the inactivation of VEGF-B leads to the expansion of WACs and increase in WAC-related gene expression, along with the whitening of BACs and a decrease in BAC-related gene expression. In contrast, inhibition of VEGF-A induces BAC development and expansion, increased energy expenditure, upregulation of BAC-related genes and

downregulation of WAC-related genes. When VEGF-B knockout and VEGF-A inhibited are were crossed, VEGF-A and VEGF-B counteract their respective regulation of a large number of genes and effectively reverse each other's effects (Jin *et al.* 2018).

4.4 Post-transcriptional regulation of genes

Non-coding RNA (ncRNA) is a general term for all functional RNAs that are not translated into proteins. In recent years, many new types of ncRNA have been discovered, and the role of some ncRNAs in gene regulatory networks has been gradually recognized. Among them, microRNAs (miRNAs) are a class of ncRNAs commonly seen in eukaryotic cells, with a length of 18~25 nt, which play key roles in the regulation of cell proliferation, differentiation, and apoptosis (Guo *et al.* 2010).

During the cold-induced browning of WACs in mice, the expression of miR-196a is significantly increased, and the expression of HOXC8 is inhibited. As a WAC gene, *HOXC8* inhibits the function of C/EBP binding and thus restricts the generation of BACs. Therefore, miR-196a/HOXC8/C/EBP can be targeted as a pathway for weight loss (Mori *et al.* 2012). Cold also increases the expression of miR-455 in mouse brown adipose cells. *In vitro* experiments have shown that miR-455 regulates the differentiation and thermogenesis of brown adipose cells. Mechanistically, miR-455 mainly activates AMPK (AMP-activated protein kinase) by acting on HIF1(hypoxia-inducible factors-1), triggering the formation of mitochondria and BACs (Zhang *et al.* 2015). miRNAs also bind to C/EBPs to regulate the production of BACs. Among them, miR-155 and C/EBP constitute a double closed-loop negative feedback mechanism and regulate the generation of BAT or beige adipose tissue (Chen *et al.* 2013). miR-378, on the other hand, regulates the browning of WACs by enhancing the binding of C/EBP to the GLUT4 promoter (Gerin *et al.* 2010). Moreover, miR-133 restricts browning of adipocytes by inhibiting the transcription of PRDM16 (Liu *et al.* 2013) and limits adipocyte differentiation by inhibiting the transcription of PPAR γ (Lee.*et al.* 2011). During exposure to cold, miR-27b expression in the adipose tissue is suppressed. Cell culture experiments have revealed that in preadipocytes undergoing browning, the suppression of miR-27b occurs at the same time as PRDM16, PPAR γ and UCP1 increase, showing that miR-27b plays a negative regulatory role in the browning of WAT (Sun *et al.* 2014).

Because microRNAs are a large family, only a small number of microRNAs have been confirmed to play a role in the browning of WACs. For example, miR-196a, miR-378 and miR-455 play positive regulatory roles in complex diseases such as obesity and type 2 diabetes, while miR-155, miR-133, miR-133 and miR-27b play negative regulatory roles (Arias *et al.* 2016) (Figure 3). With the development of RNA detection technology, more non-coding RNAs will be identified, and their regulatory mechanisms in the energy

metabolism and regulatory networks formed with the transcription factors will become more evident, providing new targets for the treatment of obesity and related metabolic diseases.

5 Problems during adipocyte differentiation

At present, we have a comprehensive understanding of adipocyte differentiation; however, several important problems remain unsolved. The first problem is the origin of adipocytes, including the formation of adipocytes in the embryo, the source of the first cells of the adipose tissue, and distribution of these cells throughout the body. Answers to these questions could enable an effective reduction in the number of adipocytes or reductions in specific adipose tissue deposits, which could prevent excessive adipose tissue deposition.

The second problem is that much work remains to be done to reveal how transcription factors promote adipocyte differentiation. Although many of the transcription factors necessary for adipocyte differentiation have been identified, such as PPAR γ , the mechanisms of action remain unclear. Elaboration of the mechanism of action of known transcription factors, functional verification of known key transcription factor target genes, and the identification of new transcription factors are the focus of current research.

Third, further research on the browning of WACs is necessary. BAT does not cause or promote metabolic disease. Instead, it consumes energy and reduces fat accumulation, making the browning of white adipocytes a new target in the treatment of metabolic diseases. However, browning of white adipocytes is very complex, involving the regulation of multiple genes and pathways, and is affected by hypoxia, exercise, cold, and other conditions. Therefore, further studies into are required and the browning of white adipocytes is expected to be an effective means to alleviate the development of metabolic diseases.

6 Conclusion

WAT, BAT and beige adipose tissue regulate the energy storage and body metabolism under the control of PPAR γ , UCP1 and PGC-1 α , which are closely implicated in the occurrence and development of obesity and related metabolic diseases. This paper reviewed the PPAR γ in white and brown adipocyte regulation and differentiation. Browning of white adipocytes is the result of the synergistic effect of various regulatory mechanisms. Among them, PPARs, PGC-1 α , PRDM16 and C/EBPs play key roles in relevant gene transcription. Moreover, the regulation of nerve and endocrine functions are accomplished through the above-mentioned transcription factors. Consequently, the browning of white fat can significantly promote systemic energy consumption and improve carbohydrate and lipid metabolism. At present, there is still a need for in-depth and detailed basic research, so that the interactions of various regulatory pathways on the browning of WAT in humans can be more clearly defined, allowing browning of white adipocytes to become an effective treatment for obesity and metabolism-related diseases.

References

- APRILE M, AMBROSIO MR, D'ESPOSITO V, BEGUINOT F, FORMISANO P, COSTA V, CICCODICOLA A: PPARG in human adipogenesis: differential contribution of canonical transcripts and dominant negative isoforms. *PPAR Res* 2014: 537865, 2014.
- ARIAS N, AGUIRRE L, FERNANDEZ-QUINTELA A, GONZÁLEZ M, LASA A, MIRANDA J, MACARULLA MT, PORTILLO MP: MicroRNAs involved in the browning process of adipocytes. *J Physiol Biochem* 72: 509-521, 2016.
- BARAK Y, NELSON MC, ONG ES, JONES YZ, RUIZ-LOZANO P, CHIEN KR, KODER A, EVANS RM: PPAR γ is required for placental, cardiac, and adipose tissue development. *Mol Cell* 4: 585-595, 1999.
- BIRSOY K, CHEN Z, FRIEDMAN J: Transcriptional regulation of adipogenesis by KLF4. *Cell Metab* 7: 339-347, 2008.
- BOSS O, FARMER SR: Recruitment of brown adipose tissue as a therapy for obesity-associated diseases. *Front Endocrinol (Lausanne)* 3: 14, 2012.
- BOSTROM P, WU J, JEDRYCHOWSKI MP, KORDE A, YE L, LO JC, RASBACH KA, BOSTRÖM EA, CHOI JH, LONG JZ, KAJIMURA S, ZINGARETTI MR, VIND BF, TU H, CINTI S, HØJLUND K, GYGI SP, SPIEGELMAN BM: A PGC1-alpha-dependent myokine that drives brown-fat-like development of white fat and thermogenesis. *Nature* 481: 463-468, 2012.
- CALL KM, GLASER T, ITO CY, BUCKLER AJ, PELLETIER J, HABER DA, ROSE EA, KRAL A, YEGER H, LEWIS WH, JONES C, HOUSMAN DE: Isolation and characterization

of a zinc finger polypeptide gene at the human chromosome 11 Wilms' tumor locus. *Cell* 60: 509-520, 1990.

CHEN Y, SIEGEL F, KIPSCHULL S, HAAS B, FRÖHLICH H, MEISTER G, PFEIFER A: MiR-155 regulates differentiation of brown and beige adipocytes via a bistable circuit. *Nat Commun* 4: 1769, 2013.

CHU DT, TAO Y: Human thermogenic adipocytes: a reflection on types of adipocyte, developmental origin, and potential application. *J Physiol Biochem* 73: 1-4, 2017.

COHEN P, LEVY JD, ZHANG Y, FRONTINI A, KOLODIN DP, SVENSSON KJ, LO JC, ZENG X, YE L, KHANDEKAR MJ, WU J, GUNAWARDANA SC, BANKS AS, CAMPOREZ JPG, JURCZAK MJ, KAJIMURA S, PISTON DW, MATHIS D, CINTI S, SHULMAN GI, SEALE P, SPIEGELMAN BM: Ablation of PRDM16 and beige adipose causes metabolic dysfunction and a subcutaneous to visceral fat switch. *Cell* 156: 304-316, 2014.

COUSIN B, CINTI S, MORRONI M, RAIMBATLT S, RICQUIER D, PENICAUD L, CASTEILLA L: Occurrence of brown adipocytes in rat white adipose tissue: Molecular and morphological characterization. *J Cell Sci* 103: 931-942, 1992.

DAPENG J, LIXING Z: Research progress on adipocyte differentiation and its regulation. *Chin J Cell Biol* 32: 690-695, 2010.

DIVELLA R, DE LUCA R, ABBATE I, NAGLIERIE, DANIELE A: Obesity and cancer: the role of adipose tissue and adipo-cytokines-induced chronic inflammation. *J Cancer* 7: 2346-2359, 2016.

EL-JACK AK, HAMM JK, PILCH, PF, FARMER SR: Reconstitution of insulin-sensitive glucose transport in fibroblasts requires expression of both PPAR γ and C/EBP α . *J Biol Chem* 274: 7946-7951, 1999.

FITZGIBBONS TP, CZECH MP: Epicardial and perivascular adipose tissues and their influence on cardiovascular disease: Basic mechanisms and clinical associations. *Am Heart Assoc* 3: e000582.

FRONTINI A, VITALI A, PERUGINI J, MURANO I, ROMITI C, RICQUIER D, CINTI S: White-to-brown transdifferentiation of omental adipocytes in patients affected by pheochromocytoma. *Biochim Biophys Acta* 1831: 950-959, 2013.

GEARING M, GOTTLICHER M, TEBOUL M, WIDMARK E, GUSTAFSSON JA: Interaction of the peroxisome-proliferator-activated receptor and retinoid X receptor. *Proc Natl Acad Sci USA* 90: 1440-1444, 1993.

GERIN I, BOMMER GT, MCCOIN CS, SOUSA KM, KRISHNAN V, MACDOUGALD OA: Roles for miRNA-378/378* in adipocyte gene expression and lipogenesis. *Am J Physiol Endocrinol Metab* 299: E198-206, 2010.

GOUDARZI F, MOHAMMADALIPOUR A, KHODADADI I, KARIMI S, MOSTOLI R, BAHABADI M, GOODARZI MT: The role of calcium in differentiation of human adipose derived stem cells to adipocytes. *Mol Biotechnol* 60: 279-289, 2018.

GUO H, INGOLIA NT, WEISSMAN JS, BARTEL DP: Mammalian microRNAs predominantly act to decrease target mRNA levels. *Nature* 466: 835-840, 2010.

HAGBERG CE, FALKEVALL A, WANG X, LARSSON E, HUUSKO J, NILSSON I, VAN MEETEREN LA, SAMEN E, LU L, VANWILDEMEERSCH M, KLAR J, GENOVE G, PIETRAS K, STONE-ELANDER S, CLAESSION-WELSH L, YLÄ-HERTTUALA S,

LINDAHL P, ERIKSSON U: Vascular endothelial growth factor b controls endothelial fatty acid uptake. *Nature* 464: 917-921, 2010.

HALLENBORG P, PETERSEN RK, KOUSKOUMVEKAKI I, NEWMAN JW, MADSEN L, KRISTIANSEN K: The elusive endogenous adipogenic PPAR γ agonists: lining up the suspects. *Prog Lipid Res* 61: 149-162, 2016.

HARMS M, SEALE P: Brown and beige fat: development, function and therapeutic potential. *Nat Med* 19: 1252-1263, 2013.

HARMS MJ, ISHIBASHI J, WANG W, LIM, HW, GOYAMA, S, SATO T, KUROKAWA M, WON KJ, SEALE P: Prdm16 is required for the maintenance of brown adipocyte identity and function in adult mice. *Cell Metab* 19: 593-604, 2016.

HARMS MJ, LIM H W, HO Y, SHAPIRA SN1, ISHIBASHI J, RAJAKUMARI S, STEGER DJ, LAZAR MA, WON KJ2, SEALE P: PRDM16 binds MED1 and controls chromatin architecture to determine a brown fat transcriptional program. *Genes Dev* 29: 298-307, 2015.

HONDARES E, GALLEGO-ESCUREDO JM, FLACHS P, FRONTINI A, CEREIJO R, GODAY A, PERUGINI J, KOPECKY P, GIRALT M, CINTI S, KOPECKY J, VILLARROYA F: Fibroblast growth factor-21 is expressed in neonatal and pheochromocytoma-induced adult human brown adipose tissue. *Metabolism* 63: 312-317, 2014.

HONG G, DAVIS B, KHATOON N, BAKER SF, BROWN J: PPAR gamma-dependent anti-inflammatory action of rosiglitazone in human monocytes: suppression of TNF alpha secretion is not mediated by PTEN regulation. *Biochem Biophys Res Comm* 303: 782-787, 2003.

IIDA S, CHEN W, NAKADAI T, OHKUMA Y, ROEDER RG: PRDM16 enhances nuclear receptor-dependent transcription of the brown fat-specific Ucp1 gene through interactions with mediator subunit MED1. *Genes Devel* 29: 308-321, 2015.

ISSEMANN I, GREEN S: Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature* 347: 645-650, 1990.

JIN H, LI D, WANG X, CHEN Y, YAO Y, ZHAO C, LU X, ZHANG S, TOGO J, JI Y, ZHANG L, FENG X, ZHENG Y: VEGFa and VEGFb play balancing roles in adipose differentiation, gene expression, and function. *Endocrinol* 159: 2036-2049, 2018.

KAJIMURA S, SEALE P, KUBOTA K, LUNSFORD E, FRANGIONI JV, GYGI SP, SPIEGELMAN BM: Initiation of myoblast to brown fat switch by a PRDM16-C/EBP-beta transcriptional complex. *Nature* 460: 1154-1158, 2009.

KAJIMURA S, SEALE P, TOMARU T, ERDJUMENT-BROMAGE H, COOPER MP, RUAS JL, CHIN S, TEMPST P, LAZAR MA, SPIEGELMAN BM: Regulation of the brown and white fat gene programs through a PRDM16/CtBP transcriptional complex. *Genes Dev* 22: 1397-1409, 2008.

KAJIMURA S, SPIEGELMAN BM, SEALE P: Brown and beige fat: physiological roles beyond heat generation. *Cell Metab* 22: 546-559, 2015.

KANG Q, SONG WX, LUO Q, TANG N, LUO J, LUO X, CHEN J, BI Y, HE BC, PARK JK, JIANG W, TANG Y, HUANG J, SU Y, ZHU GH, HE Y, YIN H, HU Z, WANG Y, CHEN L, ZUO GW, PAN X, SHEN J, VOKES T, REID RR, HAYDON RC, LUU HH, HE TC: A comprehensive analysis of the dual roles of BMPs in regulating adipogenic and osteogenic differentiation of mesenchymal progenitor cells. *Stem Cells Dev* 18: 545-559, 2009.

KARAMANLIDIS G, KARAMITRI A, DOCHERTY K, HAZLERIGG DG, LOMAX MA: C/EBP β reprograms white 3T3-L1 preadipocytes to a brown adipocyte pattern of gene expression. *J Biol Chem* 282: 24660-24669, 2007.

KERSTEN S, DESVERGNE B, WAHLI W: Roles of PPARs in health and disease. *Nature* 405: 421-424, 2000.

KIM KH, KIM SH, MIN YK, YANG HM, LEE JB, LEE MS: Acute exercise induces FGF21 expression in mice and in healthy humans. *PLoS One* 8: e63517, 2013.

KINYUI AL, LEI S: Turning WAT into BAT: a review on regulators controlling the browning of white adipocytes. *Biosci Rep* 33: 711-719, 2013.

LEE EK, LEE MJ, ABDELMOHSEN K, KIM W, KIM MM, SRIKANTAN S, MARTINDALE JL, HUTCHISON ER, KIM HH, MARASA BS, SELIMYAN R, EGAN JM, SMITH SR, FRIED SK, GOROSPE M: miR-130 suppresses adipogenesis by inhibiting peroxisome proliferator-activated receptor γ expression. *Mol Cell Biol* 31: 626-638, 2011.

LEE HS, KIM J: Constitutive expression of vascular endothelial cell growth factor (VEGF) gene family ligand and receptors on human upper and lower airway epithelial cells. *Int Forum Allergy Rhinol* 4: 8-14, 2014.

LEE JE, GE K: Transcriptional and epigenetic regulation of PPAR γ expression during adipogenesis. *Cell Biosci* 4: 29, 2014.

LEE, WS, KIM, J: Peroxisome proliferator-activated receptors and the heart: lessons from the past and future directions. *PPAR Res* 2015: 271983, 2015.

LIU HJ, LIAO HH, YANG Z, TANG QZ: Peroxisome proliferator-activated receptor- γ is critical to cardiac Fibrosis. *PPAR Res* 2016: 2198645, 2016.

LIU J, WU PH, TARR PT, LINDENBERG KS, ST-PIERRE J, ZHANG CY, MOOTHA VK, JÄGER S, VIANNA CR, REZNICK RM, CUI L, MANIERI M, DONOVAN MX, WU Z, COOPER MP, FAN MC, ROHAS LM, ZAVACKI AM, CINTI S, SHULMAN GI, LOWELL BB, KRAINIC D, SPIEGELMAN BM: Defects in adaptive energy metabolism with CNS-Linked hyperactivity in PGC-1 α null mice. *Cell* 119: 121-135, 2014.

LIU W, BI P, SHAN T, YANG X, YIN H, WANG YX, LIU N, RUDNICKI MA, KUANG S: miR-133a regulates adipocyte browning in vivo. *PLoS Genet* 9: e1003626, 2013.

LU H, GAO Z, ZHAO Z, WENG J, YE J: Transient hypoxia reprograms differentiating adipocytes for enhanced insulin sensitivity and triglyceride accumulation. *Obes (Lond)* 40: 121-128, 2015.

MAZZUCOTELLI A, VIGUERIE N, TIRABY C, ANNICOTTE JS, MAIRAL A, KLIMCAKOVA E, LEPIN E, DELMAR P, DEJEAN S, TAVERNIER G, LEFORT C, HIDALGO J, PINEAU T, FAJAS L, CLÉMENT K, LANGIN D: The transcriptional coactivator peroxisome proliferator activated receptor (PPAR) gamma coactivator-1 alpha and the nuclear receptor PPAR alpha control the expression of glycerol kinase and metabolism genes independently of PPAR gamma activation in human white adipocytes. *Diabetes* 56: 2467-2475, 2007.

MEHLEM A, PALOMBO I, WANG X, HAGBERG CE, ERIKSSON U, FALKEVALL A: Pgc-1alpha coordinates mitochondrial respiratory capacity and muscular fatty acid uptake via regulation of VEGF-b. *Diabetes* 65: 861-873, 2016.

MIROSLAVA KVANDOVA, IMA DOVINO, MIROSLAVA MAJZUNO: The Role of PPAR γ in Cardiovascular disease. *Physiol Res* 65: S343-S363, 2016.

MORI M, NAKAGAMI H, RODRIGUEZ-ARAUJO G, NIMURA K, KANEDA Y: Essential role for miR-196a in brown adipogenesis of white fat progenitor cells. *PLoS Biol* 10: e1001314, 2012.

OHNO H, SHINODA K, SPIEGELMAN BM, KAJIMURA S: PPAR gamma agonists induce a white- to-brown fat conversion through stabilization of PRDM16 protein. *Cell Metab* 15: 395-404, 2012.

OKLA M, HA JH, TEMEL RE, CHUNG S: BMP7 drives human adipogenic stem cells into metabolically active beige adipocytes. *Lipids* 50: 111-120, 2015.

PELLEGRINELLI V, CAROBBIO S, VIDAL-PUIG A: Adipose tissue plasticity: how fat depots respond differently to pathophysiological cues. *Diabetologia* 59: 1075-1088, 2016.

PENG Z, BAN K, WAWROSE RA, GOVER AG, KOZAR RA. Protection by enteral glutamine is mediated by intestinal epithelial cell peroxisome proliferator-activated receptor- γ during intestinal ischemia/reperfusion. *Shock* 43:327-333, 2015.

PETERS JM, LEE SS, LI W, WARD JM, GAVRILOVA O, EVERETT C, REITMAN ML, HUDSON LD, GONZALEZ FJ: Growth, adipose, brain, and skin alterations resulting from targeted disruption of the mouse peroxisome proliferator-activated receptor beta(delta). *Mol Cell Biol* 20: 5119-5128, 2000.

PETROVIC N, WALDEN TB, SHABALINA IG, TIMMONS JA, CANNON B, NEDERGAARD J. Chronic peroxisome proliferator-activated receptor gamma (PPAR gamma) activation of epididymally derived white adipocyte cultures reveals a population of thermogenically competent, UCP1-containing adipocytes molecularly distinct from classic brown adipocytes. *J Biol Chem* 285: 7153-7164, 2010.

PUIGSERVER P, WU Z, PARK CW, GRAVES R, WRIGHT M, SPIEGELMAN BM. A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis. *Cell* 92: 829-839, 1998.

QIAN S, HUANG H, TANG Q. Brown and beige fat: the metabolic function, induction, and therapeutic potential. *Front Med* 9: 162-172, 2015.

ROSEN ED, MACDOUGALD OA: Adipocyte differentiation from the inside out. *Nat Rev Mol Cell Biol* 7: 885-896, 2006.

ROSEN ED, SPIEGELMAN BM: Adipocytes as regulators of energy balance and glucose homeostasis. *Nature* 444: 847-853, 2006.

SANCHEZ-GURMACHES J, HUNG CM, SPARKS CA TANG Y, LI H, GUERTIN DA: PTEN loss in the Myf5 lineage redistributes body fat and reveals subsets of white adipocytes that arise from Myf5 precursors. *Cell Metab* 16: 348-362, 2012.

SATO N, KOZAR RA, ZOU L, WEATHERALL JM, ATTUWAYBI B, MOORE-OLUFEMI SD, WEISBRODT NW, MOORE FA: Peroxisome proliferator-activated receptor gamma mediates protection against cyclooxygenase2-induced gut dysfunction in a rodent model of mesenteric ischemia/reperfusion. *Shock* 24: 462-469, 2005.

SCHULZ TJ, HUANG TL, TRAN TT, ZHANG H, TOWNSEND KL, SHADRACH JL, CERLETTI M, MCDUGALL LE, GIORGADZE N, TCHKONIA T, SCHRIER D: Identification of inducible brown adipocyte progenitors residing in skeletal muscle and white fat. *Proc Natl Acad Sci USA* 108: 143-148, 2011.

SEALE P, BJORK B, YANG W, KAJIMURA S, CHIN S, KUANG S, SCIME A, DEVARAKONDA S, CONROE HM, ERDJUMENT-BROMAGE H, TEMPST P: PRDM16 controls a brown fat/skeletal muscle switch. *Nature* 454: 961-967, 2008.

SEALE P, KAJIMURA S, YANG W, CHIN S, ROHAS LM, ULDRY M, TAVERNIER G, LANGIN D, SPIEGELMAN BM: Transcriptional control of brown fat determination by PRDM16. *Cell Metab* 6: 38-54, 2007.

SHINGO K, PATRICK S, TAKUYA T, HEDIYE EB, MARCUS PC, JORGE LR, SHERRY C, PAUL T, MITCHELL AL, BRUCE MS: Regulation of the brown and white fat gene programs through a PRDM16/CtBP transcriptional complex. *Genes Dev* 22: 1397-1409, 2008.

STANFORD KI, MIDDELBECK RJ, TOWNSEND KL, AN D, NYGAARD EB, HITCHCOX KM, MARKAN KR, NAKANO K, HIRSHMAN MF, TSENG YH, GOODYEAR LJ: Brown adipose tissue regulates glucose homeostasis and insulin sensitivity. *J Clin Invest* 123: 215-223, 2013.

SUN L, TRAJKOVSKI M: MiR-27 orchestrates the transcriptional regulation of brown adipogenesis. *Metabolism* 63: 272-282, 2014.

TONG Q, HOTAMISLIGIL GS: Molecular mechanisms of adipocyte differentiation. *Rev Endocr Metab Disord* 2: 349-355, 2001.

TOWNSEND KL, LYNES M, COBURN J, PRITCHARD E, KWON YM, HUANG T, KAPLAN DL, TSENG YH: Silk-mediated sustained delivery of bone morphogenetic protein 7 (BMP7) to subcutaneous white adipose depot leads to browning and reversal of obesity. *Diabetes* 63: A64–A64, 2014.

VEIKKOLA T, KARKKAINEN M, CLAEISSON-WELSH L, ALITALO K: Regulation of angiogenesis via vascular endothelial growth factor receptors. *Cancer Res* 60: 203-212, 2000.

VELLA S, CONALDI PG, FLORIO T, PAGANO A: PPAR gamma in neuroblastoma: the translational perspectives of hypoglycemic drugs. *PPAR Res* 2016: 1-10, 2016.

VENIANT MM, SIVITS G, HELMERING J, KOMOROWSKI R, LEE J, FAN W, MOYER C, LLOYD DJ: Pharmacologic effects of FGF21 are independent of the "Browning" of white adipose tissue. *Cell Metab* 21: 731-738, 2015.

VERNOCHET C, PERES SB, DAVIS KE, MCDONALD ME, QIANG L, WANG H, FARMER SR: C/EBP α and the corepressors CtBP1 and CtBP2 regulate repression of select visceral white adipose genes during induction of the brown phenotype in white adipocytes by peroxisome proliferator-activated receptor γ agonists. *Mol Cell Biol* 29: 4714-4728, 2009.

VICTOR NA, WANDERI EW, GAMBOA J, ZHAO X, ARONOWSKI J, DEININGER K, LUST WD, LANDRETH GE, SUNDARARAJAN S: Altered PPAR γ expression and activation after Transient focal ischemia in rats. *Eur J Neurosci* 24: 1653-1663, 2006.

WALDEN TB, HANSEN IR, TIMMONS JA, CANNON B, NEDERGAARD J: Recruited vs. nonrecruited molecular signatures of brown, "brite," and white adipose tissues. *Am J Physiol Endocrinol Metab* 302: E19-E31, 2012.

WANG R, LIAN X, ZHANG L, WANG ZK, SIVASAKTHIVEL S, LIU PF, CAI JH, ZHU WL: Adaptive thermogenesis of the liver in tree shrew (*Tupaia belangeri*) during cold acclimation. *Anim Biol* 61: 385-401, 2011.

ZHANG H, GUAN M, TOWNSEND KL, HUANG TL, AN D, YAN X, XUE R, SCHULZ TJ, WINNAY J, MORI M, HIRSHMAN MF: MicroRNA-455 regulates brown adipogenesis via a novel HIF1 α AMPK-PGC1 signaling network. *Embo Rep* 16: 1378-1393, 2015.

ZHANG Y, LI H: Three important transcription factors related to lipogenesis and adipogenesis in mammal. *J Northeast Agric Univ* 17: 62-75, 2010.

Table 1. Comparison of the characteristics of the three types of adipocytes in humans

Characteristic	WACs	BACs	Beige
Distribution	Subcutaneous, perivisceral	Baby shoulder blade, around the kidneys	Adult neck, clavicle, armpit, around the spine, within WACs
Content	Majority of the body	Few and decreases with age	Few
Origin	Myf5 ⁻	Myf5 ⁺	Myf5 ⁺ /Myf5 ⁻
Molecular marker	TCF21	ZIC1	HOXC9, TMEM26
Function	Store energy	Generate heat	Generate heat
Gene transcription	PPAR γ 2, C/EBP β	PPAR γ 2, C/EBP β , PRDM16	PPAR γ , PRDM16, PGC-1 α

BACs, brown adipose cells; WACs, white adipocyte cells; Beige, beige adipose tissue; C/EBP β , CCAAT enhancer-binding protein β ; HOXC9, homologous box C9; Myf5, myogenic factor 5; PPAR, peroxisome proliferator-activated receptor; PRDM16, positive regulatory domain zinc-finger protein 16; TCF21, transcription factor 21; TMEM26, tumor microenvironment metastasis 26; ZIC1, zinc-finger structure of cerebellum 1.

Figure legends

Figure 1. Schematic representation of the domain structures of PPAR γ in humans.

AF-1, N-terminal ligand-independent transcriptional activation domain (A/B); DBD, DNA-binding domain (C); HD, transcriptional activity regulatory domain (D); LBD/AF-2, ligand-binding domain/activation function 2 (E/F).

Figure 2. Schematic of adipocyte differentiation.

MSCs, mesenchymal stem cells; Myf5, myogenic regulatory factor 5; PPAR γ , peroxisome proliferator-activated receptor γ ; C/EBP, CCAAT enhancer-binding protein; PGC-1 α , PPAR γ coactivator-1 α ; MyoD/MyoG, myodifferentiation factor; WACs, white adipocyte cells; Beige, beige adipose tissue; BACs, brown adipocyte cells; β 3-AR, β 3-adrenergic receptors; NPY, neuropeptide Y; PRMD16, positive regulatory domain zinc-finger protein 16; FGF-21, fibroblast growth factor 21; VEGF, vascular endothelial growth factor.

Figure 3. Role of miRNAs in the browning of white adipose tissue.

WACs, white adipocyte cell; Beige, beige adipose tissue; PPAR γ , peroxisome proliferator-activated receptor γ ; PRDM16, positive regulatory domain zinc-finger protein 16; C/EBP β , CCAAT/enhancer-binding protein β .



