

DORIANA

Supplemento agli

ANNALI DEL MUSEO CIVICO DI STORIA NATURALE "G.
DORIA
GENOVA

VOL. VII – N. 340

30.XI.2004

RES LIGUSTICAE

CCXLIII

FRANCESCO VITALI (*)

***XYLOTRECHUS SMEI* (CASTELNAU & GORY, 1841):
ITS PRESENCE IN WESTERN PALAEARCTIC REGION
AND DESCRIPTION OF THE PUPA**

(COLEOPTERA, CERAMBYCIDAE)

Since 1990 a new imported longicorn, contemporarily identified by RATTI (1990) and DIOLI & VIGANÒ (1990) as *Xylotrechus stebbingi* Gahan, 1906, has been recorded in Italy (Veneto and Lombardy).

The oldest Italian specimen was collected already in Piedmont in 1982 (SAMA & COCQUEMPOT, 1995) and* after such records the species was recorded in Friuli-Venezia Giulia, Liguria, Emilia-Romagna, Tuscany, Marche, Latium, Abruzzo and Sardinia (GOBBI, 1993; SAMA & COCQUEMPOT, 1995; BRUGNOLA, 1996; VITALI, 1999; SAMA, 1999; ULIANA, 2001). In Western Palaearctic region it was found also in France, Switzerland, Germany, continental Greece, Crete, Israel and Tunisia (SAMA & COCQUEMPOT, 1995; HOLZSCHUH, 1995; PAVLÍCEK *et alii*, 1998; SAMA, 1999; KÖHLER, 2000; BRAUD *et alii*, 2002; SAMA, 2002; TEUNISSEN, 2002).

Nevertheless, the GAHAN's (1906) description of *Xylotrechus stebbingi* and *X. smei* (Castelnau & Gory, 1841), besides the illustra-

(*) Via Roma 7/12, I-16121 Genova (Italia)

tions provided by STEBBING (1914) for both species, proves the mistake of the Western Palaearctic identifications, as SAMA & COCQUEMPOT (1995) already suspected.

According to the original description, *X. stebbingi* is characterised by: "Elytra ... narrowly covered with grey pubescence at the base, marked with some small spots ... which form three interrupted bands; ... the apex also narrowly bordered with ashy-grey; ... disc [of the pronotum] with a median asperate carina".

On the contrary, according to GAHAN (1906), *X. smei* is characterised by: "greyish or yellow pubescence that ... forms bands and spots on the elytra, disposed as follows: a transverse band on each at the base, ... a short transverse spot; a narrow band which begins nears the scutellum, passes close by the suture, diverges ... and curves outwards ... an apical band, which is slightly produced forwards at the suture ... Prothorax ... raised along middle of disc ... the disc more or less infuscate in the middle, especially towards the base ... This species varies considerably in colour and to some extent also in the markings".

Apart such differences, both species are very much alike.

The description of *X. smei* is that one that corresponds to the species found in Italy. The specimens with complete pubescence are characterised by elytral pattern like *Clytus*-species and not by three transversal bands. Moreover, such pattern forms distinct bands of pubescence at anterior and posterior margin of elytra and not narrow borders; the posterior one is slightly produced forwards at the suture (in the STEBBING's illustration of *X. stebbingi* it is not even present). At last, the disc of the pronotum is simply raised along middle, without any asperate carina.

Also the distribution (DUFFY, 1968) of *X. stebbingi* (Tibet, Himalayas) and of *X. smei* (Pakistan, India, Bhutan, Myanmar, introduced for time in Tanzania) suggests as more realistic scenario the importation of this latter species, through traded timbers, from countries with harbours and more open to foreign commerce.

In fact, the larval development of *X. stebbingi* (STEBBING, 1914; DUFFY, 1968) is related only to *Quercus dilatata* Lindley, a tree species widespread at 2100-2700 m, and *Quercus lanata* Sm. in Rees (= *incana* Roxburgh *nomen nudum*), widespread at 1300-3000 m (GAMBLE, 1972). On the contrary, *X. smei* is a very polyphagous species, already recorded (STEBBING, 1914; DUFFY, 1968) from a lot of




Fig. 1 - Pupa of *Xylotrechus smei* (Castelnau & Gory, 1841), from Genoa in ventral view with particular of the 7th sternite.

tropical and Palaearctic broadleaf trees, among which genera signalised also in Italy, such as *Quercus*, *Morus* and *Ficus*.

Moreover, I had the occasion to compare two specimens from Burma, (Shwegoo Myo, X.1885, L. Fea leg; Tenasseim, Meetan, IV.1887, L. Fea leg., both preserved in the Museo Civico di Storia naturale "G. Doria", Genoa), identified by Gahan himself as *X. vicinus* (Castelnau & Gory, 1841), a junior synonymous of *X. smei*, with twenty-four specimens from the series collected in Lombardy by DIOLI & VIGANÒ (ex. Coll. V. Rosa, now in my collection): I can state that the specimens belong to the same species.

It is therefore likely that all European and Mediterranean records of *X. stebbingi* must be actually referred to *X. smei*.

Nevertheless, its relative old description, pre-imago stages of *X. smei* are still unknown. The finding of a pupa referable to such species gives me the possibility to fill this gap. The description is based on a female specimen, collected in the town of Genoa (Italy, Liguria) on 3rd July 2001 under the bark of a fig log and preserved in my collection.

Xylotrechus smei (Castelnau & Gory, 1841), pupa

Length: 17 mm.

Head rounded; cheeks short, abruptly hooked at their apex and above carinate along the cutter side, pubescent at the base; labrum longitudinally striate, with short recumbent setae; clypeus transversally striate, pubescent at sides; forehead with V-shaped carina, the tip of which reaches the level of lower margin of the eyes and the ends exceed the level of antennal supports; another feeble longitudinal carina, connected at the tip with the previous one, enlarges at the level of antennal supports, forming an oval-elongate flat tubercle and reaches the hind margin of the head. Joints of the palpi subquadrate, their end rounded. Antennae short, 10th joint reaches the front knees.

Pronotum rounded, very feebly constricted at the base, smooth, with two little cluster of 7-8 sparse short conical spinules at each side, one at the base and another a little behind the middle of lateral margin, well visible from the ventral side; mesonotum and metanotum fairly punctate.

REFERENCES

- BRAUD Y., RAMOS R. & COCQUEMPOT C., 2002 - Nouvelles observations de *Xylotrechus stebbingi* Gahan, 1906, en Europe et en Afrique du Nord (Col., Cerambycidae) - *Bull. Soc. entomol. Fr.*, Paris, 107 (5): 487-488.
- BRUGNOLA R., 1993 - Note su alcuni interessanti cerambicidi della coleotterofauna marchigiana (Italia, Marche, Pesaro - Valle del Foglia) - *Lambillionea*, Tervuren, 96, 3 (2): 571-580.
- CHEREPANOV A. I., 1988 - Cerambycidae of Northern Asia. Vol. 2.: Cerambycinae Part II - Oxonian Press Pvt. Ltd., New Delhi, 354 pp.
- DIOLI P. & VIGANÒ C., 1990 - Presenza in Valtellina di un cerambicide nuovo per la fauna italiana: *Xylotrechus stebbingi* Gahan, 1906 - *Natur. valtellin.* Morbegno, 1: 7-10.
- DUFFY E. A. J., 1968 - A monograph of the immature stages of oriental timber-beetles (Cerambycidae) - Trustees of the British Museum (Natural History), London, 435 pp. VIII pl.
- GAHAN C. J., 1906 - The Fauna of British India, including Ceylon and Birma. Vol. I: Coleoptera Cerambycidae - Today & Tomorrow's Printer and Publishers, New Delhi (Reprinted), 329 pp.
- GAMBLE J. S., 1972 - A Manual of Indian Timbers - Bishen Singh Mahendra Pal Singh Ed., Dehra Dun, 868 pp.
- GOBBI G., 1993 - Interessanti reperti di Cerambycidae italiani, specialmente delle regioni centro-meridionali - *Boll. Ass. Romana Entomol.*, Roma, 48: 83-101.
- HOLZSCHUH C., 1995 - Förstschädlinge, die in den letzten fünfzig Jahren in Österreich eingewandert sind oder eingeschleppt wurden - *Stapfia* 37, Linz, 37: 129-141.
- KÖHLER F., 2000 - Erster Nachtrag zum Verzeichnis der Käfer Deutschlands —*Entom. Nachr. Berichte*, Dresden, 44 (1): 60-84.
- PAVLÍČEK T., CHIKATUNOV V., KRAVCHENKO V., DORCHIN J. & NEVO E., 1998 - *Xylotrechus stebbingi* Gahan a new species for Israeli beetle fauna (Coleoptera: Cerambycidae) - *Mitt. internat. entomol. Ver.*, Frankfurt a. M., 23 (1/2): 74-74.
- RATTI E., 1990 - Catalogo dei Coleotteri della Laguna di Venezia. V. Cerambycidae - *Lavori Soc. Ven. Sc. Nat.*, Venezia, 15: 101-114.
- RATTI E., 1999 - Diffusione nel Veneto di *Xylotrechus stebbingi* Gahan, 1906 e segnalazione di una nuova pianta ospite (Insecta, Coleoptera, Cerambycidae) - *Lavori Soc. Ven. Sc. Nat.*, Venezia, 24: 123-124.
- SAMA G., 1999 - Aggiunte e correzioni alla fauna dei Cerambycidae d'Italia (Insecta Coleoptera Cerambycidae) - *Quad. Studi Nat. Romagna*, 11 suppl.: 41-56.
- SAMA G., 2002 - Atlas of the Cerambycidae of Europe and the Mediterranean Area. 1. Northern, Western, Central and Eastern Europe. British Isles and Continental Europe from France (excl. Corsica) to Scandinavia and Urals. - *Nakladatelstvi Kabourek, Zlin*, 173 pp., 36. pl.

-
- SAMA G. & COCQUEMPOT C., 1995 - Note sur l'extension européenne de *Xylotrechus stebbingi* Gahan, 1906 (Coleoptera, Cerambycidae, Clytini) - *L'entomologiste*, Paris, 51 (2): 71-75.
- STEBBING E. P., 1914 - Indian forest insects of economic importance. Coleoptera - Today & Tomorrow's Printers and Publishers, New Delhi, 648 pp.
- TEUNISSEN A.P.J.A., 2002 - Waarnemingen van *Xylotrechus stebbingi* in Griekenland. Observations of *Xylotrechus stebbingi* in Greece, an Asian Cerambycid beetle which became recently established in Mediterranean Area (Col: Cerambycidae – Entim. Ber., Amsterdam, 62 (2): 57-58.
- ULIANA M., 2001 - Segnalazioni faunistiche italiane, 405: *Xylotrechus stebbingi* Gahan, 1906 - *Boll. Soc. ent. ital.*, Genova, 113 (III): 271.
- VITALI F., 1999 - Nuovi dati corologici per alcune specie di cerambycidi italiani (Coleoptera, Cerambycidae) - *Doriana, Genova*, 7 (315): 1-6.

ABSTRACT

The pupa of *Xylotrechus smei* (Laporte & Gory, 1841) is described and illustrated for the first time.

The presence in West-Palaeartic region of such Oriental cerambycid, erroneously recorded by previous authors as *Xylotrechus stebbingi* Gahan, 1906, is proved; the key by Chrepanov (1988) for the identification of the pupae of Western-European *Xylotrechus*-species is adapted.

RIASSUNTO

Xylotrechus smei (Laporte & Gory, 1841): sua presenza nella regione paleartica occidentale e descrizione della pupa (Coleoptera Cerambycidae).

Viene fornita la descrizione della pupa di *Xylotrechus smei* (Laporte & Gory, 1841), specie orientale importata ed acclimatata nella regione paleartica occidentale, sinora identificata erroneamente come *Xylotrechus stebbingi* Gahan, 1906. Viene inoltre modificata la tabella di Cherepanov 1988 per l'identificazione delle pupae delle specie paleartiche occidentali di *Xylotrechus*.

