

bus griseis depressis decoratus. Scutellum sub-transversum, postice obtuse rotundatum, supra planum, densissime griseo-villosum. Elytra thorace duplo fere latiora, lineari-elongata, basi truncata, juxta scutellum utrinque gibba, intra humerum profunde impressa; humeris elevatis, callosis; lateribus pone humeros leviter sinuata, apice dehiscentia, rotundata, supra modice convexa, antice creberrime rude et profunde, apicem versus sensim subtilius punctata, pube depressa grisea et ad basin pilis rigidis immixtis parce vestita, saturate cyanea, certo situ purpureo-micantia. Corpus subtus obscure cyaneum, sub-nitidum, obsoletissime at dense punctulatum, pube grisea depressa nec non pilis longioribus vage adpersum; lateribus anticis griseo-vittatis. Pedes mediocres, toti obscure cœrulei, dense pubescentes.

544. RHAGIUM *fasciculatum* MIHI. Tab. X. f. 8.

Nigrum, flavescenti-pubescentis; antennis pedibusque brunneis; elytris sub-glabris, nitidis, apice ferrugineis, pilis brevibus depressis flavis fasciculatim vage dispersis, fascia lata obliqua antica maculisque duabus pone medium in singulo testaceis.

Longit. $6\frac{1}{2}$ lin. Lat. 2 lin.

Statura et magnitudo fere Rhagii lineati Oliv. sed pictura tota diversa.

Caput elongatum, deflexum, nigrum, antice dilatatum, basi valde constrictum, vertice depressum, canaliculatum, fronte inter antennas gibba, externe transversim sat impressa; pube depressa flavescencia ubique dense tectum, punctis quibusdam

pone oculos adpersum ; ore palpisque ferrugineis. Oculi valde prominuli , sub-rotundati , nigro-luridi. Antennæ thoracis basin paullo superantes , crassæ , brunneæ , opacæ , pube brevissima dense obductæ , nec non setulis rigidis parce adpersæ. Thorax elongatus , latitudine duplo fere longior , niger , opacus , inæqualiter dense flavo-pubescentis , obsolete remote punctatus , antice angustatus , intra basin et paullo pone apicem late et profunde constrictus , lateribus in medio vix dilatatus , spina longa sub-recurvata armatus , basi truncatus , utrinque subtiliter sinuatus , intra basin transversim obsolete carinatus , apice in medio obtuse late rotundatus , pone oculos sinuatus , in disco transversim convexus , longitudinaliter concinne canaliculatus , in fundo canaliculæ carinula vix elevata nigra nitida instructus. Scutellum elongatum , triangulare , nigrum , supra planum , dense flavo-pilosum. Elytra thorace triplo fere latiora , basi truncata , valde retusa , humeris nonnihil productis , callosis ; postice parum angustata , apice rotundata , lateribus valde inflexa , subtiliter reflexo-marginata , disco modice convexa , antice vage sed rude et profunde , posterius obsolete punctata , nigra , sub-nitida , apice ferruginea , pilis flavis brevibus fasciculatim collocatis et valde depressis ubique vage adpersa , singuli elytri medio fascia lata obliqua sinuata testacea suturam haud attingente paullo ante medium , posterius maculis duabus transversim dispositis , interiore latiore , exteriori margine connexa testaceis , in quovis elytro interne lineis duabus longitudinalibus elevatis glabris nitidissimis. Corpus subtus nigrum , nitidum fasciculatim flavo-pilosum ; ano fusco. Pedes elongati , validiusculi , brunnei , sat dense pubescentes ; tibiis tarsisque nonnihil dilutioribus.



Phytoecia puncticollis Menotr. 2. *P. azurea* Steven. 3. *P. Faldermanni*
Dy. 4. *P. diademata* Fald. 5. *P. piceosa* Fald. 6. *Agapanthia smaragdina*
Dy. 7. *A. chalybea* Fald. 8. *Phagium fasciculatum* Fald. 9. *Toxotus in-*
sitivus Boeber. 10, 11. *Toxotus persicus* ♂, ♀. Fald.