

THE ROLE OF CONNEXIN 37 POLYMORPHISM IN SPONTANEOUS ABORTION

Michal KNÍŽE¹, Jan PIŤHA², Jaroslav A. HUBACEK^{2,4}, Tomáš FAIT¹

¹Department of Obstetrics and Gynecology, Motol University Hospital, Prague, Czech Republic

²Centre for Experimental Medicine, Institute of Clinical and Experimental Medicine, Prague, Czech Republic

³Department of Cardiology, Institute of Clinical and Experimental Medicine, Prague, Czech Republic

⁴3rd Department of Internal Medicine, 1st Faculty of Medicine, Charles University, Prague, Czech Republic

Short title: CX37 POLYMORPHISM AND ABORTION

Corresponding author:

M. Kníže,

Department of Obstetrics and Gynecology,

Motol University Hospital,

Prague,

Czech Republic.

E-mail: michal.knize@gmail.com

Abstract

Among unique cardiovascular risk factors in women are complications during pregnancy, including miscarriage. Important risk factor is also genetic background. One of powerful candidate genes for cardiovascular disease of atherosclerotic origin (aCVD) is gene for connexin 37 (Cx37) with strong gene-environment interaction including smoking status, that is also strong risk factor for complications in pregnancy including spontaneous abortion (SA). We analyzed association between SA and Cx37 gene polymorphism (1019C>T; Pro319Ser) in 547 fetuses and its potential interaction with smoking status of mothers. Using genetic analyses from women from general population as controls, ORs for T allele, found in our previous studies to be protective against aCVD, were calculated. T allele carriers (fetuses), had OR 0.91 (95% CI 0.72-1.14) and no interaction with smoking was observed. In conclusion, no significant association between Cx37 polymorphism and SA was observed and no modifying effect of smoking status on this association was detected.

Key words: Connexin 37 gene, Gap junctions, Spontaneous abortion, Protective factor, Smoking, Cardiovascular disease, Candidate genes

Regarding cardiovascular disease of atherosclerotic origin (aCVD) the annual incidence of cardiovascular diseases is age-dependently increasing both in men and women. The prevalence is higher in men until midlife.(Kittnar 2020). However, women are affected similarly or more than men in older age. Despite the main cardiovascular factors are shared by both sexes, several unique cardiovascular factors were described in women as is miscarriage and other complications during pregnancy (Horn et al. 2019, Oliver- Williams et al. 2013) and metabolic disorders as gestational diabetes mellitus (Vejrazkova et al. 2017). Therefore, detection of determinants of atherosclerotic disease in women including metabolic ones may help to identify and to prevent atherosclerotic process and its clinical complications (Pithová et al. 2016). However, also better understanding of the complications in pregnancy might have similar preventative effect especially when accompanied by genetic analyses. Indeed, genetic studies could help to identify not only inherited causes of early pregnancy loss, extremely serious and emotional problem both for affected women, their partners and their physicians, but also for aCVD. One of powerful candidate genes for aCVD is gene for connexin 37 (C1019>T) (Cx37) with strong gene-environment interaction including smoking status. Connexins are the main proteins in gap junctions. Connexin 37 (Gap junction protein alpha; OMIM ID 121012; Cx37) is the main gap junction representative in the vessel wall expressed in endothelial cells, smooth muscle cells but also in monocytes (Chanson et al. 2007, Wong et al. 2006), therefore in the main cells involved in the process of atherosclerosis. Gene polymorphism-association studies have detected a link between the C1019>T single nucleotide polymorphism (SNP) in the human Cx37 gene and its protective role in cardiovascular disease. This polymorphism results in a non-conservative amino acid change in the

regulatory C-terminus of the Cx37 protein (P319>S). It was recently demonstrated that Cx37 hemichannels control the initiation of atherosclerotic plaque development by regulating ATP-dependent monocyte adhesion. (Deroutte et al.2009)

In addition to cells responsible for atherosclerosis and aCVD, Cx 37 is also present in ovary, in which gap junctions mediate metabolic cooperation between granulosa cells and oocyte. These interactions are mediated not only by paracrine factors but also by gap junctions including Cx37 (Fontana et al. 2020). Preimplantation embryos express multiple connexins and assemble them into gap junctions. The suppression of connexin expression seems to play an important role in embryo implantation in several species. The gap junction induction is one of the earliest known signs for a blastocyst-derived signal which may be involved in preparing the endometrium for implantation. It could help to regulate the invasion process by a coordinated cell death of the endometrial cells replaced by the invading trophoblast. (Grümmer and Winterhager 1998) However, despite the importance of the different connexin channels expressed in the endometrium this topic was not evaluated.

Based on these facts, we analyzed data from fetuses from women with spontaneous abortion (SA) for Cx37 gene polymorphism (1019C>T; Pro319Ser) including its potential interaction with smoking, frequent risk factor for early pregnancy loss. In particular, we analyzed if T allele of Cx37 gene, found to be protective against ischemic heart disease in our previous studies, could be protective also against SA and if this protection could be modified by smoking status.

Women with verified diagnosis of SA participated in the study after giving written informed consent. The study was approved by the Institutional ethics committees and was conducted according to the Good Clinical Practice guidelines. Participating women filled in questionnaire focused on history of SA and physical and ultrasound examination were performed. Samples of aborted fetuses from Caucasian women (n=547) aged 15–44 years (mean age 32.1±4.8 years) collected during 2008–2011 at the Gynecological and Obstetrical Clinic in Prague (Rynekrova et al. 2012, Hubacek et al. 2015) were examined. Material acquired from surgical revision of uterine cavity was subjected to a cytogenetic and histological examination. Women with anatomic abnormalities of the uterus, primary endocrinopathies (such as thyreopathy, diabetes mellitus, hyperprolactinemia), known immunologic or hematologic disease, febrile infections during the first trimester, drug abuse and abnormal cytogenetic examination of the aborted tissue were excluded from the study.

As a control group we used data from the cohort of 662 women examined from the population sample in Prague (mean age 57.9±2.8 years). Detailed questionnaire about medical history, focused also on gynecological history including history of SA was obtained (Hubacek et al. 2009).

Samples from both groups of women (meant fetuses in study group) were stored at -20 °C and DNA was isolated by standard procedures. To genotype the Cx37 rs1764391 polymorphism, oligonucleotides 5'-CTGGACCCACCCCCTCAGAATGGCCAAAGA and 5'-AGGAAGCCGTAGTGCCCTGGTGG and the restriction enzyme AasI (Fermentas, Lithuania) were

used. T allele is characterized by restriction fragments 240bp + 35bp; and C allele with the unrestricted 275bp fragment. Restriction fragments were separated via electrophoresis using 2% agarose gel. (Pitha et al. 2016, b).

We compared the prevalence of *Cx37* gene variants (TT, CT, CC) between aborted material from women with SA vs. women from general population after stratifying them by smoking status (self-reported). Continuous variables were compared by unpaired Student's t-test (between study population and control group), for categorical variables χ^2 Fischer exact test was used.

In women with spontaneous abortion 5.6 % were smokers/past smokers (mean age 27.2 ± 6.5 years) and 94.4 % were non-smokers (mean age 32.3 ± 4.6 years) (difference for age, $p < 0.001$). In the population sample 44.8 % women were smokers/past smokers (mean age 57 ± 2.7 years) and 55.2 % non-smokers (mean age 57 ± 2.7 years).

Cx37 T allele was present in 55.2 % (mean age 32.1 ± 4.7 years) of aborted fetuses, and in 57.5 % (mean age 57 ± 2.7 years) of women from general population. In smoking women at least one T allele was present in 53.4 % of the fetuses, and in 56.1 % of women from general population. In non-smoking women at least one T allele was present in 55.1 % of the fetuses, and in 59.0 % of women from general population. We compared distribution of *Cx37* gene polymorphism between fetuses from women older than 32 years with women younger than 32 years (median age). No difference in the prevalence of T allele was detected (56.9 % vs. 54.2 %, $p = 0.294$). We also tested difference in prevalence of T allele between fetuses of women with one SA ($n = 448$) and fetuses of women with more than 1 SA ($n = 115$), but also in this case no significant difference was detected (54.3 % vs. 60.0 %, $p = 0.161$). The only difference found was that women with more than one SA were older than women with one SA (34.1 ± 4.7 vs. 31.6 ± 4.8 years, $p < 0.001$). In addition, in the control group from general population, women (after exclusion of diabetics) with the history of one or more SA ($n = 51$) were compared to women without history of SA with history of pregnancy ($n = 233$) of similar age, no difference in the presence of T allele was observed (66.7 % vs. 60.1 %; $p = 0.671$).

Using general population as a control sample, ORs for T allele carriers were calculated for the whole populations under study and for smoking and non-smoking women including similar stratification to smokers/past smokers and non-smokers as in the control group. Fetuses from women suffering from SA with T allele have OR 0.91 (95% CI 0.72-1.14) and there was no interaction with smoking status of mothers, OR 0.84 (0.64-1.11) for non-smokers vs. OR 0.89 (0.42-1.9) for smokers.

In summary, in our study we found no significant differences in *Cx37* gene variability between fetuses analyzed after SA and general population of middle aged women. These results were not modified by smoking status. Therefore, in contrast to our previous findings in women with acute coronary syndrome (Pitha et al. 2016, a) T allele does not seem to be neither protective against SA, nor its effect is strongly modified by smoking status. Therefore, mechanisms leading to cardiovascular

disease and complications during pregnancy including SA seems to be different regarding mechanism mediated by gap junctions involved in differentiation processes by mediating exchanges between mother and fetus cells, affecting the maternofetal blood flow interrelationships, trophoblast invasiveness and the formation of a syncytiotrophoblast (Malassiné and Cronier 2005). Expression of some connexins, among them also Cx37, has been shown to reflect maturity of luteinized follicles in animal models and Cx43 expression was associated with better prognosis in *in vitro* fertilization (Tsai et al. 2003). Transcription factors of several connexins were detected in human embryos, e.g. Cx43 was indicated in all embryonic developing stages (Bloor et al. 2004). He and Chen (2016) demonstrated that the expression of Cx43, together with VEGF, is significantly reduced in chorionic villi and decidua in women with spontaneous abortion. This may be caused by the influence on angiogenesis of placenta and developing embryo. It was reported (Togashi et al. 2015), that mouse embryos cultured with some gap junctions inhibitors presented frequent collapses and developmental delay. Functional and structural abnormalities of Cx43 might also play an important role in heart diseases (Pecoraro et al. 2015). Because of this finding and possible similarity with Cx43, we focused on Cx37 polymorphism on spontaneous abortion. In addition, Cx37 polymorphism is also supposed to play important role in cardiovascular events. Nevertheless, we did not find an association in the case of Cx37 gene as for other studies describing different connexins polymorphisms and spontaneous abortion (He and Chen, 2016).

Regarding the (absence of) effect of smoking in our study, the idea was that adhesion properties of vessels wall caused by Cx37 gene polymorphism could favor not only macrophage accumulation in the atherosclerotic lesions but also endothelial dysfunction induced by smoking initiated by reduced nitric oxide bioavailability and further by the increased expression of adhesion molecules (Messner et al. 2014) Therefore protective effect of the Cx37 T allele might be strongly modified by smoking in atherosclerosis. But according to our data the deleterious effect of smoking on pregnancy, particularly on SA seems not to be mediated through Cx37 gene.

The limitations of our study are incomplete data in women with history of SA and focus only on one polymorphism of one gene from rather large connexin family. On the other hand, the strengths of the study include high number of study women and numerous representative control group from the population sample in Prague which could reflect real population background for our findings in contrast to cases and controls usually used for such comparisons. In addition, selecting only Cx37 gene was based on knowledge, that connexin 37 is on one hand the main protein of gap junctions in cells involved in the process of atherosclerosis, but on the other hand it is also present in maternofetal organs. Moreover, data from the Czech Republic, Croatia, Hungary and Romania indicated that the strongest protective effect of the Cx37T allele was detected in the non-smoking patients without diabetes mellitus and hypertension and that effect could be mediated through stem cells (Piřha et al. 2010, 2016a). Therefore, we have chosen Cx37 polymorphism as gene potentially covering both cardiovascular and pregnancy

complications. Another limitation is the lack of data from mothers/fathers and absence of information regarding other potential modifying factors including diabetes mellitus; however, women with these characteristics were excluded to detect only the effect of this particular gene variability and smoking.

In summary, to the best of our knowledge, only a few studies have demonstrated effect of connexins on spontaneous abortion and no studies analyzed parallel effect of particular connexin gene polymorphism on cardiovascular disease and SA. In our study, no effect of Cx37 polymorphism measured in fetuses was observed on abortion and no interaction with smoking status on this association was proved.

Conflict of interest

There is no conflict of interest.

Acknowledgements

This work was supported by Supported by the project (Ministry of Health, Czech Republic) for development of research organization 00023001 (IKEM, Prague, Czech Republic) and 00064203 (Motol University Hospital, Prague, Czech republic).

Abbreviations

aCVD	-	Cardiovascular disease of atherosclerotic origin
Cx37	-	Connexin 37
CI	-	Confidence interval
DNA	-	Deoxyribonucleic acid
OR	-	Odds ratio
SA	-	Spontaneous abortion
VEGF	-	Vascular endothelial growth factor

References

- BLOOR D J, WILSON Y, KIBSCHULL M, TRAUB O, LEESE H J, WINTERHAGER E, KIMBER SJ: Expression of connexins in human preimplantation embryos in vitro. *Reprod Biol Endocrinol* 2: 25, 2004.
- CHANSON M, KWAK, BR. Connexin37: a potential modifier gene of inflammatory disease. *J Mol Med* 85: 787–795, 2007.
- DEROUELLE JP, DESPLANTEZ T, WONG CW, ROTH I, KWAK BR, WEINGART R. Functional differences between human Cx37 polymorphic hemichannels. *J Mol Cell Cardiol* 46: 499-507, 2009.
- FONTANA J, MARTINKOVA S, PETR J, ZALMANOVÁ T, TRNKA J. Metabolic cooperation in the ovarian follicle. *Physiol Res* 69: 33-48, 2020.
- GRÜMMER R, WINTERHAGER E. Regulation of gap junction connexins in the endometrium during early pregnancy. *Cell Tissue Res* 293: 189–194, 1998.
- HE X, CHEN Q: Reduced expressions of connexin 43 and VEGF in the first-trimester tissues from women with recurrent pregnancy loss. *Reprod Biol Endocrinol* 14: 46, 2016.
- HORN J, TANZ LJ, STUART JJ, MARKOVITZ AR, SKURNIK G, RIMM EB, MISSMER SA, RICH-EDWARDS JW: Early or late pregnancy loss and development of clinical cardiovascular disease risk factors: a prospective cohort study. *BJOG* 126: 33 - 42, 2019.
- HUBACEK JA, PITHA J, ADAMKOVA V, LANSKA V, POLEDNE R. A common variant in the FTO gene is associated with body mass index in males and postmenopausal females but not in premenopausal females. Czech post-MONICA and 3PMFs studies. *Clin Chem Lab Med* 47: 387-390, 2009.
- HUBACEK JA, RYNEKROVA J, KASPAROVA D, ADAMKOVA V, HOLMES MV, FAIT T. Association of MTHFR genetic variants C677T and A1298C on predisposition to spontaneous abortion in Slavonic population. *Clin Chim Acta* 440: 104-107, 2015.
- KITTNAR O. Selected sex related differences in pathophysiology of cardiovascular system. *Physiol Res*. 69(1):21-31, 2020.
- MALASSINÉ A, CRONIER L: Involvement of gap junctions in placental functions and development. *Biochim Biophys Acta* 1719/1–2: 117-124, 2005.
- MESSNER B, BERNHARD D. Smoking and cardiovascular disease: mechanisms of endothelial dysfunction and early atherogenesis. *Arterioscler Thromb Vasc Biol* 34: 509-515, 2014.
- OLIVER-WILLIAMS CT, HEYDON EE, SMITH GC, WOOD AM: Miscarriage and future maternal cardiovascular disease: a systematic review and meta-analysis. *Heart* 99: 1636-1644, 2013.

- PECORARO M, VERRILLI V, PINTO A, POPOLO A: Role of connexin 43 in cardiovascular diseases. *Eur J Pharmacol* 768: 71-76, 2015.
- PIŤHA J, HUBÁČEK JA, PIŤHOVÁ P: The connexin 37 (1019C>T) gene polymorphism is associated with subclinical atherosclerosis in women with type 1 and 2 diabetes and in women with central obesity. *Physiol Res* 59: 1029-1032, 2010.
- PIŤHA J, KRÁLOVÁ LESNÁ I, HUBÁČEK JA, SEKERKOVÁ A, LÁNSKÁ V, ADÁMKOVÁ V, DOROBANTU M, NICOLESCU R, STEINER R, IVIĆ V, BORBELY A, PAPP Z, SANDOR GV: Smoking impairs and circulating stem cells favour the protective effect of the T allele of the connexin37 gene in ischemic heart disease – A multinational study. *Atherosclerosis* 244: 73-78, 2016a).
- PIŤHA J, PIŤHOVA P, STAVEK P, HUBACEK J, AUZKY O, NESKUDLA T, PELIKANOVA T, KVAPIL M: Gene for connexin 37 is associated with atherosclerotic changes in women with diabetes mellitus type 2. *Atherosclerosis* 252: 144-145, 2016b).
- PITHOVA P, ŠTECHOVÁ K, PIŤHA J, LÁNSKÁ V, KVAPIL M: Determinants of preclinical atherosclerosis are different in type 1 and type 2 diabetic women. *Physiol Res* 65: 219-228, 2016.
- RYNEKROVA J, KASPAROVA D, ADAMKOVA V, FAIT T, HUBACEK JA. Analysis of the potential role of Apolipoprotein E polymorphism in genetic predisposition to spontaneous abortion. *Am J Reprod Immunol* 67: 179-183, 2012.
- TOGASHI K, KUMAGAI J, SATO E, SHIRASAWA H, SHIMODA Y, MAKINO K, SATO W, KUMAZAWA Y, OMORI Y, TERADA Y: Dysfunction in gap junction intercellular communication induces aberrant behavior of the inner cell mass and frequent collapses of expanded blastocysts in mouse embryos. *J Assist Reprod Genet* 32: 6, 2015.
- TSAI MY, LAN KC, HUANG KE, HUANG FJ, KUNG FT, CHANG SY. Significance of mRNA levels of connexin37, connexin43, and connexin45 in luteinized granulosa cells of controlled hyperstimulated follicles. *Fertil Steril* 80: 1437-1443, 2003.
- VEJRAZKOVA D, VANKOVA M, LUKASOVA P, VCELAK J, CIRMANOVA V, HALUZIK M, BENDLOVA B. Specific metabolic characteristics of women with former gestational diabetes: the importance of adipose tissue. *Physiol Res* 66: 349-356, 2017.
- WONG CW, CHRISTEN T, ROTH I, CHADJICHRISTOS CE, DEROUETTE JP, FOGLIA BF, CHANSON M, GOODENOUGH DA, KWAK BR. Connexin37 protects against atherosclerosis by regulating monocyte adhesion. *Nat Med* 12: 950–954, 2006.

Table 1

Cx37 polymorphism in fetuses and population control. Smoking status in fetuses correspond with this characteristic in mothers.

Entire groups

Cx37	Controls		Abortions		OR	P
	N	%	N	%		
CC	281	42.5	239	44.8	1.00	
CT	300	45.3	239	44.8	0.93 (0.73-1.19)	0.28
TT	81	12.2	55	10.4	0.80 (0.54-1.18)	0.25
+T	381	57.5	294	55.2	0.91 (0.72-1.14)	0.41

Smokers/past smokers

CX37	Controls		Abortions		OR	P
	N	%	N	%		
CC	126	43.9	14	46.7	1.0	
CT	121	42.2	14	46.7	1.04 (0.48-2.28)	0.92
TT	40	13.9	2	6.7	0.45 (0.10-2.07)	0.29
+T	161	56.1	16	53.4	0.89 (0.42-1.90)	0.77

Neversmokers

CX37	Controls		Abortions		OR	P
	N	%	N	%		
CC	144	40.7	232	44.9	1.00	
CT	172	48.6	231	44.7	0.83 (0.63-1.11)	0.21
TT	38	10.4	54	10.4	0.88 (0.55-1.40)	0.28
+T	210	59.0	285	55.1	0.84 (0.64-1.11)	0.22