

Physiological Research Pre-Press Article

Progesterone inhibitory role on gastrointestinal motility

Mohammad Alqudah^{1,2}, Othman Al-Shboul¹, Ahmed Al_Dwairi¹, Doaa Ghazi Al-Udat¹ and Abdelrahim Alqudah³

Running title: Progesterone and gut smooth muscle

1. Department of Physiology and Biochemistry, School of Medicine, Jordan University of Science and Technology, Irbid, Jordan

2. Department of Physiology, School of Medicine and biomedical Sciences, Arabian gulf University, Manama, Bahrain

3. Department of Clinical Pharmacy and Pharmacy Practice, Faculty of Pharmacy, The Hashemite University, Zarqa, Jordan.

*Corresponding author:

Mohammad Alqudah

Associate Professor

Department of Physiology

School of Medicine and Biomedical Sciences

Arabian Gulf University

Number of figures: 2

Number of tables: 0

Summary

Progesterone is a steroidal hormone that is produced from the corpus luteum of the ovaries and from the placenta. The main function of progesterone is to promote the secretory differentiation in the endometrium of the uterus and to maintain pregnancy by inhibiting uterine contractions throughout pregnancy. Progesterone performs its actions by activating the classical progesterone nuclear receptors that affect gene transcription and by the non-classical activation of cell surface membrane receptors that accounts for the rapid actions of progesterone. Besides the reproductive roles of progesterone, it exerts functions in many tissues and systems such as the nervous system, the bone, the vascular system, and the gastrointestinal (GI) tract. This review will summarize the recent literature that investigated the role of progesterone in GI tract motility. Most literature indicates that progesterone exerts an inhibitory role on gut smooth muscle cells in part by elevating nitric oxide synthesis which induces relaxation in smooth muscle. Moreover, progesterone inhibits the signaling pathways that lead to contraction such as Rho kinase inhibition. These data serve as a quick resource for the future directions of progesterone research that could lead to better understanding and more effective treatment of gender-related GI tract motility disorders.

Key words: Progesterone, Smooth muscle, GI tract

Introduction

Progesterone is the natural progestin that is mainly produced in the corpus luteum of the ovaries and by the placenta. It is synthesized from the cholesterol derivative, pregnenolone by the actions

of the cholesterol side-chain cleavage enzyme and 3-beta-hydroxysteroid dehydrogenase (1, 2). Although the focus of progesterone and its applications have been associated with female physiology, it is present in male gonads and regulates several male physiological and pathological functions such as spermiogenesis and androgen synthesis (2, 3). Moreover, progesterone is produced in the male and female central nervous systems (CNS) and all the required enzymes and precursors for progesterone synthesis are present in CNS where it acts directly on neuronal tissues (4). Furthermore, Progesterone exerts several important functions in tissues other than the reproductive tissue such as the cardiovascular, renal, adipose, bone, and gastrointestinal tissues (3, 5). The aim of this review is to summarize the role of progesterone in gastrointestinal (GI) tract motility function and to highlight the potential therapeutic benefits and risks of progestin preparations in the GI tract.

Mechanism of progesterone actions

Progesterone produces its physiological functions by activating classical and non-classical signaling pathways (6). In the predominant classical pathway, progesterone binds to its receptor; a member of the nuclear receptor superfamily of transcription factors (6). There are two main isoforms of progesterone receptors (PRs), PR-A and PR-B. Binding of progesterone to PR induces its translocation to the nucleus where the complex binds to progesterone response elements and initiates or prevents the transcription of a wide array of proteins (7). However, the non-classical pathway produces rapid actions by activating cell surface receptors that resemble G protein-coupled receptors (8). Progestin binding to these receptors rapidly activates various intracellular signaling molecules such as elevation of intracellular calcium, extracellular signal-regulated kinases 1 / 2 (ERK1/2), mitogen-activated protein kinase (MAP) kinase and Protein

kinase B/phosphatidylinositol 3-kinases (Akt/PI3K) (figure 1) (9).

Functions of progesterone

The main reproductive function of progesterone is to promote the secretory differentiation in the endometrium of the uterus during the second half of the female sexual cycle (10). These changes aim to prepare the uterus for implantation of the fertilized ovum (10). In addition to this effect on the endometrium, progesterone decreases uterine contraction force and frequency to prevent the explosion of the newly implanted ovum and maintains a quiescent environment for fetal development (11). Additionally, it influences the activity of the intrauterine immune system through the regulation of inflammatory mediators production from T cells. This action of progesterone helps to maintain pregnancy as well by increasing the immunological protection against any potential microbial threats (12). Moreover, progesterone prepares the breasts for lactation through its action to promote proliferation and enlargement of alveolar cells of the breast (13).

In addition to the unquestionable role of progesterone in reproduction, it has numerous functions in various organs in the body. In the central nervous system, progesterone is produced at high concentrations from glial and neuronal cells (14). Progestins produced in the CNS are called neurosteroids. They modulate many neuronal functions either by upregulating or downregulating neurotransmissions and are involved in cognition, memory, neuroprotection, and myelination processes (14). Moreover, progesterone is involved in sexual and maternal behaviors (15, 16). Like in the reproductive system, neurosteroids act through classical and non-classical pathways in the CNS(17).

Progesterone plays an essential role in bone physiology; it stimulates bone formation and slows down resorption. However, its effect is influenced by estrogen, age, and the predominant signaling pathways (18).

Progesterone has been studied in association with diabetes and gestational diabetes. It inhibits glucose uptake and increases liver gluconeogenesis (19). Progesterone mediates this effect by acting on progesterone receptor membrane component 1 in the liver that could exacerbate hyperglycemia in diabetes (19). Conversely, progesterone improves neuronal glucose metabolism through the augmentation of different glucose transporters in neuronal cells (20). The latter effect may be responsible for the neuroprotective role of progesterone.

The primary effects of progesterone on myometrial smooth muscle cells' contractility suggest that it may exert similar actions on smooth muscle cells of other tissues. PRs are present in vascular smooth muscle cells and progesterone administration in humans lowers blood pressure and inhibits angiotensin II-induced contraction in vascular muscle cells (21, 22). The relaxation effect of progesterone on vascular smooth muscle is independent of vascular endothelium and triggered directly via activation of PRs present in smooth muscle cells (23). The effects of progesterone on vascular smooth muscle cells' tone involve several actions on different signaling pathways that lead to smooth muscle relaxation. One prevalent mechanism is the activation of non-classical progesterone cell surface receptors that in turn activate an inhibitory G protein and subsequently activation of several intracellular pathways that upregulate nitric oxide intracellular levels which leads to inhibition of vascular smooth muscle contraction (24). Another pathway that is responsible for the inhibitory effect of progesterone on vascular smooth muscle cells contraction is the activation of the cyclic adenosine monophosphate (cAMP) pathway which inhibits contraction by reducing myosin light chain-20 (MLC 20) phosphorylation by activating

progesterone cell surface receptors directly and involving the MAP kinase/ERK-, Akt/PI3K signaling activation (23). Moreover, progesterone induces vascular smooth muscle relaxation by reducing intracellular calcium levels through activation of membrane progesterone receptor alpha (mPR α)-dependent signaling pathways, specifically, MAP and Akt signaling (25).

Similarly, progesterone targets the smooth muscle of the gastrointestinal tract and affects the contractile apparatus there. The remainder of this review will summarize the recent advances in the role of progesterone in gastrointestinal tract contraction relaxation pathways.

Progesterone in Gastrointestinal tract motility

Gastrointestinal tract motility is altered during pregnancy as a result of plasma hormonal changes such as progesterone levels. Progesterone affects gastrointestinal motility where it enhances gastric emptying at high doses as those seen toward the end of pregnancy (26). Smooth muscle cells from the colon of women with intractable constipation and slow transit time were studied by Zuo-LiangXiao et al. to elucidate the role of progesterone in constipation. They found that smooth muscle cells from women with constipation exhibited lower contraction compared to control samples. Moreover, PRs were overexpressed in smooth muscle cells from constipation women and there was downregulation of G α _q/11 and up-regulation of G α _s, responsible for contraction and relaxation respectively (27). Moreover, progesterone could increase gastric sensitivity to inhibitory gut neurotransmitters such as calcitonin gene-related peptide (CGRP) by enhancing CGRP receptors in gastric tissue. This effect is correlated with the levels of CGRP during pregnancy and could account in addition to the reported disturbed gut motility to the increased gut irritability as CGRP is one of the main sensory neurotransmitters in the gut (28). In vitro, progesterone inhibited the resting tension of the fundus and body longitudinal muscle strips, and it inhibited the mean

contractile amplitude of body and antrum longitudinal and circular muscles (29). The effect of progesterone was mostly a direct effect on gastric smooth muscle as inhibitors of other mediated pathways did not influence the effect of progesterone on gastric motility (29).

From another point of view, progesterone decreased gastrointestinal tract inflammatory cytokines such as IL-1 β and tumor necrosis factor-alpha. Which helps to protect the gut structure and prevents apoptosis after gut inflammation (30). This observation suggests the potential use of synthetic progesterone in such situations, especially if they are accompanied by diarrhea as these inflammatory cytokines are known to enhance gut motility (31, 32). Moreover, progesterone provides neuroprotective and anti-inflammatory effects on the enteric nervous system in Parkinson's disease (PD) mice model and thus could ameliorate the associated gut motility disturbances in PD. This observation could explain the beneficial effect of female hormones in PD susceptibility (33). However, these effects of progesterone on gut motility are indirect effects through the enteric nervous system.

Recently, the role of progesterone in gastric smooth muscle motility was investigated in more detail by Al-Shboul and co-authors (34-36). Single smooth muscle cells from rat gastric tissue were used to investigate the role of progesterone in gastric motility. Both isoforms of progesterone receptors (PR), PR-A and PR-B are present in gastric smooth muscle cells suggesting a direct site of action of progesterone (35). Preincubation of the dispersed smooth muscle cells 10 minutes before the administration of acetylcholine (Ach), inhibited the induced contraction by acetylcholine. Moreover, progesterone preincubation inhibited Ach-induced Rho kinase II activity without affecting its expression level suggesting that progesterone acts rapidly and directly of gastric smooth muscle cells to inhibit the contractile activity (34). The same group showed that the effect of progesterone on Ach-induced contraction was achieved by the production of nitric oxide

(NO) by progesterone from smooth muscle cells. Nitric oxide in turn induces the formation of cyclic guanosine monophosphate (cGMP) that activates protein kinase A (PKG) that plays a key role in smooth muscle relaxation (35).

Similarly, Colon smooth muscle contractility was increased in the inflammatory bowel syndrome model induced by water immersion and animal restraint in mice and this change was NO-dependent. Progesterone administrations reversed the increased contractility by elevating NO levels (37).

Future directions

Despite the obvious role of progesterone in gastrointestinal tract function, little investigations explored the topic in a mechanistic manner. This necessitates future studies to target the detailed mechanism of action of the inhibitory effect of progesterone on gut motility. Additionally, progesterone function in the GI tract should be studied in a pathological context, such as inflammatory bowel diseases, stress, and during pregnancy and lactation.

Conclusion:

Progesterone inhibits gastrointestinal tract motility by acting directly on gut smooth muscle cell surface PRs. This leads to rapid changes in the intracellular second messengers that affect contraction relaxation pathways without the involvement of gene transcription. One of the main downstream second messengers to PRs is NO that acts on Guanylate cyclase to generate cGMP and activation of the subsequent PKG. In turn PKG acts on many molecules in the contraction relaxation pathways to decrease phosphorylation of MLC-20 and eventually inhibition of contraction Figure 2.

Figures and legends:




Figure1: progesterone mechanism of action

The classical pathway involves the nuclear progesterone receptor (nPR) dimerization and translocation to the nucleus to induce genomic effects via activating or inhibiting gene transcription. The non-classical pathway is rapid action that involves membrane progesterone receptor (mPR) and the subsequent activation of several second messengers such as elevation of intracellular calcium (Ca^{2+}), extracellular signal-regulated kinases (ERK) and Protein kinase B (AKT)




Figure 2: Effect of progesterone on gut smooth muscle contraction: Progesterone activates its cell surface receptors (mPR), PR that leads to NO production. NO production leads to the generation of cGMP from GTP and in turn activation of PKG. PKG acts to inhibit signaling pathways that provoke contraction such as RhoK and activates the signaling that leads to muscle relaxation such as MLCP. mPR: progesterone receptor, NO: Nitric oxide, cGMP: cyclic guanosine monophosphate, GTP: guanosine triphosphate, PKG: protein kinase G, RhoK: Rho kinase, MLCP: myosin light chain phosphatase

References:

1. Miller WL, Auchus RJ. The molecular biology, biochemistry, and physiology of human steroidogenesis and its disorders. *Endocr Rev.* 2011;32(1):81-151. doi:10.1210/er.2010-0013
2. Cable JK, Grider MH. Physiology, Progesterone. *StatPearls [Internet]*. 2020;
3. Oettel M, Mukhopadhyay AK. Progesterone: the forgotten hormone in men? *Aging Male.* Sep 2004;7(3):236-57. doi:10.1080/13685530400004199
4. Henderson VW. Progesterone and human cognition. *Climacteric.* 2018;21(4):333-340. doi:10.1080/13697137.2018.1476484
5. Taraborrelli S. Physiology, production and action of progesterone. *Acta obstetrica et gynecologica Scandinavica.* 2015;94:8-16.
6. Garg D, Ng SSM, Baig KM, Driggers P, Segars J. Progesterone-Mediated Non-Classical Signaling. *Trends in Endocrinology & Metabolism.* 2017/09/01/ 2017;28(9):656-668. doi:<https://doi.org/10.1016/j.tem.2017.05.006>
7. Wetendorf M, DeMayo FJ. Progesterone receptor signaling in the initiation of pregnancy and preservation of a healthy uterus. *Int J Dev Biol.* 2014;58(2-4):95-106. doi:10.1387/ijdb.140069mw
8. Valadez-Cosmes P, Vázquez-Martínez ER, Carbon M, Camacho-Arroyo I. Membrane progesterone receptors in reproduction and cancer. *Molecular and cellular endocrinology.* 2016;434:166-175.
9. Simoncini T, Genazzani AR. Non-genomic actions of sex steroid hormones. *European journal of endocrinology.* 2003;148(3):281-292.
10. Bergeron C. Morphological changes and protein secretion induced by progesterone in the endometrium during the luteal phase in preparation for nidation. *Human reproduction.* 2000;15(suppl_1):119-128.
11. Putnam CD, Brann DW, Kolbeck RC, Mahesh VB. Inhibition of uterine contractility by progesterone and progesterone metabolites: mediation by progesterone and gamma amino butyric acidA receptor systems. *Biology of reproduction.* 1991;45(2):266-272.
12. Arab H, Alharbi AJ, Oraif A, et al. The Role Of Progestogens In Threatened And Idiopathic Recurrent Miscarriage. *Int J Womens Health.* 2019;11:589-596. doi:10.2147/ijwh.S224159
13. Savouret J, Misrahi M, Milgrom E. Molecular action of progesterone. *Oxford reviews of reproductive biology.* 1988;10:293-347.
14. Tsutsui K, Sakamoto H, Ukena K. Biosynthesis and action of neurosteroids in the cerebellar Purkinje neuron. *The Journal of Steroid Biochemistry and Molecular Biology.* 2003/06/01/ 2003;85(2):311-321. doi:[https://doi.org/10.1016/S0960-0760\(03\)00229-2](https://doi.org/10.1016/S0960-0760(03)00229-2)
15. Frick KM, Kim J. Mechanisms underlying the rapid effects of estradiol and progesterone on hippocampal memory consolidation in female rodents. *Hormones and behavior.* 2018;104:100-110.
16. Kavaliers M, Bishnoi IR, Ossenkopp K-P, Choleris E. Differential effects of progesterone on social recognition and the avoidance of pathogen threat by female mice. *Hormones and Behavior.* 2021;127:104873.
17. Patel S. *The Effects of Ovarian Hormones on Memory Bias and Progesterone Receptors in Female Rats.* Concordia University; 2020.
18. Yao W, Dai W, Shahnazari M, et al. Inhibition of the progesterone nuclear receptor during the bone linear growth phase increases peak bone mass in female mice. *PLoS one.* 2010;5(7):e11410.
19. Lee SR, Choi W-Y, Heo JH, et al. Progesterone increases blood glucose via hepatic progesterone receptor membrane component 1 under limited or impaired action of insulin. *Sci Rep.* 2020;10(1):16316-16316. doi:10.1038/s41598-020-73330-7

20. Wu H, Wu Z-g, Shi W-j, et al. Effects of progesterone on glucose uptake in neurons of Alzheimer's disease animals and cell models. *Life Sciences*. 2019/12/01/ 2019;238:116979. doi:<https://doi.org/10.1016/j.lfs.2019.116979>
21. Horwitz KB, Horwitz L. Canine vascular tissues are targets for androgens, estrogens, progestins, and glucocorticoids. *The Journal of clinical investigation*. 1982;69(4):750-758.
22. Barbagallo M, Dominguez LJ, Licata G, et al. Vascular effects of progesterone: role of cellular calcium regulation. *Hypertension*. 2001;37(1):142-147.
23. Pang Y, Thomas P. Progesterone induces relaxation of human umbilical cord vascular smooth muscle cells through mPR α (PAQR7). *Molecular and Cellular Endocrinology*. 2018/10/15/ 2018;474:20-34. doi:<https://doi.org/10.1016/j.mce.2018.02.003>
24. Pang Y, Dong J, Thomas P. Progesterone increases nitric oxide synthesis in human vascular endothelial cells through activation of membrane progesterone receptor- α . *American Journal of Physiology-Endocrinology and Metabolism*. 2015;308(10):E899-E911.
25. Pang Y, Thomas P. Role of mPR α (PAQR7) in progesterone-induced Ca²⁺ decrease in human vascular smooth muscle cells. *Journal of Molecular Endocrinology*. 01 Oct. 2019 2019;63(3):199-213. doi:10.1530/jme-19-0019
26. Liu C-Y, Chen L-B, Liu P-Y, Xie D-P, Wang PS. Effects of progesterone on gastric emptying and intestinal transit in male rats. *World J Gastroenterol*. 2002;8(2):338-341. doi:10.3748/wjg.v8.i2.338
27. Xiao Z-L, Pricolo V, Biancani P, Behar J. Role of progesterone signaling in the regulation of G-protein levels in female chronic constipation. *Gastroenterology*. 2005/03/01/ 2005;128(3):667-675. doi:<https://doi.org/10.1053/j.gastro.2004.12.001>
28. Yang X, Liu R, Dong Y. Regulative effects of ovarian steroids on rat gastric motility and sensitivity. *ACTA PHYSIOLOGICA SINICA-CHINESE EDITION*-. 2006;58(3):275.
29. Wang F, Zheng T-Z, Li W, Qu S-Y, He D-Y. Action of progesterone on contractile activity of isolated gastric strips in rats. *World J Gastroenterol*. 2003;9(4):775-778. doi:10.3748/wjg.v9.i4.775
30. Chen G, Shi J-X, Qi M, Wang H-X, Hang C-H. Effects of Progesterone on Intestinal Inflammatory Response, Mucosa Structure Alterations, and Apoptosis Following Traumatic Brain Injury in Male Rats. *Journal of Surgical Research*. 2008/06/01/ 2008;147(1):92-98. doi:<https://doi.org/10.1016/j.jss.2007.05.029>
31. Nalli AD, Kumar DP, Mahavadi S, et al. Hypercontractility of intestinal longitudinal smooth muscle induced by cytokines is mediated by the nuclear factor- κ B/AMP-activated kinase/myosin light chain kinase pathway. *Journal of Pharmacology and Experimental Therapeutics*. 2014;350(1):89-98.
32. Grider J. Interleukin-1 β selectively increases substance P release and augments the ascending phase of the peristaltic reflex. *Neurogastroenterology & Motility*. 2003;15(6):607-615.
33. Jarras H. Effects of progesterone at the enteric level in a mouse model of Parkinson's disease. 2019;
34. Al-Shboul O, Mustafa A, Al-hashimi F. Non-genomic effects of progesterone on Rho kinase II in rat gastric smooth muscle cells. *Journal of Smooth Muscle Research*. 2013;49:55-62.
35. Al-Shboul OA, Mustafa AG, Omar AA, et al. Effect of progesterone on nitric oxide/cyclic guanosine monophosphate signaling and contraction in gastric smooth muscle cells Corrigendum in/10.3892/br. 2019.1251. *Biomedical reports*. 2018;9(6):511-516.
36. Al-Shboul OA, Al-Rshoud HJ, Al-Dwairi AN, et al. Changes in gastric smooth muscle cell contraction during pregnancy: effect of estrogen. *Journal of pregnancy*. 2019;2019
37. LIU H-x, PAN H, SHI Z, WANG H. Estradiol and Progesterone Regulate the Activities of Colonic Smooth Muscle in Water-immersion and Restraint Stress Mice by Modulating NO. *Progress in Modern Biomedicine*. 2013:05.