

## **Roles of Isometric Contraction Training in Promoting Neuroprotection and Angiogenesis after Stroke in Adult Rats**

**Running title:** post-stroke isometric contraction training

Chengyao Mei<sup>1#\*</sup>, Teng Ma<sup>1#</sup>

1. Pukou branch, Jiangsu People's Hospital, Nanjing city, 211800, Jiangsu province, P. R. China.

\*Corresponding author,

Chengyao Mei,

Pukou branch,

Jiangsu People's Hospital,

166 Shanghe street, Pukou District, 211800,

Nanjing city, Jiangsu province,

P. R. China.

Email: meicy666214@163.com

**# both authors contributed equally to this article, and both as the first authors.**

2 **Objective:** To explore potentially neuroprotective effects of isometric contraction  
3 training (ICT) after stroke. **Methods:** 100 rats were randomly divided into a  
4 sham-operated group and middle cerebral artery occlusion (MCAO) modeling groups.  
5 The sham group after surgery was observed for 14 days. After MCAO, some rats  
6 received isometric contraction training (ICT) which was as follows: an atraumatic  
7 tourniquet was placed around left or right hind limb to achieve hind limb ischemia for  
8 5 minutes, followed by 5 minutes of reperfusion, 4 cycles for one time, once a day,  
9 and five days per week. The MCAO modeling groups included the following four  
10 groups: i) a group only received MCAO, and was observed for seven days  
11 (MCAO-7d), ii) a group only received MCAO, and was observed for 14 days  
12 (MCAO-14d), iii) a group, after MCAO, received ICT for seven days (ICT-7d), and iv)  
13 a group, after MCAO, received ICT for 14 days (ICT-14d). Brain infarct area,  
14 behavioral outcomes, the number of neurons, apoptosis, cerebral edema and cerebral  
15 water content were assessed, respectively. The mRNA expression of vascular  
16 endothelial growth factor (VEGF) was assayed with RT-PCR, and protein expression  
17 of VEGF was quantified with western blot. **Results:** compared with MCAO controls,  
18 cerebral infarction, neurological deficits and neuronal apoptosis were reduced  
19 significantly in the ICT groups, while the number of neurons was increased. Moreover,  
20 the mRNA expression of VEGF and protein expression of VEGF were enhanced after  
21 1 and 2 weeks of ICT. **Conclusion:** ICT may promote angiogenesis and  
22 neuroprotection after ischemic stroke and this new remodeling method provide a  
23 novel strategy for rehabilitation of stroke patients.

24 **Keywords:** isometric contraction training; middle cerebral artery occlusion (MCAO);  
25 neuroprotection; angiogenesis; stroke; vascular endothelial growth factor (VEGF)

## 26 **Introduction**

27 Ischemic stroke is one of the main causes of physical disability, which brings a

28 heavy burden to individuals and society [1]. So far, tissue plasminogen activator (tPA)  
29 is the only drug approved by the US Food and Drug Administration to treat ischemic  
30 stroke, while the administration of tPA is limited by its narrow therapeutic time  
31 window of 3 to 4.5 hours from ischemic stroke onset, and a large amount of patients  
32 do not receive timely tPA. Of the 795,000 cases with stroke that occur every year in  
33 Europe, about a quarter are recurrent cases [2]. The ways of preventing stroke  
34 recurrence include the control of risk factors, antithrombotic or antiplatelet therapies,  
35 and other interventions for atherosclerotic disease [3]. Although great progress has  
36 been made in vascular reconstruction technology and drug therapy for stroke in recent  
37 years, there is still a lack of safe, effective and non-invasive strategies for preventing  
38 stroke events and reducing the disability rate of stroke patients.

39 In recent decades, with the advancement of medical technology, more and more  
40 stroke patients survived the initial injury, but most patients suffer from neurological  
41 dysfunctions such as motor, learning, memory, and cognitive dysfunction, which  
42 significantly reduce the quality of daily life [4]. The sudden decrease of focal blood  
43 flow is the main cause of the occurrence of stroke [5]. The recovery of physical  
44 functions can significantly strengthen the independent ability of stroke survivors and  
45 improve their quality of life, and many studies have shown that exercise training can  
46 improve motor function after cerebral infarction, promote functional recovery, and  
47 exert neuroprotective effects [6-9].

48 Clinical studies have demonstrated that patients with a history of angina pectoris  
49 have a smaller average infarct size and a lower mortality rate when myocardial  
50 infarction occurs [10]. Similarly, studies reveal that in subsequent stroke events,

51 patients with transient ischemic attack (TIA) tend to have better recovery than  
52 non-TIA patients, suggesting that TIA might have a neuroprotective effect [11], and  
53 ischemic tolerance might play a role in TIA patients [12].

54 Isometric contraction is a muscle contraction in which the length remains  
55 constant but the tension changes, and due to the increased muscle tension, the  
56 resistance of blood vessels passing through the muscle increases, which can lead to  
57 different degrees of blood flow blockage, thereby increasing ventricular pressure [13,  
58 14]. We know that isometric contraction training (ICT) is an intervention based on the  
59 controlled application of ischemic tolerance [15], can create a series of noninvasive,  
60 reversible and controllable ischemic events in the normal skeletal muscles far away  
61 from the original ischemic site (i.e. the heart or brain) [16], and lead to local ischemia  
62 of normal limb contractile muscles, thus increasing ventricular pressure. It has been  
63 reported that blood flow in muscles generally includes 40-50% of the maximum  
64 voluntary contraction (MCV) [17]. Therefore, that isometric handgrip exercises  
65 obtained more than 50% MVC can be defined as physical ischaemia training (PIT),  
66 and PIT is a reversible ischemia of normal skeletal muscle by tourniquet or isometric  
67 contractions.

68 Clinical studies have shown the therapeutic potential of ICT in restoring  
69 coronary blood flow through central response induced by peripheral biological effects  
70 [18-20]. Lin et al.[15] found that isometric grip exercise promoted the recruitment and  
71 growth of distal collateral circulation in patients with coronary heart disease. In  
72 addition, Lin *et al.* also found that isometric exercise resulted in increased coronary  
73 collateral blood flow during acute vascular occlusion[18], and PIT induced by ICT

74 promoted the formation of collateral circulation of distal ischemic myocardium in  
75 patients with CAD[15]. Gao *et al.* investigated the patients with coronary heart  
76 disease complicated with heart failure, and found that PIT can improve the level of  
77 vascular endothelial growth factor (VEGF) in peripheral blood and quality of life[21].

78 ICT is a type of remote ischemic post-conditioning relying on a training effect.  
79 ICT repeatedly induces temporary ischemia in distant skeletal muscles, so that  
80 vascular endothelial growth factor (VEGF) is supposed to be released in the  
81 stimulated area[21], and VEGF circulates to the distal area to promote angiogenesis  
82 and eventually form a “biological bypass”[22]. The angiogenic effect of ICT has  
83 been confirmed in several previous studies featuring animal models of myocardial  
84 infarction [23, 24].

85 The angiogenic effects of ICT may also play a role in neuroprotection after  
86 ischemic stroke. In addition, ICT may have the effect of directly enhancing the  
87 expression of VEGF, thereby further stimulating the formation of new neuronal cells  
88 [25]. There are currently limited clinical data on the roles of ICT in cerebral ischemia  
89 [24]. Hahn *et al.*[26] showed that remote per-conditioning by transient limb ischemia  
90 is simple and clinically relevant, and has a strong neuroprotective effect in the model  
91 of local cerebral ischemia reperfusion injury. Zhen *et al.* [27]found in a randomized  
92 controlled trial (RCT) of 20 patients that ICT could effectively increase the expression  
93 of VEGF and the recruitment of EPCs, and promote the formation of collateral  
94 circulation. As a cytokine, VEGF binds to and activates VEGFR on the cell membrane.  
95 The signal can be transmitted to the PI3K/AKT pathway, and the pathway and  
96 downstream factors were activated [28]. In terms of neuroprotection, the PI3K/AKT

97 pathway can prevent nerve injury via nerve growth factor (NGF)/Tropomyosin  
98 receptor kinase A (TrkA) signaling [29]. PI3K/Akt pathway is an important signaling  
99 pathway that promotes neuron survival [30, 31].

100 In the present study, we used a rat model of transient middle cerebral artery  
101 occlusion (MCAO) to test the hypothesis that ICT might have angiogenic and  
102 neuroprotective effects after a stroke. Specifically, we hypothesized that ICT-induced  
103 angiogenesis might reduce the occurrence of cerebral edema and the infarct size, and  
104 ICT might reduce cell apoptosis and promote the formation of new nervous cells.

105

## 106 **Materials and Methods**

107 All experimental procedures were approved by the Key Laboratory of Nerve  
108 Regeneration of Nantong University (Nantong, China) and were carried out in  
109 accordance with the institutional animal care guidelines of Nantong University. This  
110 study was approved by the Ethics Committee of Jiangsu Province, China, with  
111 approval No. S20141103-402.

### 112 *Study design*

113 A randomized controlled animal study was performed. The randomization was  
114 achieved with computer generated random numbers. 100 Sprague-Dawley male rats  
115 weighing 350-400g were randomly divided into a sham-operated group and MCAO  
116 modeling groups. The sham group after surgery was observed for 14 days. The  
117 MCAO modeling groups included the following four groups: i) a group only received  
118 MCAO , and was observed for seven days (MCAO-7d), ii) a group only received  
119 MCAO, and was observed for 14 days (MCAO-14d), iii) a group, after MCAO,  
120 received ICT for seven days (ICT-7d), and iv) a group, after MCAO, received ICT for  
121 14 days (ICT-14d). In each group, animals were randomly assigned to six different

122 outcomes for evaluation, and each animal only was used to evaluate one of those  
123 outcomes. All procedures and measurements were performed by an investigator who  
124 was blind to the experimental groups. The concealed animal codes were revealed only  
125 after the completion of behavioral and histological analysis.

#### 126 *Chemicals and reagents*

127 2,3,5-Triphenyltetrazolium chloride (TTC), propidium iodide (PI), cresyl violet  
128 acetate, and hemoglobin assay kit were purchased from Sigma (St. Louis, MO, USA).

129 A terminal deoxynucleotidyl transferase (TdT)-mediated dUTP-biotin nick end  
130 labeling (TUNEL) assay kit was purchased from Promega (Madison, WI, USA).

131 Mammalian protein extraction reagent (M-PER), tissue protein extraction reagent  
132 (T-PER), membrane protein extraction reagent (Mem-PER), and rabbit polyclonal  
133 antibody against vascular endothelial growth factor (VEGF) were purchased from  
134 Abcam (Cambridge, Mass, UK). Mammalian protein extraction reagent (M-PER),  
135 tissue protein extraction reagent (T-PER), membrane protein extraction reagent  
136 (Mem-PER), and Coomassie plus Bradford assay kit were purchased from Thermo  
137 Scientific Pierce (Rockford, IL).

#### 138 *MACA model construction and sham-operated processing*

139 Focal cerebral ischemia was induced by the transient MCAO on the right [32,  
140 33]. Surgeries were performed by CM (physician) and SY (biologist). Both had been  
141 trained by experienced experimentalists from the Jiangsu Key Laboratory of  
142 Neuroregeneration, Nantong University, China where the surgeries were performed.  
143 Rats lay supine after anesthesia by intraperitoneal injection of sodium pentobarbital  
144 (25 mg/kg). A heating pad was used to maintain their body temperature at  $37\pm 0.5^{\circ}\text{C}$

145 during the surgical procedure. The right common carotid artery (CCA), internal  
146 carotid artery (ICA), and external carotid artery (ECA) of each rat were surgically  
147 exposed via a neck incision. The occipital artery and the superior thyroid artery were  
148 cauterized and cut. ECA was permanently ligated using 6-0 silk thread as rostrally as  
149 possible. CCA was transiently ligated as caudally as possible and ICA was transiently  
150 ligated as rostrally as possible using microvascular clips to expose enough working  
151 space. Microscissors was used to cut the ECA near the permanent ligature. 6-0 silk  
152 thread was used to ligate around the filament insertion site in the ECA, and the  
153 ligature was kept tight enough to prevent bleeding, but loose enough to allow the  
154 filament to advance. A filament (4–0 nylon suture with rounded tip) was inserted into  
155 the CCA from the insertion site, and gently entered the ICA when the microvascular  
156 clip on the ICA was removed. The origin of middle cerebral artery (MCA) lied 18–20  
157 mm from the CCA bifurcation in rats. When the insertion was close to 18mm (marked  
158 on the filament), please pay attention to the feeling of your hand. Once a slight  
159 resistance was felt, it meant the filament reached the MCA region. Then, we tightened  
160 the ligature around the filament and started to record the time of ischemia. After  
161 90-min MCAO, the transient ligature around the filament was loosened, the nylon  
162 filament was withdrawn when the microvascular clips were added on the ICA and  
163 CCA again, then we permanently tighten the ligature around the insertion site to  
164 prevent bleeding, and the microvascular clips were removed again to allow  
165 reperfusion. Followingly, the area was moistened with sterile saline, lidocaine was  
166 applied as a topical analgesic, and the wound was closed with sutures. After the rat  
167 was awake, if signs of unstable standing, paralyzed left limb was and circling were



168 observed, and the tail suspension test suggested the rat only swung to one side, it  
169 indicated that the MCAO model was successfully established. The rats in the sham  
170 group underwent the same surgical procedures except for the insertion of the filament,  
171 After the surgery, all rats were put back into their cages where they had access to food  
172 and water.

### 173 ***Experimental interventions***

174 ICT started 24 hours after MCAO. Rats were fixed by the conventional  
175 laboratory fixator. An atraumatic tourniquet was placed around the left or right  
176 hind-limb to achieve hind limb ischemia for 5 minutes, followed by 5 minutes of  
177 reperfusion, which were conducted a total of 4 cycles for one time, once a day, and  
178 five days per week [26]. Circulatory arrest in hind limbs was confirmed by vascular  
179 Doppler ultrasound [34].

180 Rats were kept under controlled environmental conditions with an ambient  
181 temperature of  $22 \pm 1^\circ\text{C}$ , a relative humidity of 65% and a light/dark cycle of 12 hours  
182 (h), and they had free access to food and water during the whole experimental period.  
183 All efforts were made to reduce the number of animals used in our study and avoid  
184 unnecessary sufferings.

### 185 **Outcome assessment**

#### 186 ***Animal euthanasia***

187 The rats were anesthetized with an overdose of sodium pentobarbital (50 mg/kg),  
188 and then were sacrificed by transcardial perfusion.

#### 189 ***Assessment of neurobehavioral deficits***

190 Modified neurological severity score (mNSS) [35] was assessed at day 7 and 14  
191 after MCAO. The mNSS is a comprehensive score of motor, sensory, reflex and

192 balance tests, ranging from 0 to 18 (normal score 0; maximal deficit score 18), and  
193 can be classified into three levels: 13-18: severe impairment; 7-12; moderate  
194 impairment; and 1- 6; mild impairment [35].

#### 195 ***Edema***

196 The examination for vasogenic edema was performed [36]. The brains of the  
197 euthanized rats were took. After removing the pons and olfactory bulb, each brain was  
198 weighed to obtain the wet weight. After the brain was dried at 60°C for 72 h, the dry  
199 weight was obtained. Brain water content percent was calculated using the following  
200 equation: water content (%) = [(wet weight - dry weight)/wet weight]×100% .

#### 201 ***Infarct area***

202 The measurement method of the infarct area has been described in many  
203 studies[37]. We took out the brains of the euthanized rats, and cut them serially into  
204 six 2-mm sections from the frontal pole using a rat brain matrix (Sunny Instruments,  
205 Beijing, China). The sections were stained with 2% solution of 2,3,5- Triphenyl  
206 tetrazolium chloride (TTC) at 37°C for 30 min, and then fixed with 4% formaldehyde  
207 buffer solution for 1h at room temperature. The area unstained by TTC was  
208 considered as the infarct area. The infarction area was calculated by the percentage of  
209 the unstained areas in the total contralateral hemisphere. A computerized image  
210 analysis system (Leica Imaging System Ltd., Cambridge, UK) was used to analyze the  
211 percentage of TTC-stained tissue with an average of more than 6 slices.

#### 212 ***Neuron Nissl staining***

213 Brains were taken out from the skull, post-fixed in buffered 4%  
214 paraformaldehyde, dehydrated in a graded sucrose series, and cut into coronal sections

215 (20- $\mu$ m thick) from the anterior commissure to the hippocampus on a cryostat  
216 microtome. A total of 40–50 sections were installed and frozen at -40 degrees. The  
217 frozen sections were stained with conventional Nissl staining (0.1% cresyl violet  
218 solution) at 37°C for 30 min, dehydrated, and installed in dibutyl phthalate in xylene.  
219 Two technicians independently calculated the number of neurons per high-power  
220 field.

### 221 *Apoptosis*

222 TUNEL assay was used for assessing apoptosis in the frozen brain sections [38].  
223 The sections were fixed with 4% methanol-free formaldehyde. The brain slices were  
224 stored in an equilibration buffer and then covered with DNA strand breaks labeled  
225 with fluorescein-12-dUTP. Recombinant TdT was added. Finally, the following steps  
226 were carried out in a dark room. Brain sections were incubated with 2 $\times$ SSC (300 mM  
227 sodium chloride and 30 mM sodium citrate, pH 7.4) at room temperature for 15 min,  
228 and then stained with 1  $\mu$ g/ml propidium iodide (PI). Apoptotic cells were positioned  
229 as bright green cells on a red background under a scanning laser confocal microscope  
230 (Leica, Germany). Data were expressed as the ratio of apoptotic cells to total cells.

### 231 *VEGF mRNA expression*

232 In order to detect VEGF gene expression, brains were taken after euthanasia.  
233 Tissue samples were collected from two cortical regions of interest (ipsilateral and  
234 contralateral to the MCAO) and stored at -80°C until further processing.

235 The frozen rat brain tissue was took, and quickly placed into a centrifuge tube.  
236 The brain tissue was weighed, and Trizol was added in proportion to fully mix the  
237 brain tissue. The brain tissue homogenate was separated, chloroform was added to

238 each tube for shaking and standing, and finally centrifugation was performed to  
239 dissolve the total RNA of brain tissue. Reverse transcription buffer and reverse  
240 transcriptase were added, respectively, and then they were placed in a 37°C water bath  
241 for reverse transcription to generate cDNA. The primers used for polymerase chain  
242 reaction (PCR) were Sense 5'-TGCACCCACGACAGAAGGGGA-3', Antisense  
243 5'-TCACCGCCTTGGCTTGTCACAT-3' for VEGF, and Sense  
244 5'-GAGAGGGAAATCGTGCGT-3', Antisense 5'-GGAGGAAGAGGATGCGG-3'  
245 for  $\beta$ -actin. Real time (RT)-PCR was performed on an ABI StepOne PCR in a 64-well  
246 plate using a final volume of 10  $\mu$ l and the following cycle conditions: 95 °C for 10  
247 min, and then 45 cycles of 15 s at 95°C and 1 min at 60 °C. The specificity of each  
248 target amplicon was assessed by dissociation curve analysis, and the results of  
249 quantification were given according to the formula  $2^{-\Delta\Delta ct}$ , using  $\beta$ -actin as the internal  
250 standard. RT-PCR for each sample was performed in triplicate.

### 251 *VEGF protein*

252 Western blot was used to detect the expression level of VEGF protein. The  
253 protein samples were extracted from the brain tissue of the ipsilateral/contralateral  
254 hemisphere. The protein was quantified with a Coomassie plus Bradford assay kit. For  
255 each sample, an equal amount of protein was separated in 12%  
256 SDS-polyacrylamide gels, and subsequently blotted onto PVDF membranes. The  
257 membranes were incubated with VEGF antibody (1:1000) and  $\beta$ -actin (1:1000) at 4°C  
258 overnight, washed with 0.01% TBST three times, and then injected with purified  
259 donkey anti-rabbit IgG (1:5000) at room temperature for 2h. The images were  
260 scanned with Odyssey infrared imaging system (LI-COR, USA), and the results were

261 analyzed using PDQuest 7.2.0 software (Bio-Rad, USA). Integrated density was  
262 quantified by background subtraction and normalization to the b-actin signal. The  
263 results were expressed as mean standard deviation (SD) of (VEGF protein from  
264 ipsilateral/VEGF protein from contralateral hemisphere) \* 100.

## 265 **Statistical analysis**

266 Data analysis was performed with SPSS 16.0 (SPSS, Chicago, IL, USA) and  
267 Stata 14 (Stata Corporation, Texas, USA). Mean differences between the groups were  
268 compared by one-way analysis of variance (ANOVA) and Bonferroni corrected  
269 pairwise post-hoc tests in case of a significant overall F-test. Mortality across groups  
270 was examined with logistic regression followed by Bonferroni corrected pairwise  
271 comparisons in case of a significant overall Wald-test. P-values smaller than 0.05 was  
272 considered statistically significant.

273

## 274 **Results**

### 275 *Isometric Contraction Training reducing mortality*

276 The mortality of the sham-operated, MCAO-7d, MCAO-14d, ICT-7d and  
277 ICT-14d groups was 0, 45.0%, 45.0%, 30.0%, and 20.0%, respectively. As shown in  
278 Figure 1A, compared with that in the MCAO-14d group, the mortality in the ICT-14d  
279 group was significantly reduced ( $F=4.27$ ,  $P<0.05$ ), and however, there was no  
280 significant difference in mortality between the ICT-7d group and the ICT-14d group  
281 ( $P>0.05$ ).

### 282 *Isometric Contraction Training reducing edema*

283 As shown in Figure 1B, ICT reduced stroke-induced edema. The mean water

284 content percentage of the sham-operated, MCAO-7d, MCAO-14d, ICT-7d and  
285 ICT-14d groups was 74.3%, 88.4%, 87.99%, 79.8%, and 72.6%, respectively.  
286 Compared with that in the sham group, brain water content in the MCAO-7d and  
287 MCAO-14d groups both significantly increased ( $F=5.93$ ,  $p<0.01$ ). Compared with  
288 that in the MCAO-7d group, brain water content in the ICT-7d group decreased  
289 significantly ( $F=7.29$ ,  $p<0.05$ ). Compared with that in the MCAO-14d group, brain  
290 water content in the ICT-14d group decreased significantly ( $F=10.73$ ,  $p<0.05$ ).

### 291 ***Isometric Contraction Training reducing neurobehavioral deficits***

292 The modified neurological severity scores measured on the 7th and 14th days  
293 after MCAO were shown in Figure 1C. The mean mNSS of the sham-operated,  
294 MCAO-7d, MCAO-14d, ICT-7d and ICT-14d groups was 0, 4.7, 4.1, 3.1, and 1.7,  
295 respectively. Compared with that in the MCAO-7d group, the mNSS in the ICT-7d  
296 group significantly decreased ( $F=7.33$ ,  $p<0.05$ ). Compared with that in the  
297 MCAO-14d group, the mNSS in the ICT-14d group significantly decreased ( $F=9.94$ ,  
298  $p<0.05$ ). All rats in the sham-operated group performed well without neurobehavioral  
299 deficits.

### 300 ***Isometric Contraction Training reducing brain infarct area***

301 Figure 2A showed the cerebral infarction area evaluated by triphenyl tetrazolium  
302 chloride (TTC) staining on the 7th and 14th day after MCAO. Normal brain tissue  
303 was dark red with staining, while the infarct area was pale gray without staining. No  
304 infarction was observed in the sham-operated group. The infarct area in the ICT  
305 groups reduced significantly (Figure 2B). The mean infarct area percentage of the  
306 sham-operated, MCAO-7d, MCAO-14d, ICT-7d and ICT-14d groups was 0, 39.64%,

307 31.09% 30.21% and 18.39%, respectively. Compared with that in the MCAO-7d  
308 group, the infarct area in the ICT-7d group decreased significantly ( $F=4.13$ ,  $p<0.05$ ).  
309 Compared with that in the MCAO-14d group, the infarct area in the ICT-14d group  
310 decreased significantly ( $F=10.53$ ,  $p<0.01$ ). In addition, the brain infarct area of the  
311 ICT-14d group was significantly smaller than that in the ICT-7d group ( $F=6.27$ ,  
312  $p<0.05$ ).

313 *The neuroprotective effect of Isometric Contraction Training assessed by Nissl*  
314 *staining*

315 As shown in the Figure 3 of the ipsilateral brain cortex, neurons in the  
316 sham-operated group had a normal shape with Nissl bodies deeply stained, and  
317 showed an integrative and granular-like configuration, while neurons in the  
318 MCAO-7d and MCAO-14d groups were apparently hypertrophic with Nissl bodies  
319 lightly stained. Neurons in the MCAO-7d and MCAO-14d groups were sparsely  
320 distributed, and the number of visible Nissl granules reduced markedly. These  
321 findings indicated that compared with the sham-operated group, transient cerebral  
322 ischemia resulted in a significant increase in the cortical necrosis in the MCAO-7d  
323 and MCAO-14d groups. Compared with that in the MCAO-7d group, the number of  
324 neurons per visual field in the ICT-7d group increased significantly ( $F=1.71$ ,  $p<0.05$ ).  
325 Compared with that in the MCAO-14d group, the number of neurons per visual field  
326 in the ICT-14d group increased significantly ( $F=5.48$ ,  $p<0.05$ ). In addition, compared  
327 with that in the sham-operated group, the number of neurons per visual field in the  
328 ICT-14d group was significantly smaller ( $F=0.73$ ,  $p<0.05$ ). These findings indicated  
329 that ICT reduced MCAO-induced neuronal injury in the cortex.

330 ***Isometric Contraction Training inhibiting MCAO-induced neuronal apoptosis in***  
331 ***the rat brain***

332 As shown in the Figure 4, in the sham-operated group, TUNEL-positive cells  
333 were hardly observed throughout the brain. The mean TUNEL-positive cells  
334 percentage of the sham-operated, MCAO-7d, MCAO-14d, ICT-7d and ICT-14d  
335 groups was 0, 43%, 37% 30% and 17%, respectively. Compared with the MCAO-7d  
336 group, the number of TUNEL-positive cells of the cortex in the ICT-7d group was  
337 significantly lower ( $P<0.01$ ). Compared with the MCAO-14d group, the number of  
338 TUNEL-positive cells of the cortex in the ICT-14d group was significantly lower  
339 ( $F=2.96$ ,  $P<0.01$ ). In addition, compared with the ICT-7d group, the number of  
340 TUNEL-positive cells of the cortex in the ICT-14d group was significantly lower  
341 ( $F=5.43$ ,  $P<0.05$ ). These findings suggested that longer ICT could improve the  
342 recovery of brain function.

343 ***Effects of Isometric Contraction Training on the expression of VEGF in rats with***  
344 ***cerebral ischemia***

345 The expression level of VEGF protein in the ipsilateral and contralateral  
346 hemisphere on the 7th and 14th days after MCAO was evaluated by western blot  
347 (Figure 5A and 5B). Compared with that in the MCAO-7d group, VEGF protein  
348 expression in the ICT-7d group increased significantly ( $F=7.48$ ,  $p<0.05$ ). Compared  
349 with that in the MCAO-14d group, VEGF protein expression in the ICT-14d group  
350 increased significantly ( $F=10.79$ ,  $p<0.05$ ). As shown in the Figure 5C, the level of  
351 VEGF mRNA expression increased to varying degrees after ICT, and was the highest  
352 in the ICT-14d group.



353

354 **Discussion**

355 This study demonstrated that ICT reduced the infarct area, attenuated  
356 stroke-induced edema, inhibited neuronal apoptosis and improved neuroprotective  
357 recovery by activating the endogenous neuroprotective program. The possible  
358 mechanism behind these findings is that ICT promotes the expression of VEGF in the  
359 infarct area by inducing reversible ischemia in distal skeletal muscle, and VEGF in  
360 turn promotes angiogenesis and reduces nerve injury, both of which contribute to  
361 neuroprotection.

362 The cardioprotective effect of ICT has previously been evaluated in experimental  
363 [25, 39] and clinical [40-43] studies. It was also demonstrated that the beneficial  
364 effects of ICT on the ventricular myocardium were not species specific [44]. The  
365 mechanism of ICT is to mobilize endothelial progenitor cells and promote vascular  
366 remodeling by up-regulating the levels of VEGF and NO [24]. Starting from the  
367 hypothesis that both cerebral ischemia and cardiac ischemia are both ischemic  
368 diseases, we demonstrated the neuroprotective effects of ICT in the brain.

369 The concept of ICT is different from ischemic preconditioning (IPC). IPC refers  
370 to the delay of cell death after coronary artery occlusion through short ischemic  
371 training before myocardial ischemia and infarction [45]. Unlike IPC, ICT causes  
372 transient ischemia of skeletal muscle through repeated isometric contraction training,  
373 thereby promoting the formation of arterial collateral and reducing the apoptosis of  
374 nerve cells. The cardioprotective effect of short-term skeletal muscle ischemia was  
375 evaluated in our previous experimental [24, 27], and Lin *et al.* used colored

376 microspheres to measure collateral circulation blood flow in rabbits (equation:  
377  $CCBF(\%) = \text{blood flow after occlusion} / \text{blood flow before occlusion}$ ), and concluded  
378 that physiologic ischemic training of skeletal muscle may induce collateral circulation  
379 development in the myocardium [34]. Most related studies [46] have suggested that  
380 isometric contraction training could produce cardioprotection after a certain period of  
381 training (such as four weeks). Both cerebral ischemia and cardiac ischemia are  
382 ischemic disease. However, only few studies have focused on the effect of ICT on  
383 cerebral ischemia. Does ICT have neuroprotective effects? To answer this question,  
384 we designed this experiment to investigate the neuroprotective effect of ICT on the rat  
385 MCAO model.

386       The rat MCAO model produces obvious infarction induced by focal occlusion of  
387 the MCA. The general ischemia model is a common clinical case, and stroke mostly  
388 occurs in local perfusion after spontaneous recanalization [47]. In this study, we  
389 performed MCAO to transiently block the distal MCA, after 90-min MCA occlusion,  
390 the transient block was removed to mimic partial reperfusion after stroke that  
391 frequently occurs in patients [48]. Our ischemic model was highly reproducible and  
392 reliable. In addition, the injured region of the cortex was most likely detected in  
393 clinical stroke patients.

394       Using this rat model, we designed three experimental groups. The purpose of the  
395 sham group was to exclude the placebo effect or any interference from the surgical  
396 procedure itself; the MCAO group was designed to establish a model of transient  
397 MCA occlusion; and the ICT group used the same ischemic model as physiologic  
398 ischemic training, and performed daily isometric contraction training.

399 Our experiment found that ICT could effectively reduce the infarct area, improve  
400 behavioral recovery, promote neurofunctional recovery, and protect neurons from  
401 focal ischemia in rats. In addition, in the ICT group, the loss of Nissl granules was  
402 attenuated, the neuronal morphology was improved, and neuronal apoptosis was  
403 reduced, indicating that ICT can regulate MCAO-induced neuronal injury.

404 Our previous studies proved that the expressions of VEGF and VEGF mRNA  
405 increased not only in the normal skeletal muscle but also increased in remote areas  
406 [21, 24]. Ren et al. [49] suggested that endogenous VEGF far away from the ischemic  
407 areas was also up-regulated. Our results showed that compared with the sham group,  
408 VEGF and VEGF mRNA increased significantly in both ischemic and contralateral  
409 sides in the ICT group. It was reported that VEGF not only promoted angiogenic and  
410 anti-inflammatory responses, but also regulated brain function at the neurovascular  
411 interface [50]. VEGF directly affects nerve cells and their progenitors, and indirectly  
412 affects the cerebral perfusion of the central and peripheral nervous systems [51],  
413 which is consistent with our study. This may also explain why ICT has  
414 neuroprotective effects.

415 We also found that ICT reduced the apoptosis of nerve cells. VEGF might be  
416 involved in the direct attenuation of cell death in the early stage of ischemic injury  
417 and the late stage of angiogenesis. Chen [52] found that TUNEL and cleaved  
418 caspase-3-positive neurons greatly reduced in AAVH9-VEGF-transduced mice,  
419 suggesting that the neuroprotective effect of VEGF was associated with the  
420 anti-apoptotic pathway. Interestingly, VEGF is also a strong survival factor in  
421 serum-deprived endothelial cells, and it also induces PI 3'-kinase activation and Akt

422 phosphorylation. Therefore, Ang1 and VEGF have a common intracellular second  
423 messenger signaling pathway, which can prevent the apoptosis of endothelial cells  
424 under serum deprivation. Recently, Akt has been shown to promote cell survival or  
425 nitric oxide production by phosphorylating Bad and procaspase-9 or endothelial nitric  
426 oxide synthase [53]. This may be a reason for the neuroprotective effect of ICT, which  
427 needs to be investigated in future studies.

428 In addition, there are some limitations in our study. Firstly, we only studied the  
429 training under a single condition, and whether using different doses and time can  
430 improve the outcomes remains to be tested. Secondly, we only studied the two-week  
431 time window, and whether longer ICT can promote the functional recovery of rats also  
432 needs to be explored further. Thirdly, this study did not employ unbiased stereology  
433 method to count the number of alive neurons or TUNEL positive cells. Fourthly,  
434 regarding the potential mechanism of the neuroprotective effect of ICT, only VEGF  
435 was studied, and whether other factors and pathways can also promote brain  
436 protection and functional recovery by regulating the expression of VEGF needs to be  
437 further investigated. Finally, the mechanism of ICT's protective effect on  
438 neurofunctional recovery still needs to be confirmed in more studies.

439

## 440 **Conclusions**

441 In summary, our results suggested that ICT might promote neurofunctional  
442 recovery and protect neurons against focal ischemia in rats, which was manifested by  
443 the decreased animal mortality, and the reduced cerebral infarct area, brain edema and  
444 functional deficits. This phenomenon might be one of the mechanisms of neurological

445 recovery in cerebral ischemic rats. ICT might provide a promising training approach  
446 for post-stroke rehabilitation.

447

#### 448 **Statements**

##### 449 **Statement of Ethics:**

450 All experimental procedures were approved by the Key Laboratory of Nerve  
451 Regeneration of Nantong University (Nantong, China) and were carried out in  
452 accordance with the institutional animal care guidelines of Nantong University. This  
453 study was approved by the Ethics Committee of Jiangsu Province, China, with  
454 approval No. S20141103-402.

##### 455 **Funding source**

456 Thanks for the support from the Jiangsu Key ' Laboratory of Neuroregeneration Open  
457 Project (grant no. 05012079).

##### 458 **Author Contributions:**

459 (1)Chengyao Mei, Teng Ma, conceiving and designing the study;

460 (2)Chengyao Mei, Teng Ma, collecting the data;

461 (3)Chengyao Mei, Teng Ma, analyzing and interpreting the data;

462 (4)Chengyao Mei, Teng Ma, writing the manuscript;

463 (5)Chengyao Mei, Teng Ma, providing critical revisions that are important for the  
464 intellectual content;

465 (6)Chengyao Mei, Teng Ma, approving the final version of the manuscript.

##### 466 **Data Availability Statement:**

467 All data generated or analysed during this study are included in this article. Further  
468 enquiries can be directed to the corresponding author.

##### 469 **Acknowledgements**

470 We would like to express our gratitude to Jiangsu Key Laboratory of  
471 Neuroregeneration, Nantong University. We thank Professor Fei Ding for assistance  
472 with the experimental procedures.

473 **Conflict of Interest:** The authors declare that there is no conflict of interest

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

494 **References**

- 495 1. Orellana-Urzúa S, Rojas I, Líbano L, Rodrigo R. Pathophysiology of Ischemic  
496 Stroke: Role of Oxidative Stress. *Curr Pharm Des* 2020;26:4246-4260.  
497 10.2174/1381612826666200708133912.
- 498 2. Roger VL, Go AS, Lloyd-Jones DM, Adams RJ, Berry JD, Brown TM,  
499 Carnethon MR, Dai S, de Simone G, Ford ES, Fox CS, Fullerton HJ, Gillespie  
500 C, Greenlund KJ, Hailpern SM, Heit JA, Ho PM, Howard VJ, Kissela BM,  
501 Kittner SJ, Lackland DT, Lichtman JH, Lisabeth LD, Makuc DM, Marcus GM,  
502 Marelli A, Matchar DB, McDermott MM, Meigs JB, Moy CS, Mozaffarian D,  
503 Mussolino ME, Nichol G, Paynter NP, Rosamond WD, Sorlie PD, Stafford RS,  
504 Turan TN, Turner MB, Wong ND, Wylie-Rosett J. Heart disease and stroke  
505 statistics--2011 update: a report from the American Heart Association.  
506 *Circulation* 2011;123:e18-e209. 10.1161/CIR.0b013e3182009701.
- 507 3. Hankey GJ. Secondary stroke prevention. *Lancet Neurol* 2014;13:178-194.  
508 10.1016/s1474-4422(13)70255-2.
- 509 4. Kuriakose D, Xiao Z. Pathophysiology and Treatment of Stroke: Present  
510 Status and Future Perspectives. *Int J Mol Sci* 2020;21:10.3390/ijms21207609.
- 511 5. Choi WJ, Li Y, Wang RK. Monitoring Acute Stroke Progression:  
512 Multi-Parametric OCT Imaging of Cortical Perfusion, Flow, and Tissue  
513 Scattering in a Mouse Model of Permanent Focal Ischemia. *IEEE Trans Med*  
514 *Imaging* 2019;38:1427-1437. 10.1109/tmi.2019.2895779.
- 515 6. Yu Q, Li X, Li Y, Fu J, Xiao Z. Effects of combined electroacupuncture and  
516 exercise training on motor function and microtubule-associated protein 2  
517 expression in the middle and late stages of cerebral infarction in rats.  
518 *Acupunct Med* 2020;38:175-180. 10.1177/0964528419882937.
- 519 7. Saunders DH, Sanderson M, Hayes S, Johnson L, Kramer S, Carter DD, Jarvis  
520 H, Brazzelli M, Mead GE. Physical fitness training for stroke patients.  
521 *Cochrane Database Syst Rev* 2020;3:CD003316.  
522 10.1002/14651858.CD003316.pub7.
- 523 8. Prior PL, Suskin N. Exercise for stroke prevention. *Stroke Vasc Neurol*  
524 2018;3:59-68. 10.1136/svn-2018-000155.
- 525 9. Svoboda J, Litvinec A, Kala D, Pošusta A, Vávrová L, Jiruška P, Otáhal J.  
526 Strain differences in intraluminal thread model of middle cerebral artery  
527 occlusion in rats. *Physiol Res* 2019;68:37-48. 10.33549/physiolres.933958.
- 528 10. Wu J, Mamas M, Rashid M, Weston C, Hains J, Luescher T, de Belder MA,  
529 Deanfield JE, Gale CP. Patient response, treatments, and mortality for acute  
530 myocardial infarction during the COVID-19 pandemic. *Eur Heart J Qual Care*  
531 *Clin Outcomes* 2021;7:238-246. 10.1093/ehjqcco/qcaa062.
- 532 11. Wang WW, Chen DZ, Zhao M, Yang XF, Gong DR. Prior transient ischemic  
533 attacks may have a neuroprotective effect in patients with ischemic stroke.  
534 *Arch Med Sci* 2017;13:1057-1061. 10.5114/aoms.2016.63744.
- 535 12. Li S, Hafeez A, Noorulla F, Geng X, Shao G, Ren C, Lu G, Zhao H, Ding Y,  
536 Ji X. Preconditioning in neuroprotection: From hypoxia to ischemia. *Prog*

- 537 Neurobiol 2017;157:79-91. 10.1016/j.pneurobio.2017.01.001.
- 538 13. Kim DH, Lee JH, Yu SM, An CM. The Effects of Ankle Position on Torque  
539 and Muscle Activity of the Knee Extensor During Maximal Isometric  
540 Contraction. *J Sport Rehabil* 2020;29:37-42. 10.1123/jsr.2018-0145.
- 541 14. Kozłowski B, Pageaux B, Hubbard EF, St Peters B, Millar PJ, Power GA.  
542 Perception of effort during an isometric contraction is influenced by prior  
543 muscle lengthening or shortening. *Eur J Appl Physiol* 2021;121:2531-2542.  
544 10.1007/s00421-021-04728-y.
- 545 15. Lin S, Chen Y, Li Y, Li J, Lu X. Physical ischaemia induced by isometric  
546 exercise facilitated collateral development in the remote ischaemic  
547 myocardium of humans. *Clin Sci (Lond)* 2014;127:581-588.  
548 10.1042/cs20130618.
- 549 16. Zhang X, Zheng Y, Geng C, Guan J, Wang L, Zhang X, Cheng Y, Li J, Lu X.  
550 Isometric exercise promotes arteriogenesis in rats after myocardial infarction.  
551 *J Biomed Res* 2021;35:436-447. 10.7555/jbr.35.20210062.
- 552 17. Saunders E, Clark BC, Clark LA, Grooms DR. Development of a trunk motor  
553 paradigm for use in neuroimaging. *Transl Neurosci* 2020;11:193-200.  
554 10.1515/tnsci-2020-0116.
- 555 18. Lin S, Lu X, Chen S, Ye F, Zhang J, Ma Y, Li J. Human coronary collateral  
556 recruitment is facilitated by isometric exercise during acute coronary occlusion.  
557 *J Rehabil Med* 2012;44:691-695. 10.2340/16501977-0989.
- 558 19. Moreira JBN, Wohlwend M, Wisløff U. Exercise and cardiac health:  
559 physiological and molecular insights. *Nat Metab* 2020;2:829-839.  
560 10.1038/s42255-020-0262-1.
- 561 20. Stock JM, Chouramanis NV, Chirinos JA, Edwards DG. Dynamic and  
562 isometric handgrip exercise increases wave reflection in healthy young adults.  
563 *J Appl Physiol* (1985) 2020;129:709-717. 10.1152/jappphysiol.00281.2020.
- 564 21. Gao M, Lu X, Chen W, Xiao GH, Zhang Y, Yu R, Li J. Randomized clinical  
565 trial of physiological ischemic training for patients with coronary heart disease  
566 complicated with heart failure: Safety of training, VEGF of peripheral blood  
567 and quality of life. *Exp Ther Med* 2018;16:260-264. 10.3892/etm.2018.6175.
- 568 22. Ni J, Lu H, Lu X, Jiang M, Peng Q, Ren C, Xiang J, Mei C, Li J. The evolving  
569 concept of physiological ischemia training vs. ischemia preconditioning. *J*  
570 *Biomed Res* 2015;29:445-450. 10.7555/jbr.29.20140142.
- 571 23. Wan C, Li J, Yang C, Hu D, Bi S. Dynamics of endogenous endothelial  
572 progenitor cells homing modulated by physiological ischaemia training. *J*  
573 *Rehabil Med* 2015;47:87-93. 10.2340/16501977-1891.
- 574 24. Zheng Y, Xiao M, Li L, Li J, Reinhardt JD, Lu X. Remote physiological  
575 ischemic training promotes coronary angiogenesis via molecular and cellular  
576 mobilization after myocardial ischemia. *Cardiovasc Ther*  
577 2017;35:10.1111/1755-5922.12257.
- 578 25. Zheng Y, Lu X, Li J, Zhang Q, Reinhardt JD. Impact of remote physiological  
579 ischemic training on vascular endothelial growth factor, endothelial progenitor  
580 cells and coronary angiogenesis after myocardial ischemia. *Int J Cardiol*  
581 2014;177:894-901. 10.1016/j.ijcard.2014.10.034.
- 582 26. Hahn CD, Manlhiot C, Schmidt MR, Nielsen TT, Redington AN. Remote



- 583 ischemic per-conditioning: a novel therapy for acute stroke? *Stroke*  
584 2011;42:2960-2962. 10.1161/strokeaha.111.622340.
- 585 27. Zhen X, Zheng Y, Hong X, Chen Y, Gu P, Tang J, Cheng H, Yuan TF, Lu X.  
586 Physiological Ischemic Training Promotes Brain Collateral Formation and  
587 Improves Functions in Patients with Acute Cerebral Infarction. *Front Neurol*  
588 2016;7:235. 10.3389/fneur.2016.00235.
- 589 28. Ruan GX, Kazlauskas A. Axl is essential for VEGF-A-dependent activation of  
590 PI3K/Akt. *Embo j* 2012;31:1692-1703. 10.1038/emboj.2012.21.
- 591 29. Chen X, Zheng Q, Li W, Lu Y, Ni Y, Ma L, Fu Y. SOX5 induces lung  
592 adenocarcinoma angiogenesis by inducing the expression of VEGF through  
593 STAT3 signaling. *Onco Targets Ther* 2018;11:5733-5741.  
594 10.2147/ott.S176533.
- 595 30. Hou Y, Wang K, Wan W, Cheng Y, Pu X, Ye X. Resveratrol provides  
596 neuroprotection by regulating the JAK2/STAT3/PI3K/AKT/mTOR pathway  
597 after stroke in rats. *Genes Dis* 2018;5:245-255. 10.1016/j.gendis.2018.06.001.
- 598 31. Samakova A, Gazova A, Sabova N, Valaskova S, Jurikova M, Kyselovic J.  
599 The PI3k/Akt pathway is associated with angiogenesis, oxidative stress and  
600 survival of mesenchymal stem cells in pathophysiologic condition in ischemia.  
601 *Physiol Res* 2019;68:S131-s138. 10.33549/physiolres.934345.
- 602 32. Wang N, Yang L, Zhang H, Lu X, Wang J, Cao Y, Chen L, Wang X, Cong L,  
603 Li J, Wang N, Liu Z, Wang L. MicroRNA-9a-5p Alleviates Ischemia Injury  
604 After Focal Cerebral Ischemia of the Rat by Targeting ATG5-Mediated  
605 Autophagy. *Cell Physiol Biochem* 2018;45:78-87. 10.1159/000486224.
- 606 33. Lopez MS, Vemuganti R. Modeling Transient Focal Ischemic Stroke in  
607 Rodents by Intraluminal Filament Method of Middle Cerebral Artery  
608 Occlusion. *Methods Mol Biol* 2018;1717:101-113.  
609 10.1007/978-1-4939-7526-6\_9.
- 610 34. Lin A, Li J, Zhao Y, Xiao M, Xiao B, Lu X, Wan C. Effect of physiologic  
611 ischemic training on protection of myocardial infarction in rabbits. *Am J Phys*  
612 *Med Rehabil* 2011;90:97-105. 10.1097/PHM.0b013e3182017483.
- 613 35. Hong Y, Liu Q, Peng M, Bai M, Li J, Sun R, Guo H, Xu P, Xie Y, Li Y, Liu L,  
614 Du J, Liu X, Yang B, Xu G. High-frequency repetitive transcranial magnetic  
615 stimulation improves functional recovery by inhibiting neurotoxic polarization  
616 of astrocytes in ischemic rats. *J Neuroinflammation* 2020;17:150.  
617 10.1186/s12974-020-01747-y.
- 618 36. Kulkarni P, Bhosle MR, Lu SF, Simon NS, Iriah S, Brownstein MJ, Ferris CF.  
619 Evidence of early vasogenic edema following minor head impact that can be  
620 reduced with a vasopressin V1a receptor antagonist. *Brain Res Bull*  
621 2020;165:218-227. 10.1016/j.brainresbull.2020.10.001.
- 622 37. Borshchev YY, Minasian SM, Burovenko IY, Borshchev VY, Protsak ES,  
623 Semenova NY, Borshcheva OV, Galagudza MM. Effects of tetracycline on  
624 myocardial infarct size in obese rats with chemically-induced colitis. *PLoS*  
625 *One* 2019;14:e0225185. 10.1371/journal.pone.0225185.
- 626 38. Sharma R, Iovine C, Agarwal A, Henkel R. TUNEL assay-Standardized  
627 method for testing sperm DNA fragmentation. *Andrologia* 2021;53:e13738.  
628 10.1111/and.13738.

- 629 39. Xiao M, Lu X, Li J, Li L, Li Y. Physiologic ischaemic training induces  
630 endothelial progenitor cell mobilization and myocardial angiogenesis via  
631 endothelial nitric oxide synthase related pathway in rabbits. *J Cardiovasc Med*  
632 (Hagerstown) 2014;15:280-287. 10.2459/JCM.0b013e32836009fe.
- 633 40. Bøtker HE, Kharbanda R, Schmidt MR, Bøttcher M, Kaltoft AK, Terkelsen CJ,  
634 Munk K, Andersen NH, Hansen TM, Trautner S, Lassen JF, Christiansen EH,  
635 Krusell LR, Kristensen SD, Thuesen L, Nielsen SS, Rehling M, Sørensen HT,  
636 Redington AN, Nielsen TT. Remote ischaemic conditioning before hospital  
637 admission, as a complement to angioplasty, and effect on myocardial salvage  
638 in patients with acute myocardial infarction: a randomised trial. *Lancet*  
639 2010;375:727-734. 10.1016/s0140-6736(09)62001-8.
- 640 41. Cheung MM, Kharbanda RK, Konstantinov IE, Shimizu M, Frndova H, Li J,  
641 Holtby HM, Cox PN, Smallhorn JF, Van Arsdell GS, Redington AN.  
642 Randomized controlled trial of the effects of remote ischemic preconditioning  
643 on children undergoing cardiac surgery: first clinical application in humans. *J*  
644 *Am Coll Cardiol* 2006;47:2277-2282. 10.1016/j.jacc.2006.01.066.
- 645 42. Li L, Luo W, Huang L, Zhang W, Gao Y, Jiang H, Zhang C, Long L, Chen S.  
646 Remote preconditioning reduces myocardial injury in adult valve replacement:  
647 a randomized controlled trial. *J Surg Res* 2010;164:e21-26.  
648 10.1016/j.jss.2010.06.016.
- 649 43. Thielmann M, Kottenberg E, Boengler K, Raffelsieper C, Neuhaeuser M,  
650 Peters J, Jakob H, Heusch G. Remote ischemic preconditioning reduces  
651 myocardial injury after coronary artery bypass surgery with crystalloid  
652 cardioplegic arrest. *Basic Res Cardiol* 2010;105:657-664.  
653 10.1007/s00395-010-0104-5.
- 654 44. Varnavas VC, Kontaras K, Glava C, Maniotis CD, Koutouzis M, Baltogiannis  
655 GG, Papalois A, Kolettis TM, Kyriakides ZS. Chronic skeletal muscle  
656 ischemia preserves coronary flow in the ischemic rat heart. *Am J Physiol*  
657 *Heart Circ Physiol* 2011;301:H1229-1235. 10.1152/ajpheart.00232.2011.
- 658 45. Nyquist P, Georgakis MK. Remote ischemic preconditioning effects on brain  
659 vasculature. *Neurology* 2019;93:15-16. 10.1212/wnl.0000000000007724.
- 660 46. Shen M, Gao J, Li J, Su J. Effect of ischaemic exercise training of a normal  
661 limb on angiogenesis of a pathological ischaemic limb in rabbits. *Clin Sci*  
662 (Lond) 2009;117:201-208. 10.1042/cs20080212.
- 663 47. Wang Y, Liu Y, Sun K, Wei Y, Fu L, Hou Z, Yi X, Ma D, Wang W, Jin X.  
664 The differential neuroprotection of HSP70-hom gene single nucleotide  
665 polymorphisms: In vitro (neuronal hypoxic injury model) and in vivo (rat  
666 MCAO model) studies. *Gene* 2019;710:354-362. 10.1016/j.gene.2019.05.059.
- 667 48. Li J, Tao T, Xu J, Liu Z, Zou Z, Jin M. HIF-1 $\alpha$  attenuates neuronal apoptosis  
668 by upregulating EPO expression following cerebral ischemia-reperfusion  
669 injury in a rat MCAO model. *Int J Mol Med* 2020;45:1027-1036.  
670 10.3892/ijmm.2020.4480.
- 671 49. Ren L, Wei C, Li K, Lu Z. LncRNA MALAT1 up-regulates VEGF-A and  
672 ANGPT2 to promote angiogenesis in brain microvascular endothelial cells  
673 against oxygen-glucose deprivation via targetting miR-145. *Biosci Rep*  
674 2019;39:10.1042/bsr20180226.

- 675 50. Lange C, Storkebaum E, de Almodóvar CR, Dewerchin M, Carmeliet P.  
676 Vascular endothelial growth factor: a neurovascular target in neurological  
677 diseases. *Nat Rev Neurol* 2016;12:439-454. 10.1038/nrneurol.2016.88.
- 678 51. Silverman DA, Martinez VK, Dougherty PM, Myers JN, Calin GA, Amit M.  
679 Cancer-Associated Neurogenesis and Nerve-Cancer Cross-talk. *Cancer Res*  
680 2021;81:1431-1440. 10.1158/0008-5472.Can-20-2793.
- 681 52. Chen L, Lin G, Chen K, Wan F, Liang R, Sun Y, Chen X, Zhu X. VEGF  
682 knockdown enhances radiosensitivity of nasopharyngeal carcinoma by  
683 inhibiting autophagy through the activation of mTOR pathway. *Sci Rep*  
684 2020;10:16328. 10.1038/s41598-020-73310-x.
- 685 53. Shang J, Gao ZY, Zhang LY, Wang CY. Over-expression of JAZF1 promotes  
686 cardiac microvascular endothelial cell proliferation and angiogenesis via  
687 activation of the Akt signaling pathway in rats with myocardial  
688 ischemia-reperfusion. *Cell Cycle* 2019;18:1619-1634.  
689 10.1080/15384101.2019.1629774.
- 690




Figure 1 Effects of isometric contraction training on mortality, brain water content, and neurobehavioral deficits. (A) Histogram showed the comparison of mortality between groups. (B) Histogram showed the comparison of brain water content between groups. (C) Histogram showed the comparison of the modified neurological severity score (mNSS).




Figure 2 Effects of isometric contraction training on the infarct area. (A) triphenyl tetrazolium chloride (TTC) staining of brain slices. (B) Histogram showed the comparison of the infarct area between groups.





Figure 3 Nissl staining and cell counts in the cortex region. (A) Nissl staining of the cortex in different groups. (B) Histogram showed the comparison of Nissl stained cells between groups.




Figure 4 Effects of isometric contraction training on neuronal apoptosis. (A) Rat brain tissue section. (B) Representative fluorescence micrographs of TUNEL staining for the cerebral cortex of different groups. (C) Histogram showing percentage of apoptotic (TUNEL-positive) cells in the cerebral cortex in different groups.




Figure 5 Effects of Isometric contraction training on the expression of vascular endothelial growth factor (VEGF). (A) Representative Western blots from cortical tissue samples. (B) Histogram showed the VEGF protein expression in the ischemic cortex of different groups. (C) Real-time polymerase chain reaction (RT-PCR) analysis of the effect of ICT on the level of VEGF mRNA expression. Note: CC: cortex contra, CI: cortex ipsi