



# Admissible rules of Łukasiewicz logic

Emil Jeřábek\*

Institute of Mathematics of the Academy of Sciences  
Žitná 25, 115 67 Praha 1, Czech Republic, email: jerabek@math.cas.cz

March 16, 2009

## Abstract

We investigate admissible rules of Łukasiewicz multi-valued propositional logic. We show that admissibility of multiple-conclusion rules in Łukasiewicz logic, as well as validity of universal sentences in free  $MV$ -algebras, is decidable (in  $PSPACE$ ).

## 1 Introduction

Investigation of nonclassical logics usually revolves around provability of formulas. When we generalize the problem from formulas to inference rules, there arises an important distinction between *derivable* and *admissible* rules, introduced by Lorenzen [15]. A rule

$$\varphi_1, \dots, \varphi_n / \psi$$

is derivable if it belongs to the consequence relation of the logic (defined semantically, or by a proof system using a set of axioms and rules); and it is admissible if the set of theorems of the logic is closed under the rule. These two notions coincide for the standard consequence relation of classical logic, but nonclassical logics often admit rules which are not derivable. (A logic whose admissible rules are all derivable is called *structurally complete*.) For example, all superintuitionistic (si) logics admit the Kreisel–Putnam rule

$$\neg p \rightarrow q \vee r / (\neg p \rightarrow q) \vee (\neg p \rightarrow r),$$

whereas many of these logics (such as  $IPC$  itself) do not derive this rule.

The research of admissible rules was stimulated by a question of H. Friedman [5], asking whether admissibility of rules in  $IPC$  is decidable. The problem was extensively investigated in a series of papers by Rybakov, who has shown that admissibility is decidable for a large class of modal and si logics, found semantic criteria for admissibility, and obtained other results on various aspects of admissibility. His results on admissible rules in transitive modal and si logics are summarized in the monograph [21]. He also applied his method to tense logics [22, 23, 24].

---

\*Supported by grants IAA900090703 and IAA100190902 of GA AV ČR, grant 1M0545 of MŠMT ČR, and a grant from the John Templeton Foundation.

Ghilardi [7, 8] discovered the connection of admissibility to projective formulas and unification, which provided another criteria for admissibility in certain modal and si logics, and new decision procedures for admissibility in some modal and si systems. Ghilardi's results were utilized by Iemhoff [10, 11, 12] to construct an explicit basis of admissible rules for *IPC* and some other si logics, and to develop Kripke semantics for admissible rules. These results were extended to modal logics by Jeřábek [13]. We note that decidability of admissibility is by no means automatic. An artificial decidable modal logic with undecidable admissibility problem was constructed by Chagrova [1], and natural examples of bimodal logics with undecidable admissibility (or even unification) problem were found by Wolter and Zakharyashev [26]. In terms of computational complexity, admissibility in basic transitive logics is *coNE*-complete by Jeřábek [14], whereas derivability in these logics is *PSPACE*-complete.

In contrast to the situation in modal and superintuitionistic logics, only very little is known about admissibility in other nonclassical logics. Here we are particularly interested in substructural and fuzzy logics. Structural completeness of various substructural logics was investigated by Olson et al. [19] and by Cintula and Metcalfe [3]. Dzik [4] studied unification in  $n$ -contractive extensions of Hájek's Basic Logic (**BL**).

In this paper we study admissible rules of Łukasiewicz logic (**L**). We chose this logic because it is one of the three fundamental  $t$ -norm fuzzy logics, and among the three it is the only one with a nontrivial admissibility problem: Gödel–Dummett logic is a si logic and it is well-known to be structurally complete, and product logic was shown to be structurally complete too by Cintula and Metcalfe [3]. For more generality, we work with multiple-conclusion rules (cf. Shoesmith and Smiley [25]). We describe a criterion for admissibility of multiple-conclusion rules in **L**, and we show that admissibility in **L** is decidable. We also compute explicit bounds on the size of counterexamples to inadmissible rules, and use them to provide a *PSPACE*-algorithm for admissibility in **L**. Our results can be restated algebraically, namely we obtain that the universal theory of free *MV*-algebras is decidable (in *PSPACE*). We also show that **L** is 1-reducible wrt admissible rules (i.e., inadmissibility of any rule can be witnessed by a substitution using only one variable), or in algebraic terms, all free *MV*-algebras over nonempty sets of generators have the same universal theory.

For completeness, we also briefly consider the case of finite-valued Łukasiewicz logics  $\mathbf{L}_n$ . Being tabular extensions of **BL**, these logics are  $n$ -contractive, hence we easily derive from Dzik's results [4] that admissibility in  $\mathbf{L}_n$  is decidable. We provide an explicit basis of admissible rules for  $\mathbf{L}_n$  (and more generally, for any  $n$ -contractive extension of **BL**).

## 2 Preliminaries

The language of Hájek's Basic Logic (**BL**) [9] consists of *propositional formulas* built from variables  $p_n$ ,  $n \in \omega$ , using connectives  $\rightarrow$ ,  $\cdot$ , and  $\perp$ . A *substitution* is a mapping of propositional formulas to propositional formulas which commutes with all connectives. A formula  $\varphi$  is *derivable* from a set of formulas  $\Gamma$ , written as  $\Gamma \vdash_{\mathbf{BL}} \varphi$ , if there exists a finite sequence of formulas  $\varphi_1, \dots, \varphi_n$  such that  $\varphi_n = \varphi$ , and each  $\varphi_i$  is a member of  $\Gamma$ , an instance of one of

the axioms (cf. Cintula [2])

$$\begin{aligned}
&(\varphi \rightarrow \psi) \rightarrow ((\psi \rightarrow \chi) \rightarrow (\varphi \rightarrow \chi)), \\
&\varphi \rightarrow (\psi \rightarrow \varphi), \\
&(\varphi \rightarrow (\psi \rightarrow \chi)) \rightarrow (\varphi \cdot \psi \rightarrow \chi), \\
&(\varphi \cdot \psi \rightarrow \chi) \rightarrow (\varphi \rightarrow (\psi \rightarrow \chi)), \\
&\varphi \cdot (\varphi \rightarrow \psi) \rightarrow \psi \cdot (\psi \rightarrow \varphi), \\
&((\varphi \rightarrow \psi) \rightarrow \chi) \rightarrow (((\psi \rightarrow \varphi) \rightarrow \chi) \rightarrow \chi), \\
&\perp \rightarrow \varphi,
\end{aligned}$$

or it is derived from some  $\varphi_j, \varphi_k, j, k < i$  by an instance of the rule of modus ponens

$$(MP) \quad \varphi, \varphi \rightarrow \psi / \psi.$$

We can introduce other connectives as abbreviations

$$\begin{aligned}
\neg\varphi &\equiv \varphi \rightarrow \perp, \\
\varphi \wedge \psi &\equiv \varphi \cdot (\varphi \rightarrow \psi), \\
\varphi \vee \psi &\equiv ((\varphi \rightarrow \psi) \rightarrow \psi) \wedge ((\psi \rightarrow \varphi) \rightarrow \varphi), \\
\varphi \leftrightarrow \psi &\equiv (\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi), \\
\top &\equiv \neg\perp,
\end{aligned}$$

and we write  $\varphi^n = \varphi \cdot \dots \cdot \varphi$  with  $n$  occurrences of  $\varphi$  (if  $n = 0$ , we put  $\varphi^0 = \top$ ). An *extension* of **BL** is a consequence relation  $\vdash_L$  defined as  $\vdash_{\mathbf{BL}}$  above, except that we allow an extra set of additional axioms, which is required to be closed under substitution. An extension of **BL** is *n-contractive* if  $\vdash_L \varphi^n \rightarrow \varphi^{n+1}$ .

*Lukasiewicz logic* (**L**) is the extension of **BL** by the axiom schema

$$\neg\neg\varphi \rightarrow \varphi.$$

In **L** we can introduce another connective

$$\varphi \oplus \psi \equiv \neg\varphi \rightarrow \psi \equiv \neg(\neg\varphi \cdot \neg\psi).$$

Connectives in **L** are interdefinable, we have

$$\begin{aligned}
\varphi \rightarrow \psi &\equiv \neg\varphi \oplus \psi \equiv \neg(\varphi \cdot \neg\psi), \\
\varphi \cdot \psi &\equiv \neg(\neg\varphi \oplus \neg\psi) \equiv \neg(\varphi \rightarrow \neg\psi),
\end{aligned}$$

hence we can take any of the sets  $\{\rightarrow, \neg\}$ ,  $\{\rightarrow, \perp\}$ ,  $\{\cdot, \neg\}$ ,  $\{\oplus, \neg\}$  as the set of basic connectives, and define the rest as abbreviations.

If  $L$  is a logic, an *L-unifier* of a formula  $\varphi$  is a substitution  $\sigma$  such that  $\vdash_L \sigma\varphi$ . A formula which has an *L-unifier* is called *L-unifiable*. A unifier is *ground* if its range consists of constant (i.e., variable-free) formulas. An *L-unifier*  $\sigma$  of  $\varphi$  is *projective* (Ghilardi [7], cf. [6]), if

$$\varphi \vdash_L \psi \leftrightarrow \sigma\psi$$

for every formula  $\psi$ . A formula is *L-projective*, if it has a projective *L*-unifier.

A *single-conclusion rule* is an expression of the form

$$\frac{\Gamma}{\varphi},$$

also written as  $\Gamma / \varphi$ , where  $\varphi$  is a formula, and  $\Gamma$  is a finite set of formulas. A single-conclusion rule  $\Gamma / \varphi$  is *derivable* in a logic *L*, if  $\Gamma \vdash_L \varphi$ . The rule  $\Gamma / \varphi$  is *L-admissible*, written as  $\Gamma \vdash_L \varphi$ , if every common *L*-unifier of  $\Gamma$  is also an *L*-unifier of  $\varphi$ . A set *B* of *L*-admissible rules is a *basis* of *L*-admissible rules, if for every *L*-admissible rule  $\Gamma / \varphi$ , the formula  $\varphi$  is derivable from  $\Gamma$  using axioms and rules of *L*, and substitution instances of rules from *B*. We will often omit the prefix *L*- when it is clear from the context.

More generally, a *multiple-conclusion rule* is an expression of the form

$$\frac{\Gamma}{\Delta},$$

also written as  $\Gamma / \Delta$ , where  $\Gamma$  and  $\Delta$  are finite sets of formulas. (Note that  $\Gamma$  or  $\Delta$  can be both empty.) We will often write just *rule* instead of “multiple-conclusion rule”. A rule  $\Gamma / \Delta$  is derivable in *L* if  $\Gamma \vdash_L \varphi$  for some  $\varphi \in \Delta$ , and it is *L*-admissible if every common *L*-unifier of  $\Gamma$  is also an *L*-unifier of some formula  $\varphi \in \Delta$ . A set *B* of *L*-admissible rules is a *basis* of *L*-admissible rules, if every *L*-admissible rule can be inferred from *L*-derivable rules and instances of rules from *B* using weakening (from  $\Gamma / \Delta$  infer  $\Gamma, \Gamma' / \Delta, \Delta'$ ) and cut (from  $\Gamma / \Delta, \varphi$  and  $\Gamma, \varphi / \Delta$  infer  $\Gamma / \Delta$ ).  $\Gamma / \Delta$  is a *passive admissible rule* if  $\Gamma$  has no common unifier, i.e., if  $\Gamma \not\vdash_L$ .

An *MV-algebra* is a structure  $\langle A, \oplus, \neg, 0 \rangle$  which satisfies the identities

$$\begin{aligned} (x \oplus y) \oplus z &= x \oplus (y \oplus z), \\ x \oplus 0 &= x, \\ x \oplus y &= y \oplus x, \\ \neg \neg x &= x, \\ x \oplus \neg 0 &= \neg 0, \\ \neg(\neg x \oplus y) \oplus y &= \neg(\neg y \oplus x) \oplus x. \end{aligned}$$

We can define other operations on an *MV*-algebra by

$$\begin{aligned} x \rightarrow y &= \neg x \oplus y, \\ x \cdot y &= \neg(\neg x \oplus \neg y), \\ x \wedge y &= x \cdot (x \rightarrow y), \\ x \vee y &= (x \rightarrow y) \rightarrow y, \\ x \leftrightarrow y &= (x \rightarrow y) \wedge (y \rightarrow x), \\ 1 &= \neg 0. \end{aligned}$$

The operations  $\wedge, \vee$  turn *A* into a distributive lattice with bounds 0, 1, which induces a partial order  $\leq$  on *A*. We can identify propositional formulas with terms in the language

of  $MV$ -algebras in a natural way. A *valuation* in an  $MV$ -algebra  $A$  is a homomorphism  $v$  from the term algebra to  $A$ . If  $\varphi$  is a formula in the first  $k$  variables,  $a \in A^k$ , and  $v$  is the assignment such that  $v(p_i) = a_i$ , we also write  $\varphi(a) = v(\varphi)$ . A valuation  $v$  satisfies a formula  $\varphi$  if  $v(\varphi) = 1$ , and it satisfies a rule  $\Gamma / \Delta$  if  $v(\varphi) \neq 1$  for some  $\varphi \in \Gamma$ , or  $v(\varphi) = 1$  for some  $\varphi \in \Delta$ . A rule  $\Gamma / \Delta$  is *valid* in an  $MV$ -algebra  $A$ , written as  $A \models \Gamma / \Delta$ , if the rule is satisfied by every valuation in  $A$ . In other words,  $A \models \Gamma / \Delta$  if and only if the open first-order formula

$$\bigwedge_{\varphi \in \Gamma} (\varphi = 1) \rightarrow \bigvee_{\varphi \in \Delta} (\varphi = 1)$$

is valid in  $A$ . Conversely, validity of open formulas (or equivalently, universal sentences) in  $A$  can be reduced to validity of rules. Any open formula  $\Phi$  can be expressed in the conjunctive normal form as  $\Phi = \bigwedge_{i < k} \Phi_i$ , where each  $\Phi_i$  is a clause: a disjunction of atomic formulas (i.e., equations) and their negations. Then  $A \models \Phi$  iff  $A \models \Phi_i$  for each  $i < k$ , and a clause

$$\bigvee_{i < n} (\varphi_i = \psi_i) \vee \bigvee_{i < m} (\varphi'_i \neq \psi'_i)$$

is valid in  $A$  iff  $A$  validates the rule

$$\{\varphi'_i \leftrightarrow \psi'_i \mid i < m\} / \{\varphi_i \leftrightarrow \psi_i \mid i < n\}.$$

Lukasiewicz logic is algebraizable, and the variety of  $MV$ -algebras is its equivalent algebraic semantics, using the translation between propositional formulas and equations described above. We thus have (cf. [9]):

**Fact 2.1** *A rule  $\Gamma / \Delta$  is valid in all  $MV$ -algebras if and only if it is derivable in  $\mathbf{L}$ .* □

A *free  $MV$ -algebra* over a set  $X$  of generators is an  $MV$ -algebra  $F \supseteq X$  such that every mapping from  $X$  to an  $MV$ -algebra  $A$  can be uniquely extended to a homomorphism from  $F$  to  $A$ . As another corollary to algebraizability of  $\mathbf{L}$ , free  $MV$ -algebras can be described as Lindenbaum–Tarski algebras of  $\mathbf{L}$ :  $F$  consists of equivalence classes of formulas using elements of  $X$  as propositional variables modulo the equivalence relation  $\varphi \sim \psi$  iff  $\vdash_{\mathbf{L}} \varphi \leftrightarrow \psi$ , with operations defined in the natural way. Note that valuations in  $F$  correspond to substitutions whose range consists of formulas using variables from  $X$ , and a formula  $\varphi$  is satisfied under a valuation given by such a substitution  $\sigma$  if and only if  $\vdash_{\mathbf{L}} \sigma\varphi$ . We obtain the following characterization of admissibility:

**Fact 2.2** *For any rule  $\Gamma / \Delta$ , the following are equivalent.*

- (i)  $\Gamma \vdash_{\mathbf{L}} \Delta$ .
- (ii)  $\Gamma / \Delta$  is valid in all free  $MV$ -algebras.
- (iii)  $\Gamma / \Delta$  is valid in all free  $MV$ -algebras over finite sets of generators.
- (iv)  $\Gamma / \Delta$  is valid in some free  $MV$ -algebra over an infinite set of generators.

□

The *standard MV-algebra*  $[0, 1]_{\mathbf{L}}$  is the algebra  $\langle [0, 1]_{\mathbb{R}}, \oplus, \neg, 0 \rangle$ , where

$$\begin{aligned}x \oplus y &= \min\{x + y, 1\}, \\ \neg x &= 1 - x.\end{aligned}$$

Notice that the rational interval  $[0, 1]_{\mathbb{Q}}$  is a subalgebra of  $[0, 1]_{\mathbf{L}}$ . Both  $[0, 1]_{\mathbf{L}}$  and  $[0, 1]_{\mathbb{Q}}$  generate the variety of *MV-algebras*, hence we have the following strengthening of Fact 2.1: a formula is derivable in  $\mathbf{L}$  iff it is valid in  $[0, 1]_{\mathbf{L}}$  iff it is valid in  $[0, 1]_{\mathbb{Q}}$ . (In fact,  $[0, 1]_{\mathbf{L}}$  and  $[0, 1]_{\mathbb{Q}}$  generate all *MV-algebras* as a quasivariety, hence the same characterization also holds for derivability of single-conclusion rules.)

For any integer  $n > 0$ , the set  $\{0, 1/n, \dots, (n-1)/n, 1\}$  is also a subalgebra of  $[0, 1]_{\mathbf{L}}$ . The extension of  $\mathbf{BL}$  given by all formulas valid in this subalgebra is the *finite-valued Łukasiewicz logic*  $\mathbf{L}_{n+1}$ .

A description of free *MV-algebras* over finite sets of generators was given by McNaughton. Let  $n \in \omega$ . A function  $f: [0, 1]_{\mathbb{Q}}^n \rightarrow [0, 1]_{\mathbb{Q}}$  is called *piecewise linear with integer coefficients*, if there are finitely many functions  $L_j: [0, 1]_{\mathbb{Q}}^n \rightarrow [0, 1]_{\mathbb{Q}}$  such that for every  $x \in [0, 1]_{\mathbb{Q}}^n$  there exists  $j$  such that  $f(x) = L_j(x)$ , and each  $L_j(x_0, \dots, x_{n-1})$  is of the form  $\sum_{i < n} a_i x_i + b$  for some  $\vec{a}, b \in \mathbb{Z}$ . Let  $F_n$  be the *MV-algebra* of continuous piecewise linear functions  $f: [0, 1]_{\mathbb{Q}}^n \rightarrow [0, 1]_{\mathbb{Q}}$  with integer coefficients, with operations defined pointwise (i.e.,  $F_n$  is a subalgebra of the Cartesian power  $[0, 1]_{\mathbf{L}}^{[0, 1]_{\mathbb{Q}}^n}$ ).

**Theorem 2.3 (McNaughton [17])**  *$F_n$  is the free  $n$ -generated *MV-algebra*. The projection functions  $\pi_i(x_0, \dots, x_{n-1}) = x_i$  for  $i < n$  are its free generators.*  $\square$

Note that  $F_n$  are usually defined to consist of *real* functions  $f: [0, 1]_{\mathbb{R}}^n \rightarrow [0, 1]_{\mathbb{R}}$  which are continuous and piecewise linear with integer coefficients. It is easy to see that both definitions lead to isomorphic algebras, and it will be more convenient for us to work with the rational version.

In general, we will denote by  $F_{\varkappa}$  the free *MV-algebra* over  $\varkappa$  generators for every cardinal number  $\varkappa$ .

If  $f = \langle f_0, \dots, f_{k-1} \rangle$  is a  $k$ -tuple of functions  $f_i \in F_n$ , we will identify  $f$  with the corresponding function  $f: [0, 1]_{\mathbb{Q}}^n \rightarrow [0, 1]_{\mathbb{Q}}^k$ .

Since  $F_n$  is isomorphic to a Lindenbaum–Tarski algebra of  $\mathbf{L}$ , elements  $f \in F_n$  represent formulas in  $n$  variables (up to  $\mathbf{L}$ -provable equivalence). As we identify propositional formulas with terms, we may evaluate them in every *MV-algebra*. We can therefore define  $f(a) \in A$  for every  $f \in F_n$  and  $a \in A^n$ , where  $A$  is an *MV-algebra*. (In algebraic terms,  $f(a) = \bar{a}(f)$ , where  $\bar{a}: F_n \rightarrow A$  is the unique homomorphism such that  $\bar{a}(\pi_i) = a_i$  for each  $i < n$ .) Notice that this notation agrees with the literal usage of  $f$  as a function  $f: [0, 1]_{\mathbb{Q}}^n \rightarrow [0, 1]_{\mathbb{Q}}$  in the case  $A = [0, 1]_{\mathbb{Q}}$ .

We assume the reader is familiar with basic linear algebra. We identify vectors  $v \in \mathbb{Q}^n$  with  $n$ -by-1 matrices, i.e., we view them as column vectors. In particular, if  $v, w \in \mathbb{Q}^m$ , then  $v^{\top}w$  coincides with the inner product of  $v$  and  $w$ . We number coordinates of vectors, as well as rows and columns of matrices, starting from 0. If  $v \in \mathbb{Q}^m$  and  $i < m$ , we denote the  $i$ th coordinate of  $v$  by  $v_i$  (though we will also use subscript indices for many other purposes).

### 3 The finite case

We first consider the easy case of finite-valued Łukasiewicz logics. The results in fact apply more generally to all  $n$ -contractive extensions of **BL**. We rely on the following key result.

**Theorem 3.1 (Dzik [4])** *Let  $L$  be an  $n$ -contractive extension of **BL** for some  $n \in \omega$ . Then every  $L$ -unifiable formula is  $L$ -projective.*  $\square$

**Corollary 3.2** *Let  $L$  be an  $n$ -contractive extension of **BL** for some  $n \in \omega$ . A rule  $\Gamma / \Delta$  is  $L$ -admissible if and only if  $\bigwedge \Gamma$  is not  $L$ -unifiable or there exists  $\delta \in \Delta$  such that  $\Gamma \vdash_L \delta$ .*

*Proof:* Assume that  $\Gamma \sim_L \Delta$ , and let  $\sigma$  be a projective unifier of  $\bigwedge \Gamma$ . We have  $\vdash_L \sigma\Gamma$ , hence  $\vdash_L \sigma\delta$  for some  $\delta \in \Delta$ , which implies  $\Gamma \vdash_L \delta$ .  $\square$

**Lemma 3.3** *Let  $L$  be a consistent extension of **BL**. A formula  $\varphi$  is  $L$ -unifiable if and only if it is satisfiable in classical logic.*

*Proof:* Right-to-left: let  $e$  be a classical assignment such that  $e(\varphi) = 1$ , and let  $\sigma$  be the ground substitution such that  $\sigma p_i = e(p_i)$ . As **BL** can evaluate constant formulas, we have  $\vdash_L \sigma\varphi$ .

Left-to-right: let  $\sigma$  be a substitution such that  $\vdash_L \sigma\varphi$ . We may assume that  $\sigma$  is ground. As **BL** can evaluate constant formulas, we have  $\vdash_L e(\varphi)$ , where  $e$  is the classical assignment such that  $e(p_i)$  is the value of the sentence  $\sigma p_i$ . As  $L$  is consistent, we must have  $e(\varphi) = 1$ .  $\square$

Notice that the last lemma already holds for extensions of **FL<sub>w</sub>** in place of **BL**. We do not have any use for this observation.

**Corollary 3.4** *Let  $L$  be a decidable  $n$ -contractive extension of **BL**. Then admissibility in  $L$  is decidable.*  $\square$

Our main contribution in this section is a description of an explicit basis of admissible rules for  $n$ -contractive extensions of **BL**. By Corollary 3.2, this amounts to axiomatization of passive admissible rules.

**Definition 3.5** We introduce the rules

$$(CC_n) \quad \frac{\neg(p \vee \neg p)^n}{\quad}$$

and their variants

$$(CC_n^1) \quad \frac{\neg(p \vee \neg p)^n}{\perp}$$

for  $n \in \omega$ . Let  $CC = \{CC_n \mid n \in \omega\}$ ,  $CC^1 = \{CC_n^1 \mid n \in \omega\}$ .

The theorem below actually holds for all extensions of **MTL**.

**Theorem 3.6** *If  $L$  is an extension of  $\mathbf{BL}$ , then  $CC^1$  is a basis of single-conclusion passive  $L$ -admissible rules. If  $L$  is consistent, then  $CC$  is a basis of multiple-conclusion passive  $L$ -admissible rules.*

*Proof:* Clearly,  $CC_n$  are passive admissible rules by Lemma 3.3. On the other hand, assume that the rule  $\Gamma / \Delta$  is passive, i.e., the formula  $\bigwedge \Gamma$  is not unifiable. Then  $\neg \bigwedge \Gamma$  is a classical tautology. As  $\mathbf{CPC} = \mathbf{BL} + p \vee \neg p$ , there are formulas  $\psi_i$  such that

$$\vdash_L \prod_{i < n} (\psi_i \vee \neg \psi_i) \rightarrow \neg \bigwedge \Gamma$$

by the deduction theorem. Put  $\psi = \bigwedge_{i < n} (\psi_i \vee \neg \psi_i)$ . Since  $\mathbf{BL}$  proves De Morgan laws and  $\neg(p \vee \neg p) \rightarrow q \vee \neg q$ , we have  $\vdash_L \neg \psi \rightarrow \psi$ , hence  $\vdash_L \psi \vee \neg \psi \rightarrow \psi_i \vee \neg \psi_i$  for every  $i$ . Thus

$$\vdash_L (\psi \vee \neg \psi)^n \rightarrow \neg \bigwedge \Gamma$$

and

$$\Gamma \vdash_L \neg(\psi \vee \neg \psi)^n,$$

hence  $\Gamma / \Delta$  follows from an instance of  $CC_n$ . If  $\Delta \neq \emptyset$ , we can use  $CC_n^1$  instead of  $CC_n$ .  $\square$

Note that in  $n$ -contractive logics,  $CC$  is equivalent to  $CC_n$ , and  $CC^1$  is equivalent to  $CC_n^1$ .

**Corollary 3.7** *If  $L$  is an  $n$ -contractive extension of  $\mathbf{BL}$  for some  $n \in \omega$ , then  $CC_n^1$  is a basis of single-conclusion  $L$ -admissible rules. If  $L$  is consistent, then  $CC_n$  is a basis of multiple-conclusion  $L$ -admissible rules.  $\square$*

## 4 The infinite case

In this section we are going to prove our main result: a characterization of admissible rules of the infinite-valued Łukasiewicz logic which establishes their decidability.

Recall that  $\Gamma \sim_{\mathbf{L}} \Delta$  if and only if  $F_n \models \Gamma / \Delta$  for all  $n \in \omega$ . We first show that it suffices to consider only the case  $n = 1$ .

**Theorem 4.1**  *$\mathbf{L}$  is 1-reducible wrt admissible rules, i.e., for every inadmissible rule  $\Gamma / \Delta$ , there exists a substitution  $\sigma$  in only one variable such that  $\vdash_{\mathbf{L}} \sigma \Gamma$  and  $\not\vdash_{\mathbf{L}} \sigma \delta$  for each  $\delta \in \Delta$ . In algebraic terms,  $\Gamma \sim_{\mathbf{L}} \Delta$  iff  $F_1 \models \Gamma / \Delta$ .*

*Proof:* If  $\Gamma \not\sim_{\mathbf{L}} \Delta$ , there exists an  $n \in \omega$  such that  $F_n \not\models \Gamma / \Delta$ . Let  $e$  be a valuation in  $F_n$  such that  $\Gamma(e) = 1$  and  $\delta(e) \neq 1$  for all  $\delta \in \Delta$ . We can represent  $e$  by a piecewise linear function  $e: [0, 1]_{\mathbb{Q}}^n \rightarrow [0, 1]_{\mathbb{Q}}^k$ , where  $k$  is such that  $\Gamma \cup \Delta$  uses only variables  $p_0, \dots, p_{k-1}$ . We enumerate  $\Delta = \{\delta_i \mid i < r\}$ . We may assume  $r > 0$  without loss of generality.

Consider an  $i < r$ . There exists an  $x^i \in [0, 1]_{\mathbb{Q}}^n$  such that  $\delta_i(e(x^i)) < 1$ . We can write  $x^i = \langle p_0/q, \dots, p_{n-1}/q \rangle$  for some natural numbers  $\vec{p}, q$  such that  $q \geq 2$ , and we define a function  $f^i: [0, 1]_{\mathbb{Q}} \rightarrow [0, 1]_{\mathbb{Q}}^n$  by

$$f_j^i(t) = \begin{cases} \min\{p_j t, 1\} & t \leq 1/2, \\ \min\{p_j(1-t), 1\} & t \geq 1/2. \end{cases}$$



Then  $f^i \in F_1^n$ ,  $f^i(0) = f^i(1) = \vec{0}$ , and  $f^i(1/q) = x^i$ .

Let us define  $f: [0, 1]_{\mathbb{Q}} \rightarrow [0, 1]_{\mathbb{Q}}^n$  by

$$f(t) = f^i(rt - i), \quad i/r \leq t \leq (i+1)/r.$$

Notice that  $f(i/r) = \vec{0}$  is well-defined. By the construction,  $f \in F_1^n$ , and for each  $\delta \in \Delta$  there exists  $t \in [0, 1]_{\mathbb{Q}}$  such that  $\delta(e(f(t))) < 1$ . Trivially  $\gamma(e(f(t))) = 1$  for each  $t$  and  $\gamma \in \Gamma$ , hence the valuation  $e \circ f: [0, 1]_{\mathbb{Q}} \rightarrow [0, 1]_{\mathbb{Q}}^k$  witnesses that  $F_1 \not\equiv \Gamma / \Delta$ .  $\square$

**Corollary 4.2** *All free MV-algebras  $F_{\varkappa}$ ,  $\varkappa \neq 0$ , have the same universal theory.*  $\square$

We can equivalently restate the last corollary as follows: every finite partial subalgebra of  $F_{\varkappa}$  can be embedded in any  $F_{\lambda}$  ( $\lambda > 0$ ). Nevertheless, it is not possible to embed all of  $F_{\varkappa}$  in  $F_{\lambda}$  at once unless  $\varkappa \leq \lambda$ :

**Theorem 4.3** *If  $\varkappa > \lambda$  are cardinals, then  $F_{\varkappa}$  is not embeddable in  $F_{\lambda}$ .*

*Proof:* It is easy to see that it suffices to show the result for finite  $\varkappa$  and  $\lambda$ . Assume for contradiction that  $\varphi: F_{\varkappa} \rightarrow F_{\lambda}$  is an embedding, and consider the continuous piecewise linear function  $f: [0, 1]_{\mathbb{Q}}^{\lambda} \rightarrow [0, 1]_{\mathbb{Q}}^{\varkappa}$  such that  $f_i = \varphi(x_i)$ , where  $\{x_i \mid i < \varkappa\}$  are the free generators of  $F_{\varkappa}$ . We can extend  $f$  to a continuous piecewise linear function  $\hat{f}: [0, 1]_{\mathbb{R}}^{\lambda} \rightarrow [0, 1]_{\mathbb{R}}^{\varkappa}$ . As piecewise linear functions do not increase topological dimension, we have  $\dim(\text{rng}(\hat{f})) \leq \dim([0, 1]_{\mathbb{R}}^{\lambda}) = \lambda < \varkappa = \dim([0, 1]_{\mathbb{R}}^{\varkappa})$ , hence  $\hat{f}$  is not onto. Being a continuous image of a compact space,  $\text{rng}(\hat{f})$  is closed, hence there exists a point  $v \in [0, 1]_{\mathbb{Q}}^{\varkappa}$  and an  $\varepsilon > 0$  such that  $\text{rng}(\hat{f}) \cap \prod_{i < \varkappa} [v_i - \varepsilon, v_i + \varepsilon] = \emptyset$ . For each  $i < \varkappa$ , we can write  $v_i = p_i/q_i$  for some integers  $p_i, q_i$  such that  $q_i > 1/\varepsilon$ , and define  $g^i: [0, 1]_{\mathbb{Q}} \rightarrow [0, 1]_{\mathbb{Q}}$  by

$$g^i(t) = \min\{1, |p_i - q_i t|\}.$$

We have  $g^i(v_i) = 0$ , and  $g^i(u) = 1$  for all  $u \notin [v_i - \varepsilon, v_i + \varepsilon]$ . We define  $g: [0, 1]_{\mathbb{Q}}^{\varkappa} \rightarrow [0, 1]_{\mathbb{Q}}$  by

$$g(t_0, \dots, t_{\varkappa-1}) = \max\{g^i(t_i) \mid i < \varkappa\}.$$

By the construction,  $g \in F_{\varkappa}$ ,  $g(v) = 0$ , and  $g(x) = 1$  for all  $x \in \text{rng}(f)$ . Thus  $\varphi(g) = g \circ f = 1 = \varphi(1)$ , but  $g \neq 1$ , which contradicts  $\varphi$  being an embedding.  $\square$

Since we will work a lot with  $F_1$ , we introduce convenient notation for elements of  $F_1^m$ , as well as other continuous piecewise linear functions in one variable.

**Definition 4.4** If  $t_0 < t_1 < \dots < t_k$  are rational numbers and  $x_0, \dots, x_k \in \mathbb{Q}^m$ , then  $L(t_0, x_0; t_1, x_1; \dots; t_k, x_k)$  is the continuous piecewise linear function  $f: [t_0, t_k]_{\mathbb{Q}} \rightarrow \mathbb{Q}^m$  such that  $f(t_i) = x_i$ , and  $f$  is linear on each interval  $[t_i, t_{i+1}]$ .

Let  $X \subseteq \mathbb{Q}^m$ . The *convex hull*  $C(X)$  is the smallest convex subset of  $\mathbb{Q}^m$  which includes  $X$ , and the *affine hull*  $A(X)$  is the smallest affine subspace of  $\mathbb{Q}^m$  which includes  $X$ . Notice

that

$$\begin{aligned}
C(X) &= \bigcup_{\substack{Y \subset X \\ Y \text{ finite}}} C(Y), \\
A(X) &= \bigcup_{\substack{Y \subset X \\ Y \text{ finite}}} A(Y), \\
C(x_0, \dots, x_{k-1}) &= \left\{ \sum_{i < k} \alpha_i x_i \mid \alpha_i \in \mathbb{Q}, \alpha_i \geq 0, \sum_{i < k} \alpha_i = 1 \right\}, \\
A(x_0, \dots, x_{k-1}) &= \left\{ \sum_{i < k} \alpha_i x_i \mid \alpha_i \in \mathbb{Q}, \sum_{i < k} \alpha_i = 1 \right\}.
\end{aligned}$$

A set  $X \subseteq \mathbb{Q}^m$  is *anchored* if  $A(X) \cap \mathbb{Z}^m \neq \emptyset$ .

The concept of a rational piecewise linear function is straightforward enough, however the definition of  $F_1$  also involves the condition of coefficients being integers, hence we have to understand what it means. Moreover, if  $\Gamma / \Delta$  is a rule in  $m$  variables,  $0 < t_1 < \dots < t_k < 1$  and  $x_0, \dots, x_{k+1} \in [0, 1]_{\mathbb{Q}}^m$ , then the validity of  $\Gamma / \Delta$  under the valuation  $e = L(0, x_0; t_1, x_1; \dots; 1, x_{k+1})$  is completely determined by the vectors  $x_i$ , it does not depend on the parametrization of the function: if we reparametrize it as  $e' = L(0, x_0; t'_1, x_1; \dots; 1, x_{k+1})$  for some  $0 < t'_1 < \dots < t'_k < 1$ , then  $e$  satisfies  $\Gamma / \Delta$  if and only if  $e'$  does. For this reason, we will investigate the following question: given  $x_0, \dots, x_k \in \mathbb{Q}^m$ , when do there exist rational  $t_0 < \dots < t_k$  such that  $L(t_0, x_0; \dots; t_k, x_k)$  has integer coefficients?

The answer given by Lemma 4.8 involves the concept of anchoredness. We thus first provide a characterization (Lemma 4.6) of this condition, which also entails its decidability. Strictly speaking, we do not need the characterization, we could show decidability of anchoredness in another way (cf. the proof of Theorem 4.17). Nevertheless, we decided to include it because we feel that it provides a useful insight into the notion of anchoredness, which allows us to understand it better.

**Lemma 4.5** *For any  $X \in \mathbb{Q}^{m \times k}$ , there exists an  $M \in GL(m, \mathbb{Z})$  such that  $MX$  is in row-echelon form.*

*Proof:* We may assume without loss of generality that  $X \in \mathbb{Z}^{m \times k}$ . The effect of multiplication by  $M$  from left is to perform certain operations on the rows of  $X$ . In particular,  $GL(m, \mathbb{Z})$  contains all permutation matrices, whose effect is to permute the rows of  $X$ , and integer matrices which differ from the identity matrix only in one element which is not on the diagonal, whose effect is to add an integer multiple of some row of  $X$  to another row. It thus suffices to show that we can transform  $X$  into row-echelon form by a sequence of these row operations.

The proof goes by induction on  $m$ . If  $X$  is the zero matrix, there is nothing to do. Otherwise let  $j$  be the first nonzero column of  $X$ . If  $x_{i,j}$  and  $x_{i',j}$  are two nonzero elements of the  $j$ th column and  $|x_{i,j}| \leq |x_{i',j}|$ , we may add a suitable multiple of the  $i$ th row to the  $i'$ th row to reduce  $x_{i',j}$  modulo  $x_{i,j}$ . This operation makes the quantity  $\sum_i |x_{i,j}|$  strictly smaller, hence after a finite number of similar steps we reach the situation that the  $j$ th column contains only one nonzero entry  $x_{i,j}$ . We may permute the rows to ensure  $i = 0$ , and using the induction

hypothesis we apply some operations on the remaining rows to bring the rest of the matrix to row-echelon form.  $\square$

**Lemma 4.6** *The following are equivalent for any  $X \subseteq \mathbb{Q}^m$ .*

(i)  $X$  is anchored.

(ii) For every  $u \in \mathbb{Z}^m$  and  $a \in \mathbb{Q}$ , if  $u^\top x = a$  for all  $x \in X$ , then  $a \in \mathbb{Z}$ .

*Proof:* (i)  $\rightarrow$  (ii):  $\{x \in \mathbb{Q}^m \mid u^\top x = a\}$  is an affine space containing  $X$ , hence it includes  $A(X)$ , which contains an integer point  $x$ . Then  $a = u^\top x \in \mathbb{Z}$ .

(ii)  $\rightarrow$  (i): Clearly  $X \neq \emptyset$ . Since the affine space  $A(X)$  is finitely dimensional, we can find finitely many points  $x_0, \dots, x_k \in X$  such that  $A(X) = A(x_0, \dots, x_k)$ . Put  $y_i = x_i - x_k$  for  $i < k$ , and let  $Y$  be the  $m$ -by- $k$  matrix whose  $i$ th column is  $y_i$ . We have

$$A(X) = \{x_k + Yv \mid v \in \mathbb{Q}^k\}.$$

The condition (ii) then states that for every  $u \in \mathbb{Z}^m$ , if  $u^\top Y = 0$ , then  $u^\top x_k \in \mathbb{Z}$ .

By Lemma 4.5, there exists a matrix  $M \in GL(m, \mathbb{Z})$  such that  $Y' = MY$  is in row-echelon form. Put  $x' = Mx_k$ , and let  $r$  be the number of nonzero rows of  $Y'$ . Consider any  $i \geq r$ . If  $u = M^\top e_i$ , where  $e_i$  is the  $i$ th basis vector, we have  $u \in \mathbb{Z}^m$  and  $u^\top Y = e_i^\top Y' = 0$ , hence  $e_i^\top x' = u^\top x_k \in \mathbb{Z}$ . Thus,  $x'_i \in \mathbb{Z}$  for all  $i \geq r$ . As the first  $r$  rows of  $Y'$  are linearly independent, there exists  $v \in \mathbb{Q}^k$  such that  $Y'v = \langle -x'_0, \dots, -x'_{r-1}, 0, \dots, 0 \rangle$ . Then  $x' + Y'v = \langle 0, \dots, 0, x'_r, \dots, x'_{m-1} \rangle \in \mathbb{Z}^m$ , hence  $A(X) \ni x_k + Yv = M^{-1}(x' + Y'v) \in \mathbb{Z}^m$ .  $\square$

**Corollary 4.7** *Given  $x_0, \dots, x_k \in \mathbb{Q}^m$ , it is decidable whether  $\{x_0, \dots, x_k\}$  is anchored.*

*Proof:* Being anchored is r.e. by definition, and co-r.e. by Lemma 4.6, hence it is decidable. In fact, the proof of Lemmas 4.6 and 4.5 provides a more efficient explicit algorithm.  $\square$

**Lemma 4.8** *If  $x_0, \dots, x_k \in \mathbb{Q}^m$ , then the following are equivalent.*

(i) There exist rationals  $t_0 < \dots < t_k$  such that  $L(t_0, x_0; \dots; t_k, x_k)$  has integer coefficients.

(ii)  $\{x_i, x_{i+1}\}$  is anchored for each  $i < k$ .

*Proof:* (i)  $\rightarrow$  (ii): If  $f = L(t_i, x_i; t_{i+1}, x_{i+1})$  has integer coefficients, then  $f(0) \in \mathbb{Z}^m \cap A(x_i, x_{i+1})$ .

(ii)  $\rightarrow$  (i): By induction on  $k$ . If  $k = 0$ , any  $t_0$  will do, as the condition on coefficients is vacuously true. Assume  $k > 0$ , and let  $t_1 < \dots < t_k$  be such that  $L(t_1, x_1; \dots; t_k, x_k)$  has integer coefficients by the induction hypothesis. We may add a sufficiently large integer to all  $t_i$ , hence we may assume  $t_1 > 0$  without loss of generality. Choose an integer  $d > 0$  such that  $d(x_1 - x_0) \in \mathbb{Z}^m$ , and an  $\alpha \in \mathbb{Q}$  such that  $b = \alpha x_0 + (1 - \alpha)x_1 \in \mathbb{Z}^m$ . By adding an integer multiple of  $d$  to  $\alpha$  we can ensure that  $\alpha > 0$ . We write  $\alpha = p/q$ ,  $t_1 = r/s$  for some natural

numbers  $p, q, r, s$ . Since we may divide all  $t_i$  by  $dqr$ , we may assume that  $r = 1$  and  $dq \mid s$ . Let

$$a = \frac{\alpha}{t_1}(x_1 - x_0) = p \frac{s}{dq}(d(x_1 - x_0)) \in \mathbb{Z}^m,$$

$$t_0 = (1 - \alpha^{-1})t_1.$$

We have  $t_0 < t_1$ , and

$$t_1 a + b = \alpha(x_1 - x_0) + \alpha x_0 + (1 - \alpha)x_1 = x_1,$$

$$t_0 a + b = (\alpha - 1)(x_1 - x_0) + \alpha x_0 + (1 - \alpha)x_1 = x_0,$$

hence  $L(t_0, x_0; t_1, x_1; \dots; t_k, x_k)$  has integer coefficients.  $\square$

A remarkable feature of Lemma 4.8 is that the linear segments do not interact with each other: in order to find a parametrization so that  $L(t_0, x_0; \dots; t_k, x_k)$  has integer coefficients, it is sufficient to parametrize individually each  $L(t, x_i; t', x_{i+1})$  to have integer coefficients.

**Definition 4.9** A (convex) *polytope* is a set of the form  $\{x \in \mathbb{Q}^m \mid \forall i < k L_i(x) \geq 0\}$ , where  $L_i$  are linear (more precisely, affine) functions with integer (or rational, it makes no difference) coefficients.

The following is an effective version of the easy part of Theorem 2.3.

**Lemma 4.10** *Let  $\Gamma$  be a finite set of formulas in  $m$  variables closed under subformulas, and  $n = |\Gamma|$ . We can compute linear functions  $L_{j,i}$  and  $L_{j,\varphi}$  with integer coefficients of absolute value less than  $2^n$  for all  $j < 2^n$ ,  $i < n$ , and  $\varphi \in \Gamma$  such that the polytopes*

$$C_j = \{x \in [0, 1]_{\mathbb{Q}}^m \mid \forall i < n L_{j,i}(x) \geq 0\}$$

satisfy

$$\bigcup_{j < 2^n} C_j = [0, 1]_{\mathbb{Q}}^m,$$

and

$$L_{j,\varphi}(x) = \varphi(x)$$

for each  $x \in C_j$  and  $\varphi \in \Gamma$ .

*Proof:* By induction on  $n$ . We assume for simplicity that all formulas are expressed in the basis  $\{\rightarrow, \perp\}$ . If  $\Gamma$  consists only of variables or  $\perp$ , we can take  $L_{j,i} = 0$ ,  $L_{j,p_i} = x_i$ , and  $L_{j,\perp} = 0$ . Otherwise we pick a formula  $\psi \rightarrow \chi \in \Gamma$  which is not a proper subformula of any formula from  $\Gamma$ . We can apply the induction hypothesis to obtain  $\{L_{j,i} \mid j < 2^{n-1}, i < n-1\}$  and  $\{L_{j,\varphi} \mid j < 2^{n-1}, \varphi \in \Gamma \setminus \{\psi \rightarrow \chi\}\}$ . We put  $L'_{2j,i} = L'_{2j+1,i} = L_{j,i}$  for  $i < n-1$ ,  $L'_{2j,n-1} = -L'_{2j+1,n-1} = L_{j,\psi} - L_{j,\chi}$ ,  $L'_{2j,\varphi} = L'_{2j+1,\varphi} = L_{j,\varphi}$  for  $\varphi \in \Gamma \setminus \{\psi \rightarrow \chi\}$ ,  $L'_{2j,\psi \rightarrow \chi} = 1 - L_{j,\psi} + L_{j,\chi}$ ,  $L'_{2j+1,\psi \rightarrow \chi} = 1$ . It is straightforward to verify the required properties using the definition of  $\rightarrow$  in  $[0, 1]_{\mathbb{L}}$ .  $\square$

The main problem in showing decidability of admissibility in  $\mathbf{L}$  is that potential counterexamples  $L(t_0, x_0; t_1, x_1; \dots; t_k, x_k)$  to a rule  $\Gamma / \Delta$  in  $F_1$  may have arbitrary length, hence we must find a way how to shorten them. The basic idea is as follows. The formulas from  $\Gamma$  define piecewise linear functions, and the domain of each piece is a polytope, let thus consider a particular polytope  $C$  such that all formulas from  $\Gamma$  are linear on  $C$ , and a part  $L(t_i, x_i; t_{i+1}, x_{i+1}; \dots; t_j, x_j)$  of the valuation such that  $x_i, \dots, x_j \in C$ . We could try to simply replace this part with  $L(t_i, x_i; t_j, x_j)$ : since  $\Gamma(x_i) = \Gamma(x_j) = 1$ , and  $\text{rng}(L(t_i, x_i; t_j, x_j)) = C(x_i, x_j) \subseteq C$ , we have  $\Gamma(x) = 1$  for each  $x$  in the range of such a function, which is the main thing we have to preserve. However, there is no guarantee that we can reparametrize  $L(t_i, x_i; t_j, x_j)$  to have integer coefficients. For example, consider the case  $i = 1, j = 3$ , and  $x_2 = \vec{0}$ : then  $\{x_1, x_2\}$  and  $\{x_2, x_3\}$  are anchored for any  $x_1$  and  $x_3$ , but  $\{x_1, x_3\}$  need not be anchored. Fortunately, it cannot get any worse, it turns out that we can do in just two steps what we cannot do in one step: there is  $x \in C$  such that  $\{x_i, x\}$  and  $\{x, x_j\}$  are anchored. This will follow from Lemma 4.12.

**Lemma 4.11** *Let  $X$  be a nonempty convex subset of  $\mathbb{Q}^m$ . There exists a point  $x \in X$  and an open neighbourhood  $U \ni x$  such that  $A(X) \cap U \subseteq X$ .*

*Proof:* Since  $A(X)$  has finite dimension, we can find points  $x_0, \dots, x_k \in X$  such that  $A(X) = A(x_0, \dots, x_k)$ , and the  $x_i$ 's are affinely independent (i.e.,  $x_i \notin A(x_0, \dots, x_{i-1}, x_{i+1}, \dots, x_k)$  for each  $i$ ). Put

$$x = \frac{1}{k+1} \sum_{i \leq k} x_i.$$

Let  $A = \{\alpha \in \mathbb{Q}^{k+1} \mid \sum_i \alpha_i = 1\}$ , and define a mapping  $\varphi: A \rightarrow \mathbb{Q}^m$  by

$$\varphi(\alpha) = \sum_{i \leq k} \alpha_i x_i.$$

Then  $\varphi$  is a homeomorphism of  $A$  onto  $A(X)$  which maps the subset  $B = \{\alpha \in A \mid \forall i \alpha_i \geq 0\}$  onto  $C(x_0, \dots, x_k)$ . The point  $a = \langle 1/(k+1), \dots, 1/(k+1) \rangle$  is in the interior of  $B$  (in  $A$ ), hence  $\varphi(a) = x$  is in the interior of  $C(x_0, \dots, x_k) \subseteq X$  relative to  $A(X)$ .  $\square$

**Lemma 4.12** *Let  $X$  be an anchored subset of  $\mathbb{Q}^m$ , and  $x_0, \dots, x_k \in \mathbb{Q}^m$ . Then there exists  $w \in C(X)$  such that  $\{x_i, w\}$  is anchored for each  $i \leq k$ .*

*Proof:* We may assume that  $X$  is convex without loss of generality. Fix  $c \in A(X) \cap \mathbb{Z}^m$ , and let  $x \in X$  and  $U$  be as in Lemma 4.11. If  $c \in X$ , we can take  $w = c$ . Otherwise  $c \neq x$ , hence  $A(c, x)$  is a line, and its intersection with  $U$  must contain two distinct points  $y, z$ . We have  $y, z \in A(X) \cap U \subseteq X$ , and  $c \in A(y, z) \cap \mathbb{Z}^m$ .

We write  $w_\alpha = (1 - \alpha)y + \alpha z$ . Fix  $\alpha$  such that  $w_\alpha = c$ , and let  $\beta > 0$  be an integer such that  $\beta(z - y) \in \mathbb{Z}^m$ . For every  $i \leq k$ , we can find an integer  $\gamma_i > 0$  such that  $\gamma_i(w_\alpha - x_i) \in \mathbb{Z}^m$ . Then  $w_{\alpha+p\beta} + q\gamma_i(w_\alpha - x_i) \in \mathbb{Z}^m$  for every  $p, q \in \mathbb{Z}$ . If  $q \geq 0$ , we have

$$\begin{aligned} w_{\alpha+p\beta} + q\gamma_i(w_\alpha - x_i) &= (1 + q\gamma_i)w_\alpha - q\gamma_i x_i + p\beta(z - y) \\ &= (1 + q\gamma_i)w_{\alpha+p\beta/(1+q\gamma_i)} - q\gamma_i x_i \in A(x_i, w_{\alpha+p\beta/(1+q\gamma_i)}), \end{aligned}$$

hence  $\{x_i, w_{\alpha+p\beta/(1+q\gamma_i)}\}$  is anchored. Let  $\gamma > \beta$  be a common multiple of all  $\gamma_i$ . There exists an integer  $p$  such that  $\delta = \alpha + p\beta/(1 + \gamma) \in [0, 1]$ . Then  $w_\delta \in C(y, z) \subseteq X$ , and  $\{x_i, w_\delta\}$  is anchored for all  $i \leq k$ .  $\square$

Now we have all tools in order, and we can proceed to our main characterization.

**Theorem 4.13** *Let  $\Gamma, \Delta$  be finite sets of formulas in variables  $\{p_i \mid i < m\}$ , and  $\{C_j \mid j < r\}$  a set of polytopes in  $\mathbb{Q}^m$  such that*

$$\bigcup_{j < r} C_j = \{x \in [0, 1]_{\mathbb{Q}}^m \mid \Gamma(x) = 1\}.$$

*The following are equivalent.*

- (i)  $\Gamma \not\prec_{\mathbf{L}} \Delta$ .
- (ii) *There exists  $a \in \{0, 1\}^m$  such that  $\Gamma(a) = 1$  and for every  $\delta \in \Delta$  there exists a sequence  $\{j_i \mid i \leq k\}$  of indices  $j_i < r$  such that*
  - ( $\alpha$ )  *$j_i$  are pairwise distinct, in particular,  $k < r$ ,*
  - ( $\beta$ )  *$a \in C_{j_0}$ ,*
  - ( $\gamma$ )  *$C_{j_i}$  is anchored for each  $i \leq k$ ,*
  - ( $\delta$ )  *$C_{j_i} \cap C_{j_{i+1}} \neq \emptyset$  for each  $i < k$ ,*
  - ( $\varepsilon$ ) *there exists  $x \in C_{j_k}$  such that  $\delta(x) < 1$ .*

*Proof:* (i)  $\rightarrow$  (ii): We may pick  $f \in F_1^m$  such that  $\Gamma(f) = 1$  and  $\delta(f) \neq 1$  for each  $\delta \in \Delta$  by Theorem 4.1. We can represent  $f$  as  $L(t_0, x_0; \dots; t_s, x_s)$  with integer coefficient for some  $0 = t_0 < t_1 < \dots < t_s = 1$  and  $x_0, \dots, x_s \in [0, 1]_{\mathbb{Q}}^m$ . Since  $\text{rng}(f) \subseteq \bigcup_j C_j$  and the intersection of  $[t_i, t_{i+1}]$  with any  $f^{-1}[C_j]$  is a (possibly empty or degenerate) interval, we can refine the sequence of  $t_i$ 's and  $x_i$ 's to ensure that the range of each  $L(t_i, x_i; t_{i+1}, x_{i+1})$  is included in some  $C_j$ . Put  $a = x_0$ . We have  $a \in \mathbb{Z}^m$  since  $a$  is the constant coefficient of  $L(0, x_0; t_1, x_1)$ , and as  $a \in [0, 1]_{\mathbb{Q}}^m$ , we must have  $a \in \{0, 1\}^m$ .

Consider any  $\delta \in \Delta$ . There exists  $t \in [0, 1]_{\mathbb{Q}}$  such that  $\delta(f(t)) < 1$ . Let  $k < s$  be such that  $t \in [t_k, t_{k+1}]$ . For every  $i \leq k$ , pick  $j_i < r$  such that  $C(x_i, x_{i+1}) = \text{rng}(L(t_i, x_i; t_{i+1}, x_{i+1})) \subseteq C_{j_i}$ . The construction immediately implies conditions ( $\beta$ ) and ( $\varepsilon$ ). Since  $L(t_i, x_i; t_{i+1}, x_{i+1})$  has integer coefficients,  $C_{j_i} \supseteq \{x_i, x_{i+1}\}$  is anchored by Lemma 4.8, hence ( $\gamma$ ) holds. Clearly,  $x_{i+1} \in C_{j_i} \cap C_{j_{i+1}}$ , thus ( $\delta$ ). It remains to satisfy ( $\alpha$ ). If  $j_i = j_{i'}$  for some  $i < i'$ , we may modify the sequence by replacing the subsequence  $j_i, j_{i+1}, \dots, j_{i'}$  with just  $j_i$ . This makes the sequence shorter, and conditions ( $\beta$ )–( $\varepsilon$ ) remain true, hence after finitely many steps we obtain an injective sequence  $\{j_i \mid i \leq k\}$ .

(ii)  $\rightarrow$  (i): We fix  $a \in \{0, 1\}^m$  as in (ii). If  $\Delta = \emptyset$ , then the constant  $a$  function is a valuation in  $F_1$  which refutes  $\Gamma / \Delta$ . Otherwise we enumerate  $\Delta = \{\delta_p \mid p \leq s\}$ . Consider any  $p \leq s$ , and let  $\{j_i \mid i \leq k_p\}$  be a sequence as in (ii). We put  $x_0^p = a$ , and find  $x_{2k_p+2}^p \in C_{j_{k_p}}$  such that  $\delta_p(x_{2k_p+2}^p) < 1$ . For every  $i < k_p$  we pick  $x_{2(i+1)}^p \in C_{j_i} \cap C_{j_{i+1}}$ . Since each  $C_{j_i}$  is

convex and anchored, we can find  $x_{2i+1}^p \in C_{j_i}$  such that  $\{x_{2i}^p, x_{2i+1}^p\}$  and  $\{x_{2i+2}^p, x_{2i+1}^p\}$  are anchored by Lemma 4.12. We form the sequence

$$x_0^0, x_1^0, \dots, x_{2k_0+2}^0, x_{2k_0+1}^0, \dots, x_1^0, x_0^0 = x_0^1, x_1^1, \dots, \\ \dots, x_0^{s-1} = x_0^s, x_1^s, \dots, x_{2k_s+2}^s, x_{2k_s+1}^s, \dots, x_1^s, x_0^s,$$

and relabel it as  $x_0, \dots, x_k$  to simplify the notation, where  $k = 4 \sum_{p \leq s} (k_p + 1)$ . By the construction, we have the following properties for each applicable  $i$ :

- $x_i \in [0, 1]_{\mathbb{Q}}^m$ ,  $x_0, x_k \in \{0, 1\}^m$ ,
- $\{x_i, x_{i+1}\}$  is anchored,
- for each  $\delta \in \Delta$  there exists  $i$  such that  $\delta(x_i) < 1$ ,
- $\{x_i, x_{i+1}\}$  is included in some  $C_j$ , in particular,  $\Gamma(x) = 1$  for every  $x \in C(x_i, x_{i+1})$ .

By Lemma 4.8, there exists a sequence of rational numbers  $t_0 < t_1 < \dots < t_k$  such that  $L(t_0, x_0; \dots; t_k, x_k)$  has integer coefficients. As in the proof of Lemma 4.8, we may add a sufficiently large integer to all  $t_i$  to ensure  $t_0 > 0$ , and we may divide all  $t_i$  by a sufficiently large integer to ensure  $t_k < 1$ . Since  $x_0, x_k \in \mathbb{Z}^m$ ,  $f = L(0, x_0; t_0, x_0; t_1, x_1; \dots; t_k, x_k; 1, x_k)$  also has integer coefficients. Thus  $f \in F_1^m$ , and the aforementioned properties ensure that  $\Gamma(f) = 1$  and  $\delta(f) \neq 1$  for every  $\delta \in \Delta$ , hence  $F_1 \not\equiv \Gamma / \Delta$ , and  $\Gamma \not\sim_{\mathbf{L}} \Delta$ .  $\square$

**Theorem 4.14** *Given a rule  $\Gamma / \Delta$ , it is decidable whether  $\Gamma \sim_{\mathbf{L}} \Delta$ .*

*Proof:* Using Lemma 4.10 we compute a description of a sequence of polytopes  $\{C'_j \mid j < r\}$  such that  $\bigcup_j C'_j = [0, 1]_{\mathbb{Q}}^m$ , and every formula  $\varphi \in \Gamma \cup \Delta$  is defined by a linear function on any  $C'_j$ . We put  $C_j = \{x \in C'_j \mid \forall \gamma \in \Gamma \gamma(x) = 1\}$ . Then the assumptions of Theorem 4.13 hold, it thus suffices to check whether the condition (ii) is true. We can do it by a brute-force search for possible  $a \in \{0, 1\}^m$  and  $\{j_i \mid i \leq k\}$ ,  $j_i < r$ ,  $k < r$ . We only need to check that conditions  $(\alpha)$ – $(\varepsilon)$  can be algorithmically verified. Conditions  $(\alpha)$  and  $(\beta)$  are immediate. We can verify  $(\delta)$  and  $(\varepsilon)$  by any linear programming algorithm, as we can express  $\delta$  by a linear function on  $C_{j_k}$ .

Finally, we need to verify whether  $C_{j_i}$  is anchored. This can be done as follows. We know from the theory of linear programming that  $C_{j_i}$  is the convex hull of its vertices, and each vertex can be described as the unique solution of a system of linear equations obtained from a subset of the defining inequalities of  $C_{j_i}$  by changing  $\leq$  to  $=$ . We can systematically list all such linear systems, use Gaussian elimination to check whether it has a unique solution, and if so, to compute the solution, and verify whether it satisfies the remaining inequalities from  $C_{j_i}$ . In this way we obtain the list  $x_0, \dots, x_k$  of all vertices of  $C_{j_i}$ , and then we can check whether  $\{x_0, \dots, x_k\}$  is anchored by Corollary 4.7.  $\square$

**Corollary 4.15** *The universal theory of any free MV-algebra is decidable.*  $\square$

## 4.1 Complexity

In this section we take a look on complexity issues concerning admissibility in  $\mathbf{L}$ . First, Theorem 4.14 implies that there exists a computable bound on the size (number of bits) of a counterexample to an inadmissible rule. We provide explicit estimates below. Second, we give an upper bound on the computational complexity of  $\vdash_{\mathbf{L}}$ , namely we show that it is computable in polynomial space (and therefore in exponential time).

**Definition 4.16** The *height*  $H(x)$  of a rational number  $x$  is  $\max\{|p|, |q|\}$ , where  $p, q$  are coprime integers such that  $x = p/q$ . More generally, if  $x \in \mathbb{Q}^m$ , its height  $H(x)$  is

$$\max\{|qx_i|, q \mid i < m\},$$

where  $q$  is the smallest nonzero natural number such that  $qx \in \mathbb{Z}^m$ . Notice that the natural representation of  $x$  in binary takes  $O(m \log H(x))$  bits.

**Theorem 4.17** Let  $\Gamma / \Delta$  be a rule of length  $n = \sum_{\varphi \in \Gamma \cup \Delta} |\varphi|$  in  $m$  variables. If  $\Gamma \not\vdash_{\mathbf{L}} \Delta$ , there exists a valuation  $f = L(t_0 = 0, x_0; t_1, x_1; \dots; t_k = 1, x_k) \in F_1^m$  such that  $\Gamma(f) = 1$ , for each  $\delta \in \Delta$  there is  $r \leq k$  such that  $\delta(x_r) < 1$ ,  $k = 1 + O(|\Delta|2^n) = O(n2^n)$ , for each  $r \leq k$ :  $H(x_r) = 2^{O(nm)}$ ,  $H(t_r) = 2^{O(nmk)} = 2^{O(n^2m2^n)}$ , and there exists  $a_r \in A(x_r, x_{r+1}) \cap \mathbb{Z}^m$  of height  $H(a_r) = 2^{O(nm)}$ .

*Proof:* Note that  $t_r \in [0, 1]$  and  $x_r \in [0, 1]^m$ , hence their height is just a bound on the denominator  $q$ , whereas the height of  $a_r$  coincides with its  $L^\infty$  norm  $\|a_r\|_\infty$ . Let  $\Sigma$  be the set of all subformulas of  $\Gamma \cup \Delta$ . As  $n \geq |\Sigma|$ , by Lemma 4.10 we can write

$$\{x \in [0, 1]_{\mathbb{Q}}^m \mid \Gamma(x) = 1\} = \bigcup_{j < 2^n} C_j,$$

where each  $C_j$  is described by a set of linear inequalities with integer coefficients of absolute value at most  $2^n$ . Also, every  $\delta \in \Delta$  is defined on each  $C_j$  by a linear function with integer coefficients bounded by  $2^n$ . We assume that condition (ii) of Theorem 4.13 is satisfied, and we will extract explicit bounds from the proof of the implication (ii)  $\rightarrow$  (i).

We first bound  $x_{2r}$ . By the construction, we have  $x_{2r} = a \in \{0, 1\}^m$ , which has height 1, or we choose an arbitrary  $x_{2r}$  in some  $C = C_j \cap C_{j'}$ , or we choose  $x_{2r} \in C = C_j$  such that  $\delta(x_{2r}) < 1$  for some  $\delta \in \Delta$ . In the latter two cases, we may take for  $x_{2r}$  a vertex of the polytope  $C$ . Such a vertex is the unique solution of a system of linear equations obtained from a subset of the defining inequalities of  $C$  by replacing  $\geq$  with  $=$ . If we take a minimal set of these equations, then their number must be  $m$  (a larger number of equations would be linearly dependent, and a smaller number cannot have a unique solution). We can thus write  $Ax_{2r} = b$ , where  $A$  is a regular  $m$ -by- $m$  matrix, and the coefficients of  $A$  and  $b$  are integers bounded by  $2^n$ . Then  $x_{2r} = A^{-1}b$ , and by Cramer's rule the height (i.e., common denominator) of  $x_{2r}$  is a divisor of  $\det(A)$ . We thus have  $H(x_{2r}) \leq |\det(A)| \leq m!2^{nm} = 2^{O(nm)}$ .

In order to get a better bound, we exploit the structure of the defining inequalities of  $C$  using extension variables. Let us denote the original variables of  $L_{j,i}$  and  $L_{j,\varphi}$  by  $u_i$  to avoid clashes with the vectors  $x_r$ . We assume for simplicity  $C = C_j$ , the case of  $C = C_j \cap C_{j'}$  is



similar. We introduce new variables  $u_\varphi$  for all  $\varphi \in \Sigma$ , where we identify  $u_i = u_{p_i}$ . If  $v \in \mathbb{Q}^\Sigma$ , let us write  $v \upharpoonright m = \langle v_0, \dots, v_{m-1} \rangle$ . We put

$$C'_j = \{v \in \mathbb{Q}^\Sigma \mid v \upharpoonright m \in C_j, \forall \varphi \in \Sigma v_\varphi = L_{j,\varphi}(v \upharpoonright m)\}.$$

Then  $C'_j$  is also a bounded polytope, its projection to the first  $m$  coordinates is  $C_j$ , and for each  $w \in C_j$  there exists a unique  $v \in C'_j$  such that  $v \upharpoonright m = w$ . It follows easily that vertices of  $C_j$  are exactly the points  $v \upharpoonright m$  where  $v$  is a vertex of  $C'_j$ , so let  $x'$  be a vertex of  $C'_j$  such that  $x' \upharpoonright m = x_{2r}$ . We can describe  $C'_j$  by the following set of inequalities and equations (assuming for simplicity that formulas are expressed in the basis  $\{\perp, \rightarrow\}$ ):  $0 \leq u_i \leq 1$  for each  $i < m$ ,  $u_\gamma \geq 1$  for each  $\gamma \in \Gamma$ ,  $u_\perp = 0$ , and for each  $\varphi \rightarrow \psi \in \Sigma$ , either  $u_\varphi \geq u_\psi$ ,  $u_{\varphi \rightarrow \psi} = 1 - u_\varphi + u_\psi$ , or  $u_\varphi \leq u_\psi$ ,  $u_{\varphi \rightarrow \psi} = 1$ , depending on  $j$ . By the same reasoning as above, we can write  $Ax' = b$ , where  $A$  is a regular  $|\Sigma|$ -by- $|\Sigma|$  matrix, all coefficients of  $A$  and  $b$  are integers with absolute value at most 1, and there are at most 3 nonzero coefficients in every row of  $A$ . Thus  $H(x_{2r}) \leq H(x') \leq |\det(A)| \leq 3^{|\Sigma|} = 2^{O(n)}$ .

Now we proceed to bound  $x_{2r+1}$ . Recall that this point is chosen so that  $\{x_{2r}, x_{2r+1}\}$  and  $\{x_{2r+2}, x_{2r+1}\}$  are anchored, and  $x_{2r+1} \in C_j$  for certain  $j$ , where we know that  $C_j$  is anchored. Let  $\{u_0, \dots, u_l\}$  be a maximal affinely independent set of vertices of  $C_j$ . Clearly  $l \leq m$ , and by the same reasoning as above, we have  $H(u_i) = 2^{O(n)}$ . Since  $C_j$  is the convex hull of its vertices, we must have  $A(C_j) = A(u_0, \dots, u_l)$ . For each  $i < l$ , let  $v_i$  be a multiple of  $u_i - u_l$  such that  $v_i \in \mathbb{Z}^m$  and  $H(v_i) = 2^{O(n)}$ . We have  $A(C_j) = \{u_l + Vz \mid z \in \mathbb{Q}^l\}$ , where  $V$  is the  $m$ -by- $l$  integer matrix whose  $i$ th column is  $v_i$ . There exists a point  $c = u_l + Vz \in A(C_j) \cap \mathbb{Z}^m$ . We may add any integer vector to  $z$ , hence we may assume that  $\|z\|_\infty \leq 1$ , thus  $\|Vz\|_\infty \leq \sum_i \|v_i\|_\infty = 2^{O(n)}$  and  $H(c) = \|c\|_\infty = 2^{O(n)}$ .

Put  $x = \frac{1}{l+1} \sum_{i \leq l} u_i$ . Note that the denominator of  $x$  is at most the product of the denominators of the  $u_i$ 's times  $l+1$ , hence  $H(x) = 2^{O(nm)}$ . We need to find an  $\varepsilon > 0$  such that  $A(C_j) \cap U \subseteq C_j$ , where  $U$  is the Euclidean ball around  $x$  with radius  $\varepsilon$ . The interior of  $C(u_0, \dots, u_l)$  relative to  $A(C_j)$  is the set  $\{\sum_i \alpha_i u_i \mid \alpha_i > 0, \sum_i \alpha_i = 1\}$ , hence a suitable  $\varepsilon$  is

$$\varepsilon = \min_{i \leq l} d(x, A(u_0, \dots, u_{i-1}, u_{i+1}, \dots, u_l)) = \frac{1}{l+1} \min_{i \leq l} d(u_i, A(u_0, \dots, u_{i-1}, u_{i+1}, \dots, u_l)),$$

where  $d$  denotes the Euclidean ( $L^2$ ) distance. By symmetry, it suffices to find a lower bound on

$$d(u_l, A(u_0, \dots, u_{l-1})) = d(u_l - u_0, S),$$

where  $S$  is the linear span of  $\{u_1 - u_0, \dots, u_{l-1} - u_0\}$ . For each  $i = 1, \dots, l-1$ , let  $w_i$  be a multiple of  $u_i - u_0$  such that  $w_i \in \mathbb{Z}^m$  and  $H(w_i) = 2^{O(n)}$ . Let  $W$  be the  $m$ -by- $(l-1)$  integer matrix whose columns are the vectors  $w_i$ . As  $w_i$  are linearly independent,  $W^\top W$  is regular, and we may define  $P = W(W^\top W)^{-1}W^\top$ . We have  $P = P^\top$ ,  $PW = W$ , and  $P = WX$  for some matrix  $X$ , hence  $P$  is the orthogonal projection on  $S$ . Thus

$$\begin{aligned} d(u_l - u_0, S)^2 &= \|(u_l - u_0) - P(u_l - u_0)\|_2^2 = (u_l - u_0)^\top (E - P^\top)(E - P)(u_l - u_0) \\ &= (u_l - u_0)^\top (E - P)(u_l - u_0). \end{aligned}$$

Since  $W$  is an integer matrix, we have  $dP \in \mathbb{Z}^{m \times m}$  for  $d = |\det(W^T W)| = 2^{O(nm)}$  by Cramer's rule. Since also  $H(u_l - u_0) = 2^{O(n)}$ , the denominator of  $d(u_l - u_0, S)^2$  is  $2^{O(nm)}$ . As  $u_i$  are affinely independent, we must have  $d(u_l - u_0, S) > 0$ , hence we obtain

$$\varepsilon = 2^{-O(nm)}.$$

Therefore we can find  $y \neq z$ ,  $y, z \in C_j \cap A(c, x)$  such that  $H(y), H(z) = 2^{O(nm)}$ . Then, using the notation of Lemma 4.12, we have  $H(\alpha) = 2^{O(nm)}$ ,  $\beta = 2^{O(nm)}$ ,  $\gamma_i = 2^{O(n)}$ ,  $\gamma = 2^{O(nm)}$ , and  $H(\delta) = 2^{O(nm)}$ , hence the height of  $x_{2r+1} = w_\delta$  is  $2^{O(nm)}$ .

Also  $p = 2^{O(nm)}$ , hence the height of the integer point  $a_{2r} = w_{\alpha+p\beta} + \gamma(w_\alpha - x_{2r}) \in A(x_{2r}, x_{2r+1})$  is  $2^{O(nm)}$ , and similarly for  $a_{2r+1}$ .

It remains to analyze the inductive argument in Lemma 4.8. In order to get a decent bound, we modify the induction hypothesis: we require  $0 < t_0 < \dots < t_k < 1$ , and  $1/t_0 \in \mathbb{Z}$ . Consider the induction step. Using the notation of Lemma 4.8, we have  $d = 2^{O(nm)}$  and  $H(\alpha) = 2^{O(nm)}$ . We may ensure  $\alpha > 2$ . What happens to the  $t_r$ 's is that we first divide all  $t_r$  by  $dq = 2^{O(nm)}$ , and we put  $t_0 = (1 - \alpha^{-1})t_1$ . Then  $t_0 \geq t_1/2$ , hence the extra conditions  $t_0 > 0$  and  $t_k < 1$  are satisfied. Since  $t_0 = (p - q)/ps$ , it suffices to divide all  $t_r$  by  $p - q = 2^{O(nm)}$  in order to satisfy the condition  $t_0^{-1} \in \mathbb{Z}$ . All in all, we see that  $\max_r H(t_r)$  is multiplied by at most  $2^{O(nm)}$  in each step, hence the final  $H(t_r)$  is  $2^{O(knm)}$ . (In fact, the same argument shows  $H(\langle t_0, \dots, t_k \rangle) = 2^{O(knm)}$ .)  $\square$

**Theorem 4.18** *Admissibility in  $\mathbf{L}$  is in PSPACE. More precisely, we can test admissibility of multiple-conclusion rules in  $\mathbf{L}$  using space  $O(n^2m)$ , where  $n$  is the total size of the input, and  $m \leq n$  is the number of variables.*

*Proof:* Let us first describe a nondeterministic polynomial-space algorithm. The basic idea is to nondeterministically search for a counterexample  $f$  as in Theorem 4.17, but there are several caveats. First, the numbers  $t_r$  have exponential size. Fortunately we do not really need them, it suffices to construct the  $x_r$ 's and the integer points  $a_r$  witnessing that  $\{x_r, x_{r+1}\}$  is anchored. Second,  $k$  is exponentially large, hence we cannot write down a full description of  $f$ . Instead we search for it piece by piece, we only need to remember two successive  $x_r$  at a time, as all the conditions are local. Third, we need somehow to certify that  $\Gamma(x) = 1$  for all  $x \in C(x_r, x_{r+1})$ . By the proof of Theorem 4.17, we may require that  $x_r, x_{r+1} \in C$  for a suitable polytope  $C$  from Lemma 4.10 such that all  $\gamma \in \Gamma$  are linear on  $C$ ; then it suffices to verify  $\Gamma(x_r) = \Gamma(x_{r+1}) = 1$ .

More precisely, let  $\Sigma$  be the closure of  $\Gamma \cup \Delta$  under subformulas, and  $n_\Sigma = |\Sigma| \leq n$ . Let  $\{C'_j \mid j < 2^{n_\Sigma}\}$  be the sequence of polytopes from Lemma 4.10 for  $\Sigma$ . An inspection of the proof shows that given  $j < 2^{n_\Sigma}$  in binary, we can compute the sets of linear functions  $\{L_{j,i} \mid i < n_\Sigma\}$  and  $\{L_{j,\varphi} \mid \varphi \in \Sigma\}$  in space linear in the size of the output, i.e.,  $O(n^2m)$ . Putting  $C_j = \{x \in C'_j \mid \Gamma(x) = 1\}$ , we see that we can compute inequalities defining  $C_j$  within the same space bound: they consist of

$$\{L_{j,i} \geq 0 \mid i < n_\Sigma\} \cup \{x_i \geq 0, 1 - x_i \geq 0 \mid i < m\} \cup \{L_{j,\gamma} - 1 \geq 0 \mid \gamma \in \Gamma\}.$$

Consider the algorithm in Figure 1. Notice that the test  $a \in A(x, y)$  can be implemented

```

 $D \leftarrow \Delta$ 
nondeterministically guess  $x \in \{0, 1\}^m$ 
for  $i \leftarrow 1, \dots, O(n2^n)$  do:
  nondeterministically guess  $j < 2^{n^\Sigma}$ 
  if  $x \notin C_j$  then REJECT
  if  $\exists \gamma \in \Gamma L_{j,\gamma}(x) < 1$  then REJECT
  if  $L_{j,\delta}(x) < 1$  for some  $\delta \in D$ , then  $D \leftarrow D \setminus \{\delta\}$ 
  if  $D = \emptyset$  then ACCEPT
  nondeterministically guess  $y \in [0, 1]_{\mathbb{Q}}^m$  with  $H(y) = 2^{O(nm)}$ 
  if  $y \notin C_j$  then REJECT
  nondeterministically guess  $a \in \mathbb{Z}^m$  with  $H(a) = 2^{O(nm)}$ 
  if  $a \notin A(x, y)$  then REJECT
   $x \leftarrow y$ 
REJECT

```

Figure 1: A nondeterministic algorithm for  $\not\prec_{\mathbf{L}}$

as follows. If  $x = y$ , then it is equivalent to  $a = x$ . Otherwise we find the least  $i$  such that  $x_i \neq y_i$ , compute  $\alpha = (a_i - x_i)/(y_i - x_i)$ , and check whether  $a_j = (1 - \alpha)x_j + \alpha y_j$  for all  $j$ .

It follows from Theorems 4.17 and 4.13 that there exists an accepting computation path if and only if  $\Gamma \not\prec_{\mathbf{L}} \Delta$ . The space requirements of the algorithm are dominated by  $O(nm^2)$  to store the vectors  $x, y, a$ , and  $O(n^2m)$  to perform the operations with  $C_j$ . In total, the algorithm works in space  $O(n^2m) \subseteq O(n^3)$ .

By Savitch's theorem, we can construct a deterministic algorithm working in space  $O(n^6)$ . We can obtain a better bound if we write down the algorithm explicitly and analyze it. We consider the recursive procedure in Figure 2. We see that  $Path(x, y, k, D)$  accepts if and only if there exists a sequence  $x_0, \dots, x_k \in [0, 1]_{\mathbb{Q}}^m$  with  $H(x_r) = 2^{O(nm)}$  such that  $x_0 = x$ ,

```

function  $Path(x, y, k, D)$ :
if  $k = 1$ :
  for every  $j < 2^{n^\Sigma}$  and  $a \in \mathbb{Z}^m$  such that  $H(a) = 2^{O(nm)}$ :
    if  $x \in C_j \wedge y \in C_j \wedge a \in A(x, y)$ 
       $\wedge \forall \gamma \in \Gamma L_{j,\gamma}(x) = 1 \wedge \forall \delta \in D L_{j,\delta}(x) < 1$ 
    then ACCEPT
  REJECT
for every  $D' \subseteq D$  and  $z \in [0, 1]_{\mathbb{Q}}^m$  such that  $H(z) = 2^{O(nm)}$ :
  if  $Path(x, z, \lfloor k/2 \rfloor, D') \wedge Path(z, y, \lceil k/2 \rceil, D \setminus D')$  then ACCEPT
REJECT

```

Figure 2: A deterministic subprocedure for  $\not\prec_{\mathbf{L}}$

$x_k = y$ , for each  $r < k$ ,  $\{x_r, x_{r+1}\}$  has an anchor of height  $2^{O(nm)}$  and is contained in some  $C_j$ ,  $\Gamma(x_r) = 1$ , and for each  $\delta \in D$  there is  $r < k$  such that  $\delta(x_r) < 1$ . Using Theorems 4.17 and 4.13 again, we have

$$\Gamma \not\sim_{\mathbf{L}} \Delta \Leftrightarrow \exists a \in \{0, 1\}^m \text{Path}(a, a, O(n2^n), \Delta).$$

Each recursive call of *Path* needs local storage  $O(nm^2)$ , and the depth of recursion is  $O(n)$ , hence the recursion needs space  $O(n^2m^2)$ . The base case  $k = 1$  needs  $O(nm^2)$  bits to store  $a$ , and  $O(n^2m)$  to perform operations with  $C_j$ . Therefore the total space requirements of the algorithm are  $O(n^2m^2)$ .

We can reduce the space further by observing that in Theorem 4.17 we only need  $H(x_r) = 2^{O(nm)}$  for odd  $r$ , whereas for even  $r$  we have a better bound  $H(x_r) = 2^{O(n)}$ . We can modify *Path* so that all recursive calls except the deepest one are performed with  $z$  being an ‘‘even point’’. This brings down the space requirement of the recursive phase to  $O(n^2m)$ . The base case also fits into this bound as  $m \leq n$ , hence the total space used by the algorithm is  $O(n^2m)$ .  $\square$

**Remark 4.19** We can also devise a *PSPACE* algorithm for  $\sim_{\mathbf{L}}$  by an exhaustive search for the sequences  $\{j_i \mid i \leq k\}$  from Theorem 4.13 instead of the sequence  $\{x_r \mid r \leq k\}$ . We can use the estimates of Theorem 4.17 to implement space-efficient tests for  $C_{j_i} \cap C_{j_{i+1}} \neq \emptyset$  and anchoredness of  $C_{j_i}$ . If we further employ a log-space algorithm for undirected connectivity (Reingold [20]) and space-efficient formula evaluation similar to Lynch [16], we can obtain in this way an algorithm for  $\sim_{\mathbf{L}}$  working in space  $O(nm^2 + n \log n)$ , which is slightly better than the bound of Theorem 4.18. We omit the details.

**Corollary 4.20** *The universal theory of any free MV-algebra is decidable in PSPACE.*

*Proof:* The case of  $F_0$  is trivial, let thus  $\varkappa > 0$ . If  $\varphi(\vec{x})$  is an open formula in the language of *MV*-algebras, we can write  $\varphi$  as  $\psi(t_0 = s_0, \dots, t_{k-1} = s_{k-1})$  for some propositional formula  $\psi$ , and terms  $t_i, s_i$  in variables  $\vec{x}$ . We identify  $t_i, s_i$  with the corresponding formulas of  $\mathbf{L}$ . Write  $\alpha^0 = \alpha$ ,  $\alpha^1 = \neg\alpha$ . Using Corollary 4.2, we have

$$\begin{aligned} F_{\varkappa} \models \forall \vec{x} \varphi(\vec{x}) &\Leftrightarrow \forall e: k \rightarrow 2 \left( \psi(e) = 0 \Rightarrow F_{\varkappa} \models \bigvee_{i < k} (t_i = s_i)^{e(i)} \right) \\ &\Leftrightarrow \forall e: k \rightarrow 2 (\psi(e) = 0 \Rightarrow \{t_i \leftrightarrow s_i \mid e(i) = 1\} \sim_{\mathbf{L}} \{t_i \leftrightarrow s_i \mid e(i) = 0\}), \end{aligned}$$

which is decidable in *PSPACE* by Theorem 4.18.  $\square$

## 5 Open problems

We have provided a solution to the most obvious question concerning admissibility in  $\mathbf{L}$ , namely whether it is decidable. Nevertheless, many problems in this area remain open. First, we did not obtain any information on bases of  $\mathbf{L}$ -admissible rules.

**Problem 5.1** *Does  $\mathbf{L}$  have a finite basis of admissible rules? Describe an explicit basis of  $\mathbf{L}$ -admissible rules.*

Another set of questions concerns unification in  $\mathbf{L}$ , and projectivity. In many superintuitionistic and modal logics, as well as  $n$ -contractive extensions of  $\mathbf{BL}$ , every formula has a finite basis of unifiers, which are the most general unifiers of some projective formulas. Moreover, projective formulas have a transparent semantic characterization. Let us say that a *projective approximation* of a formula  $\varphi$  is a finite set  $\Pi$  of projective formulas such that  $\varphi \sim_{\mathbf{L}} \Pi$ , and  $\pi \vdash_{\mathbf{L}} \varphi$  for each  $\pi \in \Pi$ .

**Problem 5.2** *Does every formula have a projective approximation in  $\mathbf{L}$ ? Give a description of  $\mathbf{L}$ -projective formulas. What is the unification type of  $\mathbf{L}$ ?*

With regards to computational complexity of admissibility in  $\mathbf{L}$ , we have shown a *PSPACE* upper bound, but we only have a trivial *coNP* lower bound given by the complexity of the set of  $\mathbf{L}$ -tautologies (Mundici [18]).

**Problem 5.3** *Is  $\sim_{\mathbf{L}}$  PSPACE-complete? Is it in coNP, or at least in the polynomial hierarchy?*

Finally, our analysis of admissible rules in  $\mathbf{L}$  heavily relied on the relatively transparent structure of free *MV*-algebras given by McNaughton's theorem. It seems much more difficult to generalize it to weaker fuzzy logics.

**Problem 5.4** *Is admissibility decidable in  $\mathbf{BL}$  or  $\mathbf{MTL}$ ?*

## References

- [1] Alexander V. Chagrov, *A decidable modal logic with the undecidable admissibility problem for inference rules*, Algebra and Logic 31 (1992), no. 1, pp. 53–55.
- [2] Petr Cintula, *Short note: on the redundancy of axiom (A3) in BL and MTL*, Soft Computing 9 (2005), no. 12, p. 942.
- [3] Petr Cintula and George Metcalfe, *Structural completeness in fuzzy logics*, Notre Dame Journal of Formal Logic, to appear.
- [4] Wojciech Dzik, *Unification of some substructural logics of BL-algebras and hoops*, Reports on Mathematical Logic 43 (2008), pp. 73–83.
- [5] Harvey M. Friedman, *One hundred and two problems in mathematical logic*, Journal of Symbolic Logic 40 (1975), no. 2, pp. 113–129.
- [6] Silvio Ghilardi, *Unification through projectivity*, Journal of Logic and Computation 7 (1997), no. 6, pp. 733–752.
- [7] ———, *Unification in intuitionistic logic*, Journal of Symbolic Logic 64 (1999), no. 2, pp. 859–880.

- [8] ———, *Best solving modal equations*, *Annals of Pure and Applied Logic* 102 (2000), no. 3, pp. 183–198.
- [9] Petr Hájek, *Metamathematics of fuzzy logic*, Kluwer, Dordrecht, 1998.
- [10] Rosalie Iemhoff, *On the admissible rules of intuitionistic propositional logic*, *Journal of Symbolic Logic* 66 (2001), no. 1, pp. 281–294.
- [11] ———, *Intermediate logics and Visser’s rules*, *Notre Dame Journal of Formal Logic* 46 (2005), no. 1, pp. 65–81.
- [12] ———, *On the rules of intermediate logics*, *Archive for Mathematical Logic* 45 (2006), no. 5, pp. 581–599.
- [13] Emil Jeřábek, *Admissible rules of modal logics*, *Journal of Logic and Computation* 15 (2005), no. 4, pp. 411–431.
- [14] ———, *Complexity of admissible rules*, *Archive for Mathematical Logic* 46 (2007), no. 2, pp. 73–92.
- [15] Paul Lorenzen, *Einführung in die operative Logik und Mathematik*, *Grundlehren der mathematischen Wissenschaften* vol. 78, Springer, 1955 (in German).
- [16] Nancy Lynch, *Log space recognition and translation of parenthesis languages*, *Journal of the Association for Computing Machinery* 24 (1977), no. 4, pp. 583–590.
- [17] Robert McNaughton, *A theorem about infinite-valued sentential logic*, *Journal of Symbolic Logic* 16 (1951), no. 1, pp. 1–13.
- [18] Daniele Mundici, *Satisfiability in many-valued sentential logic is NP-complete*, *Theoretical Computer Science* 52 (1987), no. 1–2, pp. 145–153.
- [19] Jeffrey S. Olson, James G. Raftery, and Clint J. van Alten, *Structural completeness in substructural logics*, *Logic Journal of the IGPL* 16 (2008), no. 5, pp. 453–495.
- [20] Omer Reingold, *Undirected connectivity in log-space*, *Journal of the Association for Computing Machinery* 55 (2008), no. 4.
- [21] Vladimir V. Rybakov, *Admissibility of logical inference rules*, *Studies in Logic and the Foundations of Mathematics* vol. 136, Elsevier, 1997.
- [22] ———, *Logical consecutions in discrete linear temporal logic*, *Journal of Symbolic Logic* 70 (2005), no. 4, pp. 1137–1149.
- [23] ———, *Logical consecutions in intransitive temporal linear logic of finite intervals*, *Journal of Logic and Computation* 15 (2005), no. 5, pp. 663–678.

- [24] —————, *Linear temporal logic with Until and Before on integer numbers, deciding algorithms*, in: Computer Science – Theory and Applications (D. Grigoriev, J. Harrison, and E. A. Hirsch, eds.), Lecture Notes in Computer Science vol. 3967, Springer, 2006, pp. 322–333.
- [25] D. J. Shoesmith and Timothy J. Smiley, *Multiple-conclusion logic*, Cambridge University Press, 1978.
- [26] Frank Wolter and Michael Zakharyashev, *Undecidability of the unification and admissibility problems for modal and description logics*, ACM Transactions on Computational Logic 9 (2008), no. 4.