

Kybernetika

VOLUME 43 (2007), NUMBER 2

The Journal of the Czech Society for
Cybernetics and Information Sciences

Published by:

Institute of Information Theory
and Automation of the AS CR, v.v.i.

Editor-in-Chief:

Milan Mareš

Managing Editors:

Karel Sladký
Lucie Fajfrová

Editorial Board:

Jiří Anděl, Sergej Čelikovský, Marie
Demlová, Petr Hájek, Jan Flusser, Martin
Janžura, Jan Ježek, George Klir, Ivan
Kramosil, Tomáš Kroupa, Friedrich Liese,
Jean-Jacques Loiseau, František Matúš,
Radko Mesiar, Jiří Outrata, Jan Štecha,
Olga Štěpánková, Igor Vajda, Jiřina
Vejnarová, Miloslav Vošvrda, Pavel Zítek

Editorial Office:

Pod Vodárenskou věží 4, 182 08 Praha 8

Kybernetika is a bi-monthly international journal dedicated for rapid publication of high-quality, peer-reviewed research articles in fields covered by its title.

Kybernetika traditionally publishes research results in the fields of Control Sciences, Information Sciences, System Sciences, Statistical Decision Making, Applied Probability Theory, Random Processes, Fuzziness and Uncertainty Theories, Operations Research and Theoretical Computer Science, as well as in the topics closely related to the above fields.

The Journal has been monitored in the Science Citation Index since 1977 and it is abstracted/indexed in databases of Mathematical Reviews, Current Mathematical Publications, Current Contents ISI Engineering and Computing Technology.

Kybernetika. Volume 43 (2007)

ISSN 0023-5954, MK ČR E 4902.

Published bimonthly by the Institute of Information Theory and Automation of the Academy of Sciences of the Czech Republic, Pod Vodárenskou věží 4, 182 08 Praha 8. — Address of the Editor: P. O. Box 18, 182 08 Prague 8, e-mail: kybernetika@utia.cas.cz. — Printed by PV Press, Pod vrstevnicí 5, 140 00 Prague 4. — Orders and subscriptions should be placed with: MYRIS TRADE Ltd., P. O. Box 2, V Štíhlách 1311, 142 01 Prague 4, Czech Republic, e-mail: myris@myris.cz. — Sole agent for all “western” countries: Kubon & Sagner, P. O. Box 34 01 08, D-8 000 München 34, F.R.G.

Published in June 2007.

© Institute of Information Theory and Automation of the AS CR, v.v.i., Prague 2007.

ON THE STRUCTURE OF CONTINUOUS UNINORMS

PAWEŁ DRYGAŚ

Uninorms were introduced by Yager and Rybalov [13] as a generalization of triangular norms and conorms. We ask about properties of increasing, associative, continuous binary operation U in the unit interval with the neutral element $e \in [0, 1]$. If operation U is continuous, then $e = 0$ or $e = 1$. So, we consider operations which are continuous in the open unit square. As a result every associative, increasing binary operation with the neutral element $e \in (0, 1)$, which is continuous in the open unit square may be given in $[0, 1]^2$ or $(0, 1)^2$ as an ordinal sum of a semigroup and a group. This group is isomorphic to the positive real numbers with multiplication. As a corollary we obtain the results of Hu, Li [7].

Keywords: uninorms, continuity, t -norms, t -conorms, ordinal sum of semigroups

AMS Subject Classification: 06F05, 03E72, 03B52

1. INTRODUCTION

Uninorms were introduced by Yager and Rybalov [13] as a generalization of triangular norms and conorms. However similar operations were considered in [3] and [4]. In [6] Fodor, Yager and Rybalov examined a general structure of uninorms. For example, the frame structure of uninorms and characterization of representable uninorms are presented.

In this paper we consider a more general class of operations than uninorms, i.e. operations from the class $\mathcal{U}(e) = \{U : [0, 1]^2 \rightarrow [0, 1] : U \text{ is an increasing, associative binary operation with the neutral element } e\}$ for $e \in [0, 1]$, where we omit the assumption about the commutativity. We ask about properties of continuous operation U in $\mathcal{U}(e)$ where $e \in [0, 1]$. If operation U is continuous then $e = 0$ or $e = 1$ (cf. [3]). So, we consider operations which are continuous in the open unit square. The structure of operations continuous on another subset of unit square we can find in [6, 11, 12].

First, in the Section 2 we present the notion of uninorms and the frame structure of uninorms. Next we present the construction of ordinal sum of semigroups. In Section 4 we present properties of the operation which is continuous in $(0, 1)^2$.

As a result every operation in $\mathcal{U}(e)$ with $e \in (0, 1)$, which is continuous in the open unit square may be given in $[0, 1]^2$ or $(0, 1)^2$ as an ordinal sum of a semigroup and a group. This group is isomorphic to the positive real numbers with multiplication.

Moreover this operation is commutative beyond from two points at the most. As a corollary we obtain results of Hu, Li [7] and Fodor, Yager, Rybalov [6].

2. NOTION OF UNINORMS

We discuss the structure of binary operations $U : [0, 1]^2 \rightarrow [0, 1]$.

Definition 1. (Yager and Rybalov [13]) An operation U is called a uninorm if it is commutative, associative, increasing and has the neutral element $e \in [0, 1]$.

Uninorms are generalizations of triangular norms (case $e = 1$) and triangular conorms (case $e = 0$). In the case $e \in (0, 1)$ a uninorm U is composed by using a triangular norm and a triangular conorm.

Theorem 1. (Fodor, Yager and Rybalov [6]) If a uninorm U has the neutral element $e \in (0, 1)$, then there exist a triangular norm T and a triangular conorm S such that

$$U = \begin{cases} T^* \text{ in } [0, e]^2, \\ S^* \text{ in } [e, 1]^2, \end{cases} \quad (1)$$

where

$$\begin{cases} T^*(x, y) = \varphi^{-1}(T(\varphi(x), \varphi(y))), \quad \varphi(x) = x/e, & x, y \in [0, e], \\ S^*(x, y) = \psi^{-1}(S(\psi(x), \psi(y))), \quad \psi(x) = (x - e)/(1 - e), & x, y \in [e, 1]. \end{cases} \quad (2)$$

Lemma 1. (Fodor, Yager and Rybalov [6]) If U is increasing and has the neutral element $e \in (0, 1)$ then

$$\min \leq U \leq \max \text{ in } A(e) = [0, e] \times (e, 1] \cup (e, 1] \times [0, e). \quad (3)$$

Furthermore, if U is associative, then $U(0, 1), U(1, 0) \in \{0, 1\}$.

Theorem 2. (Li and Shi [10]) Let $e \in (0, 1)$. If T is an arbitrary triangular norm and S is an arbitrary triangular conorm then formula (1) with $U = \min$ or $U = \max$ in $A(e)$ gives uninorms.

Remark 1. Uninorms from Theorem 2 are not continuous in some points such that one of the variables is equal to the neutral element.

Example 1. (Fodor, Yager and Rybalov [6]) Formula

$$U(x, y) = \begin{cases} 0, & \text{if } x = 0 \text{ or } y = 0, \\ \frac{xy}{(1-x)(1-y)+xy}, & \text{if } x > 0 \text{ and } y > 0 \end{cases}$$

gives a uninorm with $e = \frac{1}{2}$, $T(x, y) = \frac{xy}{2-(x+y-xy)}$, $S(x, y) = \frac{x+y}{1+xy}$, $x, y \in [0, 1]$. This uninorm is continuous apart from the points $(0, 1)$ and $(1, 0)$.




Fig. 1. Frame structure of uninorm U with neutral element e .

Theorem 3. (Czogala and Drewniak [3]) If a uninorm is continuous then $e = 0$ or $e = 1$.

3. REMARK ABOUT THE ORDINAL SUM THEOREM

In this section we consider the ordinal sum and dual ordinal sum of semigroups. Next we present the characterization of continuous t -norms and t -conorms by using the ordinal sum theorem. Additional information about the ordinal sum of semigroups one may find in [1, 2, 5, 8, 9, 12].

Theorem 4. (Clifford [1], Climescu [2]) If (X, F) , (Y, G) are disjoint semigroups then $(X \cup Y, H)$ is a semigroup, where H is given by

$$H(x, y) = \begin{cases} F(x, y), & \text{if } x, y \in X, \\ G(x, y), & \text{if } x, y \in Y, \\ x, & \text{if } x \in X, y \in Y, \\ y, & \text{if } x \in Y, y \in X. \end{cases} \quad (4)$$

By duality we obtain

Theorem 5. (Drewniak and Drygaś [5]) If (X, F) , (Y, G) are disjoint semigroups, then $(X \cup Y, H)$ is a semigroup, where H is given by

$$H(x, y) = \begin{cases} F(x, y), & \text{if } x, y \in X, \\ G(x, y), & \text{if } x, y \in Y, \\ y, & \text{if } x \in X, y \in Y, \\ x, & \text{if } x \in Y, y \in X. \end{cases} \quad (5)$$




Fig. 2. Ordinal sum (left) and dual ordinal sum (right) of semigroups (X, F) and (Y, G) .

For our consideration it will be useful to remember the characterization of continuous t -norms or t -conorms by using ordinal sum theorems.

Theorem 6. (Klement, Mesiar and Pap [9], p.128, Sander [12]) Operation $T : [0, 1]^2 \rightarrow [0, 1]$ is continuous, associative, increasing, with the neutral element $e = 1$ iff there exists a family $\{(a_k, b_k)\}_{k \in A}$ (where $A \subset \mathbb{Q} \cap [0, 1]$) of nonempty, pairwise disjoint, open subintervals of $[0, 1]$ such that the operations $T_k = T|_{[a_k, b_k]^2}$ are continuous, increasing, associative with Archimedean property, neutral element b_k and T is given by

$$T(x, y) = \begin{cases} T_k(x, y), & \text{for } (x, y) \in (a_k, b_k]^2, \\ \min(x, y), & \text{otherwise.} \end{cases} \quad (6)$$

Moreover, the operation T is commutative.

Theorem 7. (Klement, Mesiar and Pap [9], p.130) Operation $S : [0, 1]^2 \rightarrow [0, 1]$ is continuous, associative, increasing, with the neutral element $e = 0$ iff there exists a family $\{(a_k, b_k)\}_{k \in A}$ (where $A \subset \mathbb{Q} \cap [0, 1]$) of nonempty, pairwise disjoint, open subintervals of $[0, 1]$ such that the operations $S_k = S|_{[a_k, b_k]^2}$ are continuous, increasing, associative with Archimedean property, neutral element a_k and S is given by

$$S(x, y) = \begin{cases} S_k(x, y), & \text{for } (x, y) \in [a_k, b_k)^2, \\ \max(x, y), & \text{otherwise.} \end{cases} \quad (7)$$

Moreover, the operation S is commutative.

4. MAIN RESULTS

In Theorems 6 and 7 a characterization of continuous operations in the class $\mathcal{U}(1)$ and $\mathcal{U}(0)$ respectively is given. Moreover, if operation in the class $\mathcal{U}(e)$ is continuous,

then $e = 0$ or $e = 1$ (see Theorem 3). Thus, we ask about the structure of operations in the class $\mathcal{U}(e)$ which are continuous in the open unit square for $e \in (0, 1)$.

Lemma 2. Let $e \in (0, 1)$. If operation $U \in \mathcal{U}(e)$ is continuous in $(0, 1)^2$ then operation $U|_{[0, e]^2}$ is isomorphic to a continuous t -norm and $U|_{[e, 1]^2}$ is isomorphic to a continuous t -conorm.

Proof. First we prove that operation $U|_{[e, 1]^2}$ is continuous. The operator U is continuous in $(0, 1)^2$. From this we obtain the continuity of the operation $U|_{[e, 1]^2}$ in $[e, 1]^2$. Moreover $U(x, y) \geq \max(x, y)$ for $x, y \in [e, 1]$ and $U(x, 1) = U(1, x) = 1$ for $x \in [e, 1]$. Let $x, y \in [e, 1]$, then $1 \geq U(x, y) \geq \max(x, y)$, $\lim_{x \rightarrow 1} \max(x, y) = 1$ and $\lim_{y \rightarrow 1} \max(x, y) = 1$. It means that $\lim_{x \rightarrow 1} U(x, y) = 1$ and $\lim_{y \rightarrow 1} U(x, y) = 1$, i. e. functions $U(x, t)$ and $U(t, y)$, $t \in [e, 1]$ are continuous for all $x, y \in [e, 1]$. This implies continuity of the operation $U|_{[e, 1]^2}$. It means, that $U|_{[e, 1]^2}$ is a continuous, associative, increasing operation with neutral element e , then it is isomorphic to a continuous t -conorm.

In similar way we obtain that the operation $U|_{[0, e]^2}$ is isomorphic to a continuous t -norm. \square

Lemma 3. Let $e \in (0, 1)$ and $U \in \mathcal{U}(e)$. If there exists $a \in [0, e)$ such that $U(x, y) = x$ for $x \in (a, e)$, $y \in (e, 1)$ or $U(x, y) = y$ for $x \in (e, 1)$, $y \in (a, e)$ then U is not continuous in $(0, 1)^2$.

Proof. Let $U(x, y) = x$ for $x \in (a, e)$, $y \in (e, 1)$. Take $s \in (e, 1)$ and let $f(t) = U(t, s)$, $t \in [0, 1]$. We have $f(t) = U(t, s) = t < e$ for $t \in (a, e)$ and $f(e) = s > e$. It means, that the function f is not continuous at the point e . This implies, that U is not continuous in $(0, 1)^2$.

In similar way as above we obtain the second part of Lemma. \square

In the next part of this paper we need the following lemmas

Lemma 4. (Klement, Mesiar and Pap [9]) Let $J = [a, b]$ and $F : J^2 \rightarrow J$ be associative, increasing operation with the neutral element b . If $x \in J$ is an idempotent element of operation F and functions $f(t) = F(x, t)$, $h(t) = F(t, x)$, $t \in J$ are continuous in J then $F(x, y) = F(y, x) = \min(x, y)$ for $y \in J$.

Lemma 5. Let $J = [a, b]$ and $F : J^2 \rightarrow J$ be associative, increasing operation with the neutral element a . If $x \in J$ is an idempotent element of operation F and functions $f(t) = F(x, t)$, $h(t) = F(t, x)$, $t \in J$ are continuous in J then $F(x, y) = F(y, x) = \max(x, y)$ for $y \in J$.

Lemma 6. Let $e \in (0, 1)$ and $U \in \mathcal{U}(e)$ be continuous in $(0, 1)^2$. If there exists $b \in (0, e)$ such that $U(b, y) = b$ for $y \in (b, e)$ or $U(x, b) = b$ for $x \in (b, e)$ then $U(x, y) = U(y, x) = \min(x, y)$ for $x \in [0, b]$ and $y \in [b, 1]$.




Fig. 3. The operation U from the Lemma 6.

Proof. Let $x \in [0, b]$ and $y \in (e, 1)$. For all $t \in (b, e)$ we have $U(b, t) = b$. By the continuity of the operation U we have $U(b, b) = b$. This means that b is an idempotent element of the continuous operation $U|_{[0, e]^2}$ and by Lemma 4 we have $U(b, t) = U(t, b) = \min(t, b)$ for $t \in [0, e]$. Hence, by monotonicity of U we have $U(s, t) = \min(s, t)$ for $s \in [0, b]$, $t \in [b, e]$.

Suppose that there exists $z \in (e, 1)$ such that $U(b, z) \geq e$. By continuity of the operation U and condition $U(b, e) = b$ there exists $w \in (e, z]$ such that $U(b, w) = e$. Then

$$b = U(b, e) = U(b, U(b, w)) = U(U(b, b), w) = U(b, w) = e,$$

which is a contradiction. Therefore $U(b, y) < e$ for all $y \in (e, 1)$. By continuity of the operation U and condition $U(e, y) = y$ there exists $v \in (b, e)$ such that $U(v, y) = e$. Therefore for all $x \leq b$ we have

$$U(x, y) = U(\min(x, v), y) = U(U(x, v), y) = U(x, U(v, y)) = U(x, e) = x.$$

By commutativity of the operation $U|_{[0, e]^2}$ we obtain $U(y, x) = x$ for $x \in [0, b]$ and $y \in [b, e]$. In similar way as above we obtain $U(y, x) = \min(x, y)$ for $x \in [0, b]$, $y \in [b, 1]$. If we assume that $U(x, b) = b$ for $x \in (b, e)$ then the proof is analogous. \square

By duality we obtain

Lemma 7. Let $e \in (0, 1)$ and $U \in \mathcal{U}(e)$ be continuous in $(0, 1)^2$. If there exists $a \in (e, 1)$, such that $U(a, y) = a$ for $y \in (e, a)$ or $U(x, a) = a$ for $x \in (e, a)$ then $U(x, y) = U(y, x) = \max(x, y)$ for $x \in [a, 1]$ and $y \in (0, a]$.

Lemma 8. (cf. Hu and Li [7]) Let $e \in (0, 1)$ and $U \in \mathcal{U}(e)$ be continuous in $(0, 1)^2$. Then there exist idempotent elements $a \in [0, e)$ and $b \in (e, 1]$ such that operations $U|_{(a, e]^2}$ and $U|_{[e, b)^2}$ are strictly increasing. Moreover $a = 0$ or $b = 1$.




Fig. 4. The operation $U \in \mathcal{U}(e)$ from Lemma 8.

Proof. By Lemma 2 operation $U|_{[0,e]^2}$ is isomorphic to a continuous t -norm. By Theorem 6 there exists a countably family of intervals $(a_k, b_k) \subset [0, e]$ such that $U|_{[0,e]^2}$ is an ordinal sum of semigroups $T_k = U|_{[a_k, b_k]^2}$ with Archimedean property or $T_k = \min$.

Suppose that there does not exist such $a \in [0, e)$ that $U|_{[a,e]^2}$ is a semigroup with Archimedean property. Then there exists $r \in [0, e)$ such that $U|_{[r,e]^2} = \min$ or for every neighborhood of the point e there exists k such that interval (a_k, b_k) is included in that neighborhood, i. e. there exists an increasing subsequence $\{b_{k_n}\}$ of sequence $\{b_k\}$ convergent to e . So, we construct the sequence of idempotent elements $\{c_n\}$, e. g. $c_n = e - \frac{1}{n + \lceil \frac{1}{e-r} \rceil} \in [r, e)$ in the first case, and $c_n = b_{k_n}$ in the second case. According to (6) we have $U(c_n, y) = c_n$ for all $y \in (c_n, e)$. By Lemma 6, $U(x, y) = x$ for $x \in [0, c_n]$ and $y \in (e, 1)$. It implies that $U(x, y) = x$ for $x \in [0, e) = \bigcup_{n=1}^{\infty} [0, c_n]$ and $y \in (e, 1)$. Now, by Lemma 3, operation U is not continuous in $(0, 1)^2$, which is a contradiction. So, there exists $a \in [0, e)$ such that $U|_{[a,e]^2}$ is isomorphic to a continuous Archimedean t -norm. Moreover a is an idempotent element of operation U and the zero element of operation $U|_{[a,e]^2}$.

Now we show that $U|_{(a,e]^2}$ is strictly increasing. Suppose that it is not. It means that $U|_{[a,e]^2}$ is isomorphic to the Łukasiewicz t -norm T_L . By continuity of U there exist $p \in (a, e)$ and $w \in (e, 1)$ such that $U(p, w) = e$. By the fact that $U|_{[a,e]^2}$ is isomorphic to T_L (all elements from (a, e) are zero divisors, where zero element is equal to a) it follows that $U(p, q) = U(q, p) = a$ for some $q \in (a, e)$ and by monotonicity of operation U and because $U(a, a) = a$ we have $U(t, p) = a$ for all $t \in [a, q]$. Therefore $U(t, U(p, w)) = U(t, e) = t$ and $U(U(t, p), w) = U(a, w)$. By associativity of U we have $U(a, w) = t$ for all $t \in [a, q]$, which leads to a contradiction. Thus $U|_{(a,e]^2}$ is strictly increasing.

In similar way we prove that there exists idempotent element $b \in (e, 1]$, which is the zero element of $U|_{[e,b]^2}$, such that $U|_{[e,b]^2}$ is strictly increasing.

Suppose that $a > 0$ and $b < 1$. Since $U(a, y) = a$ for all $y \in (a, e)$, Lemma 6 implies that $U(x, y) = \min(x, y)$ for $x \in [0, a]$ and $y \in (e, 1)$. Similarly, since b is the

zero element of $U|_{[e,b]^2}$, Lemma 7 implies that $U(x, y) = \max(x, y)$ for $x \in (0, e)$ and $y \in [b, 1]$. Therefore $U(x, y) = x$ and $U(x, y) = y$ for $x \in (0, a]$ and $y \in [b, 1]$, which is a contradiction.

Accordingly $a = 0$ or $b = 1$. \square

Lemma 9. Let $e \in (0, 1)$ and $U \in \mathcal{U}(e)$ be continuous in $(0, 1)^2$. If there exists $a \in [0, e)$ such that operations $U|_{(a,e]^2}$ and $U|_{[e,1]^2}$ are strictly increasing then the operation $U|_{(a,1)^2}$ is strictly increasing.

Proof. To show, that $U|_{(a,1)^2}$ is strictly increasing we must show that U is strictly increasing on the set $(a, e] \times [e, 1] \cup [e, 1] \times (a, e]$. By Lemma 2 operations $U|_{[0,e]^2}$ and $U|_{[e,1]^2}$ are commutative. Let $x, y \in (a, e]$, $x < y$ and $z \in [e, 1]$. Suppose that $U(x, z) = U(y, z)$. Then $z > e$ because $U(x, e) = x < y = U(y, e)$.

If $U(x, z) = U(y, z) < e$ then by continuity of U and inequality $U(e, z) = z > e$ there exists $s \in (x, e)$ such that $U(s, z) = e$. Then

$$\begin{aligned} x &= U(x, e) = U(x, U(s, z)) = U(U(x, s), z) = U(U(s, x), z) = U(s, U(x, z)) \\ &= U(s, U(y, z)) = U(U(s, y), z) = U(U(y, s), z) = U(y, U(s, z)) = U(y, e) = y, \end{aligned}$$

which is a contradiction.

If $U(x, z) = U(y, z) \geq e$ then, by continuity of U and condition $U(x, e) = x$, $x < y \leq e$, there exists $c \in (e, z]$ such that $U(x, c) = y$. From $U(y, e) = y \leq e \leq U(y, z)$, there exists $d \in [e, z]$ such that $U(y, d) = e$. Thus $U(e, z) = z$ and

$$\begin{aligned} z &= U(e, z) = U(U(y, d), z) = U(y, U(d, z)) = U(y, U(z, d)) \\ &= U(U(x, c), U(z, d)) = U(x, U(c, U(z, d))) = U(x, U(U(c, z), d)) \\ &= U(x, U(U(z, c), d)) = U(x, U(z, U(c, d))) = U(x, U(z, U(d, c))) \\ &= U(U(x, z), U(d, c)) = U(U(y, z), U(d, c)) = U(y, U(z, U(d, c))) \\ &= U(y, U(U(z, d), c)) = U(y, U(U(d, z), c)) = U(y, U(d, U(z, c))) \\ &= U(U(y, d), U(z, c)) = U(e, U(z, c)) = U(z, c). \end{aligned}$$

Moreover operation $U|_{[e,1]^2}$ is strictly increasing and $z, c \in (e, 1)$. This leads to a contradiction. Therefore U is strictly increasing with respect to the first variable in the $(a, e] \times [e, 1]$.

Now let $x, y \in [e, 1]$, $x < y$ and $z \in (a, e]$. Suppose that $U(z, x) = U(z, y)$. Then $z < e$ because $U(e, x) = x < y = U(e, y)$.

If $U(z, x) = U(z, y) > e$ then, by continuity of U and inequality $U(z, e) = z < e$, there exists $s \in (e, x)$ such that $U(z, s) = e$. Therefore

$$\begin{aligned} x &= U(e, x) = U(U(z, s), x) = U(z, U(s, x)) = U(z, U(x, s)) = U(U(z, x), s) \\ &= U(U(z, y), s) = U(z, U(y, s)) = U(z, U(s, y)) = U(U(z, s), y) = U(e, y) = y, \end{aligned}$$

which is a contradiction.

If $U(z, x) = U(z, y) \leq e$ then, by continuity of U and condition $U(e, y) = y$, $e \leq x < y$, there exists $c \in (z, e)$ such that $U(c, y) = x$. From $U(e, x) = x > e \geq U(z, x)$ there exists $d \in [z, e]$ such that $U(d, x) = e$. Therefore

$$\begin{aligned}
 z &= U(z, e) = U(z, U(d, x)) = U(U(z, d), x) = U(U(d, z), x) \\
 &= U(U(d, z), U(c, y)) = U(d, U(z, U(c, y))) = U(d, U(U(z, c), y)) \\
 &= U(d, U(U(c, z), y)) = U(d, U(c, U(z, y))) = U(U(d, c), U(z, y)) \\
 &= U(U(c, d), U(z, x)) = U(U(U(c, d), z), x) = U(U(c, U(d, z)), x) \\
 &= U(U(c, U(z, d)), x) = U(U(U(c, z), d), x) = U(U(c, z), U(d, x)) \\
 &= U(U(c, z), e) = U(c, z).
 \end{aligned}$$

Moreover, operation $U|_{(a,e]^2}$ is strictly increasing and $z, c \in (a, e)$. This leads to a contradiction. Thus U is strictly increasing with respect to second variable on $(a, e] \times [e, 1)$.

In a similar way we prove that U is strictly increasing on $[e, 1) \times (a, e]$. \square

Theorem 8. Let $e \in (0, 1)$ and $U \in \mathcal{U}(e)$ be continuous in $(0, 1)^2$. If there exists an idempotent element $a \in [0, e)$ of U such that operations $U|_{(a,e]^2}$ and $U|_{[e,1)^2}$ are strictly increasing, then operation $U|_{[0,1)^2}$ is an ordinal sum of continuous semigroup $U|_{[0,a]^2}$ with the neutral element a and continuous group $U|_{(a,1)^2}$ with Archimedean property and the neutral element e .

Proof. By Lemma 2, the operation $U|_{[0,e]^2}$ is isomorphic to a continuous t -norm and, since a is an idempotent element of this operation, $U|_{[0,a]^2}$ is also isomorphic to a continuous t -norm. By Lemma 9, operation $U|_{(a,1)^2}$ is strictly increasing and therefore it is isomorphic to the real numbers with addition. Now, taking into account Lemma 6 we have that $U|_{[0,1)^2}$ is an ordinal sum of the semigroup $U|_{[0,a]^2}$ and the group $U|_{(a,1)^2}$. \square

Similarly, we obtain the following results:

Lemma 10. Let $e \in (0, 1)$ and $U \in \mathcal{U}(e)$ be continuous in $(0, 1)^2$. If there exists $b \in (e, 1]$ such that operations $U|_{(0,e]^2}$ and $U|_{[e,b)^2}$ are strictly increasing then the operation $U|_{(0,b)^2}$ is strictly increasing.

Theorem 9. Let $e \in (0, 1)$ and $U \in \mathcal{U}(e)$ be continuous in $(0, 1)^2$. If there exists an idempotent element $b \in (e, 1]$ of U such that operations $U|_{(0,e]^2}$ and $U|_{[e,b)^2}$ are strictly increasing then operation $U|_{(0,1)^2}$ is a dual ordinal sum of continuous group $U|_{(0,b)^2}$ with Archimedean property and the neutral element e and continuous semigroup $U|_{[b,1]^2}$ with the neutral element b .

So, we have the characterization of this operation in the open unit square. Now we ask about it's structure on the boundary.

Lemma 11. Let $e \in (0, 1)$ and $U \in \mathcal{U}(e)$ be continuous in $(0, 1)^2$. If there exists an idempotent element $a \in [0, e]$ of U such that operations $U|_{(a,e]^2}$ and $U|_{[e,1]^2}$ are strictly increasing then there exist idempotent elements $c, d \in [0, a]$ of operation U such that

$$U(x, 1) = \begin{cases} x, & \text{if } x \in [0, c), \\ 1, & \text{if } x \in (c, 1], \\ x \text{ or } 1, & \text{if } x = c, \end{cases} \quad (8)$$

$$U(1, x) = \begin{cases} x, & \text{if } x \in [0, d), \\ 1, & \text{if } x \in (d, 1], \\ x \text{ or } 1, & \text{if } x = d. \end{cases} \quad (9)$$

Moreover $c = d$.

Proof. By the Lemma 1, $U(0, 1) = 0$ or $U(0, 1) = 1$. If $U(0, 1) = 1$ then by monotonicity of U we have $U(x, 1) = 1$ for $x \in [0, 1]$. Therefore we obtain (8) for $c = 0$. Moreover 0 is an idempotent element of the operation U .

If $U(0, 1) = 0$ then by Theorem 9 the semigroup $U|_{(a,1)^2}$ is isomorphic to the real numbers with addition. Thus we have $\lim_{y \rightarrow 1} U(x, y) = 1$ for $x \in (a, 1)$ and by monotonicity of the operation U we obtain $U(x, 1) = 1$ for $x \in (a, 1]$. Let $x \in (0, a]$. First we will prove that $U(x, 1) = x$ or $U(x, 1) = 1$. Suppose that there exists $z \in (0, a]$ such that $z < U(z, 1) < 1$ and let $w = U(z, 1)$.

If $w \in (a, 1)$ then for $y \in (e, 1)$, by associativity of U and strictly monotonicity of $U|_{(a,1)^2}$, we obtain

$$\begin{aligned} w &= U(z, 1) = U(z, U(y, 1)) = U(z, U(1, y)) \\ &= U(U(z, 1), y) = U(w, y) > U(w, e) = w, \end{aligned}$$

which is a contradiction.

If $w \in (z, a]$ then by the conditions $U(0, w) = 0$, $U(e, w) = w$ and continuity of $U|_{[0,e]^2}$ there exists $v \in (0, e)$ such that $U(v, w) = z$ and by associativity of U , we obtain

$$\begin{aligned} w &= U(z, 1) = U(U(v, w), 1) = U(U(v, U(z, 1)), 1) \\ &= U(U(v, z), U(1, 1)) = U(U(v, z), 1) = U(v, U(z, 1)) = U(v, w) = z, \end{aligned}$$

which is a contradiction. Therefore $U(x, 1) = x$ or $U(x, 1) = 1$ for $x \in [0, 1]$.

Thus, for $c = \inf\{x \in [0, a] : U(x, 1) = 1\}$ we obtain (8), moreover $c \in [0, a]$.

Let $x \in (0, c)$, $y \in (c, e]$ then we have

$$\begin{aligned} U(x, y) &= U(y, x) = U(y, U(x, 1)) = U(U(y, x), 1) \\ &= (U(x, y), 1) = U(x, U(y, 1)) = U(x, 1) = x = \min(x, y). \end{aligned}$$

By monotonicity of U and inequality $U|_{[0,e]^2} \leq \min$ we obtain $U(c, y) = c$ for $y \in (c, e)$. By above and continuity of U we have $U(c, c) = c$, i.e. c is an idempotent element of operation U . Similarly we prove (9).

To prove that $c = d$ suppose that $d < c$. Then there exists $y \in (d, c)$ such that $U(1, y) = 1$ and $U(y, 1) = y$. Taking $z \in (d, y)$ we have $U(1, z) = 1$ and

$$y = U(y, 1) = U(y, U(1, z)) = U(U(y, 1), z) = U(y, z) \leq U(e, z) = z < y,$$

which is a contradiction, thus $d \geq c$.

If we suppose that $d > c$ then there exists $y \in (c, d)$ such that $U(1, y) = y$ and $U(y, 1) = 1$. Taking $z \in (y, d)$ we have

$$z = U(1, z) = U(U(y, 1), z) = U(y, U(1, z)) = U(y, z) \leq U(y, e) = y < z,$$

which is a contradiction. Thus $c = d$. \square

Lemma 12. Let $e \in (0, 1)$ and $U \in \mathcal{U}(e)$ be continuous in $(0, 1)^2$. If there exists an idempotent element $b \in (e, 1]$ of U such that operations $U|_{(0, e]^2}$ and $U|_{[e, b]^2}$ are strictly increasing then there exist idempotent elements $p, q \in [b, 1]$ of operation U such that

$$U(x, 0) = \begin{cases} 0, & \text{if } x \in [0, p), \\ x, & \text{if } x \in (p, 1], \\ 0 \text{ or } x, & \text{if } x = p, \end{cases} \quad (10)$$

$$U(0, x) = \begin{cases} 0, & \text{if } x \in [0, q), \\ x, & \text{if } x \in (q, 1], \\ 0 \text{ or } x, & \text{if } x = q. \end{cases} \quad (11)$$

Moreover $p = q$.




Fig. 5. Operation $U \in \mathcal{U}(e)$ continuous in the open unit square with $a > 0$.

As a results of our considerations we obtain

Theorem 10. Let $e \in (0, 1)$ and $U \in \mathcal{U}(e)$ be continuous in $(0, 1)^2$. Then one of the following two cases holds:

- (i) There exist idempotent elements $a \in [0, e)$ and $c \in [0, a]$ of operation U such that $U|_{[0,1]^2}$ is an ordinal sum of continuous semigroup $U|_{[0,a]^2}$ with the neutral element a and continuous group $U|_{(a,1)^2}$ with Archimedean property and the neutral element e and conditions (8) and (9) hold.
- (ii) There exist idempotent elements $b \in (e, 1]$ and $p \in [b, 1]$ of operation U , such that $U|_{(0,1]^2}$ is a dual ordinal sum of continuous semigroup $U|_{[b,1]^2}$ with the neutral element b and continuous group $U|_{(0,b)^2}$ with Archimedean property and the neutral element e and conditions (10) and (11) hold.




Fig. 6. Operation $U \in \mathcal{U}(e)$ continuous in the open unit square with $b < 1$.

Proof. By Lemma 8 there exist $a \in [0, e)$ and $b \in (e, 1]$ ($a = 0$ or $b = 1$) such that $U|_{(a,b)^2}$ is strictly increasing (Lemma 9 and 10).

If $b = 1$ then by Theorem 8 and Lemma 11 we obtain (i).

If $a = 0$ then by Theorem 9 and Lemma 9 we obtain (ii). □

Remark 2. Operation U in the previous theorem is commutative in the set

(i) $[0, 1]^2 \setminus \{(c, 1), (1, c)\},$

(ii) $[0, 1]^2 \setminus \{(0, p), (p, 0)\}.$

5. CONCLUSION

By the above consideration we obtain the following results known from the papers [6] and [7]

Theorem 11. (Hu and Li [7], Theorem 4.5) Let $e \in (0, 1)$ and U be a uninorm which is continuous in $(0, 1)^2$. Then U can be represented as follows:

$$(i) \quad U(x, y) = \begin{cases} eT(\frac{x}{e}, \frac{y}{e}), & \text{if } x, y \in [0, a], \\ h^{-1}(h(x) + h(y)), & \text{if } x, y \in (a, 1), \\ x, & \text{if } x \in [0, a], y \in (a, 1) \text{ or } x \in [0, c], y = 1, \\ y, & \text{if } x \in (a, 1), y \in [0, a] \text{ or } x = 1, y \in [0, c], \\ 1, & \text{if } x \in (c, 1], y = 1 \text{ or } x = 1, y \in (c, 1], \\ x \text{ or } y, & \text{if } x = c, y = 1 \text{ or } x = 1, y = c, \end{cases}$$

where $a \in [0, e)$, $c \in [0, a]$, $U(c, c) = c$, function $h : [a, 1] \rightarrow [-\infty, +\infty]$ is strict and $h(a) = -\infty$, $h(e) = 0$, $h(1) = +\infty$;

$$(ii) \quad U(x, y) = \begin{cases} e + (1 - e)S(\frac{x-e}{1-e}, \frac{y-e}{1-e}), & \text{if } x, y \in [b, 1], \\ h^{-1}(h(x) + h(y)), & \text{if } x, y \in (0, b), \\ y, & \text{if } x \in (0, b), y \in [b, 1] \text{ or } x = 0, y \in (p, 1], \\ x, & \text{if } x \in [b, 1], y \in (0, b) \text{ or } x \in (p, 1], y = 0, \\ 0, & \text{if } x = 0, y \in [0, p) \text{ or } x \in [0, p), y = 0, \\ x \text{ or } y, & \text{if } x = p, y = 0, \text{ or } x = 0, y = p, \end{cases}$$

where $b \in (e, 1]$, $p \in [b, 1]$, $U(p, p) = p$, function $h : [0, b] \rightarrow [-\infty, +\infty]$ is strict and $h(0) = -\infty$, $h(e) = 0$, $h(b) = +\infty$.

Theorem 12. (Fodor, Yager and Rybalkov [6]) Let $e \in (0, 1)$ and U be a uninorm continuous without the points $(0, 1)$ and $(1, 0)$. Then operations $U|_{(0,e]^2}$ and $U|_{[e,1]^2}$ are strictly increasing and

$$U(x, y) = \begin{cases} h^{-1}(h(x) + h(y)), & \text{for } (x, y) \in [0, 1]^2 \setminus \{(0, 1), (1, 0)\}, \\ 0 \text{ or } 1, & \text{elsewhere,} \end{cases} \quad (12)$$

where $h : [0, 1] \rightarrow [-\infty, +\infty]$ is an increasing bijection such that $h(e) = 0$.

Proof. Operation $U|_{(0,1)^2}$ is continuous. Suppose that in Theorem 10 the condition (i) holds, i. e. there exists $a \in [0, e)$, such that operation $U|_{(a,1)^2}$ is strictly increasing. By Lemma 11 there exists $c \in [0, a]$ such that (8) holds.

Suppose that $c < a$, then for $x \in (c, a)$ and $y \in (e, 1)$ we have $U(x, y) = \min(x, y) = x$ and $U(x, 1) = 1$. It means that U is not continuous at the points $(x, 1)$, $x \in (c, a)$. Therefore $c = a$.

Suppose now, that $a > 0$. By Lemma 11 we have $U(x, 1) = x$ for $x \in [0, a]$ and $U(x, 1) = 1$ for $x \in (a, 1]$. It means that the point $(a, 1)$ is a point of discontinuity of the operation U , which leads to a contradiction. Thus $a = 0$. Now, directly by the above theorem, we obtain (12). \square

(Received April 17, 2006.)

REFERENCES

-
- [1] A. H. Clifford: Naturally totally ordered commutative semigroups. *Amer. J. Math.* **76** (1954), 631–646.
 - [2] A. C. Climescu: Sur l'équation fonctionnelle de l'associativité. *Bull. Ecole Polytechn.* **1** (1946), 1–16.
 - [3] E. Czogała and J. Drewniak: Associative monotonic operations in fuzzy set theory. *Fuzzy Sets and Systems* **12** (1984), 249–269.
 - [4] J. Dombi: Basic concepts for a theory of evaluation: The aggregative operators. *European J. Oper. Res.* **10** (1982), 282–293.
 - [5] J. Drewniak and P. Drygaś: Ordered semigroups in constructions of uninorms and nullnorms. In: *Issues in Soft Computing Theory and Applications* (P. Grzegorzewski, M. Krawczak, and S. Zadrozny, eds.), EXIT, Warszawa 2005, pp. 147–158.
 - [6] J. Fodor, R. Yager, and A. Rybalov: Structure of uninorms. *Internat. J. Uncertain. Fuzziness Knowledge-Based Systems* **5** (1997), 411–427.
 - [7] S.-K. Hu and Z.-F. Li: The structure of continuous uninorms. *Fuzzy Sets and Systems* **124** (2001), 43–52.
 - [8] S. Jenei: A note on the ordinal sum theorem and its consequence for the construction of triangular norm. *Fuzzy Sets and Systems* **126** (2002), 199–205.
 - [9] E. P. Klement, R. Mesiar, and E. Pap: *Triangular Norms*. Kluwer Academic Publishers, Dordrecht 2000.
 - [10] Y.-M. Li and Z.-K. Shi: Remarks on uninorm aggregation operators. *Fuzzy Sets and Systems* **114** (2000), 377–380.
 - [11] M. Mas, M. Monserrat, and J. Torrens: On left and right uninorms. *Internat. J. Uncertain. Fuzziness Knowledge-Based Systems* **9** (2001), 491–507.
 - [12] W. Sander: Associative aggregation operators. In: *Aggregation Operators* (T. Calvo, G. Mayor, and R. Mesiar, eds), Physica-Verlag, Heidelberg 2002, pp. 124–158.
 - [13] R. Yager and A. Rybalov: Uninorm aggregation operators. *Fuzzy Sets and Systems* **80** (1996), 111–120.

Paweł Drygaś, Institute of Mathematics, University of Rzeszów, ul. Rejtana 16a, 35-310 Rzeszów. Poland.

e-mail: paweldr@univ.rzeszow.pl