

Kybernetika

VOLUME 43 (2007), NUMBER 1

The Journal of the Czech Society for
Cybernetics and Information Sciences

Published by:

Institute of Information Theory
and Automation of the AS CR, v.v.i.

Editor-in-Chief:

Milan Mareš

Managing Editors:

Karel Sladký
Lucie Fajfrová

Editorial Board:

Jiří Anděl, Sergej Čelikovský, Marie
Demlová, Petr Hájek, Jan Flusser, Martin
Janžura, Jan Ježek, George Klir, Ivan
Kramosil, Tomáš Kroupa, Friedrich Liese,
Jean-Jacques Loiseau, František Matúš,
Radko Mesiar, Jiří Outrata, Jan Štecha,
Olga Štěpánková, Igor Vajda, Jiřina
Vejnarová, Miloslav Vošvrda, Pavel Zítek

Editorial Office:

Pod Vodárenskou věží 4, 182 08 Praha 8

Kybernetika is a bi-monthly international journal dedicated for rapid publication of high-quality, peer-reviewed research articles in fields covered by its title.

Kybernetika traditionally publishes research results in the fields of Control Sciences, Information Sciences, System Sciences, Statistical Decision Making, Applied Probability Theory, Random Processes, Fuzziness and Uncertainty Theories, Operations Research and Theoretical Computer Science, as well as in the topics closely related to the above fields.

The Journal has been monitored in the Science Citation Index since 1977 and it is abstracted/indexed in databases of Mathematical Reviews, Current Mathematical Publications, Current Contents ISI Engineering and Computing Technology.

Kybernetika. Volume 43 (2007)

ISSN 0023-5954, MK ČR E 4902.

Published bimonthly by the Institute of Information Theory and Automation of the Academy of Sciences of the Czech Republic, Pod Vodárenskou věží 4, 182 08 Praha 8. — Address of the Editor: P. O. Box 18, 182 08 Prague 8, e-mail: kybernetika@utia.cas.cz. — Printed by PV Press, Pod vrstevnicí 5, 140 00 Prague 4. — Orders and subscriptions should be placed with: MYRIS TRADE Ltd., P. O. Box 2, V Štíhlách 1311, 142 01 Prague 4, Czech Republic, e-mail: myris@myris.cz. — Sole agent for all “western” countries: Kubon & Sagner, P. O. Box 34 01 08, D-8 000 München 34, F.R.G.

Published in March 2007.

© Institute of Information Theory and Automation of the AS CR, v.v.i., Prague 2007.

WEAK LAW OF LARGE NUMBERS FOR I.I.D. FUZZY RANDOM VARIABLES

DUG HUN HONG AND KYUNG TAE KIM

In this paper, weak laws of large numbers for sum of independent and identically distributed fuzzy random variables are obtained.

Keywords: fuzzy number, fuzzy random variable, weak law of large numbers

AMS Subject Classification: 60F05, 60B12

1. INTRODUCTION

The celebrated Feller weak law of large numbers provided a necessary and sufficient condition in the i.i.d. case in [1] as follows:

Theorem 1.1. (Feller) Let $\{X_n\}$ be i.i.d. random variables and $S_n = \sum_{i=1}^n X_i$. Suppose that $EX_1 I_{[|X_1| \leq n]} \rightarrow u$ as $n \rightarrow \infty$. Then $S_n/n \rightarrow u$ in probability as $n \rightarrow \infty$ if and only if $n \Pr\{|X_1| > n\} \rightarrow 0$ as $n \rightarrow \infty$.

Let $a > 0$. A positive measurable function f on $[a, \infty)$ varies regularly at infinity with exponent ρ , $\infty < \rho < \infty$, denote $f \in RV(\rho)$, if and only if

$$\frac{f(tx)}{f(t)} \rightarrow x^\rho \text{ as } t \rightarrow \infty \text{ for all } x > 0.$$

If $\rho = 0$ the function is slowly varying at infinity; $f \in SV$.

Recently, Gut [3] provided the following more general Feller law concerning regularly and slowly varying functions.

Theorem 1.2. (Gut) Let $\{X_n\}$ be i.i.d. random variables and $S_n = \sum_{i=1}^n X_i$. Further, let, for $x > 0$, $b \in RV(1/\rho)$, for some $\rho \in (0, 1]$, that is, let $b(x) = x^{1/\rho} l(x)$, where $l \in SV$. Set $b_n = b(n)$. Then $(1/b_n)(S_n - nEX_1 I_{[|X_1| \leq b_n]}) \rightarrow 0$ in probability as $n \rightarrow \infty$ if and only if $n \Pr\{|X_1| > b_n\} \rightarrow 0$ as $n \rightarrow \infty$.

The author should simply think c (constant), $\log x$ or $\log \log x$ each time the function $l(x)$ appears.

The concept of a fuzzy random variable was introduced as a natural generalization of random sets in order to represent relationships between the outcomes of a random experiment and inexact data. By inexactness here we mean non-statistical inexactness due to the subjectivity and imprecision of human knowledge. Limit theorems for sums of independent fuzzy random variables have received much attentions because of its usefulness in several applied fields. This paper concerns with weak law of large numbers which is the one of limit theorems. Many authors have studied the strong laws of large numbers for sums of independent fuzzy random variables. For example, see Hong and Kim [4], Hong [5], Inoue [6], Joo and Kim [8], Kim [10], Klement et al. [11], Kruse [12], Miyakoshi and Shimbo [13], Uemura [16] and so on [7, 14]. On the other hand, weak laws of large numbers for fuzzy random variables have been studied only by Joo [9] and Taylor et al. [15]. The purpose of this paper is to obtain Feller, Gut [3] weak law of large numbers for sums of independent and identically distributed (i.i.d.) fuzzy random variables.

2. PRELIMINARIES

In this section, we describe some basic concepts of fuzzy numbers. Let R denote the real line. A fuzzy number is a fuzzy set $\tilde{u} : R \rightarrow [0, 1]$ with the following properties;

1. \tilde{u} is normal, i. e., there exists $x \in R$ such that $\tilde{u}(x) = 1$.
2. \tilde{u} is upper semicontinuous.
3. $\text{supp } \tilde{u} = \text{cl}\{x \in R | \tilde{u}(x) > 0\}$ is compact.
4. \tilde{u} is a convex fuzzy set, i. e., $\tilde{u}(\lambda x + (1 - \lambda)y) \geq \min(\tilde{u}(x), \tilde{u}(y))$ for $x, y \in R$ and $\lambda \in [0, 1]$.

We denote the family of all fuzzy numbers by $F(R)$. For a fuzzy set \tilde{u} , the α -level set of \tilde{u} is defined by

$$L_\alpha \tilde{u} = \begin{cases} \{x | \tilde{u}(x) \geq \alpha\}, & 0 < \alpha \leq 1, \\ \text{supp } \tilde{u}, & \alpha = 0. \end{cases} \quad (1)$$

Then it follows that \tilde{u} is a fuzzy number if and only if $L_1 \tilde{u} \neq \emptyset$ and $L_\alpha \tilde{u}$ is a closed bounded interval for each $\alpha \in [0, 1]$. From this characterization of fuzzy numbers, a fuzzy number, \tilde{u} is completely determined by the endpoints of the intervals $L_\alpha \tilde{u} = [u_\alpha^1, u_\alpha^2]$.

Theorem 2.1. (Goetschel and Voxman [2]) For $\tilde{u} \in F(R)$, denote $u^1(\alpha) = u_\alpha^1$ and $u^2(\alpha) = u_\alpha^2$ by considering them as functions of $\alpha \in [0, 1]$. Then the following hold:

1. u^1 is a bounded non-decreasing function on $[0, 1]$.
2. u^2 is a bounded non-increasing function on $[0, 1]$.
3. $u^1(1) \leq u^2(1)$.
4. u^1 and u^2 are left continuous on $[0, 1]$ and right continuous at 0.
5. If v^1 and v^2 satisfy above (1)–(4), then there exists a unique $\tilde{v} \in F(R)$ such that $v_\alpha^1 = v^1(\alpha)$, $v_\alpha^2 = v^2(\alpha)$, for all $\alpha \in [0, 1]$.

The above theorem implies that we can identify a fuzzy number \tilde{u} with the parameterized representation $\{(u_\alpha^1, u_\alpha^2) \mid 0 \leq \alpha \leq 1\}$, where u^1 and u^2 satisfy (1)–(4) of Theorem 2.1. Suppose now that $\tilde{u}, \tilde{v} \in F(R)$ are fuzzy numbers whose representations are $\tilde{u} = \{(u_\alpha^1, u_\alpha^2) \mid 0 \leq \alpha \leq 1\}$ and $\tilde{v} = \{(v_\alpha^1, v_\alpha^2) \mid 0 \leq \alpha \leq 1\}$, respectively. If we define

$$\begin{aligned} (\tilde{u} + \tilde{v})(z) &= \sup_{x+y=z} \min(\tilde{u}(x), \tilde{v}(y)), \\ (\lambda\tilde{u})(z) &= \begin{cases} \tilde{u}(z/\lambda), & \lambda \neq 0, \\ \tilde{0}, & \lambda = 0, \end{cases} \end{aligned}$$

where $\tilde{0} = I_{\{0\}}$ is the indicator function of $\{0\}$, then

$$\begin{aligned} \tilde{u} + \tilde{v} &= \{(u_\alpha^1 + v_\alpha^1, u_\alpha^2 + v_\alpha^2) \mid 0 \leq \alpha \leq 1\} \\ \lambda\tilde{u} &= \{(\lambda u_\alpha^1, \lambda u_\alpha^2) \mid 0 \leq \alpha \leq 1\} \text{ for } \lambda > 0. \end{aligned}$$

Now, we define the metric d_∞ on $F(R)$ by

$$d_\infty(\tilde{u}, \tilde{v}) = \sup_{0 \leq \alpha \leq 1} d_H(L_\alpha \tilde{u}, L_\alpha \tilde{v}),$$

where d_H denotes Hausdorff metric on closed subsets of R which admits in our particular case the form

$$d_H(L_\alpha \tilde{u}, L_\alpha \tilde{v}) = \max(|u_\alpha^1 - v_\alpha^1|, |u_\alpha^2 - v_\alpha^2|).$$

Also, the norm $\|\tilde{u}\|$ of fuzzy number \tilde{u} will be defined as

$$\|\tilde{u}\| = d_\infty(\tilde{u}, \tilde{0}) = \max(|u_0^1|, |u_0^2|).$$

Let (Ω, \mathcal{A}, P) denotes a complete probability space. For a fuzzy number valued function $\tilde{X} : \Omega \rightarrow F(R)$ and a subset B of R , $\tilde{X}^{-1}(B)$ denotes the fuzzy subset of Ω defined by

$$\tilde{X}^{-1}(B)(\omega) = \sup_{x \in B} \tilde{X}(\omega)(x)$$

for every $\omega \in \Omega$. The function $\tilde{X} : \Omega \rightarrow F(R)$ is called a fuzzy random variables if for every closed subset B of R , the fuzzy set $\tilde{X}^{-1}(B)$ is measurable when considered as a function from Ω to $[0, 1]$. If we denote $\tilde{X}(\omega) = \{(X_\alpha^1(\omega), X_\alpha^2(\omega)) \mid 0 \leq \alpha \leq 1\}$, then it is well known that \tilde{X} is a fuzzy random variable if and only if for each $\alpha \in [0, 1]$, X_α^1 and X_α^2 are random variables in the usual sense.

A fuzzy random variable $\tilde{X}(\omega) = \{(X_\alpha^1(\omega), X_\alpha^2(\omega)) \mid 0 \leq \alpha \leq 1\}$, is called integrable if for each $\alpha \in [0, 1]$, X_α^1 and X_α^2 are integrable, equivalently, $\int \|\tilde{X}\| dP < \infty$. In this case, the expectation of \tilde{X} is the fuzzy number $E\tilde{X}$ defined by

$$E\tilde{X} = \int \tilde{X} dP = \left\{ \left(\int X_\alpha^1 dP, \int X_\alpha^2 dP \right) \mid 0 \leq \alpha \leq 1 \right\}.$$

Let $\{\tilde{X}_n\}$ be a sequence of fuzzy random variables. For $\alpha \in [0, 1]$ $\{X_{n\alpha}^1\}$, $\{X_{n\alpha}^2\}$ are sequences of independent and identically distributed random variables. Then $\{\tilde{X}_n\}$ is called a sequence of independent and identically distributed fuzzy random variables.

3. MAIN RESULTS

We define $\tilde{X}I_{[\|\tilde{X}\|\leq t]}$ as

$$\tilde{X}I_{[\|\tilde{X}\|\leq t]}(\omega) = \begin{cases} \tilde{X}(\omega) & \text{if } \|\tilde{X}(\omega)\| \leq t, \\ I_{\{0\}} & \text{otherwise.} \end{cases}$$

Theorem 3.1. Let $\{\tilde{X}_n\}$ be a sequence of independent and identically distributed fuzzy random variables and $\tilde{S}_n = \sum_{i=1}^n \tilde{X}_i$. Suppose that for a fuzzy number $\tilde{u} \in F(R)$, $d_\infty(\mathbb{E}\tilde{X}_1 I_{[\|\tilde{X}_1\|\leq n]}, \tilde{u}) \rightarrow 0$, then

$$d_\infty\left(\frac{\tilde{S}_n}{n}, \tilde{u}\right) \rightarrow 0 \text{ in probability} \quad (2)$$

iff

$$nP\{\|\tilde{X}_1\| > n\} \rightarrow 0. \quad (3)$$

To prove our main result, we need some lemmas.

Lemma 3.2. (Kim [10]) For each $\tilde{u} \in F(R)$ and each $\varepsilon > 0$, there exists a partition $0 = \alpha_0 < \alpha_1 < \dots < \alpha_r = 1$ of $[0, 1]$ such that

$$\max(u_{\alpha_k}^1 - u_{\alpha_{k-1}^+}^1, u_{\alpha_{k-1}^+}^2 - u_{\alpha_k}^2) < \varepsilon, \quad k = 1, 2, \dots, r,$$

where $u_{\alpha^+}^i$ denotes the right-hand limit of u^i at α , $i = 1, 2$.

Lemma 3.3. (Hong [5]) For a sequence $\{\tilde{u}_n\} \in F(R)$ and a continuous $\tilde{u} \in F(R)$, suppose that for each $\alpha \in [0, 1]$, $d_H(L_\alpha \tilde{u}_n, L_\alpha \tilde{u}) \rightarrow 0$. Then we have $d_\infty(\tilde{u}_n, \tilde{u}) \rightarrow 0$.

Lemma 3.4. Let $\{\tilde{X}_n\}$ be a sequence of independent and identically distributed fuzzy random variables. Let $\tilde{X}'_{jn} = \tilde{X}_j I_{[\|\tilde{X}_j\|\leq n]}$ and let $S'_n = \sum_{j=1}^n \tilde{X}'_{jn}$. If $nP\{\|\tilde{X}_1\| > n\} \rightarrow 0$, then for each $\alpha \in [0, 1]$, $(1/n)((S'_n)_\alpha^1 - (\mathbb{E}S'_n)_\alpha^1) \rightarrow 0$ in probability.

Proof. By the assumption, for $\alpha \in [0, 1]$,

$$\begin{aligned}
\mathbb{E} \left(\left(\frac{\tilde{S}'_n}{n} \right)_\alpha^1 - \left(\frac{\mathbb{E}\tilde{S}'_n}{n} \right)_\alpha^1 \right)^2 &= \frac{1}{n^2} \sum_{j=1}^n \mathbb{E}((X'_{jn})_\alpha^1 - \mathbb{E}(X'_{jn})_\alpha^1)^2 \\
&\leq \frac{1}{n^2} \sum_{j=1}^n \mathbb{E}((X'_{jn})_\alpha^1)^2 \\
&= \frac{1}{n} \mathbb{E}((X'_{1n})_\alpha^1)^2 \\
&= \frac{1}{n} \sum_{j=1}^n \int_{[j-1 < \|\tilde{X}_1\| \leq j]} ((X'_{1n})_\alpha^1)^2 \\
&\leq \frac{1}{n} \sum_{j=1}^n \int_{[j-1 < \|\tilde{X}_1\| \leq j]} \|\tilde{X}_1\|^2 \\
&\leq \frac{1}{n} \sum_{j=1}^n j^2 [P\{\|\tilde{X}_1\| > j-1\} - P\{\|\tilde{X}_1\| > j\}] \\
&= \frac{1}{n} [P\{\|\tilde{X}_1\| > 0\} - n^2 P\{\|\tilde{X}_1\| > n\}] \\
&\quad + \sum_{j=1}^{n-1} ((j+1)^2 - j^2) P\{\|\tilde{X}_1\| > j\} \\
&\leq \frac{3}{n} \left[1 + \sum_{j=1}^{n-1} j P\{\|\tilde{X}_1\| > j\} \right] \\
&= o(1),
\end{aligned}$$

which completes the proof. \square

Proof of Theorem 3.1. (\Leftarrow) Set $\tilde{X}'_{jn} = \tilde{X}_j I_{[\|\tilde{X}_j\| \leq n]}$ for $1 \leq j \leq n$ and $\tilde{S}'_n = \sum_{j=1}^n \tilde{X}'_{jn}$. Then, for each $n \geq 2$, $\{\tilde{X}'_{jn}, 1 \leq j \leq n\}$ are i.i.d. and for $\varepsilon > 0$,

$$P\{d_\infty(\tilde{S}_n/n, \tilde{S}'_n/n) > \varepsilon\} \leq P\{\tilde{S}_n \neq \tilde{S}'_n\} = P\{\cup_{j=1}^n [\tilde{X}_j \neq \tilde{X}'_{jn}]\} \leq nP\{\|X_1\| > n\},$$

so that (2) entails

$$d_\infty(\tilde{S}_n/n, \tilde{S}'_n/n) \rightarrow 0 \text{ in probability.} \quad (4)$$

Next, we verify that $d_\infty(\tilde{S}'_n/n, \mathbb{E}\tilde{S}'_n/n) = d_\infty(\tilde{S}'_n/n, \mathbb{E}\tilde{X}_1 I_{[\|\tilde{X}_1\| \leq n]}) \rightarrow 0$ in probability. By Lemma 3.2, there exists a partition $0 = \alpha_0 < \alpha_1 < \dots < \alpha_k = 1$ satisfying $|u_{\alpha_i}^1 - u_{\alpha_{i-1}}^1| < \varepsilon/4$ for all i , and since $(\mathbb{E}\tilde{X}_1 I_{[\|\tilde{X}_1\| \leq n]})_\alpha^1$ converges to $u^1(\alpha)$ uniformly with respect to α , there exist $N_1 > 0$ such that for any $n \geq N_1$, $|(\mathbb{E}\tilde{X}_1 I_{[\|\tilde{X}_1\| \leq n]})_\alpha^1 - u_\alpha^1| < \varepsilon/4$ for all α . Now, since $nP\{|X_{1\alpha}^1| > n\} \leq nP\{\|X_1^1\| > n\} \rightarrow 0$, by Lemma 3.4 there exist $N_2 > 0$ such that for any $n \geq N_2$, $P\{|(\frac{\tilde{S}'_n}{n})_{\alpha_i}^1 - (\frac{\mathbb{E}\tilde{S}'_n}{n})_{\alpha_i}^1| >$

$\frac{\varepsilon}{4}\} \leq \varepsilon/k$ for all $i = 1, 2, \dots, k$. Here, we note that, for $n \geq \max\{N_1, N_2\} = N$, if $\sup_i |(\tilde{S}'_n/n)_{\alpha_i}^1 - (\mathbf{E}\tilde{X}_1 I_{[\|\tilde{X}_1\| \leq n]})_{\alpha_i}^1| \leq \varepsilon/4$, then $\sup_{\alpha} |(\tilde{S}'_n/n)_{\alpha}^1 - (\mathbf{E}\tilde{X}_1 I_{[\|\tilde{X}_1\| \leq n]})_{\alpha}^1| \leq \varepsilon$. It is because for $\alpha \in (\alpha_{i-1}, \alpha_i]$

$$\begin{aligned} (\tilde{S}'_n/n)_{\alpha}^1 - \left(\mathbf{E}\tilde{X}_1 I_{[\|\tilde{X}_1\| \leq n]}\right)_{\alpha}^1 &\leq (\tilde{S}'_n/n)_{\alpha_i}^1 - \left(\mathbf{E}\tilde{X}_1 I_{[\|\tilde{X}_1\| \leq n]}\right)_{\alpha_{i-1}^+}^1 \\ &\leq \left(\mathbf{E}\tilde{X}_1 I_{[\|\tilde{X}_1\| \leq n]}\right)_{\alpha_i}^1 - \left(\mathbf{E}\tilde{X}_1 I_{[\|\tilde{X}_1\| \leq n]}\right)_{\alpha_{i-1}^+}^1 + \frac{\varepsilon}{4} \\ &\leq u^1(\alpha_i) - u^1(\alpha_{i-1}^+) + \frac{\varepsilon}{4} + \frac{\varepsilon}{4} + \frac{\varepsilon}{4} \\ &= \varepsilon, \end{aligned}$$

and similarly we have

$$\left(\mathbf{E}\tilde{X}_1 I_{[\|\tilde{X}_1\| \leq n]}\right)_{\alpha}^1 - \left(\frac{\tilde{S}'_n}{n}\right)_{\alpha}^1 \leq \varepsilon,$$

and hence it follows. Now, for $n \geq N$

$$\begin{aligned} &P \left\{ \sup_{\alpha} \left| \left(\frac{\tilde{S}'_n}{n}\right)_{\alpha}^1 - \left(\frac{\mathbf{E}\tilde{S}'_n}{n}\right)_{\alpha}^1 \right| > \varepsilon \right\} \\ &\leq P \left\{ \sup_i \left| \left(\frac{\tilde{S}'_n}{n}\right)_{\alpha_i}^1 - \left(\frac{\mathbf{E}\tilde{S}'_n}{n}\right)_{\alpha_i}^1 \right| > \frac{\varepsilon}{4} \right\} \\ &\leq \sum_{i=1}^k P \left\{ \left| \left(\frac{\tilde{S}'_n}{n}\right)_{\alpha_i}^1 - \left(\frac{\mathbf{E}\tilde{S}'_n}{n}\right)_{\alpha_i}^1 \right| > \frac{\varepsilon}{4} \right\} \\ &\leq k \frac{\varepsilon}{k} = \varepsilon. \end{aligned}$$

Hence, we proved $\sup_{\alpha} |(\tilde{S}'_n/n)_{\alpha}^1 - (\mathbf{E}\tilde{X}_1 I_{[\|\tilde{X}_1\| \leq n]})_{\alpha}^1| \rightarrow 0$ in probability. Similarly, it can be proved $\sup_{\alpha} |(\tilde{S}'_n/n)_{\alpha}^2 - (\mathbf{E}X_1 I_{[\|X_1\| \leq n]})_{\alpha}^2| \rightarrow 0$ in probability. Therefore, we obtain

$$d_{\infty}(\tilde{S}'_n/n, \mathbf{E}\tilde{S}'_n/n) \rightarrow 0 \quad \text{in probability.} \quad (5)$$

Now by (3) and (4), we have that

$$\begin{aligned} &P \left\{ d_{\infty} \left(\frac{\tilde{S}_n}{n}, \tilde{u} \right) > \varepsilon \right\} \\ &\leq P \left\{ d_{\infty} \left(\frac{\tilde{S}_n}{n}, \frac{\tilde{S}'_n}{n} \right) + d_{\infty} \left(\frac{\tilde{S}'_n}{n}, \frac{\mathbf{E}\tilde{S}'_n}{n} \right) + d_{\infty} \left(\frac{\mathbf{E}\tilde{S}'_n}{n}, \tilde{u} \right) > \varepsilon \right\} \\ &\leq P \left\{ d_{\infty} \left(\frac{\tilde{S}_n}{n}, \frac{\tilde{S}'_n}{n} \right) > \frac{\varepsilon}{3} \right\} + P \left\{ d_{\infty} \left(\frac{\tilde{S}'_n}{n}, \frac{\mathbf{E}\tilde{S}'_n}{n} \right) > \frac{\varepsilon}{3} \right\} + P \left\{ d_{\infty} \left(\frac{\mathbf{E}\tilde{S}'_n}{n}, \tilde{u} \right) > \frac{\varepsilon}{3} \right\} \rightarrow 0 \end{aligned}$$

which completes the proof. \square

(\implies) If (1) holds, then $S_{n0}^1/n \rightarrow u_0^1$ in probability and $S_{n0}^2/n \rightarrow u_0^2$ in probability. Then the classical WLLN (see [1]) implies

$$nP\{|X_{10}^1| > n\} \rightarrow 0 \quad \text{and} \quad nP\{|X_{10}^2| > n\} \rightarrow 0.$$

Therefore,

$$\begin{aligned} nP\{\|\tilde{X}_1\| > n\} &= nP\{\max(|X_{10}^1|, |X_{10}^2|) > n\} \\ &\leq n(P\{|X_{10}^1| > n\} + P\{|X_{10}^2| > n\}) \rightarrow 0. \end{aligned}$$

Theorem 3.5. Let $\{\tilde{X}_n\}$ be a sequence of independent and identically distributed fuzzy random variables and $\tilde{S}_n = \sum_{i=1}^n \tilde{X}_i$. Let $b(x) = x^{1/\rho}l(x)$ for some $\rho \in (0, 1]$, where $l \in SV$ and $\lim_{x \rightarrow \infty} l(x) = \infty$ for $\rho = 1$. Set $b_n = b(n)$. Suppose that $(1/b_n)d_\infty(nE\tilde{X}_1 I_{\|\tilde{X}_1\| \leq b_n}, \tilde{0}) \rightarrow 0$ as $n \rightarrow \infty$ then

$$(1/b_n)d_\infty(\tilde{S}_n, \tilde{0}) \rightarrow 0 \quad \text{in probability as } n \rightarrow \infty \quad (6)$$

iff

$$nP\{\|\tilde{X}_1\| > b_n\} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (7)$$

To prove above result, we need the following lemma.

Lemma 3.6. (Gut [3]) Let $\{X_n\}$ be a sequence of i.i.d. random variables. Let $X'_j = X_j I_{|X_j| \leq b_n}$ for $1 \leq j \leq n$ and let $S'_n = \sum_{j=1}^n X'_j$. Let $b(x) = x^{1/\rho}l(x)$ for some $\rho \in (0, 1]$, where $l \in SV$ and set $b_n = b(n)$. If $nP\{|X_1| > b_n\} \rightarrow 0$ as $n \rightarrow \infty$, then $\frac{1}{b_n}(S'_n - ES'_n) \rightarrow 0$ in probability as $n \rightarrow \infty$.

Proof of Theorem 3.5. (\Leftarrow) Set $\tilde{X}'_j = \tilde{X}_j I_{\|\tilde{X}_j\| \leq b_n}$ for $1 \leq j \leq n$ and $\tilde{S}'_n = \sum_{j=1}^n \tilde{X}'_j$. Then, for each $n \geq 2$, $\{\tilde{X}'_j, 1 \leq j \leq n\}$ are i.i.d. and for $\varepsilon > 0$,

$$P\{d_\infty(\tilde{S}_n/n, \tilde{S}'_n/n) > \varepsilon\} \leq P\{\tilde{S}_n \neq \tilde{S}'_n\} = P\{\cup_{j=1}^n [\tilde{X}_j \neq \tilde{X}'_j]\} \leq nP\{\|\tilde{X}_1\| > b_n\},$$

so that (6) entails

$$(1/b_n)d_\infty(\tilde{S}_n, \tilde{S}'_n) \rightarrow 0 \quad \text{in probability as } n \rightarrow \infty. \quad (8)$$

Next, we verify that $(1/b_n)d_\infty(\tilde{S}'_n, \tilde{0}) \rightarrow 0$ in probability as $n \rightarrow \infty$. We note that

$$\begin{aligned} d_\infty(\tilde{S}'_n, \tilde{0}) &\leq |(\tilde{S}'_n)_0^1| + |(\tilde{S}'_n)_0^2| \\ &\leq \left[|(\tilde{S}'_n)_0^1 - (nE\tilde{X}_1 I_{\|\tilde{X}_1\| \leq b_n})_0^1| + d_\infty(nE\tilde{X}_1 I_{\|\tilde{X}_1\| \leq b_n}, \tilde{0}) \right] \\ &\quad + \left[|(\tilde{S}'_n)_0^2 - (nE\tilde{X}_1 I_{\|\tilde{X}_1\| \leq b_n})_0^2| + d_\infty(nE\tilde{X}_1 I_{\|\tilde{X}_1\| \leq b_n}, \tilde{0}) \right] \\ &\leq 2[d_H(L_0\tilde{S}'_n, L_0(nE\tilde{X}_1 I_{\|\tilde{X}_1\| \leq b_n})) + d_\infty(nE\tilde{X}_1 I_{\|\tilde{X}_1\| \leq b_n}, \tilde{0})]. \end{aligned}$$

Therefore, by Lemma 3.6 and the assumption, we obtain

$$(1/b_n) d_\infty(\tilde{S}'_n, \tilde{0}) \rightarrow 0 \text{ in probability as } n \rightarrow \infty. \quad (9)$$

Now by (7), and (8), we have that

$$\begin{aligned} & P \left\{ (1/b_n) d_\infty(\tilde{S}_n, \tilde{0}) > \varepsilon \right\} \\ & \leq P \left\{ (1/b_n) d_\infty(\tilde{S}_n, \tilde{S}'_n) + (1/b_n) d_\infty(\tilde{S}'_n, \tilde{0}) > \varepsilon \right\} \\ & \leq P \left\{ (1/b_n) d_\infty(\tilde{S}_n, \tilde{S}'_n) > \frac{\varepsilon}{2} \right\} + P \left\{ (1/b_n) d_\infty(\tilde{S}'_n, \tilde{0}) > \frac{\varepsilon}{2} \right\} \rightarrow 0. \end{aligned}$$

(\implies) If (5) holds, then $S_{n0}^1/b_n \rightarrow 0$ in probability and $S_{n0}^2/b_n \rightarrow 0$ in probability. Then

$$(1/b_n)(S_{n1}^i - nEX_{10}^i I_{[|X_{10}^i| \leq b_n]}) \rightarrow 0, \quad i = 1, 2$$

in probability as $n \rightarrow \infty$, since

$$EX_{10}^i I_{[|X_{10}^i| \leq b_n]} \leq d_\infty(E(\tilde{X}_1 I_{[|\tilde{X}_1| \leq b_n]}), \tilde{0}), \quad i = 1, 2.$$

Then Theorem 1.2 implies

$$nP\{|X_{10}^1| > b_n\} \rightarrow 0 \quad \text{and} \quad nP\{|X_{10}^2| > b_n\} \rightarrow 0.$$

Therefore,

$$\begin{aligned} & nP\{\|\tilde{X}_1\| > b_n\} = nP\{\max(|X_{10}^1|, |X_{10}^2|) > b_n\} \\ & \leq n(P\{|X_{10}^1| > b_n\} + P\{|X_{10}^2| > b_n\}) \rightarrow 0 \end{aligned}$$

as $n \rightarrow \infty$, which completes the proof. \square

Remark 3.7. In Theorem 3.5, if $d_\infty(E\tilde{X}_1 I_{[|\tilde{X}_1| \leq b_n]}, \tilde{u}) \rightarrow 0$ for a fuzzy number $\tilde{u} \in F(\mathbb{R})$ or $\|E\tilde{X}_1 I_{[|\tilde{X}_1| \leq b_n]}\|$, $n = 1, 2, \dots$, are bounded then the assumption $(1/b_n) d_\infty(nE\tilde{X}_1 I_{[|\tilde{X}_1| \leq b_n]}, \tilde{0}) \rightarrow 0$ as $n \rightarrow \infty$ holds since $n/b_n \rightarrow 0$ as $n \rightarrow \infty$.

Example 3.8. Let $\tilde{u} \in F(\mathbb{R})$ be fixed and let $\{Y_n\}$ be a sequence of i.i.d. random variables with common density

$$f(x) = \begin{cases} \frac{c}{x^2 \log|x|} & \text{for } |x| \geq 2, \\ 0 & \text{otherwise,} \end{cases}$$

where c is a normalizing constants. We define $(\tilde{X}_n(\omega))(x) = \tilde{u}(x - Y_n(\omega))$, i. e., $\tilde{X}_n(\omega)$ is the translation of \tilde{u} by $Y_n(\omega)$ in the real axis. Then

$$\tilde{X}_{n\alpha}^1(\omega) = u_\alpha^1 + Y_n(\omega) \quad \text{and} \quad \tilde{X}_{n\alpha}^2(\omega) = u_\alpha^2 + Y_n(\omega).$$

We note that

$$\begin{aligned} nP\{\|\tilde{X}_1\| > n\} &= nP\{\max(|X_{10}^1|, |X_{10}^2|) > n\} \\ &\leq n(P\{|u_0^1 + Y_1| > n\} + P\{|u_0^2 + Y_1| > n\}) \rightarrow 0 \end{aligned}$$

and

$$\begin{aligned} &d_H(L_\alpha \mathbb{E} \tilde{X}_1 I_{\{\|\tilde{X}_1\| \leq n\}}, L_\alpha \tilde{u}) \\ &= \max \left(\left| (\mathbb{E} \tilde{X}_1 I_{\{\|\tilde{X}_1\| \leq n\}})_\alpha^1 - u_\alpha^1 \right|, \left| (\mathbb{E} \tilde{X}_1 I_{\{\|\tilde{X}_1\| \leq n\}})_\alpha^2 - u_\alpha^2 \right| \right) \\ &= \left| \int_{\{\|\tilde{X}_1\| \leq n\}} (u_\alpha^1 + Y_1) dP - u_\alpha^1 \right| + \left| \int_{\{\|\tilde{X}_1\| \leq n\}} (u_\alpha^2 + Y_1) dP - u_\alpha^2 \right| \\ &\leq 2 \left| \int_{\{\|\tilde{X}_1\| \leq n\}} Y_1 dP \right| + \|\tilde{u}\| P\{\|\tilde{X}_1\| > n\} \\ &\leq 2 \int_{[n-u_0^1, n+u_0^1]} |Y_1| dP + 2 \int_{[n-u_0^2, n+u_0^2]} |Y_1| dP + \|\tilde{u}\| P\{\|\tilde{X}_1\| > n\} \\ &\leq 4 \int_{[n-\|\tilde{u}\|, n+\|\tilde{u}\|]} |Y_1| dP + \|\tilde{u}\| P\{\|\tilde{X}_1\| > n\} \\ &\leq 4(n + \|\tilde{u}\|) P\{|Y_1| > n - \|\tilde{u}\|\} + \|\tilde{u}\| P\{\|\tilde{X}_1\| > n\} \\ &\leq 4(n - \|\tilde{u}\|) P\{|Y_1| > n - \|\tilde{u}\|\} + 8\|\tilde{u}\| P\{\|\tilde{X}_1\| > n - \|\tilde{u}\|\} + \|\tilde{u}\| P\{\|\tilde{X}_1\| > n\} \\ &\rightarrow 0. \end{aligned}$$

Then we have

$$d_\infty \left(\frac{\tilde{S}_n}{n}, \tilde{u} \right) \rightarrow 0 \text{ in probability}$$

by the criteria in Theorem 3.1.

Example 3.9. Let $\rho \in (0, 1]$, and suppose that $\{Y_n\}$ is a sequence of i.i.d. random variables with common density

$$f(x) = \begin{cases} \frac{\rho}{2|x|^{1+\rho}} & \text{for } |x| \geq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Then $nP\{|Y_1| > (n \log n)^{1/\rho}\} \rightarrow 0$ as $n \rightarrow \infty$ (Example 1.3 [3]). We define $(\tilde{X}_n(\omega))(x) = \tilde{u}(x - Y_n(\omega))$, i.e., $\tilde{X}_n(\omega)$ is the translation of \tilde{u} by $Y_n(\omega)$ in the real axis. Then

$$\tilde{X}_{n\alpha}^1(\omega) = u_\alpha^1 + Y_n(\omega) \quad \text{and} \quad \tilde{X}_{n\alpha}^2(\omega) = u_\alpha^2 + Y_n(\omega).$$

We can easily check that $nP\{\|\tilde{X}_1\| > (n \log n)^{1/\rho}\} \rightarrow 0$. Next we consider that

$$\begin{aligned}
& d_H \left(L_\alpha \mathbf{E} \tilde{X}_1 I_{\|\tilde{X}_1\| \leq (n \log n)^{1/\rho}}, 0 \right) \\
&= \max \left(\left| (\mathbf{E} \tilde{X}_1 I_{\|\tilde{X}_1\| \leq (n \log n)^{1/\rho}})_\alpha^1 \right|, \left| (\mathbf{E} \tilde{X}_1 I_{\|\tilde{X}_1\| \leq (n \log n)^{1/\rho}})_\alpha^2 \right| \right) \\
&= \left| \int_{\|\tilde{X}_1\| \leq (n \log n)^{1/\rho}} (u_\alpha^1 + Y_1) dP \right| + \left| \int_{\|\tilde{X}_1\| \leq (n \log n)^{1/\rho}} (u_\alpha^2 + Y_1) dP \right| \\
&\leq 2 \left| \int_{\|\tilde{X}_1\| \leq (n \log n)^{1/\rho}} Y_1 dP \right| + 2\|\tilde{u}\| \\
&\leq 4 \int_{[(n \log n)^{1/\rho} - \|\tilde{u}\|, (n \log n)^{1/\rho} + \|\tilde{u}\|]} |Y_1| dP + 2\|\tilde{u}\| \\
&\leq 4((n \log n)^{1/\rho} + \|\tilde{u}\|)P\{|Y_1| > (n \log n)^{1/\rho} - \|\tilde{u}\|\} + 2\|\tilde{u}\|,
\end{aligned}$$

and hence we have that $(1/(n \log n)^{1/\rho}) d_\infty(n\mathbf{E}\tilde{X}_1 I_{\|\tilde{X}_1\| \leq (n \log n)^{1/\rho}}, \tilde{0}) \rightarrow 0$ as $n \rightarrow \infty$. Then we have

$$d_\infty \left(\frac{\tilde{S}_n}{(n \log n)^{1/\rho}}, \tilde{0} \right) \rightarrow 0 \text{ in probability}$$

by the criteria in Theorem 3.5.

ACKNOWLEDGEMENT

This work was supported by Grant No. R05-2004-000-10304-0(2004) from the Korea Science and Engineering Foundation.

(Received July 22, 2005.)

REFERENCES

- [1] Y. S. Chow and H. Teicher: Probability Theory. Second edition. Springer-Verlag, New York 1998.
- [2] R. Goetschel and W. Voxman: Elementary fuzzy calculus. *Fuzzy Sets and Systems* 18 (1986), 31–43.
- [3] A. Gut: An extension of the Kolmogorov–Feller weak law of large numbers with an application to the St. Petersburg game. *J. Theoret. Probab.* 17 (2004), 769–779.
- [4] D. H. Hong and H. J. Kim: Marcinkiewicz-type law of large numbers for fuzzy random variables. *Fuzzy Sets and Systems* 64 (1994), 387–393.
- [5] D. H. Hong: A convergence of fuzzy random variables. *Kybernetika* 39 (2003), 275–280.
- [6] H. Inoue: A strong law of large numbers for fuzzy random sets. *Fuzzy Sets and Systems* 41 (1991), 285–291.
- [7] S. Y. Joo, S. S. Lee, and Y. H. Yoo: A strong law of large numbers for stationary fuzzy random variables. *J. Korean Statist. Soc.* 30 (2001), 153–161.
- [8] S. Y. Joo and Y. K. Kim: Kolmogorov’s strong law of large numbers for fuzzy random variables. *Fuzzy Sets and Systems* 120 (2001), 499–503.

- [9] S. Y. Joo: Weak laws of large numbers for fuzzy random variables. *Fuzzy Sets and Systems* 147 (2004), 453–464.
- [10] Y. K. Kim: A strong law of large numbers for fuzzy random variables. *Fuzzy Sets and Systems* 111 (2000), 319–323.
- [11] E. P. Klement, M. L. Puri, and D. A. Ralescu: Limit theorems for fuzzy random variables. *Proc. Roy. Soc. London Ser. A* 407 (1986), 171–182.
- [12] R. Kruse: The strong law of large numbers for fuzzy random variables. *Inform. Sci.* 28 (1982), 233–241.
- [13] M. Miyakoshi and M. Shimbo: A strong law of large numbers for fuzzy random variables. *Fuzzy Sets and Systems* 12 (1984), 133–142.
- [14] I. Molchanov: Strong laws of large numbers for random upper semicontinuous. *J. Math. Anal. Appl.* 235 (1999), 349–355.
- [15] R. L. Taylor, L. Seymour, and Y. Chen: Weak laws of large numbers for fuzzy random sets. *Nonlinear Anal.* 47 (2001), 1245–1256.
- [16] T. Uemura: A law of large numbers for random sets. *Fuzzy Sets and Systems* 59 (1993), 181–188.

Dug Hun Hong, Department of Mathematics, Myongji University, Yongin Kyunggido 449-728. South Korea.

e-mail: dhhong@mju.ac.kr

Kyung Tae Kim, Department of Electronics and Electrical Information Engineering, Kyungwon University, Sunnam Kyunggido. South Korea.

e-mail: ktkim@kyungwon.ac.kr