

PITMAN EFFICIENCIES OF L_p -GOODNESS-OF-FIT TESTS

JAN BEIRLANT AND LÁSZLÓ GYÖRFI

Pitman efficiencies are used to describe the problem of choice of the number of classes in L_p -goodness-of-fit tests ($p \geq 1$) based on histogram density estimates. We consider the case where the number of classes increases with the sample size.

1. INTRODUCTION

Let X_1, X_2, \dots be i.i.d. real valued random variables. Consider the problem of testing the simple null hypothesis H_0 that the X_i 's are distributed according to some distribution μ_0 with a continuous distribution function F_0 , versus a simple hypothesis H_1 . Without loss of generality we may assume that μ_0 is the uniform distribution on $[0, 1]$, otherwise one transforms the data by F_0 . Let μ_1 be the distribution of X_i 's under H_1 .

Let μ_n be the empirical distribution for the sample X_1, X_2, \dots, X_n :

$$\mu_n(A) = \frac{\#\{i; X_i \in A, 1 \leq i \leq n\}}{n}$$

and let $\mathcal{P}_n = \{A_{n,j}, j = 1, 2, \dots\}$ be a uniform partition of $[0, 1]$ of interval size $h_n > 0$ ($k_n = 1/h_n$ is integer).

In this paper we consider the L_p -goodness-of-fit test statistics:

$$J_{k,n}^{(p)} = \sum_{j=1}^{k_n} |\mu_n(A_{n,j}) - \mu_0(A_{n,j})|^p.$$

In case no confusion is possible we abbreviate $J_{k,n}^{(p)}$ to J_n . The case $p = 2$ corresponds to a χ^2 -statistic, while $p = 1$ was studied by Györfi and van der Meulen [4] and Beirlant, Györfi and Lugosi [1]. These authors introduced these statistics in the context of L_p -errors for the histogram density estimator

$$f_n(x) = \mu_n(A_{n,j})/h_n \quad (x \in A_{n,j}).$$

Indeed,

$$J_n = h_n^{p-1} \int_0^1 |f_n - 1|^p.$$

The aim of this paper is to prove some new results with respect to the Pitman efficiency of any pair of tests induced by two different values of p , hence providing some new insight in the number of classes to be taken relatively from one L_p -test with respect to another. This note can also be considered as an addition to the work of Quine and Robinson [6] who performed this same program for the L_2 -test and the likelihood ratio test for uniformity.

2. MAIN RESULT

The Pitman efficiency is defined along a sequence of neighboring alternatives, assuming that

$$\mu_1(A) = \mu_0(A) + \nu_n \tau(A)$$

where $\nu_n > 0$ and $\nu_n \rightarrow 0$ as $n \rightarrow \infty$, and τ is a signed measure with density g . Obviously

$$\tau([0, 1]) = \int_0^1 g(x) dx = 0.$$

The technique in the sequel is based on a Poissonization argument developed in Beirlant, Györfi and Lugosi [1] and Beirlant and Mason [2] as a generalization of the work of Morris [5]. When using this technique \tilde{N} will denote a Poisson(n) random variable independent of the data, and $\tilde{\Pi}_n$ be the empirical Poisson measure:

$$\tilde{\Pi}_n(A) = \#\{i; X_i \in A, 1 \leq i \leq \tilde{N}\},$$

leading to the auxiliary test statistics

$$\tilde{J}_n = n^{-p} \sum_{j=1}^{k_n} |\tilde{\Pi}_n(A_{n,j}) - n\mu_0(A_{n,j})|^p.$$

Furthermore let us introduce sequences of normalizing constants $E_{\ell,n}$, $V_{\ell,n}$, $\ell = 0, 1$ such that if $nh_n \rightarrow \infty$ and $h_n \rightarrow 0$ as $n \rightarrow \infty$, we have under H_0

$$\frac{J_n - E_{0,n}}{\sqrt{V_{0,n}}} \xrightarrow{\mathcal{D}} N(0, 1),$$

and under $H_{1,n}$

$$\frac{J_n - E_{1,n}}{\sqrt{V_{1,n}}} \xrightarrow{\mathcal{D}} N(0, 1).$$

In the sequel N stands for a standard normal random variable.

In the same way as in Beirlant, Györfi and Lugosi [1] or in Beirlant and Mason [2] one can prove the following result:

Theorem 1. Assume that for some $\delta > 0$, $\int_0^1 |g(x)|^{4+\delta} dx < \infty$. If $h_n \rightarrow 0$ and $nh_n \rightarrow \infty$ as $n \rightarrow \infty$, and $\nu_n^2 = O(h_n^{-1/2}n^{-1})$ then under H_0 and $H_{1,n}$ respectively

$$\frac{J_n - E_{l,n}}{\sqrt{V_{l,n}}} \xrightarrow{\mathcal{D}} N(0, 1), \quad (l = 0, 1)$$

where

$$\lim_{n \rightarrow \infty} \frac{V_{1,n}}{V_{0,n}} = 1$$

and

$$\lim_{n \rightarrow \infty} \frac{E_{1,n} - E_{0,n}}{\sqrt{V_{0,n}}} = \frac{c(p)}{2} \nu_n^2 n \sqrt{h_n} \int_0^1 g^2(x) dx + o(1),$$

where

$$c(p) = \frac{pE(|N|^p)}{\sqrt{\text{Var}(|N|^p)}}.$$

The limit results mentioned in the preceding theorem now can be used to derive Pitman efficiencies of the L_p -tests under consideration for sequences of alternatives considered in the introduction. To this end, as in Quine and Robinson [6] we will suppose that the number of cells $k = k(n)$ will be induced by a function k which, when taken as a function of the continuous variable x , is regularly varying, that is that for some q , $k(ax)/k(x) \rightarrow a^q$ as $x \rightarrow \infty$, for all $a > 0$. We consider then tests of the hypothesis H_0 using J_n chosen in such a way that that the power of the size α test under $H_{1,n}$ tends to β ($\alpha < \beta < 1$) as n tends to ∞ . Let J'_n be another statistic and n' a sequence such that the power of the size α test based on $J'_{n'}$ under $H_{1,n'}$ also tends to β as $n' \rightarrow \infty$. Then if the limit of n'/n exists and is the same for all such sequences n' , we call it the Pitman efficiency of J_n with respect to J'_n and write

$$PE(J_n, J'_n) = \lim n'/n.$$

Specifically, we choose $\nu_n = \nu(n)$ such that

$$\lim_{n \rightarrow \infty} \frac{E_{1,n} - E_{0,n}}{\sqrt{V_{0,n}}} \rightarrow b > 0$$

and we take n' such that the same limit relation holds with the same constant b when using the other test based on J'_n and when $k(n)$ is replaced by $k'(n')$. Here the role of J and J' is played by considering two different values of p . With the method of proof used in Section 2 of Quine and Robinson [6] the following result now follows from Theorem 1:

Theorem 2. Under the conditions of Theorem 1 and assuming that both $k(n)$ and $k'(n)$ are regularly varying sequences of numbers of intervals with indices of regular variation q and q' in $[0, 1]$, then

$$PE(J_{k,n}^{(p_1)}, J_{k',n}^{(p_2)}) = \left(\frac{c^2(p_1)}{c^2(p_2)} c \right)^{\frac{1}{2-q}} \quad (1 \leq p_1, p_2 < \infty)$$

if $q = q'$ and $k'(n)/k(n) \rightarrow c \in (0, \infty)$, and

$$PE \left(J_{k,n}^{(p_1)}, J_{k',n}^{(p_2)} \right) = \infty$$

if $k'(n)/k(n) \rightarrow \infty$.

Since for any $a > 1$ one obtains that

$$E(|N|^a) = \frac{2^{a/2}}{\sqrt{\pi}} \Gamma \left(\frac{a+1}{2} \right)$$

one finds that

$$c(p) = \frac{p}{\sqrt{\frac{\sqrt{\pi}\Gamma(p+1/2)}{\Gamma^2(p/2)} - 1}}.$$

For large p , using Stirling's formula one gets

$$c(p) \sim p2^{-p/2} \quad (p \rightarrow \infty).$$

Finally Lemma 1 in the Appendix states that $c(p)$ has a unique maximum at $p = 2$. Figure 1 provides a graph of this function, showing that for any test J'_n using a value p_2 different from 2 one has to choose $k'(n)/k(n) \rightarrow c^2(p_2)/2 < 1$ as $n \rightarrow \infty$ in order to have Pitman efficiency 1 with respect to the χ^2 test.

Fig. 1.

3. PROOF OF THEOREM 1

It is shown in Theorem 3.1 in Beirlant and Mason [2] that

$$\frac{J_n - E_{0,n}}{\sqrt{V_{0,n}}} \xrightarrow{D} N(0, 1)$$

if $h_n \rightarrow 0$ and $nh_n \rightarrow \infty$ with

$$E_{0,n} = n^{-p} \sum_{j=1}^{k_n} E_0 \left(|\tilde{\Pi}_n(A_{n,j}) - n\mu_0(A_{n,j})|^p \right)$$

and

$$V_{0,n} = n^{-2p} \text{Var}(|N|^p) n^p \sum_{j=1}^{k_n} \mu_0^p(A_{n,j}) = n^{-p} h_n^{p-1} \text{Var}(|N|^p).$$

From Lemma 2.2 in Beirlant and Mason [2] it follows that

$$E_0 \left(|\tilde{\Pi}_n(A_{n,j}) - n\mu_0(A_{n,j})|^p (n\mu_0(A_{n,j}))^{-p/2} \right) - E(|N|^p) = O(1/\sqrt{n})$$

as $n \rightarrow \infty$ so that then

$$\begin{aligned} E_{0,n} &= \sum_{j=1}^{k_n} E \left(|N| \sqrt{\mu_0(A_{n,j})/n} \right) + O \left(\frac{1}{\sqrt{n}} (nh_n)^{-p/2} h_n^{p-1} \right) \\ &= E(|N|^p) h_n^{-1+p/2} n^{-p/2} + O \left(\frac{1}{\sqrt{n}} (nh_n)^{-p/2} h_n^{p-1} \right) \\ &= E(|N|^p) h_n^{-1+p/2} n^{-p/2} + O \left(\frac{1}{\sqrt{n}} (nh_n)^{-p/2} h_n^{p-1} \right) \\ &=: E_{0,n}^* + b_n. \end{aligned}$$

The proof of the given limit result under $H_{1,n}$ asks for a little bit more care. Again the technique used in the proof of Theorem 3.1 in Beirlant and Mason [2] can be applied given that the following conditions are satisfied:

$$\max_{j=1, \dots, k_n} \mu_1(A_{n,j}) \rightarrow 0 \tag{1}$$

$$n \min_{j=1, \dots, k_n} \mu_1(A_{n,j}) \rightarrow \infty \tag{2}$$

$$\max_{j=1, \dots, k_n} \mu_1^p(A_{n,j}) / \sum_{j=1}^{k_n} \mu_1^p(A_{n,j}) \rightarrow 0. \tag{3}$$

First, using Hölder’s inequality we obtain that

$$\begin{aligned} \max_{j=1,\dots,k_n} \mu_1(A_{n,j}) &= h_n + \nu_n \max_{j=1,\dots,k_n} \int_{A_{n,j}} g(x) \, dx \\ &= h_n + O(n^{-1/2}h_n^{1/2-1/4}) \left(\int_0^1 g^2(x) \, dx \right)^{1/2} \\ &= h_n + O(n^{-1/2}h_n^{1/4}) = o(1) \quad (n \rightarrow \infty). \end{aligned}$$

Next, to prove (2) remark that

$$n \min_{j=1,\dots,k_n} \mu_1(A_{n,j}) \geq nh_n \left(1 - \frac{\nu_n}{h_n} \max_{j=1,\dots,k_n} \left| \int_{A_{n,j}} g(x) \, dx \right| \right).$$

Now, using Hölder’s inequality again, we obtain

$$\begin{aligned} \frac{\nu_n}{h_n} \left| \int_{A_{n,j}} g(x) \, dx \right| &\leq \frac{\nu_n}{h_n} h_n^{1-\frac{1}{4+\delta}} \left(\int_0^1 g^{4+\delta}(x) \, dx \right)^{1/(4+\delta)} \\ &= O\left(h_n^{-1/4-1/(4+\delta)} n^{-1/2} \right) \end{aligned}$$

which tends to zero as $n \rightarrow \infty$, hence finishing the proof of (2). The proof of (3) goes along similar lines.

With the method of proof used in the proof of Theorem in Beirlant, Györfi and Lugosi [1] or Theorem 3.1 in Beirlant and Mason [2] one can now check that under the given conditions

$$\frac{J_n - E_{1,n}}{\sqrt{V_{1,n}}} \xrightarrow{\mathcal{D}} N(0, 1),$$

where the expressions for $E_{1,n}$ and $V_{1,n}$ will now be specified.

First,

$$\begin{aligned} V_{1,n} &= n^{-p} \text{Var}(|N|^p) \sum_{j=1}^{k_n} \mu_{1,n}^p(A_{n,j}) \\ &= n^{-p} \text{Var}(|N|^p) \sum_{j=1}^{k_n} \left(h_n + O(n^{-1/2}h_n^{-1/4}) \int_{A_{n,j}} g(x) \, dx \right)^p \end{aligned}$$

which as in the proof of (2) leads to

$$V_{1,n} = n^{-p} h_n^{-1+p} \text{Var}(|N|^p) (1 + o(1))$$

as $n \rightarrow \infty$, from which we get that

$$\lim_{n \rightarrow \infty} \frac{V_{1,n}}{V_{0,n}} = 1.$$

Secondly, introducing the notation

$$\psi_p(a) = E(|N + a|^p)$$

one obtains with the help of a straightforward extension of Lemma 2.2(a) in Beirlant and Mason [2] that

$$\begin{aligned} E_{1,n} &= n^{-p} \sum_{j=1}^{k_n} E_1(|\tilde{\Pi}_n(A_{n,j}) - n\mu_0(A_{n,j})|^p) \\ &= n^{-p} \sum_{j=1}^{k_n} E_1(|\tilde{\Pi}_n(A_{n,j}) - n\mu_1(A_{n,j}) + n\mu_1(A_{n,j}) - n\mu_0(A_{n,j})|^p) \\ &= n^{-p} \sum_{j=1}^{k_n} E_1(|\tilde{\Pi}_n(A_{n,j}) - n\mu_1(A_{n,j}) + n\nu_n\tau(A_{n,j})|^p) \\ &= \sum_{j=1}^{k_n} \left(\frac{\mu_1(A_{n,j})}{n}\right)^{p/2} \left\{ E \left(\left| N + \sqrt{\frac{n}{\mu_1(A_{n,j})}} \nu_n\tau(A_{n,j}) \right|^p \right) \right\} \\ &\quad + O(1/\sqrt{n}) \\ &= \sum_{j=1}^{k_n} \left(\frac{\mu_1(A_{n,j})}{n}\right)^{p/2} \psi_p \left(\sqrt{\frac{n}{\mu_1(A_{n,j})}} \nu_n\tau(A_{n,j}) \right) \\ &\quad + O \left(\frac{1}{\sqrt{n}} n^{-p/2} h_n^{-1+p/2} \right) \\ &=: E_{1,n}^* + c_n. \end{aligned}$$

Using the same method as in the derivation of (2) one first shows that

$$\frac{c_n}{\sqrt{V_{0,n}}} = O(1/\sqrt{nh_n})$$

as $n \rightarrow \infty$. Next, by Lemma 2 in the Appendix

$$\begin{aligned}
E_{1,n}^* - E_{0,n}^* &= \sum_{j=1}^{k_n} \left(\frac{\mu_1(A_{n,j})}{n} \right)^{p/2} \psi_p \left(\sqrt{\frac{n}{\mu_1(A_{n,j})}} \nu_n \tau(A_{n,j}) \right) \\
&\quad - \sum_{j=1}^{k_n} \left(\frac{\mu_0(A_{n,j})}{n} \right)^{p/2} \psi_p(0) \\
&= \sum_{j=1}^{k_n} \left(\frac{\mu_1(A_{n,j})}{n} \right)^{p/2} \left[\psi_p \left(\sqrt{\frac{n}{\mu_1(A_{n,j})}} \nu_n \tau(A_{n,j}) \right) - \psi_p(0) \right] \\
&\quad + \psi_p(0) \sum_{j=1}^{k_n} \left[\left(\frac{\mu_1(A_{n,j})}{n} \right)^{p/2} - \left(\frac{\mu_0(A_{n,j})}{n} \right)^{p/2} \right] \\
&= \sum_{j=1}^{k_n} \left(\frac{\mu_1(A_{n,j})}{n} \right)^{p/2} \frac{pE(|N|^p)}{2} \left(\sqrt{\frac{n}{\mu_1(A_{n,j})}} \nu_n \tau(A_{n,j}) \right)^2 [1 + o(1)] \\
&\quad + \psi_p(0) \sum_{j=1}^{k_n} \frac{p}{2n} \left(\frac{\mu_0(A_{n,i})}{n} \right)^{p/2-1} \nu_n \tau(A_{n,i}) [1 + o(1)] \\
&= \frac{pE(|N|^p)}{2} \sum_{j=1}^{k_n} \left(\frac{\mu_1(A_{n,j})}{n} \right)^{p/2-1} \nu_n^2 \tau(A_{n,j})^2 [1 + o(1)].
\end{aligned}$$

Hence

$$\begin{aligned}
\frac{E_{1,n}^* - E_{0,n}^*}{\sqrt{V_{0,n}}} &= \frac{pE(|N|^p)}{2\sqrt{V_p(0)}} \nu_n^2 n \sqrt{h_n} \sum_{j=1}^{k_n} \left(\frac{\mu_0(A_{n,j})}{h_n} \right)^{p/2-1} \left(\frac{\tau(A_{n,j})}{h_n} \right)^2 h_n + o(1) \\
&= \frac{c(p)}{2} \nu_n^2 n \sqrt{h_n} \int_0^1 g(x)^2 dx + o(1).
\end{aligned}$$

4. APPENDIX

Lemma 1. The function $c : [1, \infty) \rightarrow (0, \infty)$ has a unique maximum at $p = 2$.

Proof. Setting $h(p) = E(|N|^{2p}) / (E(|N|^p))^2$ we get $c^2(p) = \frac{p^2}{h(p)-1}$.

Now

$$g(q) = \left(\frac{d}{dq} E(|N|^q) \right) / E(|N|^q) = \frac{\ln 2}{2} + \frac{1}{2} \Psi\left(\frac{q+1}{2}\right)$$

where $\Psi(z) = \Gamma'(z)/\Gamma(z)$ denotes the logarithmic derivative of the gamma function.

We find that

$$\begin{aligned}
h'(p) &= 2h(p)(g(2p) - g(p)) \\
&= h(p) \left(\Psi(p+1/2) - \Psi\left(\frac{p+1}{2}\right) \right).
\end{aligned}$$

On the other hand, the numerator of the expression for the derivative of c^2 is equal to

$$p(2h(p) - 2 - ph'(p)) = p \left(\left[2 - p \left(\Psi(p + 1/2) - \Psi \left(\frac{p+1}{2} \right) \right) \right] h(p) - 2 \right)$$

so that it remains to show that

$$1 - \frac{p}{2} \left(\Psi(p + 1/2) - \Psi \left(\frac{p+1}{2} \right) \right) \left\{ \begin{matrix} > \\ = \\ < \end{matrix} \right\} \frac{1}{h(p)} = \frac{\Gamma^2((p+1)/2)}{\sqrt{\pi}\Gamma(p+1/2)}$$

as

$$p \left\{ \begin{matrix} < \\ = \\ > \end{matrix} \right\} 2.$$

To this end remark that

$$h(p) = \exp \left\{ \int_0^{p/2} (\Psi(v + 1/2 + p/2) - \Psi(v + 1/2)) dv \right\},$$

so that it suffices to prove that

$$1 - \frac{p}{2} d_{p/2}(p/2) \left\{ \begin{matrix} > \\ = \\ < \end{matrix} \right\} \exp \left\{ - \int_0^{p/2} d_{p/2}(v) dv \right\}$$

as

$$p \left\{ \begin{matrix} < \\ = \\ > \end{matrix} \right\} 2,$$

with

$$d_q(u) = \Psi(u + q + 1/2) - \Psi(u + 1/2).$$

As $d_1(u) = 1/u$, one immediately checks the equality in case $p = 2$. From Gauss' expression for the logarithmic derivative Ψ of the gamma function one obtains that if $u > 0$ and $u + q > 0$

$$d_q(u) = \int_0^1 \frac{x^{u-1/2}(1-x^q)}{1-x} dx,$$

from which one derives that

- for any q , d_q is a decreasing function of u ,
- $d_{q_1} \leq d_{q_2}$ if $q_1 < q_2$,
- $d'_{q_1} \geq d'_{q_2}$ if $q_1 < q_2$.

From these three properties of d_q the result now follows since they imply that

$$q \mapsto qd_q(q) + \exp \left\{ - \int_0^q d_q(v) dv \right\}$$

is a decreasing function in $q > 1/2$. □

Lemma 2. As $a \downarrow 0$

$$\psi_p(a) = \psi_p(0) + pE(|N|^p) \frac{a^2}{2} (1 + o(1)).$$

Proof. One easily checks that $\psi_p'(0) = 0$ and $\psi_p''(0) = pE(|N|^p)$. □

(Received March 3, 1994.)

REFERENCES

-
- [1] J. Beirlant, L. Györfi and G. Lugosi: On the asymptotic normality of the L_1 - and L_2 -errors in histogram density estimation. *Canad. J. Statist.* (to appear 1995).
 - [2] J. Beirlant and D. M. Mason: On the asymptotic normality of L_p norms of empirical functionals. *Mathem. Methods Statist.* (to appear 1995).
 - [3] L. Devroye and L. Györfi: *Nonparametric Density Estimation: The L_1 -View*. Wiley, New York 1985.
 - [4] L. Györfi and E. C. van der Meulen: A consistent goodness-of-fit test based on the total variation distance. In: *Nonparametric Functional Estimation and Related Topics* (G. Roussas, ed.), Kluwer Academic Publishers, Dordrecht 1991, pp. 631–645.
 - [5] C. Morris: Central limit theorems for multinomial sums. *Ann. Statist.* *3* (1975), 165–188.
 - [6] M. P. Quine and J. Robinson: Efficiencies of chi-square and likelihood ratio goodness-of-fit tests. *Ann. Statist.* *13* (1985), 727–742.

Dr. Jan Beirlant, Department of Mathematics, Katholieke Universiteit Leuven, 200B Celestijnenlaan, B-3000 Leuven. Belgium.

Dr. László Györfi, Department of Mathematics, Technical University Budapest, H-1521 Budapest, Stoczek u. 2. Hungary.