

Kybernetika

VOLUME 43 (2007), NUMBER 6

The Journal of the Czech Society for
Cybernetics and Information Sciences

Published by:

Institute of Information Theory
and Automation of the AS CR, v.v.i.

Editor-in-Chief:

Milan Mareš

Editorial Board:

Managing Editors:

Karel Sladký
Lucie Fajfrová

Jiří Anděl, Sergej Čelikovský, Marie
Demlová, Petr Hájek, Jan Flusser, Martin
Janžura, Jan Ježek, George Klir, Ivan
Kramosil, Tomáš Kroupa, Friedrich Liese,
Jean-Jacques Loiseau, František Matúš,
Radko Mesiar, Jiří Outrata, Jan Štecha,
Olga Štěpánková, Igor Vajda, Jiřina
Vejnarová, Miloslav Vošvrda, Pavel Zítek

Editorial Office:

Pod Vodárenskou věží 4, 182 08 Praha 8

Kybernetika is a bi-monthly international journal dedicated for rapid publication of high-quality, peer-reviewed research articles in fields covered by its title.

Kybernetika traditionally publishes research results in the fields of Control Sciences, Information Sciences, System Sciences, Statistical Decision Making, Applied Probability Theory, Random Processes, Fuzziness and Uncertainty Theories, Operations Research and Theoretical Computer Science, as well as in the topics closely related to the above fields.

The Journal has been monitored in the Science Citation Index since 1977 and it is abstracted/indexed in databases of Mathematical Reviews, Current Mathematical Publications, Current Contents ISI Engineering and Computing Technology.

Kybernetika. Volume 43 (2007)

ISSN 0023-5954, MK ČR E 4902.

Published bimonthly by the Institute of Information Theory and Automation of the Academy of Sciences of the Czech Republic, Pod Vodárenskou věží 4, 182 08 Praha 8. — Address of the Editor: P. O. Box 18, 182 08 Prague 8, e-mail: kybernetika@utia.cas.cz. — Printed by PV Press, Pod vrstevnicí 5, 140 00 Prague 4. — Orders and subscriptions should be placed with: MYRIS TRADE Ltd., P. O. Box 2, V Štíhlách 1311, 142 01 Prague 4, Czech Republic, e-mail: myris@myris.cz. — Sole agent for all “western” countries: Kubon & Sagner, P. O. Box 34 01 08, D-8 000 München 34, F.R.G.

Published in December 2007.

© Institute of Information Theory and Automation of the AS CR, v.v.i., Prague 2007.

ON THE TANGENTIAL VELOCITY ARISING IN A CRYSTALLINE APPROXIMATION OF EVOLVING PLANE CURVES

SHIGETOSHI YAZAKI

In a crystalline algorithm, a tangential velocity is used implicitly. In this short note, it is specified for the case of evolving plane curves, and is characterized by using the intrinsic heat equation.

Keywords: tangential velocity, intrinsic heat equation, crystalline algorithm, admissible polygonal curve

AMS Subject Classification: 34A26, 34A34, 35K65, 53A04, 53C80, 53C44, 65L20, 65M12, 65N12

1. INTRODUCTION

It is well known that the tangential velocity of evolution of plane curves causes stability from a numerical point of view. In this short note, the tangential velocity in a crystalline algorithm is specified, and it is compared with the one in other schemes by using the intrinsic heat equation. Our goal is simple. In the crystalline algorithm, the tangential velocity equals the quantity which is described as a negative of partial derivatives of the normal velocity with respect to the arc-length and its division by the curvature.

Following Mikula and Ševčovič [12], our target curves are defined as follows. We consider an embedded, orientable and closed plane curve Γ which is parameterized by a smooth function $\mathbf{x} : \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}^2$ such that $\Gamma = \text{Image}(\mathbf{x}) = \{\mathbf{x}(u); u \in \mathbb{R}/\mathbb{Z}\}$ and $|\partial_u \mathbf{x}| > 0$. Here and hereafter, we denote $\partial_\xi(\cdot) = \partial(\cdot)/\partial\xi$. The unit tangent vector can be defined as $\mathbf{T} = \partial_u \mathbf{x} / |\partial_u \mathbf{x}| = \partial_s \mathbf{x}$, where s is the arc-length parameter and $ds = |\partial_u \mathbf{x}| du$, and the unit inward normal vector is defined by $\mathbf{N} = \mathbf{T}^\perp$, where $(a, b)^\perp = (-b, a)$. The curvature in the direction \mathbf{N} is denoted by k , and the Frenet's formulae are $\partial_s \mathbf{T} = k \mathbf{N}$ and $\partial_s \mathbf{N} = -k \mathbf{T}$. Let ν be the angle of \mathbf{T} , i. e., $\mathbf{T} = (\cos \nu, \sin \nu)$ and $\mathbf{N} = (-\sin \nu, \cos \nu)$. Our problem is as follows: For a given initial curve $\Gamma^0 = \text{Image}(\mathbf{x}^0) = \Gamma$, find a family of curve $\{\Gamma^t\}_{t \geq 0}$, $\Gamma^t = \{\mathbf{x}(u, t); u \in \mathbb{R}/\mathbb{Z}\}$ which starts from $\mathbf{x}(u, 0) = \mathbf{x}^0(u)$ for $u \in \mathbb{R}/\mathbb{Z}$ and evolves according to the normal velocity

$$v = \beta(k, \nu). \tag{1}$$

Information of β determines movement of curves, and in many physical phenomena β includes k and ν . In general, β may also be a function of \mathbf{x} and L^t , etc., other than k and ν , where $L^t = \int_{\Gamma^t} ds = \int_0^1 |\partial_u \mathbf{x}| du$ is the total length of the curve Γ^t . The normal velocity ν is the normal component of the following evolution equation of a solution \mathbf{x} :

$$\partial_t \mathbf{x} = \beta \mathbf{N} + \alpha \mathbf{T}, \quad \mathbf{x}(\cdot, 0) = \mathbf{x}^0(\cdot). \tag{2}$$

Our main result is that, in a crystalline algorithm, the tangential velocity α is used essentially and implicitly such as:

$$\alpha = -\frac{\partial_s \beta}{k}. \tag{3}$$

This equation follows from the equality $\partial_t \nu = 0$, where $\partial_t \nu$ is described as

$$\partial_t \nu = \partial_s \beta + \alpha k, \quad \nu(\cdot, 0) = \nu^0(\cdot).$$

See [12, (3.8)].

In the crystalline algorithm, the class of polygonal curves is restricted to an admissible class and this restriction corresponds to the equality $\partial_t \nu = 0$. The admissibility is a powerful concept which may handle singular anisotropic energy such as crystalline energy (see references in the next section). We note that in the three dimensional crystalline algorithm for mean curvature flow has not been successfully constructed in the sense that the class of polyhedron is unclear. However, convergence of crystalline algorithm for the three dimensional Gauss curvature flow has been proved [16, 17].

In the next section, we will construct evolution of polygonal curves under the assumption $\partial_t \nu = 0$. In Section 3, the tangential velocity (3) will be compared with the one in other known schemes. In the last Section 4, some remarks on numerical scheme will be mentioned.

2. CRYSTALLINE ALGORITHM

A crystalline algorithm describes the motion of polygonal curves in an admissible class, and the admissible polygonal curves are the polygonal curves with each tangential angles belonging to a prescribed set. Let the prescribed set be $\mathcal{T} = \{\eta_1 < \eta_2 < \dots < \eta_J < \eta_1 + 2\pi\}$, where J is a natural number $J \geq 3$ and $\eta_{j+1} - \eta_j < \pi$ is satisfied for all $j = 1, 2, \dots, J$ ($\eta_{J+1} = \eta_1$). On a smooth curve $\Gamma = \{\mathbf{x}(u); u \in \mathbb{R}/\mathbb{Z}\}$, we distribute N points $\mathbf{x}_i = \mathbf{x}(u_i)$ ($u_1 < u_2 < \dots < u_N < u_1 + 1$), which satisfy two conditions: (1) the tangential angle $\nu_i = \nu(u_i)$ of \mathbf{x}_i belongs to \mathcal{T} , i.e., $(\cos \nu_i, \sin \nu_i) = \partial_s \mathbf{x}(\nu_i)$ and $\nu_i \in \mathcal{T}$ hold for $i = 1, 2, \dots, N$; (2) For $i = 1, 2, \dots, N$, there exists j such that $\{\nu_i, \nu_{i+1}\} = \{\eta_j, \eta_{j+1}\}$ holds ($\nu_{N+1} = \nu_1$). Let l_i be the straight line passing through \mathbf{x}_i in the direction $\mathbf{T}_i = (\cos \nu_i, \sin \nu_i)$, and let \mathbf{y}_i be the intersection point of l_{i-1} and l_i for all i . Thus a closed polygonal curve $\mathcal{P} = \bigcup_{i=1}^N [\mathbf{y}_i, \mathbf{y}_{i+1}]$ is constructed ($\mathbf{y}_{N+1} = \mathbf{y}_1$). This \mathcal{P} is called \mathcal{T} -admissible polygonal curve. The length of the i th edge $d_i = |\mathbf{y}_{i+1} - \mathbf{y}_i|$ is given by

$$d_i = (\cot \xi_i + \cot \xi_{i+1})h_i - h_{i+1} \operatorname{cosec} \xi_{i+1} - h_{i-1} \operatorname{cosec} \xi_i \tag{4}$$

for $i = 1, 2, \dots, N$, where $\xi_i = \nu_i - \nu_{i-1}$ and $h_i = \mathbf{x}_i \cdot \mathbf{N}_i$ ($\nu_0 = \nu_N$, $\mathbf{N}_i = \mathbf{T}_i^\perp$). And the i th vertex \mathbf{y}_i is given by

$$\mathbf{y}_i = h_i \mathbf{N}_i + \frac{h_{i-1} - h_i \cos \xi_i}{\sin \xi_i} \mathbf{T}_i \tag{5}$$

for $i = 1, 2, \dots, N$.

Crystalline algorithm is formulated as the following problem: For a given initial \mathcal{T} -admissible N -sided polygonal curve $\mathcal{P}^0 = \bigcup_{i=1}^N [\mathbf{y}_i^0, \mathbf{y}_{i+1}^0] = \mathcal{P}$, find a family of \mathcal{T} -admissible polygonal curves $\{\mathcal{P}^t\}_{t \geq 0}$, $\mathcal{P}^t = \bigcup_{i=1}^N [\mathbf{y}_i(t), \mathbf{y}_{i+1}(t)]$ which starts from $\mathbf{y}_i(0) = \mathbf{y}_i^0$ for $i = 1, 2, \dots, N$ and evolves according to the i th normal velocity

$$v_i = \beta(k_i, \nu_i), \quad i = 1, 2, \dots, N, \tag{6}$$

where k_i is the i th discretized curvature, which is called crystalline curvature if $k_i = c_i/d_i$ for some geometric quantity c_i depending on \mathcal{T} . Since the problem is considered in the admissible class, the i th tangential angle ν_i does not depend on time t , i.e., $\partial_t \nu_i = 0$ holds for all i . Moreover, from (4), we have the evolution equations as a system of ordinary differential equations:

$$\dot{d}_i = (\cot \xi_i + \cot \xi_{i+1})v_i - v_{i+1} \operatorname{cosec} \xi_{i+1} - v_{i-1} \operatorname{cosec} \xi_i \tag{7}$$

for $i = 1, 2, \dots, N$, where $v_i = \dot{h}_i = \dot{\mathbf{x}}_i \cdot \mathbf{N}_i$. Here and hereafter, we denote $\dot{(\cdot)} = d(\cdot)/dt$. It is easy to check that (7) is equivalent to the evolution equations of vertices:

$$\dot{\mathbf{y}}_i = v_i \mathbf{N}_i + \frac{v_{i-1} - v_i \cos \xi_i}{\sin \xi_i} \mathbf{T}_i \tag{8}$$

for $i = 1, 2, \dots, N$. Therefore, it is possible to interpret the crystalline approximation as follows. Let s_i be the arc-length defined as $s_i = s + \int_u^{u_i} |\partial_u \mathbf{x}| du$. By putting $\beta(s) = \beta(k(s), \nu(s))$ and $v_i = \beta(s_i)$, we have $v_{i-1} = \beta(s_i - a_i) = v_i - \partial_s \beta(s_i) a_i + o(a_i)$ ($a_i = s_i - s_{i-1}$). Hence it holds that

$$\frac{v_{i-1} - v_i \cos \xi_i}{\sin \xi_i} = \beta(s_i) \frac{1 - \cos \xi_i}{\sin \xi_i} - \frac{\partial_s \beta(s_i)}{\sin \xi_i} a_i + \frac{o(a_i)}{\sin \xi_i} \rightarrow -\frac{\partial_s \beta(s_i)}{k(s_i)}$$

as $s_{i-1} \rightarrow s_i$. Here we have used $\partial_s \nu = k$ and $\xi_i = \nu_i - \nu_{i-1} = \int_{s_{i-1}}^{s_i} k(s) ds$. This corresponds to (3) at $s = s_i$.

This crystalline algorithm was introduced by J. E. Taylor [13, 14] and S. Angenent and M. E. Gurtin [1] at the end of 1980's. Since then crystalline curvature flow equation has been studied under various kinds of evolution law by several authors. We refer the reader, other than the above pioneer works, to the surveys by Taylor, Cahn and Handwerker [15] and the books by Gurtin [6] for a geometric and physical background. Besides this crystalline strategy, other strategies by subdifferential and level-set method have been extensively studied. See Giga [4] and references therein. Recently, the admissibility is extended as an essentially admissibility. See, e.g., Hontani, Giga, Giga and Deguchi [8] and Yazaki [19].

3. COMPARISON WITH OTHER SCHEMES

In this section, the tangential velocity (3) is compared with the one in other schemes by using the intrinsic heat equation for a solution \mathbf{x} :

$$\partial_t \mathbf{x} = \theta_1^{-1} \partial_s (\theta_2^{-1} \partial_s \mathbf{x}), \quad \mathbf{x}(\cdot, 0) = \mathbf{x}^0(\cdot),$$


where $\theta_1 \theta_2 = k/\beta$. Hence, the tangential velocity α is described as $\alpha = \theta_1^{-1} \partial_s \theta_2^{-1}$.

We will sketch the known results. Dziuk [3] studied a numerical scheme for the case $\beta(k) = k$ with $\alpha = 0$, i. e., $\theta_1 = \theta_2 = 1$. Kimura [10] proposed a uniform redistribution scheme in the case $\beta(k) = k$ by using a nontrivial tangential velocity α which satisfies $\theta_1 \theta_2 = 1$ and discretizes the conditions $\int_0^1 \alpha \, du = 0$ and $|\partial_u \mathbf{x}| = L^t$. At this stage, the relation between these conditions and θ_1, θ_2 is not clear. Another scheme for the case $\beta(k) = k$ was proposed by Deckelnick [2], who used a nontrivial $\alpha = -\partial_u (|\partial_u \mathbf{x}|^{-1})$ which satisfies $\theta_1 = |\partial_u \mathbf{x}| = 1/\theta_2$. Mikula and Ševćovič [11] generalized these results for the case $\beta(k) = |k|^{m-1}k$ by putting $\alpha = \partial_s |k|^{m-1}/2$ and $\theta_1 = \theta_2 = \sqrt{k/\beta(k)}$. This result was improved in Mikula and Ševćovič [12] for the general case $v = \beta(k, \nu)$. In this, θ_1 and θ_2 satisfy $\alpha = \theta_1^{-1} \partial_s \theta_2^{-1}$ and $\partial_s \alpha = \bar{k}v - kv$, where $\bar{f} = \int_{\Gamma^t} f \, ds/L^t$ is the average of f .

Applying the crystalline case (3) for $\beta(k) = |k|^{m-1}k$, we obtain $\alpha = -\bar{m} \partial_s |k|^{m-1}$, $\bar{m} = m/(m-1)$ if $m \neq 1$, and $\alpha = -\partial_s \log |k|$ if $m = 1$; and $\theta_1 = k/\beta^2, \theta_2 = \beta$.

4. REMARKS ON NUMERICAL SCHEMES

Discretizing (7) (or (8)) in time and using the adapted time step control, we obtain a sequence of \mathcal{T} -admissible polygonal curves $\mathcal{P}^0 = \mathcal{P}_0, \mathcal{P}_1, \mathcal{P}_2, \dots$, where \mathcal{P}_j approximates \mathcal{P}^{t_j} ($0 = t_0 < t_1 < t_2 < \dots$). The following figure is a numerical scheme in the case $\beta(k) = |k|^{m-1}k$ with $m = 0.1$.



We can observe that vertices are concentrated at sharp corners. But scheme does not break down, since the elimination of some edges and renumbering are done. (In this figure, the last partition number is $N = 36$ starting with the initial partition number $N = 380$.) In other words, the solution is extended in several times while the solution curve keeps admissibility. However, in general, preservation of admissibility is open problems. Indeed, in the case where $\beta(k, \nu) = a(\nu)|k|^{m-1}k$, there exist $m \in (0, 1)$, $a(\cdot)$ and \mathcal{P}^0 such that the admissibility collapses in finite time, i. e., we have the examples that admissible nonconvex polygonal curve becomes nonadmissible in finite time. See, e. g., Hirota, Ishiwata and Yazaki [7]. This is contrast result to the following: For any $m \geq 1$, $a(\cdot)$ and \mathcal{P}^0 if the anisotropy is symmetric, then the solution curve keeps admissibility. See Giga and Giga [5].

We can also observe that no swallow tail occurs because of the adapted time step control. Then we can continue the scheme until the maximal existence time. Moreover several convergence results are known until the maximal existence time. See, e. g., [5, 18] and references therein. At this stage, before the maximal existence time, it is open whether \mathcal{P}^t becomes convex or not. For instance, the following non-convexified phenomenon is known: In the case where $\beta(k, \nu) = a(\nu)|k|^{m-1}k$, there exist $m \in (0, 1)$, $a(\cdot)$ and \mathcal{P}^0 such that nonconvex solution curve shrinks homothetically, i. e., there exists a nonconvex self-similar solution polygonal curve. See Ishiwata, Ushijima, Yagisita and Yazaki [9].

ACKNOWLEDGEMENT

The author would like to thank the organizers of the First Slovak-Japan Workshop on Computational Mathematics which was held while the author was visiting Czech Republic within the period from August 18 to September 19, 2006; this visit was sponsored by Czech Technical University in Prague, Faculty of Nuclear Sciences and Physical Engineering within the Jindřich Nečas Center for Mathematical Modeling (Project of the Czech Ministry of Education, Youth and Sports, project No. LC 06052). Furthermore, the author is grateful to the anonymous referee for her or his comments and suggestions. The present research is/was supported by Grant-in-Aid for Encouragement of Young Scientists No. 17740063.

(Received November 30, 2006.)

REFERENCES

-
- [1] S. Angenent and M. E. Gurtin: Multiphase thermomechanics with interfacial structure, 2. Evolution of an isothermal interface. *Arch. Rational Mech. Anal.* *108* (1989), 323–391.
 - [2] K. Deckelnick: Weak solutions of the curve shortening flow. *Calc. Var. Partial Differential Equations* *5* (1997), 489–510.
 - [3] G. Dziuk: Convergence of a semi discrete scheme for the curve shortening flow. *Math. Models Methods Appl. Sci.* *4* (1994), 589–606.
 - [4] Y. Giga: Anisotropic curvature effects in interface dynamics. *Sūgaku* *52* (2000), 113–127; English transl., *Sugaku Expositions* *16* (2003), 135–152.
 - [5] M.-H. Giga and Y. Giga: Crystalline and level set flow – convergence of a crystalline algorithm for a general anisotropic curvature flow in the plane. *GAKUTO Internat. Ser. Math. Sci. Appl.* *13* (2000), 64–79.
 - [6] M. E. Gurtin: *Thermomechanics of Evolving Phase Boundaries in the Plane*. Clarendon Press, Oxford 1993.
 - [7] C. Hirota, T. Ishiwata, and S. Yazaki: Numerical study and examples on singularities of solutions to anisotropic crystalline curvature flows of nonconvex polygonal curves. In: *Advanced Studies in Pure Mathematics (ASPM); Proc. MSJ-IRI 2005 “Asymptotic Analysis and Singularity”*, Sendai 2005 (to appear).
 - [8] H. Hontani, M.-H. Giga, Y. Giga, and K. Deguchi: Expanding selfsimilar solutions of a crystalline flow with applications to contour figure analysis. *Discrete Appl. Math.* *147* (2005), 265–285.
 - [9] T. Ishiwata, T. K. Ushijima, H. Yagisita, and S. Yazaki: Two examples of nonconvex self-similar solution curves for a crystalline curvature flow. *Proc. Japan Academy, Ser. A* *80* (2004), 8, 151–154.

- [10] M. Kimura: Accurate numerical scheme for the flow by curvature. *Appl. Math. Letters* **7** (1994), 69–73.
- [11] K. Mikula and D. Ševčovič: Solution of nonlinearly curvature driven evolution of plane curves. *Appl. Numer. Math.* **31** (1999), 191–207.
- [12] K. Mikula and D. Ševčovič: Evolution of plane curves driven by a nonlinear function of curvature and anisotropy. *SIAM J. Appl. Math.* **61** (2001), 1473–1501.
- [13] J. E. Taylor: Constructions and conjectures in crystalline nondifferential geometry. In: *Proc. Conference on Differential Geometry, Rio de Janeiro, Pitman Monographs Surveys Pure Appl. Math.* **52** (1991), pp. 321–336, Pitman, London 1991.
- [14] J. E. Taylor: Motion of curves by crystalline curvature, including triple junctions and boundary points. *Diff. Geom.: Partial Diff. Eqs. on Manifolds (Los Angeles 1990)*, *Proc. Sympos. Pure Math.*, **54** (1993), Part I, pp. 417–438, AMS, Providence.
- [15] J. E. Taylor, J. Cahn, and C. Handwerker: Geometric models of crystal growth. *Acta Metall.* **40** (1992), 1443–1474.
- [16] T. K. Ushijima and H. Yagisita: Approximation of the Gauss curvature flow by a three-dimensional crystalline motion. In: *Proc. Czech–Japanese Seminar in Applied Mathematics 2005; Kuju Training Center, Oita, Japan, September 15–18, 2005, COE Lecture Note 3, Faculty of Mathematics, Kyushu University (M. Beneš, M. Kimura, and T. Nakaki, eds.)*, 2006, pp. 139–145.
- [17] T. K. Ushijima and H. Yagisita: Convergence of a three-dimensional crystalline motion to Gauss curvature flow. *Japan J. Indust. Appl. Math.* **22** (2005), 443–459.
- [18] T. K. Ushijima and S. Yazaki: Convergence of a crystalline approximation for an area-preserving motion. *Journal of Comput. and Appl. Math.* **166** (2004), 427–452.
- [19] S. Yazaki: Motion of nonadmissible convex polygons by crystalline curvature. *Publ. Res. Inst. Math. Sci.* **43** (2007), 155–170.

*Shigetoshi Yazaki, Faculty of Engineering, University of Miyazaki, 1-1 Gakuen Kibanadai Nishi, Miyazaki 889-2192. Japan.
e-mail: yazaki@cc.miyazaki-u.ac.jp*