

Kybernetika

VOLUME 39 (2003), NUMBER 3

The Journal of the Czech Society for
Cybernetics and Information Sciences

Published by:

Institute of Information Theory
and Automation of the Academy
of Sciences of the Czech Republic

Editor-in-Chief:

Milan Mareš

Managing Editors:

Karel Sladký

Editorial Board:

Jiří Anděl, Marie Demlová, Petr Hájek,
Martin Janžura, Jan Ježek, Radim Jiroušek,
George Klir, Ivan Kramosil, Friedrich Liese,
Jean-Jacques Loiseau, František Matúš,
Radko Mesiar, Jiří Outrata, Jan Štecha,
Olga Štěpánková, Igor Vajda, Pavel Zítek,
Pavel Žampa

Editorial Office:

Pod Vodárenskou věží 4, 182 08 Praha 8

Kybernetika is a bi-monthly international journal dedicated for rapid publication of high-quality, peer-reviewed research articles in fields covered by its title.

Kybernetika traditionally publishes research results in the fields of Control Sciences, Information Sciences, System Sciences, Statistical Decision Making, Applied Probability Theory, Random Processes, Fuzziness and Uncertainty Theories, Operations Research and Theoretical Computer Science, as well as in the topics closely related to the above fields.

The Journal has been monitored in the Science Citation Index since 1977 and it is abstracted/indexed in databases of Mathematical Reviews, Current Mathematical Publications, Current Contents ISI Engineering and Computing Technology.

Kybernetika. Volume 39 (2003)

ISSN 0023-5954, MK ČR E 4902.

Published bi-monthly by the Institute of Information Theory and Automation of the Academy of Sciences of the Czech Republic, Pod Vodárenskou věží 4, 182 08 Praha 8. — Address of the Editor: P. O. Box 18, 182 08 Prague 8, e-mail: kybernetika@utia.cas.cz. — Printed by PV Press, Pod vrstevnicí 5, 140 00 Prague 4. — Orders and subscriptions should be placed with: MYRIS TRADE Ltd., P. O. Box 2, V Štíhlách 1311, 142 01 Prague 4, Czech Republic, e-mail: myris@myris.cz. — Sole agent for all “western” countries: Kubon & Sagner, P. O. Box 34 01 08, D-8 000 München 34, F.R.G.

Published in June 2003.

© Institute of Information Theory and Automation of the Academy of Sciences of the Czech Republic, Prague 2003.

***T*-EQUIVALENCES GENERATED BY SHAPE FUNCTION ON THE REAL LINE**

DUG HUN HONG

This paper is devoted to give a new method of generating *T*-equivalence using shape function and finding the exact calculation formulas of *T*-equivalence induced by shape function on the real line. Some illustrative examples are given.

Keywords: fuzzy number, fuzzy relation, *t*-norm, *T*-equivalence, shape function

AMS Subject Classification: 26A21, 03E02

1. INTRODUCTION

For the fuzzy set-theoretical modelling of verbal quantities and computing with these quantities, it appears useful to part the class of real numbers into fuzzy equivalence classes. Jacas and Recasens [8] considered the idea of generating fuzzy numbers as equivalence classes of a *T*-indistinguishability operator based on a scale function. The theoretical approach suggested in [10] and further developed in [11] indicates that partitions based on the concept of a shape function can be especially significant. De Baets et al [2] and Marková [12] characterized that the shapes by means of which *T*-equivalences can be generated, are based on the knowledge of idempotents of the *T*-addition of fuzzy numbers.

In this paper, we give a new method of generating *T*-equivalence using shape function and finding the exact calculation formulas of *T*-equivalence induced by shape function on the real line. Some illustrative examples are given.

2. PRELIMINARIES

Definition 1. (Jacas and Recasens [8]) A fuzzy number is a mapping $A : R \rightarrow [0, 1]$ such that there exists $a \in R$ with $A(a) = 1$ and A is increasing on $(-\infty, a]$ and A is decreasing on $[a, \infty)$.

Definition 2. (De Baets and Mesiar [3]) Consider a *t*-norm T . A binary fuzzy relation E on an universe X is called a *T*-equivalence on X if and only if it is reflexive, symmetric and *T*-transitive, i. e. if and only if for any (x, y, z) in X^3 :

- (i) $E(x, x) = 1$;

- (ii) $E(x, y) = E(y, x)$;
- (iii) $T(E(x, y), E(y, z)) \leq E(x, z)$.

Definition 3. (Jacas and Recasens [8]) A scale is a continuous non-decreasing surjective monotonic mapping $S : R \rightarrow R$.

Definition 4. A shape is a non-increasing mapping $\phi : R^+ \rightarrow [0, 1]$ such that $\phi(0) = 1$.

Definition 5. A mapping $d : X^2 \rightarrow [0, \infty]$ is called a pseudo-metric on X if and only if for any (x, y, z) in X^3

- (i) $d(x, x) = 0$;
- (ii) $d(x, y) = d(y, x)$;
- (iii) $d(x, z) \leq d(x, y) + d(y, z)$.

It is called a metric if it moreover satisfies, for any $(x, y) \in X^2$

- (iv) $d(x, y) = 0 \Leftrightarrow x = y$.

Consider a scale s , then the mapping $d_s : R^2 \rightarrow R^+$ defined by

$$d_s(x, y) = |s(x) - s(y)|$$

is a pseudo-metric on R . Now consider a shape ϕ , then we construct the binary fuzzy relation $E_{s,\phi}$ as follows:

$$E_{s,\phi}(x, y) = \phi(|s(x) - s(y)|).$$

Definition 6. A generator (or source of vagueness) g is a scale such that $g(0) = 0$.

A function $T : [0, 1] \times [0, 1] \rightarrow [0, 1]$ is said to be a triangular norm [9, 14] (t -norm for short) iff T is symmetric, associative, non-decreasing in each argument, and $T(x, 1) = x$ for all $x \in [0, 1]$, and, in general, $T(x_1, \dots, x_n) = T(T(\dots T(T(x_1, x_2), x_3), \dots, x_{n-1}), x_n)$. Some well-known continuous t -norms are the minimum operator T_M , the algebraic product T_P and the Lukasiewicz t -norm T_L defined by $T_L(x, y) = \max(x + y - 1, 0)$. The minimum operator T_M is the strongest (greatest) t -norm. The weakest (smallest) t -norm T_W is defined by

$$T_W(x, y) = \begin{cases} \min(x, y) & \text{if } \max(x, y) = 1, \\ 0, & \text{elsewhere.} \end{cases}$$

We will call t -norm T is Archimedean if and only if T is continuous and $T(x, x) < x$ for all $x \in (0, 1)$. Every Archimedean t -norm T is representable by a continuous and decreasing function $f : [0, 1] \rightarrow [0, \infty]$ with $f(1) = 0$ and

$$T(x_1, \dots, x_n) = f^{[-1]}(f(x_1) + \dots + f(x_n))$$

for all $x_i \in [0, 1]$, $1 \leq i \leq n$, where $f^{[-1]}$ is the pseudo-inverse of f , defined by

$$f^{[-1]}(y) = \begin{cases} f^{-1}(y) & \text{if } y \in [0, f(0)], \\ 0 & \text{if } y \in [f(0), \infty]. \end{cases}$$

The function f is the additive generator of T . If $T = T_P$, then $f(x) = \log x^{-1}$ and if $T = T_L$, then $f(x) = 1 - x$.

For arbitrary fuzzy numbers A_i , $i = 1, \dots, n$, $n \in N$, on the real line, their T -sum is defined by means of the extension principle as follows:

$$A_1 \oplus_T \dots \oplus_T A_n(z) = \sup_{x_1 + \dots + x_n = z} T(A_1(x_1), \dots, A_n(x_n)), \quad z \in R.$$

Definition 7. Let J be a finite or countable set. Let $\{T_i | i \in J\}$ be a collection of t -norms and $\{(a_i, b_i) | i \in J\}$ a collection of disjoint intervals in $[0, 1]$. We call ordinal sum of t -norms $\{T_i | i \in J\}$ to the following t -norm :

$$T(x, y) = \begin{cases} a_i + (b_i - a_i)T_i\left(\frac{x - a_i}{b_i - a_i}, \frac{y - a_i}{b_i - a_i}\right) & \text{whenever } (x, y) \in (a_i, b_i)^2 \\ & = (a_i, b_i) \times (a_i, b_i), \\ \min(x, y) & \text{otherwise,} \end{cases}$$

which is denoted by $T = (\langle a_i, b_i, T_i \rangle | i \in J)$, and only if all T_i are generated, then equivalently it can be used $T = (\langle a_i, b_i, f_i \rangle | i \in J)$ where f_i is the additive generator of T_i .

The following theorem gives a general classification of continuous t -norms [9].

Theorem 1. (Ling [9]) Let T be a continuous t -norm. Then T is Archimedean or T -min or T is an ordinal sum of Archimedean t -norms.

3. T -EQUIVALENCE GENERATED BY SHAPES

Consider a generator g and a shape ϕ , and the fuzzy relation $E_{g,\phi}$, which is always reflexive and symmetric. Let T be a t -norm and $\phi_n = \phi \oplus_T \dots \oplus_T \phi$ (n -fold T -sum of ϕ). Then $\phi_n(x) \leq \phi_{n+1}(x)$ for any $x \in R$ and for $n \in N$, the natural numbers. Hence the limit always exists. Let $\lim_{n \rightarrow \infty} \phi_n \equiv \phi^*$. We also note that if we define $|\phi| : R \rightarrow [0, 1]$ such that $|\phi|(z) = \phi(|z|)$ and $|\phi|_n = |\phi| \oplus_T \dots \oplus_T |\phi|$, then $\lim_{n \rightarrow \infty} |\phi|_n \equiv |\phi|^* = |\phi^*|$.

Theorem 2. For a continuous t -norm T , a generator g and a shape ϕ , the fuzzy relation E_{f,ϕ^*} is a T -equivalence on R .

Proof. We only need to show that for any $a, b, y \in R$

$$T(E_{g,\phi^*}(a, y), E_{g,\phi^*}(y, b)) \leq E_{g,\phi^*}(a, b),$$

or equivalently

$$T(|\phi|^*(g(y) - g(a)), |\phi|^*(g(b) - g(y))) \leq |\phi|^*(g(b) - g(a)). \tag{1}$$

By the continuity of the t -norm T , we have

$$\begin{aligned} & T(|\phi|^*(g(y) - g(a)), |\phi|^*(g(b) - g(y))) \\ &= \lim_{n \rightarrow \infty} T(|\phi|_n(g(y) - g(a)), |\phi|_n(g(b) - g(y))) \end{aligned}$$

and

$$\begin{aligned} & T(|\phi|_n(g(y) - g(a)), |\phi|_n(g(b) - g(y))) \\ &= T \left(\sup_{x_1 + \dots + x_n = g(y) - g(a)} T(|\phi|(x_1), \dots, |\phi|(x_n)), \right. \\ & \quad \left. \sup_{x_{n+1} + \dots + x_{2n} = g(b) - g(y)} T(|\phi|(x_{n+1}), \dots, |\phi|(x_{2n})) \right) \\ &= \sup_{\substack{x_1 + \dots + x_n = g(y) - g(a) \\ x_{n+1} + \dots + x_{2n} = g(b) - g(y)}} T(T(|\phi|(x_1), \dots, |\phi|(x_n)), T(|\phi|(x_{n+1}), \dots, |\phi|(x_{2n}))) \\ &\leq \sup_{x_1 + \dots + x_{2n} = g(b) - g(a)} T(|\phi|(x_1), \dots, |\phi|(x_{2n})) \\ &= |\phi|_{2n}(g(b) - g(a)) \end{aligned}$$

where the second equality comes from the continuity of T and the inequality comes from non-decreasing property of T , hence equation (1) is proved since $\lim_{n \rightarrow \infty} |\phi|_{2n}(g(b) - g(a)) = |\phi|^*(g(b) - g(a))$. \square

The following theorem is due to B. De Baets et al [2]. Here, we give a new proof using the idea of Theorem 2.

Theorem 3. (De Baets et al [2]) Consider a t -norm T , a generator g and a shape ϕ . Let $H = \{|g(u) - g(v)| | (u, v) \in R^2\}$. If for any $x \in H$, $\phi \oplus_T \phi(x) = \phi(x)$, then the fuzzy relation $E_{g,\phi}$ is a T -equivalence on R .

Proof. Define ϕ_0 as follows :

$$\phi_0(x) = \begin{cases} \phi(x) & \text{if } x \in H, \\ \inf\{\phi(w) | w < x, w \in H\} & \text{if } x \notin H. \end{cases}$$

Then ϕ_0 is a shape with $E_{g,\phi}(x, y) = E_{g,\phi_0}(x, y)$ for $(x, y) \in R^2$. We can also show that for any $x \in R$, $\phi_0 \oplus_T \phi_0(x) = \phi_0(x)$. It is because $\phi_0 \oplus_T \phi_0(x) \geq \phi_0(x)$ is always true and for $x \notin H$, $w \in H$ and $w < x$,

$$\begin{aligned} \phi_0 \oplus_T \phi_0(x) &\leq \phi_0 \oplus \phi_0(w) \\ &= \phi(w) \end{aligned}$$

and hence

$$\begin{aligned} \phi_0 \oplus_T \phi_0(x) &\leq \inf\{\phi(w) | w < x, w \in H\} \\ &= \phi_0(x). \end{aligned}$$

We now note that $\phi_0 = \phi_0^*$ and can prove that E_{g, ϕ_0} is a T -equivalence on R according to the exactly same method as Theorem 1 without the assumption of continuity of T using $\phi_0 \oplus_T \phi_0 = \phi_0$. This completes the proof. \square

Recently, many authors [5, 6, 7, 13] studied facts about T -sums of shape function and their limits.

Theorem 4. (Hong and Hwang [6], Hong and Ro [7], Mesiar [11]) Consider a continuous Archimedean t -norm T with additive generator f and a shape ϕ . If $f \circ \phi$ is convex, then

$$\phi_n(x) = f^{[-1]} \left(n f \circ \phi \left(\frac{x}{n} \right) \right).$$

Theorem 5. (Hong and Hwang [5]) Consider a continuous Archimedean t -norm T with additive generator f and a shape ϕ . If $f \circ \phi$ is convex, then $\phi^*(0) = 1$ and for $x > 0$,

$$\lim_{n \rightarrow \infty} \phi_n(x) = \phi^*(x) = f^{[-1]}(x f'_-(1) \phi'_+(0)).$$

Definition 8. Consider $(a, b) \in R, a \neq b$, then $\phi_{(a,b)}$ is the linear transformation defined by

$$\phi_{(a,b)}(x) = \frac{x - a}{b - a}$$

Note that the inverse mapping $\phi_{(a,b)}^{-1}$ of $\phi_{(a,b)}$ is given by $\phi_{(a,b)}^{-1}(x) = a + (b - a)x$.

Definition 9. Consider a fuzzy quantity A and $(a, b) \in [0, 1]^2, a < b$.

- (i) The fuzzy quantity $A^{[a,b]}$ is defined as $A^{[a,b]} = \text{tr} \circ \phi_{(a,b)} \circ A$, i. e. $A^{[a,b]}(x) = \text{tr}((A(x) - a)/(b - a))$, where $\text{tr} : R \rightarrow [0, 1]$ is defined by

$$\text{tr}(x) = \begin{cases} 0, & \text{if } x < 0, \\ x, & \text{if } 0 \leq x \leq 1, \\ 1, & \text{if } x > 1. \end{cases}$$

- (ii) The fuzzy quantity $A_{[a,b]}$ is defined by

$$A_{[a,b]}(x) = \begin{cases} \phi_{(a,b)}^{-1}(A(x)), & \text{if } A(x) > 0, \\ 0, & \text{elsewhere.} \end{cases}$$

We need the following result to generalize Theorem 5 to arbitrary continuous t -norm.

Theorem 6. (De Baets and Marková [1]) Consider an ordinal sum of continuous t -norm $T = (\langle a_i, b_i, f_i \rangle | i \in I)$ written in such a way that $\bigcup_{\lambda \in I} [a_i, b_i] = [0, 1]$ and a shape ϕ . If $f_i \circ \phi^{[a_i, b_i]}$ is convex for all $i \in I$, then

$$\phi_n(x) = \sup_{i \in I} \left\{ (\phi_n^{T_i, [a_i, b_i]})_{[a_i, b_i]}(x) \right\}$$

where $\phi_n^{T_i, [a_i, b_i]}(x) = f_i^{[-1]}(n f_i \circ \phi^{[a_i, b_i]}(\frac{x}{n}))$.

Theorem 5 can be easily generalized to arbitrary ordinal sums of continuous t -norm T .

Theorem 7. Consider an ordinal sums of continuous t -norm $T = (\langle a_i, b_i, f_i \rangle | i \in I)$ written in such a way that $\bigcup_{\lambda \in I} [a_i, b_i] = [0, 1]$ and a shape ϕ . If $f_i \circ \phi^{[a_i, b_i]}$ is convex for all $i \in I$, then

$$\begin{aligned} \phi^*(x) &= \lim_{n \rightarrow \infty} \phi_n(x) \\ &= \sup_{i \in I} \left\{ (\phi^{T_i, [a_i, b_i]})_{[a_i, b_i]}(x) \right\}, \end{aligned}$$

where $\phi^{T_i, [a_i, b_i]}(x) = \lim_{n \rightarrow \infty} \phi_n^{T_i, [a_i, b_i]}(x) = f_i^{[-1]}(x(f_i)'_-(1) (\phi^{[a_i, b_i]})'_+(0))$.

4. EXAMPLES

Example 1. Consider the product t -norm T_P with additive generator $f(x) = \log x^{-1}$, and a generator g and a shape function ϕ defined by $\phi(x) = \max\{1 - x, 0\}$. Then, by Theorem 5 (or see [5]), $\phi^*(x) = e^{-x}$, and hence $E_{g, \phi^*}(x, y) = e^{-|g(x) - g(y)|}$ is a T -equivalence on R .

Example 2. Consider the Lukasiewicz t -norm T_L with additive generator $f(x) = 1 - x$, and generator g and a shape function ϕ defined by $\phi(x) = \max\{1 - x, 0\}$. Then, by Theorem 5 (or see [5]), $\phi^*(x) = \phi(x)$, and hence $E_{g, \phi^*}(x, y) = \max\{1 - |g(x) - g(y)|, 0\}$ is a T -equivalence on R .

Example 3. Consider the ordinal sums $T = (\langle 0, \frac{1}{3}, \log x^{-1} \rangle, \langle \frac{1}{3}, 1, 1 - x \rangle)$, a generator g and a shape function ϕ defined by $\phi(x) = \max\{1 - x, 0\}$. Then, by Theorem 7, $\phi^*(x) = \max\{1 - x, \frac{1}{3}\}$, and hence $E_{g, \phi^*}(x, y) = \max\{1 - |g(x) - g(y)|, \frac{1}{3}\}$ is a T -equivalence on R .

Example 4. Consider the ordinal sums $T = (\langle 0, \frac{1}{3}, 1 - x \rangle, \langle \frac{1}{3}, 1, \log x^{-1} \rangle)$, a generator g and a shape function ϕ defined by $\phi(x) = \max\{1 - x, 0\}$. Then, by Theorem 7, $\phi^*(x) = \frac{1}{3} + \frac{2}{3}e^{-\frac{3}{2}x}$ since $f^{T_P, [\frac{1}{3}, 1]}(x) = e^{-\frac{3}{2}x}$ and $f^{T_L, [0, \frac{1}{3}]}(x) = 1$. Hence $E_{g, \phi^*}(x, y) = \frac{1}{3} + \frac{2}{3}e^{-\frac{3}{2}|g(x) - g(y)|}$ is a T -equivalence on R .

Example 5. Consider the ordinal sums $T = (\langle 0, \frac{1}{3}, \log x^{-1} \rangle, \langle \frac{1}{3}, 1, 1 - x \rangle)$, a generator g and a shape function ϕ defined by

$$\phi(x) = \begin{cases} 1 & \text{if } x = 0, \\ \frac{1}{3}(1 - x) & \text{if } 0 < x \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Then, by Theorem 7,

$$\phi^*(x) = \begin{cases} 1 & \text{if } x = 0, \\ \frac{1}{3}e^{-x} & \text{otherwise,} \end{cases}$$

since

$$f^{T_L, [\frac{1}{3}, 1]}(x) = \begin{cases} 1 & \text{if } x = 0, \\ 0 & \text{otherwise,} \end{cases}$$

and $f^{T_P, [0, \frac{1}{3}]}(x) = e^{-x}$. Hence

$$E_{g, \phi^*}(x, y) = \begin{cases} 1 & \text{if } x = y, \\ \frac{1}{3}e^{-|g(x) - g(y)|} & \text{otherwise,} \end{cases}$$

is a T -equivalence on R .

Example 6. Consider the ordinal sums $T = (\langle 0, \frac{1}{3}, 1 - x \rangle, \langle \frac{1}{3}, 1, \log x^{-1} \rangle)$, a generator g and a shape function ϕ defined by

$$\phi(x) = \begin{cases} 1 & \text{if } x = 0, \\ \frac{1}{3}(1 - x) & \text{if } 0 < x \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Then, by Theorem 7,

$$\phi^*(x) = \begin{cases} 1 & \text{if } x = 0, \\ \frac{1}{3}(1 - x) & \text{if } |x| \leq 1, \\ 0 & \text{otherwise,} \end{cases}$$

since

$$f^{T_P, [\frac{1}{3}, 1]}(x) = \begin{cases} 1 & \text{if } x = 0, \\ 0 & \text{otherwise,} \end{cases}$$

and $f^{T_L, [0, \frac{1}{3}]}(x) = 1 - x$. Hence

$$\phi_{g, \phi^*}(x, y) = \begin{cases} 1 & \text{if } g(x) = g(y), \\ \frac{1}{3}(1 - |g(x) - g(y)|) & \text{if } |g(x) - g(y)| \leq 1, \\ 0 & \text{otherwise,} \end{cases}$$

is a T -equivalence on R .

ACKNOWLEDGEMENTS

This work was supported by grant R01-2000-000-00011-0 from the Korea Science & Engineering Foundation.

(Received September 3, 2002.)

REFERENCES

-
- [1] B. De Baets and A. Marková: Analytical for the addition of fuzzy intervals. *Fuzzy Sets and Systems* *91* (1997), 203–213.
 - [2] B. De Baets, M. Mareš, and R. Mesiar: T-partition of the real line generated by impotent shapes. *Fuzzy Sets and Systems* *91* (1997), 177–184.
 - [3] B. De Baets and R. Mesiar: Pseudo-metrics and T-equivalence. *J. Fuzzy Math.* *5* (1997), 471–481.
 - [4] M. A. Harrison: *Introduction to Switching and Automata Theory*. McGraw–Hill, New York 1965.
 - [5] D. H. Hong and S. Y. Hwang: On the convergence of T-sum of L-R fuzzy numbers. *Fuzzy Sets and Systems* *63* (1994), 175–180.
 - [6] D. H. Hong and C. Hwang: A T-sum bound of L-R fuzzy numbers. *Fuzzy Sets and Systems* *91* (1997), 239–252.
 - [7] D. H. Hong and P. Ro: The law of large numbers for fuzzy numbers with bounded supports. *Fuzzy Sets and Systems* *116* (2000), 269–274.
 - [8] J. Jacas and J. Recasens: Fuzzy numbers and equality relations. In: *Proc. 2nd IEEE International Conference on Fuzzy Systems, San Francisco 1993*, pp. 1298–1301.
 - [9] C. Ling: Representation of associative function. *Publ. Math. Debrecen* *12* (1965), 189–212.
 - [10] M. Mareš and R. Mesiar: Composition of shape generators. *Acta Mathematica et Informatica Universitatis Ostraviensis* *4* (1996), 37–46.
 - [11] M. Mareš and R. Mesiar: Fuzzy quantities and their aggregation, in aggregation operations. In: *New Trends and Applications*. Physica–Verlag, Heidelberg 2002, pp. 291–352.
 - [12] A. Marková–Stupňanová: Idempotents of T-addition of fuzzy numbers. *Tatra Mt. Math. Publ.* *12* (1997), 67–72.
 - [13] R. Mesiar: A note to T-sum of L-R fuzzy numbers. *Fuzzy Sets and Systems* *87* (1996), 259–261.
 - [14] B. Schweizer and A. Sklar: *Probabilistic Metric Spaces*. North–Holland, New York 1983.

*Dr. Dug Hun Hong, School of Mechanical and Automotive Engineering, Catholic University of Daegu, Kyungbuk 712–702. South Korea.
e-mail: dhhong@cuth.cataegu.ac.kr*