

# Kybernetika

VOLUME 39 (2003), NUMBER 2

The Journal of the Czech Society for  
Cybernetics and Information Sciences

Published by:

Institute of Information Theory  
and Automation of the Academy  
of Sciences of the Czech Republic

Editor-in-Chief:

Milan Mareš

Managing Editors:

Karel Sladký

Editorial Board:

Jiří Anděl, Marie Demlová, Petr Hájek,  
Martin Janžura, Jan Ježek, Radim Jiroušek,  
George Klir, Ivan Kramosil, Friedrich Liese,  
Jean-Jacques Loiseau, František Matúš,  
Radko Mesiar, Jiří Outrata, Jan Štecha,  
Olga Štěpánková, Igor Vajda, Pavel Zítek,  
Pavel Žampa

Editorial Office:

Pod Vodárenskou věží 4, 182 08 Praha 8

*Kybernetika* is a bi-monthly international journal dedicated for rapid publication of high-quality, peer-reviewed research articles in fields covered by its title.

*Kybernetika* traditionally publishes research results in the fields of Control Sciences, Information Sciences, System Sciences, Statistical Decision Making, Applied Probability Theory, Random Processes, Fuzziness and Uncertainty Theories, Operations Research and Theoretical Computer Science, as well as in the topics closely related to the above fields.

The Journal has been monitored in the Science Citation Index since 1977 and it is abstracted/indexed in databases of Mathematical Reviews, Current Mathematical Publications, Current Contents ISI Engineering and Computing Technology.

---

*Kybernetika*. Volume 39 (2003)

ISSN 0023-5954, MK ČR E 4902.

Published bi-monthly by the Institute of Information Theory and Automation of the Academy of Sciences of the Czech Republic, Pod Vodárenskou věží 4, 182 08 Praha 8. — Address of the Editor: P. O. Box 18, 182 08 Prague 8, e-mail: kybernetika@utia.cas.cz. — Printed by PV Press, Pod vrstevnicí 5, 140 00 Prague 4. — Orders and subscriptions should be placed with: MYRIS TRADE Ltd., P. O. Box 2, V Štíhlách 1311, 142 01 Prague 4, Czech Republic, e-mail: myris@myris.cz. — Sole agent for all “western” countries: Kubon & Sagner, P. O. Box 34 01 08, D-8 000 München 34, F.R.G.

Published in April 2003.

© Institute of Information Theory and Automation of the Academy of Sciences of the Czech Republic, Prague 2003.

## CONTINUOUS EXTENSION OF ORDER-PRESERVING HOMOGENEOUS MAPS

ANDREW D. BURBANKS, COLIN T. SPARROW AND ROGER D. NUSSBAUM

Maps  $f$  defined on the interior of the standard non-negative cone  $K$  in  $\mathbb{R}^N$  which are both homogeneous of degree 1 and order-preserving arise naturally in the study of certain classes of Discrete Event Systems. Such maps are non-expanding in Thompson's part metric and continuous on the interior of the cone. It follows from more general results presented here that all such maps have a homogeneous order-preserving continuous extension to the whole cone. It follows that the extension must have at least one eigenvector in  $K - \{0\}$ . In the case where the cycle time  $\chi(f)$  of the original map does not exist, such eigenvectors must lie in  $\partial K - \{0\}$ .

*Keywords:* discrete event systems, order-preserving homogeneous maps

*AMS Subject Classification:* 93B27, 06F05

### 1. INTRODUCTION

We study classes of maps motivated originally by applications to Discrete Event Systems. We first consider a general setting, and then apply our results to a particular class of maps.

In Section 2, we consider the general case of maps which are defined on the *interior* of a closed cone  $K_1$ , taking values in another closed cone  $K_2$ , which are continuous and order-preserving with respect to the usual partial orderings induced by  $K_1$  and  $K_2$ . We examine the problem of extending these maps, in a natural way, to the whole of  $K_1$ . We give conditions, in considerable generality, (both finite- and infinite-dimensional) under which a natural extension exists and is continuous. (There are also interesting examples in which some of the conditions do not hold and the extension is not everywhere continuous.)

Specifically, we consider order-preserving continuous maps  $f$  from the interior of  $K_1$  to  $K_2$ , where  $K_1$  satisfies a geometrical condition (which generalizes the notion of a polyhedral cone),  $K_2$  satisfies a weak version of normality,  $f$  satisfies a weak version of homogeneity of degree 1. Our results show that all such maps have a continuous extension to the whole of  $K_1$ . We state these general results without proofs; the interested reader is referred to [1].

In Section 3, we examine the case of maps  $f$  defined on the interior of the standard non-negative cone  $K$  in  $\mathbb{R}^N$  which are both homogeneous of degree 1 and order-

preserving. Our results imply that such maps always have a continuous extension. We examine briefly connections between properties of the extension and existence of the cycle-time vector for such maps.

## 2. GENERAL RESULTS

We first give some basic definitions.

**Definition 2.1.** A closed cone (with vertex at zero) in a topological vector space  $X$  is a closed convex subset  $K \subset X$  such that (1)  $K \cap (-K) = \{0\}$  and (2)  $\lambda K \subseteq K$  for all real  $\lambda \geq 0$ . The cone structure induces a partial ordering: We write  $x \leq_K y$  if  $y - x \in K$  (or simply  $x \leq y$ , if  $K$  is obvious from the context). If  $K$  has non-empty interior,  $\overset{\circ}{K} \neq \emptyset$ , we write  $x \ll y$  if  $y - x \in \overset{\circ}{K}$ .

In what follows, we usually let  $K_1$  be a closed cone in a Hausdorff topological vector space  $X_1$  with  $\overset{\circ}{K}_1 \neq \emptyset$  and let  $K_2$  be a closed cone in a topological vector space (t.v.s.)  $X_2$ , and consider maps  $f : \overset{\circ}{K}_1 \rightarrow K_2$  that are both (1) *continuous* and (2) *order-preserving* in the sense that, for all  $x, y \in \overset{\circ}{K}_1$ ,

$$x \leq_{K_1} y \quad \text{implies} \quad f(x) \leq_{K_2} f(y).$$

**Definition 2.2.** When considering the extension of such maps to points  $x \in \partial K_1$ , on the boundary of  $K_1$ , it will be natural to consider sequences  $\langle x_n \in \overset{\circ}{K}_1 : n \geq 1 \rangle$  such that  $x_n \rightarrow x$  as  $n \rightarrow \infty$  and  $x_n \gg x$  for all  $n$ . We will call such sequences *allowable*.

Assuming that  $\overset{\circ}{K}_1 \neq \emptyset$ , if we take any  $u \in \overset{\circ}{K}_1$  and define  $x_n := x + n^{-1}u$ , then  $\lim_{n \rightarrow \infty} x_n = x$  and  $x_n \gg x$  for all  $n \geq 1$  (by convexity). So such a sequence always exists.

### 2.1. Natural extension

Under the assumption (*Condition A*) given below, namely that the images under  $f$  of decreasing sequences must converge, it follows that a natural extension exists.

**Definition 2.3.** Let  $f : \overset{\circ}{K}_1 \rightarrow K_2$  where  $K_1, K_2$  are closed cones in Hausdorff topological vector spaces  $X_1, X_2$ , respectively. Suppose that if  $x_1 \geq x_2 \geq \dots \geq x_k \geq \dots$  is any decreasing sequence in  $\overset{\circ}{K}_1$ , then the sequence  $\langle f(x_j) : j \geq 1 \rangle$  converges in  $K_2$ . If  $f, K_1$ , and  $K_2$  satisfy these conditions, then we shall say that *Condition A is satisfied*.

Condition A holds in some useful cases:

**Lemma 2.4.** Let  $K_1$  be a closed cone in a Hausdorff t.v.s.  $X_1$  with  $\overset{\circ}{K}_1 \neq \emptyset$ . Assume that  $K_2$  is a closed cone in a *finite-dimensional* Hausdorff t.v.s.  $X_2$ . Let  $f : \overset{\circ}{K}_1 \rightarrow K_2$  be a continuous order-preserving map. Then Condition A holds.

Note that there are interesting closed cones  $K$  which are *not* finite-dimensional, but which have the property that decreasing sequences converge. This motivates the following definition.

**Definition 2.5.** Let  $K$  be a closed cone in a Hausdorff t.v.s.  $X$ . We say that  $K$  has the *monotone convergence property* if whenever  $\langle y_j : j \geq 1 \rangle$  is a sequence in  $K$  and  $y_{j+1} \leq y_j$  for all  $j \geq 1$ , there exists  $y \in K$  with  $\lim_{j \rightarrow \infty} y_j = y$ .

**Example 2.6.** Any finite-dimensional closed cone has the monotone convergence property.

**Example 2.7.** Let  $H$  be a real Hilbert space and  $X$  be the set of bounded self-adjoint linear maps  $A : H \rightarrow H$ . Equip  $X$  with the strong operator topology: if  $\langle A_j : j \geq 1 \rangle$  is a sequence in  $X$ , then  $A_j \rightarrow A$  in the strong operator topology if and only if  $\|A_j(x) - A(x)\| \rightarrow 0$  as  $j \rightarrow \infty$  for all  $x \in H$ . Let  $K$  be the cone of nonnegative-definite bounded self-adjoint operators in  $X$ , so that  $\langle Ax, x \rangle \geq 0$  for all  $x \in H$ . It is a standard result that  $K$  has the monotone convergence property (see, for example, [8]).

**Example 2.8.** Let  $(S, \mathcal{M}, \mu)$  be a measure space and let  $X = L^1(S, \mathcal{M}, \mu)$  denote the usual Banach space of  $\mu$ -integrable real-valued maps. Let  $K$  denote the closed cone in  $X$  of maps which are greater than or equal to zero  $\mu$ -almost everywhere (two maps in  $X$  being identified if they agree  $\mu$ -almost everywhere). The monotone convergence theorem from real analysis implies that  $K$  has the monotone convergence property.

We use the monotone convergence property and the previous remarks to generalize Lemma 2.4 to the case where  $X_2$  need not be finite-dimensional:

**Lemma 2.9.** Let  $K_1$  be a closed cone in a Hausdorff t.v.s.  $X_1$  with  $\overset{\circ}{K}_1 \neq \emptyset$ . Let  $K_2$  be a closed cone in a Hausdorff t.v.s.  $X_2$  and suppose that  $K_2$  satisfies the monotone convergence property. Let  $f : \overset{\circ}{K}_1 \rightarrow K_2$  be a continuous order-preserving map. Then Condition A is satisfied.

We now state our first extension theorem:

**Theorem 2.10.** Let  $K_1$  be a closed cone with non-empty interior in a Hausdorff t.v.s.  $X_1$ . Let  $K_2$  be a closed cone in a Hausdorff t.v.s.  $X_2$ . Assume that  $f : \overset{\circ}{K}_1 \rightarrow K_2$  is order-preserving and continuous and that Condition A is satisfied. Suppose that  $x \in K_1$  and that  $\langle x_n \in \overset{\circ}{K}_1 : n \geq 1 \rangle$  is an allowable sequence with  $\lim_{n \rightarrow \infty} x_n = x$ . Then:

(a) There exists  $z = z_x \in K_2$  such that  $\lim_{n \rightarrow \infty} f(x_n) = z_x$ .

(b) Further, if  $\langle y_n \in \overset{\circ}{K}_1 : n \geq 1 \rangle$  is another allowable sequence such that  $\lim_{n \rightarrow \infty} y_n = x$ , then  $\lim_{n \rightarrow \infty} f(y_n) = \lim_{n \rightarrow \infty} f(x_n) = z_x$ .

(c) Further, if we define  $F(x) := z_x$  for all  $x \in K_1$ , then  $F(x) = f(x)$  for all  $x \in \overset{\circ}{K}_1$ , i. e.,  $F$  is an extension of  $f$ . If  $x \in K_1$  and  $\langle x_n \in \overset{\circ}{K}_1 : n \geq 1 \rangle$  is any sequence such that  $\lim_{n \rightarrow \infty} x_n = x$  and  $x_{n+1} \ll x_n$  for all  $n \geq 1$ , then  $\lim_{n \rightarrow \infty} F(x_n) = F(x)$ . (In fact, if  $\langle y_n \in \overset{\circ}{K}_1 : n \geq 1 \rangle$  is any sequence such that  $y_n \geq x$  for all  $n \geq 1$  and  $\lim_{n \rightarrow \infty} y_n = x$ , it follows that  $\lim_{n \rightarrow \infty} F(y_n) = F(x)$ .)

That the above limits all exist and take the same value, namely  $F(x)$  as defined in the theorem, is our intended meaning of the term ‘‘natural’’ extension.

**Remark 2.11.** The extended map  $F$  is also order-preserving since, if  $x, y \in K_1$  and  $x \leq y$ , we may take  $u \in \overset{\circ}{K}_1$  and note that

$$\begin{aligned} F(x) &= \lim_{n \rightarrow \infty} f(x + n^{-1}u) \\ &\leq \lim_{n \rightarrow \infty} f(y + n^{-1}u) = F(y). \end{aligned}$$

Theorem 2.10 also implies that  $F : K_1 \rightarrow K_2$  is continuous at  $0 \in K_1$ . However, as we show in detail in [1], the map  $F$  need not be everywhere continuous, even if  $K_1$  and  $K_2$  are finite-dimensional closed cones.

## 2.2. Continuity of the extension

The aim of this section is to give some further conditions on  $f$ ,  $K_1$ , and  $K_2$ , in considerable generality, which ensure that  $F : K_1 \rightarrow K_2$  is continuous.

We begin with a geometrical condition on  $K_1$  which generalizes the polyhedral property of a cone.

**Definition 2.12.** Let  $K$  be a closed cone in a Hausdorff t.v.s.  $X$ . If  $x \in K$ , we shall say that  $K$  satisfies Condition G at  $x$  if, whenever  $\langle x_n \in K : n \geq 1 \rangle$  is a sequence in  $K$  with  $\lim_{n \rightarrow \infty} x_n = x$  and  $\lambda < 1$ , there exists an integer  $n_*$  such that  $\lambda x \leq x_n$  for all  $n \geq n_*$ . We shall say simply that  $K$  satisfies Condition G if  $K$  satisfies Condition G at  $x$  for all  $x \in K$ .

In [1], we exhibit a map  $f : \overset{\circ}{K}_1 \rightarrow K_2$  where  $K_1$  is a normal cone for which Condition G does not hold. The map is the harmonic mean  $(A^{-1} + B^{-1})^{-1}$  of two

nonnegative-definite symmetric real  $2 \times 2$  matrices  $A, B$ . This is a map from the cone  $C \times C = K_1$  to  $C = K_2$ , where  $C$  is the cone of such matrices. Its extension is order-preserving, homogeneous of degree 1 and continuous on the interior of  $K_1$ , but is *not* continuous at certain points on the boundary  $\partial K_1$ .

**Definition 3.12.** Recall that a closed cone  $K$  in a Hausdorff t.v.s.  $X$  is called *polyhedral* if there exist continuous linear functionals  $\varphi_j : X \rightarrow \mathbb{R}$ ,  $1 \leq j \leq N < \infty$ , such that

$$K = \{x \in X : \varphi_j(x) \geq 0 \text{ for } 1 \leq j \leq N\}.$$

**Lemma 2.14.** Let  $K$  be a closed cone in a Hausdorff t.v.s.  $X$ . It follows that  $K$  is polyhedral if and only if  $K$  is finite-dimensional and satisfies Condition G.

We are grateful to Cormac Walsh, who pointed-out the converse part of the lemma.

The next condition that we will need to ensure continuity of  $F$  is a weak form of *homogeneity*.

**Definition 2.15.** We say that  $f : \overset{\circ}{K}_1 \rightarrow K_2$  is *homogeneous (of degree 1)* if for all  $x \in \overset{\circ}{K}$  and  $\lambda > 0$ ,

$$f(\lambda x) = \lambda f(x).$$

**Definition 2.16.** Let  $K_1$  be a closed cone with  $\overset{\circ}{K}_1 \neq \emptyset$  in a Hausdorff t.v.s.  $X_1$  and let  $K_2$  be a closed cone in a Hausdorff t.v.s.  $X_2$ . Let  $f : \overset{\circ}{K}_1 \rightarrow K_2$ . We say that  $f$  satisfies *Condition WH* at  $x \in K_1$  if, for every real  $\alpha$ ,  $0 < \alpha < 1$ , there exists  $\delta > 0$  and an open neighbourhood  $V$  of  $x$  in  $X_1$  such that for all  $y \in V \cap \overset{\circ}{K}_1$  and for all real  $\lambda \in [1 - \delta, 1]$ ,

$$f(\lambda y) \geq \alpha f(y).$$

If  $f$  satisfies Condition WH at  $x$  for every  $x \in K_1$ , then we shall say simply that  $f$  satisfies *Condition WH* or that  $f$  is *weakly homogeneous*.

**Example 2.17.** If  $f : \overset{\circ}{K}_1 \rightarrow K_2$  is continuous, it is easy to prove that  $f$  satisfies Condition WH at every  $x \in \overset{\circ}{K}_1$  such that  $f(x) \in \overset{\circ}{K}_2$ .

**Example 2.18.** If there exists  $\delta_* > 0$  and a map  $\varphi : [1 - \delta_*, 1] \rightarrow (0, \infty)$  with  $\lim_{\lambda \rightarrow 1^-} \varphi(\lambda) = 1$  such that

$$f(\lambda y) \geq \varphi(\lambda) f(y),$$

for all  $y \in \overset{\circ}{K}_1$  and  $\lambda \in [1 - \delta_*, 1]$ , then  $f$  satisfies Condition WH.

The final condition that we shall need is a variant of *normality* for the cone  $K_2$ . We call this the *weak normality* (WN) or *sandwich* condition.

**Definition 2.19.** A cone  $K$  in a normed linear space, not necessarily finite-dimensional, is said to be *normal* if there exists a constant  $M$  such that  $\|x\| \leq M\|y\|$  whenever  $0 \leq_K x \leq_K y$ .

**Definition 2.20.** Let  $K_2$  be a closed cone in a Hausdorff t.v.s.  $X_2$ . We shall say that  $K_2$  *satisfies Condition WN* (or *is weakly normal*) if, whenever  $\langle x_n \rangle$ ,  $\langle y_n \rangle$ , and  $\langle z_n \rangle$  are sequences in  $K_2$  with  $0 \leq x_n \leq y_n \leq z_n$  for all  $n \geq 1$  and  $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} z_n = y$ , for some  $y \in K_2$ , it follows that  $\lim_{n \rightarrow \infty} y_n$  exists and equals  $y$ .

**Lemma 2.21.** Let  $K_2$  be a closed cone in a normed linear space. Then  $K_2$  is normal if and only if  $K_2$  satisfies Condition WN.

We now state our second extension theorem; we use the above additional conditions to ensure that the extended map  $F$  is sequentially continuous:

**Theorem 2.22.** Let  $K_1$  be a closed cone with  $\overset{\circ}{K}_1 \neq \emptyset$  in a Hausdorff t.v.s.  $X_1$ ,  $K_2$  a closed cone in a Hausdorff t.v.s.  $X_2$  and  $f : \overset{\circ}{K}_1 \rightarrow K_2$  a continuous order-preserving map. Assume that Condition A is satisfied, that  $K_1$  satisfies Condition G at some  $x \in \partial K_1$ , that  $f$  satisfies Condition WH at  $x$  and that  $K_2$  satisfies Condition WN. Define the extension  $F$  as in Theorem 2.10. Then  $F$  is sequentially continuous at  $x$ .

**Corollary 2.23.** Let  $K_1$  be a closed polyhedral cone with  $\overset{\circ}{K}_1 \neq \emptyset$  in a Hausdorff t.v.s.  $X_1$ . Let  $K_2$  be a closed cone in a Hausdorff t.v.s.  $X_2$  and assume either that (i)  $K_2$  is finite-dimensional or, more generally, that (ii)  $K_2$  has the monotone convergence property and satisfies Condition WN. Let  $f : \overset{\circ}{K}_1 \rightarrow K_2$  be continuous and order-preserving and satisfy Condition WH on  $K_1$ . Then  $f$  has a sequentially continuous extension  $F : K_1 \rightarrow K_2$  that is order-preserving and satisfies Condition WH on  $K_1$ .

### 3. THE POSITIVE CONE

In this section, we focus on maps  $f$  defined on the interior of the standard non-negative cone  $K = \mathbb{R}_+^N$  in  $\mathbb{R}^N$  which are both homogeneous of degree 1 and order-preserving. In this case, the partial ordering induced by the cone is exactly the usual partial product ordering on  $\mathbb{R}^N$ . These maps arise naturally in the study of certain classes of Discrete Event Systems. Such maps are non-expanding in Thompson's part metric and continuous on the interior of the cone, so that continuity is no longer required as an explicit assumption.

First, we work in a more general setting. Recall the definition of Thompson's part metric:

**Definition 3.1.** Let  $v, y$  denote elements of a cone  $K$ . We say that  $v$  and  $y$  are comparable if there exist reals  $\alpha > 0$  and  $\beta > 0$  with  $\alpha y \leq_K v \leq_K \beta y$ . The notion of comparability divides  $K$  into disjoint equivalence classes called the *components* or *parts* of  $K$ ; for  $v \in K$  we let  $K(v)$  denote the set of points that are comparable with  $v$ .

**Definition 3.2.** Let  $v, y$  denote comparable elements of a cone  $K$ . We define the positive real quantities,

$$\begin{aligned} M(v/y) &:= \inf\{\beta > 0 : v \leq \beta y\}, \\ \bar{d}(v, y) &:= \max\{\log M(v/y), \log M(y/v)\}. \end{aligned}$$

The restriction of  $\bar{d}$  to each part  $K(v)$  is a metric, called *Thompson's (part) metric*. By defining  $\bar{d}(v, y) := +\infty$  if  $v, y$  are non-comparable elements of  $K$ , we extend  $\bar{d}$  to the whole of  $K$  (but only the restrictions to each part are actually metrics).

**Remark 3.3.** Let  $K_1$  be a normal cone with  $\overset{\circ}{K}_1 \neq \emptyset$  in a Banach space  $X_1$  and  $K_2$  be a normal cone in a Banach space  $X_2$ . Let  $\bar{d}_1$  denote the part metric on  $\overset{\circ}{K}_1$ . It is known that  $\bar{d}_1$  gives the same topology on  $\overset{\circ}{K}_1$  as the topology induced by the norm on  $X_1$ . See Proposition 1.1 in [7] for a proof and references to the literature. See also [9]. If  $f : \overset{\circ}{K}_1 \rightarrow K_2$  is homogeneous of degree 1 and order-preserving, then there exists  $v \in K_2$  such that  $f(x)$  is comparable to  $v$  in  $K_2$  for all  $x \in \overset{\circ}{K}_1$  (i. e., the interior maps to a single part). If  $K_2(v)$  is the part corresponding to  $v$  and  $\bar{d}_2$  is the part metric on  $K_2(v)$ , then

$$\bar{d}_2(f(x_1), f(x_2)) \leq \bar{d}_1(x_1, x_2),$$

for all  $x_1, x_2 \in \overset{\circ}{K}_1$ , i. e.,  $f$  is non-expanding on  $\overset{\circ}{K}_1$  in the part metric. It follows automatically that  $f$  is continuous as a map from  $\overset{\circ}{K}_1$  to  $K_2$ . Thus, in this case we do not need to assume explicitly that  $f : \overset{\circ}{K}_1 \rightarrow K_2$  is continuous.

### 3.1. Topical maps on the positive cone

Let  $K_1 = K_2 = K := \mathbb{R}_+^N$ , the standard positive cone in  $\mathbb{R}^N$ . Maps  $f : \overset{\circ}{K} \rightarrow \overset{\circ}{K}$  that are order-preserving and homogeneous of degree 1 are a specific example of the general case above; they are non-expanding with respect to the part metric on  $\overset{\circ}{K}$ . In fact, if homogeneity holds, then being order-preserving is equivalent to being non-expanding, see [3]. Hence these maps are continuous on  $\overset{\circ}{K}$ . Such maps are sometimes called “topical” in the literature and are of interest for certain classes of discrete event systems. They may be viewed as the image under the bijection (component-wise exponentiation)  $\exp : \mathbb{R}^N \rightarrow \overset{\circ}{K}$  of maps  $g : \mathbb{R}^N \rightarrow \mathbb{R}^N$  that are *additively* homogeneous and order-preserving and, consequently, non-expanding in



the supremum ( $\ell_\infty$ ) norm. To each such map  $g$  there corresponds a map  $f : \overset{\circ}{K} \rightarrow \overset{\circ}{K}$  with

$$f(x) = (\exp \circ g \circ \log)(x).$$

With suitable modifications of the proofs to take advantage of full, rather than weak, homogeneity, our results imply the following corollary. This result was proved previously, by more direct means involving the comparison of various component-wise limits, by [2].

**Corollary 3.4.** All homogeneous order-preserving maps  $f : \overset{\circ}{K} \rightarrow \overset{\circ}{K}$  with  $K = \mathbb{R}_+^N$  have an extension  $F : K \rightarrow K$  that is homogeneous, order-preserving, and continuous on the whole of  $K$ .

**Remark 3.5.** We can also show, see [2], that the extended map  $F$  is non-expanding in the extended part metric, mapping each part of  $K$  to a part.

**Corollary 3.6.** Let  $f : \overset{\circ}{K} \rightarrow \overset{\circ}{K}$ , with  $K = \mathbb{R}_+^N$ , be an order-preserving homogeneous map with the order-preserving homogeneous (and continuous) extension  $F : K \rightarrow K$ . Then  $F$  has at least one eigenvector in  $K - \{0\}$ .

*Proof.* Let  $\Pi \subset K$  denote the intersection of the positive cone with the surface of the ( $\ell_2$ ) unit hyper-sphere,

$$\Pi := \{x \in K : \|x\|_2 = 1\},$$

and let  $\pi$  denote the projection (normalization),

$$\pi : K - \{0\} \rightarrow \Pi, \quad x \mapsto \frac{x}{\|x\|_2}.$$

We have seen that  $f$  has a continuous extension  $F$  to the whole cone  $K$ . If  $F(x) = 0$  for some  $x \in \Pi$  then, by definition,  $F$  has an eigenvector with eigenvalue 0 and we are done. This happens, for example, in the case  $f : \overset{\circ}{\mathbb{R}}_+^2 \rightarrow \overset{\circ}{\mathbb{R}}_+^2$  with  $f(x_1, x_2) := \sqrt{x_1 x_2}(1, 1)$ , where all  $x \in \partial\Pi$  are mapped to the vertex 0. Suppose, on the other hand, that  $F(x) \neq 0$  for all  $x \in \Pi$ , then the projected map  $\pi \circ F : \Pi \rightarrow \Pi$  is well-defined and continuous on  $\Pi$ . Further,  $\Pi$  is homeomorphic to a compact convex set. Hence, by Brouwer's fixed point theorem,  $\pi \circ F$  has at least one fixed point in  $\Pi$ . By homogeneity, it follows that  $F$  itself must have at least one eigenvector in  $K - \{0\}$ .  $\square$

In fact, we can prove a stronger result:

**Lemma 3.7.** Let  $C$  be a closed cone with non-empty interior in a finite dimensional Banach space  $X$ . Let  $F : C \rightarrow C$  be a continuous order-preserving homogeneous (of degree one) map that maps the interior of  $C$  into itself. Then  $F$  has an eigenvector in  $C - \{0\}$  with nonzero eigenvalue.

*Proof.* (Outlined.) If  $u \in \overset{\circ}{C}$ , there exists  $c > 0$  such that  $F(u) \geq cu$ . Take  $a > 1$  and define  $g(x) := (a/c)F(x)$ , so that  $g(u) \geq au$  and the sequence  $\langle g^m(u) : m \geq 1 \rangle$  is unbounded in norm. Theorem 2.1 in [6] then implies that there exists  $x \in C$  with  $\|x\| = 1$  and  $t \geq 1$  such that  $g(x) = tx$ . (The assumption of compactness of the map  $g$ , in the statement of that theorem, can be weakened.)

It follows that  $F(x) = (tc/a)x$ . A simple limiting and compactness argument, letting  $a \rightarrow 1$ , shows that  $F$  has an eigenvector with eigenvalue not less than  $c$ . In fact, if we define

$$c := \sup\{k : \exists x \in C, \|x\| = 1, F(x) \geq kx\},$$

then we deduce that there exists an eigenvector of  $F$  with eigenvalue  $c$ . □

**Definition 3.8.** From the viewpoint of applications, a natural question is whether the map  $f : \overset{\circ}{K} \rightarrow \overset{\circ}{K}$  has a *cycle-time vector*, defined formally by

$$\chi(f) := \lim_{k \rightarrow \infty} (f^k(x))^{1/k}.$$

If this limit exists for some  $x \in \overset{\circ}{K}$ , then it follows, from the fact that  $f$  is non-expanding, that it exists for all  $y \in \overset{\circ}{K}$  and takes the same value everywhere. Thus the cycle time is naturally regarded as a property of the map itself.

Existence of an eigenvector  $x \in \overset{\circ}{K}$  in the interior with, say,  $f(x) = \lambda x$  for some  $\lambda > 0$ , implies directly the existence of the cycle time with  $\chi(f) = \lambda \mathbf{1}$ , where  $\mathbf{1} := (1, 1, \dots, 1)$ . Thus our above result establishes the following corollary:

**Corollary 3.9.** If  $\chi(f)$  does not exist, for  $f : \overset{\circ}{K} \rightarrow \overset{\circ}{K}$  homogeneous and order-preserving on the positive cone  $K$ , then the extended map  $F$  has at least one eigenvector in  $\partial K - \{0\}$  with nonzero eigenvalue and there are no eigenvectors in  $\overset{\circ}{K}$ .

The cycle-time vector is known to exist for certain classes of maps in general dimension  $N$ . Specifically, a nonlinear hierarchy of such maps may be built from simple maps by closure under a finite set of operations, see [4]. The cycle-time also exists for all order-preserving homogeneous maps with  $N = 1, 2$ .

However,  $\chi(f)$  need not exist in general for  $N \geq 3$ , as illustrated by a family of maps introduced by [5]: consider a sequence of reals  $\langle a^k \in [0, 1] : k \geq 1 \rangle$  and let

$\sigma^k := \sum_{j=1}^k a^j$  with  $\sigma^0 := 0$ . Then there exists a homogeneous order-preserving map  $f : \overset{\circ}{K} \rightarrow \overset{\circ}{K}$  with  $K := \mathbb{R}_+^3$ , such that for all  $k \geq 0$ ,

$$f^k(1, 1, 1) = (1, \exp(\sigma^k), \exp(k)).$$

For suitable choices of the sequence  $\langle a^k \rangle$ , we can arrange that the sequence  $\langle \sigma^k/k \rangle$  does not converge and, hence, that  $\chi(f)$  does not exist.

Construction of a particular family of such maps, for which there is no cycle time, reveals that the projected map  $\pi \circ F$  fixes a continuum of points on one edge of  $\Pi$ .

#### 4. CONCLUSION

We have presented results on the continuous extension of maps defined on the interior of closed cones, firstly in considerable generality and secondly applied to the case of the standard positive cone.

Our general results have applications in other areas, particularly to operator-valued means, where the existence of continuous extensions is a natural question. In this case, it is also relevant to consider maps which are not order-preserving. For further discussions, see [1].

For the positive cone, we have shown that all homogeneous order-preserving maps have a continuous extension to the boundary of the cone and that non-existence of the cycle time restricts the fixed-point set of the extended map. It would be interesting to have a characterization of the possible fixed-point sets and to clarify connections between existence of a cycle time and properties of these sets.

#### ACKNOWLEDGEMENTS

During this work, A. Burbanks was supported partially by EPSRC GR/L42742, by the research network ALAPEDES, and by BRIMS (Hewlett-Packard Laboratories, Bristol, UK). R. Nussbaum supported partially by NSF DMS 0070829.

(Received April 23, 2002.)

#### REFERENCES

- 
- [1] A. D. Burbanks, R. D. Nussbaum, and C. T. Sparrow: Extension of order-preserving maps on a cone. *Proc. Roy. Soc. Edinburgh Sect. A* 133 (2003), 35–59.
  - [2] A. D. Burbanks and C. T. Sparrow: All Monotone Homogeneous Functions (on the Positive Cone) Admit Continuous Extension. Technical Report No. 1999-13, Statistical Laboratory, University of Cambridge 1999.
  - [3] M. G. Crandall and L. Tartar: Some relations between nonexpansive and order preserving mappings. *Proc. Amer. Math. Soc.* 78 (1980), 385–390.
  - [4] S. Gaubert and J. Gunawardena: A Nonlinear Hierarchy for Discrete Event Systems. Technical Report No. HPL-BRIMS-98-20, BRIMS, Hewlett-Packard Laboratories, Bristol 1998.
  - [5] J. Gunawardena and M. Keane: On the Existence of Cycle Times for Some Nonexpansive Maps. Technical Report No. HPL-BRIMS-95-003, BRIMS, Hewlett-Packard Laboratories, Bristol 1995.

- [6] R. D. Nussbaum: Eigenvectors of Nonlinear Positive Operators and the Linear Krein–Rutman Theorem (Lecture Notes in Mathematics 886). Springer Verlag, Berlin 1981, pp. 309–331.
- [7] R. D. Nussbaum: Finsler structures for the part-metric and Hilbert’s projective metric, and applications to ordinary differential equations. *Differential and Integral Equations* 7 (1994), 1649–1707.
- [8] F. Riesz and B. Sz.-Nagy: *Functional Analysis*. Frederick Ungar Publishing Company, New York 1955.
- [9] A. C. Thompson: On certain contraction mappings in a partially ordered vector space. *Proc. Amer. Math. Soc.* 14 (1963), 438–443.

*Dr. Andrew D. Burbanks, Laboratory for Advanced Computation in the Mathematical Sciences, School of Mathematics, University of Bristol, University Walk, Bristol BS8 1TW. United Kingdom.*

*e-mail: A.Burbanks@bristol.ac.uk*

*Dr. Colin T. Sparrow, Mathematical Institute, University of Warwick, Coventry CV4 7AL. United Kingdom.*

*e-mail: csparrow@maths.warwick.ac.uk*

*Prof. Dr. Roger D. Nussbaum, Department of Mathematics, Hill Center, Rutgers University, 110 Frelinghuysen Road, Piscataway, NJ 08854-8019. U.S.A.*

*e-mail: nussbaum@math.rutgers.edu*