



Contributions to Soil Zoology in Central Europe I

Proceedings of the 7th Central European Workshop on Soil Zoology

held in České Budějovice, Czech Republic, April 14-16, 2003

Edited by

Karel Tajovský, Jiří Schlaghamerský & Václav Pižl

Institute of Soil Biology, Academy of Sciences of the Czech Republic,
České Budějovice, Czech Republic

2005

ISBN 80-86525-04-X

Content

Epigaeic activity of eight common Collembola species in spruce forests and clearings after bark beetle outbreaks in the National Park Šumava, Czech Republic (Brůhová J. & Rusek J.) ♦ Dispersal of *Micranurida sensillata* (Collembola, Neanuridae) by feral domestic pigeons (Christian E.) ♦ Mites (Acarina, Gamasida) in littoral zone of Jakubov fishponds (Slovakia) (Fend'a P. & Schniererová E.) ♦ Energetic budget of three species of bibionid larvae (Frouz J., Šustr V. & Kalčík J.) ♦ Nematodes in clay colliery spoil dumps and experimentally introduced strips of meadow soil (Háněl L.) ♦ Using actinomycete-specific PCR primers to characterize the bacterial communities of wormcasts (Hill P., Krištúfek V., Feijoo A.M. & Gallego G.) ♦ Ants (Hymenoptera, Formicidae) as an important part of the epigaeic fauna in Carpathian oak-hornbeam forests (Holecová M., Lukáš J., Havířová M. & Harakařová E.) ♦ Can earthworm gut passage modify clay minerals structure: preliminary results (Hušák J., Frouz J. & Had J.) ♦ Ruins of medieval castles as refuges of interesting land snails in the landscape (Juřičková L. & Kučera T.) ♦ Niche differentiation among epigaeic Collembola in primeval Carpathian beech and ash forests in Uzhans'ky National Park (Ukraine) (Kapruš I.J. & Steržinská M.) ♦ Abundance and diversity of soil Collembola (Hexapoda) assemblages in the Zádielska Valley, Slovak Karst (Slovakia) (Kostúrová N., Kováč L. & Miklisová N.) ♦ Terrestrial arthropods of the Domica Cave system and the Ardovská Cave (Slovak Karst) - principal microhabitats and diversity (Kováč L., Mock A., Luptáčik P., Košel V., Fend'a P., Svatoň J. & Mašán P.) ♦ Does the cellular fatty acid and enzyme content of cave bacteria affect the feeding preferences of *Enchytraeus crypticus* (Oligochaeta, Enchytraeidae)? (Krištúfek V., Elhotová D., Šustr V., Lasák R., Kováč L., Mock A. & Luptáčik P.) ♦ Floods – as the factor of degradation and recovery of araneocoenoses (Krumpálová Z.) ♦ Soil oribatid mite communities (Acari: Oribatida) across a terrain depression in an arable field in the East-Slovakian Lowland (Luptáčik P. & Miklisová D.) ♦ Structure and seasonal dynamics of

oribatid communities (Acari: Oribatida) inhabiting moss covers on granite boulders in a mountain spruce forest (Materna J.) ♦ Terrestrial arthropods inhabiting caves near Veľký Folkmar (Čierna hora Mts., Slovakia) (Mock A., Luptáčik P., Fend'a P., Svatoň J., Országh I. & Krumpál M.) ♦ Interesting and rare saprotrophic microfungi isolated from excrements and other substrata in the Domica and Ardovská Caves (Slovak Karst National Park, Slovakia) (Nováková A.) ♦ Feeding sources of invertebrates in Ardovská cave and Domica cave systems – preliminary results (Nováková A., Elhotová D., Krištúfek V., Lukešová A., Hill P., Kováč L., Mock A. & Luptáčik P.) ♦ Earthworms in an arable field-forest ecotone (Pižl V. & Zeithaml J.) ♦ Subterranean traps in two high alpine habitats (Querner P. & Gereben-Krenn B.-A.) ♦ Protura and Diplura succession on chemical factory deposits (Rusek J.) ♦ The community of small annelids (Enchytraeidae, Tubificidae, Aeolosomatidae) in the first phase of meadow restoration on arable land (Schlaghamerský J.) ♦ The effect of lead and zinc on the moss mite *Archegozetes longisetosus* Aoki (Acari, Oribatida) under laboratory conditions (Seniczak A., Seniczak S. & Dlugosz J.) ♦ Soil Gamasida (Acari) of the ecotone between the Scots pine and birch forests in Tuchola Forest, Poland (Seniczak S., Kaczmarek S. & Seniczak A.) ♦ Old mine underground galleries as the habitat for mites (Acari) (Skubala P., Madej G., Solarz K. & Klys G.) ♦ Two dwellers of one habitat - two strategies for the different microhabitats (Smrž J.) ♦ Influence of grass sowing on oribatid mite communities (Acari: Oribatida) in initial successive stages during rich meadow recovery (Starý J.) ♦ Species representation of Carabidae (Coleoptera) registered in agricultures of the school farm in Telč (Šťastná P. & Bezděk J.) ♦ Check List of Collembola in the Nature Reserve Stroomdalladschap Drentsche Aa (Netherlands) (Sterzyńska M. & Ehrnsberger R.) ♦ Epigaeic spiders (Araneae) of non-forest and forest habitats of the Poloniny National Park, Slovakia (Svatoň J., Gajdoš P. & Krumpálová Z.) ♦ Development of soil fauna in meadows restored on arable land – Initial phases of successional development (Tajovský K., Pižl V., Starý J., Balík V., Frouz J., Schlaghamerský J., Háněl L. & Rusek J.) ♦ Notes to ecological characteristic of chosen protozoan groups in leaf-litter and mineral layer of soil (Tirjaková E. & Mrva M.) ♦ Heat-extractor – an indispensable tool for soil zoological studies (Tuf I.H. & Tvardík D.) ♦ Survival under water – comparative study of millipedes (Diplopoda), centipedes (Chilopoda) and terrestrial isopods (Oniscidea) (Tufová J. & Tuf I.H.) ♦ To what extent do different farming systems influence biodiversity and composition of collembolan communities? (Uteseny K.) ♦ New digitizing tools (digital microscope and HP photo scanner) and their use in biological laboratory (Weyda F.) ♦ Centipedes in the spruce forests of the Moravskoslezské Beskydy Mountains, Czech Republic (Wytrwal J., Tajovský K.) ♦ Nematode community structure in horse-grazed and ungrazed dry grass meadows in eastern Austria (Zolda P.) (221 pages in total)