

Studies on Soil Fauna in Central Europe

Proceedings of the 6th Central European Workshop on Soil Zoology

held in České Budějovice, Czech Republic, April 23-25, 2001

Edited by

Karel Tajovský, Vladimír Balík & Václav Pižl


Institute of Soil Biology, Academy of Sciences of the Czech Republic,
České Budějovice, Czech Republic

2002

ISBN 80-86525-00-7

Content

Balík, V.: Testate amoebae (Protozoa) in soils of restored flowery meadows in the Bílé Karpaty Mts. (Czech Republic) ■ Błoszyk, J., Bajerlein, D., Skoracka, A. & Bajaczyk, R.: *Uropoda orbicularis* (Müller, 1776) (Acari: Uropodina) as an example of a mite adapted to synanthropic habitats ■ Bryja, V. & Kula, E.: Succession of epigeic spider communities in substitute birch forests between 1988 and 1998 ■ Christian, E.: Copepoda from forest floor habitats in Vienna ■ Dmowska, E. & Makulec-Ilieva, K.: Influence of antimicrobial feed additives on nematode community structure: Primary field experiment ■ Elhottová, D., Frouz, J., Křišťůfek, V., Lukešová, A., Nováková, A. & Tříška, J.: Potential sources of polyunsaturated fatty acids for saprophagous soil invertebrates ■ Fojtová, H.: The influence of water regime on soil-surface fauna in floodplain forest ■ Háněl, L.: Fauna of soil nematodes and other soil micro-mesofauna in spruce clearings in the Šumava Mts., Czech Republic ■ Háněl, L.: Short-term effects of grass mixtures on nematodes inhabiting agricultural soil in the Bílé Karpaty Mts., Czech Republic - a preliminary study ■ Holecová, M. & Sukupová, J.: Weevils (Coleoptera, Curculionoidea) as a part of the oak-hornbeam forest epigaeon ■ Hubert, J.: Abundant and frequent oribatid mites (Acari: Oribatida) in different plots on two meadows ■ Košel, V.: What do we know on terrestrial Tricladida (Turbellaria) from the former Czechoslovakia? ■ Kováč, L., Magdová, J. & Miklisová, D.: Introductory monitoring of Collembola (Hexapoda) in the Pieniny Mts. (Western Carpathians) ■ Kováč, L., Mock, A., Luptáček, P. & Palacios-Vargas, J.G.: Distribution of *Eukoenia spelaea* (Peyerimhoff, 1902) (Arachnida, Palpigradida) in the Western Carpathians with remarks on its biology and behaviour ■ Krivolutsky, D.A. & Lebedeva, N.V.: The oribatid mites in the bird feathers ■ Krumpálová, Z. & Bartoš, D.: Lycosid spiders (Araneae) of the oak forests of the Malé Karpaty Mts. near Modra ■ Luptáček, P. & Miklisová, D.: Soil Oribatida (Acarina) communities across a land depression in an arable field ■ Madej, G., Błoszyk, J., Stachowiak, M. & Bajaczyk, R.: Structure of pioneer communities of mesostigmatid mites (Arachnida, Acari) at initial stages of ecological succession on coal mining ■ Mock, A. & Tajovský, K.: Genus *Melogona* Cook, 1895 (Diplopoda: Chordeumatida: Chordeumatidae) in Slovakia ■ Nováková, A. & Pižl, V.: Mycotic flora in vermicultures and intestines of *Eisenia andrei* ■ Pižl, V.: Earthworm communities in spruce forest soils – effects of some natural and anthropogenic factors ■ Pokarzhevskii, D., Van Straalen, N.M. & Zaboev, D.P.: New approaches to the trophic structure of soil ecosystems ■ Rusek, J.: Impact of boric acid on Collembola, Protura and Pauropoda in a mountain elm forest in the Šumava National Park, Czech Republic ■ Skoracka, A., Skoracki, M., Błoszyk, J. & Stachowiak, M.: The seasonal abundance of



Mesostigmata (Acari) in compost ■ Skoracka, A., Skoracki, M., Błoszyk, J. & Stachowiak, M.: The structure of dominance and frequency of Mesostigmata (Acari) in compost ■ Skubała, P.: Development of oribatid mite fauna (Acari, Oribatida) in a sedimentation tank ■ Smrž, J.: The excrement analysis - the useful tool for the biological and autecological studies in soil zoology ■ Šobotník, J. & Weyda, F.: Ontogeny of tarsal glands in *Reticulitermes lucifugus* (Isoptera: Rhinotermitidae) ■ Starý, J.: Changes of oribatid mite communities (Acari: Oribatida) during primary succession on colliery spoil heaps near Sokolov, North-West Bohemia, Czech Republic ■ Sterzyńska, M.: Collembola of the hydrogenic soils of Biebrza River valley ■ Szymkowiak, P.: The soil mites Gamasina (Acari, Gamasida) in the Carpathian beech forest: abundance, spatial organization and influence of two environmental factors ■ Tajovský, K.: Soil macrofauna (Diplopoda, Chilopoda, Oniscidea) in a pine forest disturbed by wildfire ■ Tirjaková, E., Mrva, M. & Hlúbiková, D.: Ciliophora nad Rhizopoda in soil, leaf-litter and mosses of oak-hornbeam forests in the Malé Karpaty Mts. (Western Slovakia) ■ Tuf, I.H.: Contribution to the knowledge of vertical distribution of soil macrofauna (Chilopoda, Oniscidea) in floodplain forest * Tufová, J.: Development of millipede assemblages in floodplain forests after summer flood ■ Urbánková, M. & Schlaghamerský, J.: The impact of inundation on floodplain forest Brachycera (Diptera) associated with oak logs ■ Weyda, F.: Scientific digital photography and its application to modern zoological research ■ Weyda, F., Šobotník, J. & Nebesářová, J.: Ontogeny of fat body in *Prorethinosia simplex* (Isoptera: Rhinotermitidae) ■ Krysiak, D., Błoszyk, J. & Bajaczyk, R.: The influence of an urban agglomeration on the dispersion and the abundance of the mites of the suborder Uropodina (Acari: Mesostigmata) in Poznań (285 pages in total)