

V. MÜLLER, PRAGUE

Let  $A$  be a linear associative algebra. By [3] it is always possible to define a topology on  $A$  which makes of  $A$  a locally convex algebra with separately continuous multiplication (i.e.  $x_\alpha, x, y \in A, x_\alpha \rightarrow x$  implies  $x_\alpha y \rightarrow xy, yx_\alpha \rightarrow yx$ ).

On the other hand (cf. [3]) in general it is not possible to introduce a topology on  $A$  which makes of  $A$  a locally convex algebra with jointly continuous multiplication (i.e.  $x_\alpha \rightarrow x, y_\beta \rightarrow y \Rightarrow x_\alpha y_\beta \rightarrow xy$ ). The aim of this note is to exhibit two examples which continue these investigations.

In the first example we construct a commutative algebra which admits no topology. This gives a negative answer to the question raised in [2]. In the second example we construct a topological algebra which admits no locally convex topology.

All algebras in this paper will be complex (this condition, however, is not essential).

We say that an algebra  $A$  is topologizable (topologizable as a locally convex algebra) if there exists a topology on  $A$  which makes of  $A$  a topological (locally convex) algebra with jointly continuous multiplication.

It is easy to see that an algebra  $A$  is topologizable if and only if there exists a system  $\mathcal{V}$  of subset of  $A$  (zero-neighbourhoods in  $A$ ) satisfying

- (1)  $\bigcap_{V \in \mathcal{V}} V = \{0\}$
- (2)  $\lambda V \subset V$  for every  $v \in \mathcal{V}$  and complex number  $\lambda, |\lambda| \leq 1$
- (3) each  $V \in \mathcal{V}$  is absorbent
- (4) for every  $V \in \mathcal{V}$  there exists  $W \in \mathcal{V}$  such that  $W + W \subset V$
- (5) for every  $W \in \mathcal{V}$  there exists  $W \in \mathcal{V}$  such that  $W \cdot W \subset V$ .

For basic properties of topological algebras see e.g. [1].

**THEOREM 1.** *There exists a commutative algebra which is not topologizable.*

**PROOF.** Denote by  $N$  the set of all positive integers and by  $\mathcal{F}$  the set of all sequences  $f = \{f_j\}_{j=1}^\infty$  of positive integers. Consider the linear space  $A$  of all formal linear combinations of elements  $c, x_i$  ( $i \in N$ ) and  $a_f$  ( $f \in \mathcal{F}$ ). We define the multiplication in  $A$  by

$$\begin{aligned} cz = zc = 0 & & \text{for every } z \in A, \\ x_i x_j = 0 & & (i, j \in N), \\ a_f a_{f'} = 0 & & (f, f' \in \mathcal{F}), \\ x_n a_f = a_f x_n = f_n \cdot c & & (n \in N, f \in \mathcal{F}). \end{aligned}$$

Clearly these relations define uniquely a multiplication on  $A$  which makes of  $A$  a commutative algebra (for the associative law note that the product of any three of the basis elements is equal to zero).

We prove that  $A$  is not topologizable. Suppose on the contrary that there exists a system  $\mathcal{V}$  of zero-neighbourhoods in  $A$  satisfying (1) - (5). Let  $V, W \in \mathcal{V}$  satisfy  $c \notin V$  and  $W \cdot W \subset V$ .

For  $n = 1, 2, \dots$  choose  $s_n > 0$  such that  $x_n \in s_n \cdot W$ . Let  $f = \{f_n\}_{n=1}^{\infty}$  be a sequence of positive integers  $f_n$  with  $f_n > n \cdot s_n$ . Then  $a_f \in r \cdot W$  for some  $r > 0$ . We have

$$c = \frac{1}{f_n}(x_n \cdot a_f) = \frac{r \cdot s_n}{f_n} \left( \frac{x_n}{s_n} \cdot \frac{a_f}{r} \right) \in \frac{r s_n}{f_n} \cdot W \cdot W \subset \frac{r s_n}{f_n} V.$$

Since  $c \notin V$  we have

$$\frac{r s_n}{f_n} > 1 \quad \text{and} \quad r > \frac{f_n}{s_n} > n \quad (n \in N),$$

a contradiction.

*Remark.* Let  $X$  be a linear space of infinite dimension and let  $\mathcal{L}(X)$  be the algebra of all linear mappings acting in  $X$ . By [3],  $\mathcal{L}(X)$  can not be topologized as a locally convex algebra. Using analogous method as in example 1 it is possible to show that  $\mathcal{L}(X)$  is not topologizable. In fact even the algebra of all finite-dimensional operators in  $X$  is not topologizable.

**THEOREM 2.** *There exists a commutative topological algebra which is not topologizable as a locally convex algebra.*

**PROOF.** Let  $K$  be an uncountable set. Denote by  $\mathcal{D}$  the set of all functions  $d : N \times K \rightarrow N$ . For  $d \in \mathcal{D}, n \in N$  and  $k \in K$  we shall write shortly  $d_{nk}$  instead of  $d(n, k)$ .

Clearly for every  $d \in \mathcal{D}$  and  $n \in N$  there exists a subset  $K_{d,n} \subset K$  and a positive integer  $d_n$  such that  $\text{card}K_{d,n} = d_n$  and  $d_{nk} = d_n$  for every  $k \in K_{d,n}$ . Let  $A$  be the linear space of all (finite) linear combinations of elements  $c, x_{nk}$  ( $n \in N, k \in K$ ),  $a_d$  ( $d \in \mathcal{D}$ ) and  $y_{dnk}$  ( $d \in \mathcal{D}, n \in N, k \in K_{d,n} \subset K$ ).

We define the multiplication in  $A$  by

$$\begin{aligned} cz = zc &= 0 & (z \in A), \\ y_{dnk}z &= zy_{dnk} = 0 & (z \in A, d \in \mathcal{D}, n \in N, k \in K_{d,n}), \\ a_d a_{d'} &= 0 & (d, d' \in \mathcal{D}), \\ x_{nk} \cdot x_{n'k'} &= 0 & (n, n' \in N, k, k' \in K), \\ x_{nk} \cdot a_d &= a_d \cdot x_{nk} = \begin{cases} d_n y_{dnk} & (d \in \mathcal{D}, n \in N, k \in K_{d,n}) \\ 0 & (k \notin K_{d,n}). \end{cases} \end{aligned}$$

Clearly  $A$  is a commutative algebra. To define the topology on  $A$  we shall need the following notations:

Let  $\mathcal{L}$  be the set of all complex valued functions  $\lambda : k \mapsto \lambda_k$  defined on  $K$  with a finite support. For  $\lambda \in \mathcal{L}$  and  $i \in \{0, 1, 2, \dots\}$  define

$$m_i(\lambda) = \min_{\substack{M \subset K \\ \text{card}M=i}} \max_{j \in K-M} |\lambda_j|.$$

Clearly  $\max_{j \in K} |\lambda_j| = m_0(\lambda) \geq m_1(\lambda) \geq \dots$  and  $\text{card}\{j \in K, |\lambda_j| > m_i(\lambda)\} \leq i$ .

LEMMA 3. Let  $\lambda, \mu \in \mathcal{L}$  and let  $s, t \in \{0, 1, 2, \dots\}$ . Then

$$m_{s+t}(\lambda + \mu) \leq m_s(\lambda) + m_t(\mu)$$

where  $\lambda + \mu \in \mathcal{L}$  is defined by  $(\lambda + \mu)_k = \lambda_k + \mu_k$  ( $k \in K$ ).

PROOF. Suppose  $j \in K, |\lambda_j + \mu_j| > m_s(\lambda) + m_t(\mu)$ . Then either  $|\lambda_j| > m_s(\lambda)$  or  $|\mu_j| > m_t(\mu)$ . Since

$$\text{card}\{j, |\lambda_j + \mu_j| > m_s(\lambda) + m_t(\mu)\} \leq \text{card}\{j, |\lambda_j| > m_s(\lambda)\} + \text{card}\{j, |\mu_j| > m_t(\mu)\} \leq s + t, \blacksquare$$

we conclude that  $m_{s+t}(\lambda + \mu) \leq m_s(\lambda) + m_t(\mu)$ .

$$\text{For } \lambda \in \mathcal{L} \text{ define } h(\lambda) = \sum_{i=0}^{\infty} (i+1)m_i(\lambda).$$

LEMMA 4. If  $\lambda, \mu \in \mathcal{L}$  then

$$h(\lambda + \mu) \leq 4[h(\lambda) + h(\mu)].$$

PROOF. We have

$$\begin{aligned} h(\lambda + \mu) &= \sum_{r=0}^{\infty} (2r+1)m_{2r}(\lambda + \mu) + \sum_{r=0}^{\infty} (2r+2)m_{2r+1}(\lambda + \mu) \leq \\ &\leq \sum_{r=0}^{\infty} (2r+1)[m_r(\lambda) + m_r(\mu)] + \sum_{r=0}^{\infty} (2r+2)[m_r(\lambda) + m_{r+1}(\mu)] \leq \\ &\leq \sum_{r=0}^{\infty} (4r+3)[m_r(\lambda) + m_r(\mu)] \leq 4[h(\lambda) + h(\mu)]. \end{aligned}$$

(continuation of the proof of Theorem 2):

Let  $u \in A$ , i.e.  $u$  can be expressed as

$$(6) \quad u = \alpha c + \sum_{n \in N} \sum_{k \in K} \beta_{nk} X_{nk} + \sum_{d \in \mathcal{D}} \gamma_d a_d + \sum_{d \in \mathcal{D}} \sum_{n \in N} \sum_{k \in K_{dn}} \delta_{dnk} y_{dnk}$$

where  $\alpha, \beta_{nk}, \gamma_d, \delta_{dnk}$  are complex numbers such that only a finite number of them is non-zero. For  $u$  of form (6) define

$$f(u) = |\alpha| + \sum_{n \in N} h(\{\beta_{nk}\}_{k \in K}) + \sum_{d \in \mathcal{D}} |\gamma_d| + \sum_{d \in \mathcal{D}} \sum_{n \in N} \frac{2}{d_n + 1} h(\{\delta_{dnk}\}_{k \in K})$$

(we put formally  $\delta_{dnk} = 0$  for  $k \in K - K_{dn}$ ).

The function  $f : A \rightarrow \langle 0, \infty \rangle$  has the following properties:

- a)  $u \in A, u \neq 0 \Rightarrow f(u) \neq 0$
- b)  $f(\varepsilon u) = |\varepsilon|f(u)$  for each complex number  $\varepsilon$  and  $u \in A$
- c)  $f(u + u') \leq 4[f(u) + f(u')]$
- d)  $f(u, u') \leq 8f(u)f(u')$ .

The first two properties are evident, property c) follows from Lemma 4. To prove d) suppose that  $u, u' \in A$  are of form (6) (i.e.  $u' = \alpha'c + \sum_n \sum_k \beta'_{nk}x_{nk} + \dots$ ). Then

$$\begin{aligned} f(uu') &= f\left(\sum_{d,n} \sum_{k \in K_{dn}} d_n y_{dnk} (\beta_{nk} \gamma'_d + \beta'_{nk} \gamma_d)\right) = \\ &= \sum_{d,n} \frac{2d_n}{d_{n+1}} h\left(\{\beta_{nk} \gamma'_d + \beta'_{nk} \gamma_d\}_{k \in K_{dn}}\right) \leq \\ &\leq 8 \sum_{d,n} \left[ |\gamma'_d| h\left(\{\beta_{nk}\}_{k \in K_{dn}}\right) + |\gamma_d| h\left(\{\beta'_{nk}\}_{k \in K_{dn}}\right) \right] \leq 8f(u)f(u'). \end{aligned}$$

Let  $V = \{u \in A, f(u) < 1\}$  and  $\mathcal{V} = \{tV, t \in (0, \infty)\}$ . Then  $\mathcal{V}$  satisfies conditions (1) – (5) so  $A$  with the topology given by  $\mathcal{V}$  is a topological algebra.

Let  $M \subset A$  be the subspace generated by the elements of form  $c - \frac{1}{d_n} \sum_{k \in K_{dn}} y_{dnk}$ ,

$d \in \mathcal{D}, n \in N$ . Clearly  $M$  is a two-sided ideal in  $A$ .

Let  $u \in A$  be of form (6). If  $\beta_{nk} \neq 0$  for some  $n \in N, k \in K$  or  $\gamma_d \neq 0$  for some  $d \in \mathcal{D}$  then  $(u + tV) \cap M = \phi$  for a suitable  $\epsilon > 0$ , so  $u \notin \bar{M}$ . Similarly,  $u \notin \bar{M}$  if  $\delta_{dnk} \neq \delta_{dnk'}$  for some  $d, n, k, k'$ . Finally, if  $u = \alpha c - \sum_{d,n} \sum_{k \in K_{dn}} \varepsilon_{dn} y_{dnk}$  and  $\alpha \neq \sum_{d,n} d_n \varepsilon_{dn}$  we have  $u \notin \bar{M}$  as  $f\left(\frac{1}{d_n} \sum_{k \in K_{dn}} y_{dnk}\right) = 1$  ( $d \in \mathcal{D}, n \in N$ ).

Hence  $M$  is a closed ideal in  $A$  and  $c \notin M$ . Let  $B = A/M$  and let  $\pi : A \rightarrow B$  be the canonical homomorphism. Then  $B$  is a topological algebra and  $\pi(c) \neq 0$ .

We prove that  $B$  is not topologizable as a locally convex algebra. Suppose on the contrary that there exists a system  $\mathcal{W}$  of convex zero-neighbourhoods in  $B$  satisfying (1) – (5). We shall need the following lemma:

**LEMMA 5.** *For every  $W \in \mathcal{W}$  there exists  $d \in \mathcal{D}$  and  $n \in N$  such that  $\pi(y_{dnk}) \in W$  for every  $k \in K_{dn}$ .*

**PROOF.** Let  $W \in \mathcal{W}$ . Suppose on the contrary that for every  $d \in \mathcal{D}$  and  $n \in N$  there exists  $k \in K_{dn}$  with  $\pi(y_{dnk}) \notin W$ . Let  $W' \in \mathcal{W}$  satisfy  $W'W' \subset W$ . For  $n \in N$  and  $k \in K$  choose  $s_{nk} > 0$  such that  $\pi(x_{nk}) \in s_{nk}W'$ .

Choose  $d = \{d_{nk}\}_{n \in N} \in \mathcal{D}$  such that  $d_{nk} > ns_{nk}$  ( $n \in N, k \in K$ ). Then  $a_d \in rW'$  for some  $r > 0$ .

We supposed that for every  $n \in N$  there exists  $k \in K_{dn}$  such that  $\pi(y_{dnk}) \notin W$ . On the other hand we have

$$\pi(y_{dnk}) = \frac{1}{d_n} \pi(x_{nk}) \pi(a_d) \in \frac{1}{d_n} s_{nk} W' r W' \subset \frac{s_{nk} r}{d_n} W.$$

So  $s_{nk} r / d_n > 1$ ,  $r > d_n / s_{nk} > n$  for every  $n \in N$  which is a contradiction.

(continuation of the proof of Theorem 2):

Let  $W \in \mathcal{W}$ . Let  $d \in \mathcal{D}$  and  $n \in N$  be given by Lemma 5. Then

$$\pi(c) = \frac{1}{d_n} \sum_{k \in K_{dn}} \pi(y_{dnk})$$

and  $\pi(y_{dnk}) \in W$  for every  $k \in K_{dn}$ . Since  $W$  is convex and  $\text{card}K_{dn} = d_n$  we have  $\pi(c) \in W$  for every  $W \in \mathcal{W}$ , a contradiction with condition (1).

Problem: Is it possible to construct separable algebras with properties of Theorem 1 (Theorem 2)?

#### REFERENCES

- [1] W.Żelazko, Selected topics in topological algebras, Aarhus University Lecture Notes, Series No 31 (1971).
- [2] W.Żelazko, On certain open problems in topological algebras, Rend. Sem. Mat. Fis. Milano (in print).
- [3] W.Żelazko, Example of an algebra which is non-topologizable as a locally convex topological algebra, Proc. Amer. Math. Soc. (in print).

Institute of Mathematics ČSAV  
Žitná 25, 115 67 Praha 1  
Czechoslovakia