

URYSOHN'S LEMMA, GLUING LEMMA AND CONTRACTION*
MAPPING THEOREM FOR FUZZY METRIC SPACES

E. ROJA, M. K. UMA, Salem, G. BALASUBRAMANIAN, Chennai

(Received November 11, 2006)

Abstract. In this paper the concept of a fuzzy contraction* mapping on a fuzzy metric space is introduced and it is proved that every fuzzy contraction* mapping on a complete fuzzy metric space has a unique fixed point.

Keywords: fuzzy contraction mapping, fuzzy continuous mapping

MSC 2000: 54A40, 03E72

1. INTRODUCTION

The theory of fuzzy sets was introduced by Zadeh in 1965 [7]. Since then many authors (Zi-ke 1982 [8], Erceg 1979 [1], George and Veeramani 1994 [2], Kaleva and Seikkala 1984 [5]) have introduced the concept of a fuzzy metric space in different ways. In this paper we follow the definition of a metric space given by George and Veeramani [2] since the topology induced by the fuzzy metric according to the definition of George and Veeramani [2] is Hausdorff. Motivated by the concept of a metric space, Urysohn's lemma and gluing lemma are studied. Based on the concept of a fuzzy contraction mapping [6], the fuzzy contraction* mapping theorem is established.

2. PRELIMINARIES

Definition 1 [4]. A binary operation $*$: $[0, 1] \times [0, 1] \rightarrow [0, 1]$ is a continuous t -norm if $*$ satisfies the following conditions:

1. $*$ is associative and commutative,
2. $*$ is continuous,

3. $a * 1 = a$ for all $a \in [0, 1]$,
4. $a * b \leq c * d$ whenever $a \leq c$ and $b \leq d$, ($a, b, c, d \in [0, 1]$).

Definition 2 [2]. The triple $(X, M, *)$ is said to be a fuzzy metric space if X is an arbitrary set, $*$ is a continuous t -norm and M is a fuzzy set on $X^2 \times (0, \infty)$ satisfying the following conditions:

1. $M(x, y, t) > 0$,
2. $M(x, y, t) = 1$ if and only if $x = y$,
3. $M(x, y, t) = M(y, x, t)$,
4. $M(x, y, t) * M(y, z, s) \leq M(x, z, t + s)$, $x, y, z \in X$ and $t, s > 0$,
5. $M(x, y, \cdot): X^2 \times (0, \infty) \rightarrow [0, 1]$ is continuous, $x, y, z \in X$ and $t, s > 0$.

Remark 1 [2]. $M(x, y, t)$ can be thought of as the degree of nearness between x and y with respect to t . We identify $x = y$ with $M(x, y, t) = 1$, for $t > 0$ and $M(x, y, t) = 0$ with $x = \infty$ or $y = \infty$.

Remark 2 [2]. In a fuzzy metric space $(X, M, *)$, whenever $M(x, y, t) > 1 - r$ for x, y in X , $t > 0$, $0 < r < 1$, we can find a t_0 , $0 < t_0 < 1$ such that $M(x, y, t_0) > 1 - r$.

Definition 3 [4]. A sequence $\{x_n\}$ in a fuzzy metric space $(X, M, *)$ is said to be a Cauchy sequence if for each ε , $0 < \varepsilon < 1$ and $t > 0$ there exists $n_0 \in \mathbb{N}$ such that $M(x_n, x_m, t) > 1 - \varepsilon$ for all $n, m \geq n_0$.

Definition 4 [2]. Let $(X, M, *)$ be a fuzzy metric space. We define the open ball $B(x, r, t)$ with centre $x \in X$ and radius r , $0 < r < 1$, $t > 0$ as

$$B(x, r, t) = \{y \in X : M(x, y, t) > 1 - r\}.$$

Definition 5 [4]. Let $(X, M, *)$ be a fuzzy metric space. Define $T = \{A \subset X : x \in A \text{ if and only if there exist } r, t > 0, 0 < r < 1 \text{ such that } B(x, r, t) \subset A\}$. Then T is topology on X . This topology is called the topology induced by the fuzzy metric.

Then by Theorem 3.11 of (George and Veeramani 1994 [2]) we know that a sequence $x_n \rightarrow x$ (x_n converges to x) if and only if $M(x_n, x, t) \rightarrow 1$ as $n \rightarrow \infty$.

Definition 6 [2]. A fuzzy metric space is said to be complete if every Cauchy sequence is convergent.

Notation. $M_A(x, y, t)$ denotes the degree of nearness between x and y with respect to t when $x, y \in A$.

3. URYSOHN'S LEMMA AND GLUING LEMMA

Proposition 1 (Urysohn's Lemma). *Let $(X, M, *)$ be a fuzzy metric space. Let T be a topology on X induced by the fuzzy metric. Let A and B be distinct members of r . Then there exists a fuzzy continuous function $f: X \rightarrow [0, 1]$ such that $f = 0$ on A and $f = 1$ on B .*

Proof. Define a function $f: X \rightarrow [0, 1]$ by

$$f(x) = \frac{1 - M_A(x, x, t)}{M_B(x, x, t) - M_A(x, x, t)}.$$

Note that $M_B(x, x, t) - M_A(x, x, t) \neq 0$ for any $x \in X$. If $x \in A$, $M_A(x, x, t) = 1$, then $f(x) = 0$. If $x \in B$, $M_B(x, x, t) = 1$, then $f(x) = 1 - M_A(x, x, t)/1 - M_A(x, x, t) = 1$. Since $M(x, y, t)$ is fuzzy continuous (George and Veeramani 1994 [2]), f is fuzzy continuous. □

Proposition 2 (Gluing Lemma). *Let $(X, M, *)$ and $(Y, M, *)$ be two fuzzy metric spaces. Let $U_i, i \in I$ be members of fuzzy induced topology T on X such that $\bigcup_{i \in I} U_i = X$. Assume that there exists a fuzzy continuous function [3] $f_i: U_i \rightarrow Y$ for each $i \in I$ with the property that $f_i(x) = f_j(x)$ for all $x \in U_i \cap U_j$ and $i, j \in I$. Then the function $f: X \rightarrow Y$ defined by $f(x) = f_i(x)$ if $x \in U_i$ is well defined and fuzzy continuous on X .*

Proof. Let $x, y \in X$. Since f_i is continuous for given $r \in (0, 1), t > 0$ we can find $r_0 \in (0, 1), t/4 > 0$ such that $M(x, y, t_0) > 1 - r_0$ implies $M(f_i(x), f_i(y), t/2) > 1 - r$. Now $M(x, y, t/4) > 1 - r_0$. Let $x \in U_i, y \in U_j$ for some $i \neq j$. Let $x_i \in U_i \cap U_j$. Then

$$\begin{aligned} M(f(x), f(y), t/2) &> M(f(x), f(x_i), t/4) * M(f(x_i), f(y), t/4) \\ &= M(f_i(x), f_i(x_i), t/4) * M(f_j(x_i), f_j(y), t/4) \\ &> (1 - r) * (1 - r) = 1 - r. \end{aligned}$$

Therefore f is fuzzy continuous. □

4. FUZZY CONTRACTION * MAPPING

Definition 7. Let $(X, M, *)$ be a fuzzy metric space. A function $f: X \rightarrow X$ is called a fuzzy contraction* mapping if $M(x, y, t) \geq 1 - (1 - r^2)$ for all $0 < 1 - r^2 < 1$. Then $M(f(x), f(y), t) \geq 1 - (1 - r_0^2)$ for each $x, y \in X$ for some $1 - r_0^2 < 1 - r^2, 1$.

Example 1. Consider the fuzzy metric space $(\mathbb{R}, M, *)$, where \mathbb{R} is the set of all real numbers and $M(x, y, t) = t/(t + |x - y|)$. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ and define $f(x) = x/2$. Then $M(x, y, t) = t/(t + |x - y|) \geq 1 - (1 - r^2)$, $t > 0$, $0 < 1 - r^2 < 1$ where $1 - r^2 \geq |x - y|/(t + |x - y|)$. Then

$$\begin{aligned} M(f(x), f(y), t) &= \frac{t}{t + |(x/2) - (y/2)|} \\ &= \frac{1 - (|(x/2) - (y/2)|)}{t + |(x/2) - (y/2)|} \geq 1 - (1 - r_0^2) \end{aligned}$$

where

$$1 - r_0^2 \geq \frac{|(x/2) - (y/2)|}{t + |(x/2) - (y/2)|}.$$

Further,

$$\begin{aligned} (1 - r^2) - (1 - r_0^2) &\geq \frac{|x - y|}{t + |x - y|} - \frac{(|(x/2) - (y/2)|)}{t + |(x/2) - (y/2)|} \\ &\geq \frac{|x - y|}{t + |x - y|} - \frac{\frac{1}{2}|x - y|}{t + \frac{1}{2}|x - y|} \\ &\geq \frac{|x - y|(t + \frac{1}{2}|x - y|) - \frac{1}{2}|x - y|(t + |x - y|)}{(t + |x - y|)(t + \frac{1}{2}|x - y|)} \\ &\geq \frac{|x - y|t - \frac{1}{2}(|x - y|t)}{(t + |x - y|)(t + \frac{1}{2}|x - y|)} \\ &\geq \frac{(t/2)|x - y|}{(t + |x - y|)(t + \frac{1}{2}|x - y|)} = 0, \end{aligned}$$

which implies that f is a fuzzy contraction* by Definition 7.

Definition 8. A mapping from a fuzzy metric space X to a fuzzy metric space Y is said to be fuzzy continuous* if for given $1 - r^2, t > 0$, $0 < 1 - r^2 < 1$ we can find $1 - r_0^2 \in (0, 1)$, $t_0 > 0$ such that $M(x, y, t_0) > 1 - (1 - r_0^2)$ implies $M(f(x), f(y), t/2) > 1 - (1 - r^2)$.

Proposition 3. Every fuzzy contraction* mapping on a fuzzy metric space is fuzzy continuous*.

Proof. Let $f: X \rightarrow X$ be a fuzzy contraction* mapping. Therefore for $x, y \in X$, given $1 - r^2 \in (0, 1)$, $t > 0$, we can find $1 - r_0^2 \in (0, 1)$, $t/4 > 0$ such that $1 - r^2 = (1 - (1 - r_0^2)) * (1 - (1 - r_0^2))$. Now $M(x, y, t/4) > 1 - (1 - r_0^2)$ implies $M(f(x), f(y), t/4) > 1 - (1 - s^2) > 1 - (1 - r_0^2)$ where $1 - s^2 \in (0, (1 - r_0^2))$ (since f is a fuzzy contraction* mapping). Let $x_1 \in X$. Then

$$\begin{aligned} M(f(x), f(y), t/2) &> M(f(x), f(x_1), t/4) * M(f(x_1), f(y), t/4) \\ &> (1 - (1 - r_0^2)) * (1 - (1 - r_0^2)) > (1 - (1 - r^2)), (1 - r^2) \in (0, 1) \end{aligned}$$

which implies that f is a fuzzy continuous* mapping. \square

Remark 3. The converse need not be true as the following example shows.

Example 2. Consider the fuzzy metric space $(\mathbb{R}, M, *)$ [2] where \mathbb{R} is the set of all real numbers and

$$M(x, y, t) = \frac{t}{t + |x - y|}.$$

Let $f: \mathbb{R} \rightarrow \mathbb{R}$ and define $f(x) = x^2$. Then

$$M(x, y, t) = \frac{t}{t + |x - y|} \geq 1 - (1 - r^2)$$

where $(1 - r^2) \geq |x - y|/(t + |x - y|)$. Then

$$\begin{aligned} M(f(x), f(y), t/2) &= \frac{(t/2)}{(t/2) + |x^2 - y^2|} = \frac{t}{t + 2(|x^2 - y^2|)} \\ &\geq 1 - (1 - r_0^2) \text{ where } 1 - r_0^2 \geq \frac{2|x^2 - y^2|}{t + 2(|x^2 - y^2|)} \end{aligned}$$

which implies that f is a fuzzy continuous* mapping. However, $M(f(x), f(y), t) = t/(t + |x^2 - y^2|) \geq 1 - (1 - s^2)$ where $1 - s^2 \geq |x^2 - y^2|/(t + |x^2 - y^2|)$ since

$$\begin{aligned} (1 - s^2) - (1 - r^2) &\geq \frac{|x^2 - y^2|}{t + |x^2 - y^2|} - \frac{|x - y|}{t + |x - y|} \\ &\geq \frac{(t + |x - y|)|x^2 - y^2| - (t + |x^2 - y^2|)|x - y|}{(t + |x^2 - y^2|)(t + |x - y|)} \\ &\geq \frac{t(|x^2 - y^2| - |x - y|)}{(t + |x^2 - y^2|)(t + |x - y|)} \begin{cases} \geq 0 & \text{if } x, y \text{ are integers} \\ \leq 0 & \text{if } x, y \text{ are not integers} \end{cases} \end{aligned}$$

and consequently, f is not a fuzzy contraction* mapping.

Proposition 4. *Every fuzzy contraction* mapping on a complete fuzzy metric space [2] has a unique fixed point.*

Proof. Let f be a fuzzy contraction* mapping on a complete fuzzy metric space $(X, M, *)$. □

Uniqueness part. If possible let $x_0 \neq y_0$ be two fixed points of f . Then we have

$$\begin{aligned} x_0 &= f^1(x_0) = f^2(x_0) = f^3(x_0) = \dots = f^n(x_0), \\ y_0 &= f^1(y_0) = f^2(y_0) = f^3(y_0) = \dots = f^n(y_0) \quad \text{for each } n \in \mathbb{N}. \end{aligned}$$

Now

$$\begin{aligned} M(x_0, y_0, t) &= M(f^n(x_0), f^n(y_0), t) \geq 1 - (1 - r^2)/k^n \\ &> M(x_0, y_0, t) \quad (= 1 - (1 - r^2)) \end{aligned}$$

where $k > 1$, a contradiction, hence $x_0 = y_0$. Therefore the fixed points are unique.

Existence part. Let $x_1 = f(x_0)$, $x_2 = f(x_1)$, \dots , $x_n = f(x_{n-1}) = f^{n-1}(x_1)$. Then

$$\begin{aligned} M(x_n, x_{n+1}, t) &= M(f^{n-1}(x_1), f^{n-1}(x_2), t) \geq 1 - (1 - r^2)/k^{n-1} \\ &\geq 1 - \frac{1}{1 - s^2} \quad \text{for some } \frac{1}{1 - s^2} \in (0, 1). \end{aligned}$$

Therefore,

$$(A) \quad M(x_n, x_{n+1}, t) \geq 1 - \frac{1}{1 - s^2}.$$

For a given $t' = (m - n)t > 0$, $\varepsilon > 0$, choose n_0 such that $1/n_0 < \varepsilon$. Then for $m \geq n \geq n_0$,

$$\begin{aligned} M(x_n, x_m, t') &\geq M(x_n, x_{n+1}, t) * M(x_{n+1}, x_{n+2}, t) * \dots * M(x_{m-1}, x_m, t) \\ &\geq (1 - (1 - s^2)^{-1}) * (1 - (1 - s^2)^{-1}) * \dots * (1 - (1 - s^2)^{-1}) \\ &\geq 1 - \frac{1}{n} \quad \text{for some } \frac{1}{n} \in (0, 1) \geq 1 - \varepsilon \end{aligned}$$

and hence $\{x_n\}$ is a Cauchy sequence. Since X is a complete metric space, this sequence converges to, say, $z_0 \in X$. Now we assert that z_0 is a fixed point of f . Consider $n \in \mathbb{N}$ for $0 < 1 - r^2 < 1$, $t > 0$. Then we have

$$\begin{aligned} M(f(z_0), z_0, t) &\geq M(f(z_0), f(x_0), t/n + 1) * M(f(x_0), f^2(x_0), t/n + 1) * \dots \\ &\quad * M(f^n(x_0), z_0, t/n + 1), \end{aligned}$$

and since f is a fuzzy contraction* mapping, this is for $k > 1$ and $1/(1 - s_n^2) \in (0, 1)$ greater than or equal to

$$\begin{aligned} & (1 - (1 - s_n^2)) * (1 - (1 - r^2)) * (1 - (1 - r^2)/k) * \dots \\ & * (1 - (1 - r^2)/k^{n-1}) * M(f^n(x_0), z_0, t/n + 1) \\ & \geq (1 - (1 - r^2)/k^{n+p}) * M(f^n(x_0), z_0, t/n + 1) \end{aligned}$$

for some $p \in \mathbb{N}$. Taking limit on both sides as $n \rightarrow \infty$ we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} M(f(z_0), z_0, t) & \geq \lim_{n \rightarrow \infty} (1 - (1 - r^2)/k^{n+p}) * \lim_{n \rightarrow \infty} M(f^n(x_0), z_0, t/n + 1) \\ & \Rightarrow M(f(z_0), z_0, t) \geq 1 * 1 \Rightarrow f(z_0) = z_0. \end{aligned}$$

Acknowledgement. The authors thank the referee for his/her comments and suggestions.

References

- [1] *M. A. Erceg*: Metric spaces in fuzzy set theory. *J. Math. Anal. Appl.* *69* (1979), 205–230. [zbl](#)
- [2] *A. George, P. Veeramani*: On some results in fuzzy metric spaces. *Fuzzy Sets Syst.* *64* (1994), 395–399. [zbl](#)
- [3] *A. George, P. Veeramani*: Some theorems in fuzzy metric spaces. *J. Fuzzy Math.* *3* (1995), 933–940. [zbl](#)
- [4] *A. George, P. Veeramani*: On some results of analysis for fuzzy metric spaces. *Fuzzy Sets Syst.* *90* (1997), 365–368. [zbl](#)
- [5] *O. Kaleva, S. Seikkala*: On fuzzy metric spaces. *Fuzzy Sets Syst.* *12* (1984), 215–229. [zbl](#)
- [6] *G. Rajendran, G. Balasubramanian*: Fuzzy contraction mapping theorem for fuzzy metric spaces. *Bull. Calcutta Math. Soc.* *94* (2002), 453–458. [zbl](#)
- [7] *L. A. Zadeh*: Fuzzy sets. *Inform and Control* *8* (1965), 338–353. [zbl](#)
- [8] *Zi-Ke, Deng*: Fuzzy pseudo metric spaces. *J. Math. Anal. Appl.* *86* (1982), 74–95. [zbl](#)

Authors' addresses: *E. Roja, M. K. Uma*, Department of Mathematics, Sri Sarada College for Women, Salem-636 016, Tamilnadu, Indie, e-mail: rpbalan@sancharnet.in; *G. Balasubramanian*, Ramanujan Institute for Advanced Study in Mathematics, University of Madras, Chennai-600 005, Tamilnadu, Indie.