

ON MINIMAL IDEALS IN SEMIGROUPS WITH RESPECT TO
THEIR SUBSETS, I

IMRICH ABRHAN, Bratislava

(Received September 13, 1994, revised February 5, 1996)

Abstract. In the paper, the following concept are defined:

- (i) a minimal left (right, two-sided) ideal with respect to a subset B of a semigroup S ,
 - (ii) a kernel with respect to a subset B of a semigroup S ,
- and their basic properties are investigated.

Keywords: minimal left (right, two-sided) ideal with respect to a subset B of a semigroup S , kernel with respect to a subset B of a semigroup S , partial group

MSC 1991: 20M10, 20M12

In many papers concerning the algebraic theory of semigroups, properties of the following types of ideals in semigroups are investigated:

- 1) the minimal left (right, two-sided) ideals (see for example [3], [5], [6], [7], [8], [9], [11]);
- 2) the 0-minimal left (right, both-sided) ideals (see for example [4]);
- 3) the minimal quasi-ideals (see for example [12]);
- 4) the simple left (right, two-sided) ideals (see for example [8], [10]).

In this paper, the following concepts are defined:

- a) a minimal left (right, two-sided) ideal with respect to a subset B of a semigroup S ;
- b) a kernel with respect to a subset B of a semigroup S .

An example of two semigroups, each satisfying exactly one of the following two properties, is given:

- a) S_1 does not contain any minimal left (right, two-sided) ideal (it does not have a kernel), and it contains infinitely many mutually different subsets such that with

respect to each of them S_1 contains minimal left (right, two-sided) ideals and the kernel.

b) S_2 contains at least one minimal left (right, two-sided) ideal, hence it contains the kernel, nonetheless it does not contain any simple left (right, two-sided) ideal and contains infinitely many mutually different subsets such that with respect to each of them S_2 has minimal, left (right, two-sided) ideals (none of them is a minimal left (right, two-sided) ideal of S) and with respect to each of them it also has the kernel.

Let S be a semigroup and let $\emptyset \neq B \subseteq S$. In this paper, basic properties of a minimal left (right, two-sided) ideal with respect to the set B of the semigroup S (under certain conditions on a subset B of a semigroup S) are investigated. The main result of this paper is Theorem 3, which is a generalization of Corollary 9 (see [3]).

After the basic assertions on minimal left (right, two-sided) ideals with respect to a set B of a semigroup S , some well known corollaries will be given, e.g. on minimal left, on 0-minimal right (if the semigroup S is a semigroup with the zero 0) and on simple left (if the semigroup S contains the kernel) ideals of a semigroup S .

Throughout the paper, the following notation will be used:

$X \subset Y$ will mean that X is a proper subset of the set Y (to distinguish it from $X \subseteq Y$ which means either $X \subset Y$ or $X = Y$).

Let S be a semigroup and let $\emptyset \neq A \subseteq S$. $L(A)$ ($R(A)$, $J(A)$) is the left (right, two-sided) ideal generated by A . If $a \in S$ and $A = \{a\}$, then instead of $L(\{a\})$ we will write $L(a)$.

\mathcal{L} (\mathcal{R} , \mathcal{J}) is the Green \mathcal{L} -equivalence (\mathcal{R} -equivalence, \mathcal{J} -equivalence) on S (see [1]).

S/\mathcal{L} (S/\mathcal{J} , S/\mathcal{R}) is the set of all \mathcal{L} -classes (\mathcal{J} -classes, \mathcal{R} -classes) which belong to the equivalence \mathcal{L} (\mathcal{J} , \mathcal{R}) on S .

L_a (J_a , R_a) is the element of S/\mathcal{L} (S/\mathcal{J} , S/\mathcal{R}) containing the element $a \in S$.

\leq is a partial ordering on S/\mathcal{L} (S/\mathcal{J} , S/\mathcal{R}) (see [1]). We will write $R_a < R_b$ provided $R_a \leq R_b$ and $R_a \neq R_b$.

$NL(A)$ ($N(A)$, $NR(A)$) will denote the set of all elements $x \in S$ such that for each $a \in A$: $L_a \not\leq L_x$ ($J_a \not\leq J_x$, $R_a \not\leq R_x$) (see [13]).

L_B (R_B) will denote the set $\cup\{L_b \mid b \in B\}$ ($\cup\{R_b \mid b \in B\}$).

\overline{A} is the set $S \setminus A$.

We will use the following assertion: Let S be a semigroup and let $\emptyset \neq A \subseteq S$. Then (see [13]):

If $NL(A) \neq \emptyset$, ($N(A) \neq \emptyset$, $NR(A) \neq \emptyset$), then $NL(A)$ ($N(A)$, $NR(A)$) is a left (two-sided, right) ideal in S .

In what follows the definitions of new concepts will be mostly omitted and the theorems about them will be given only for left ideals of S . Theorems on left ideals

of S will be referred to (without further notice) in case analogous theorems (concepts) concerning right (two-sided) ideals of S should be used.

Definition 1. Let S be a semigroup and let $\emptyset \neq B \subseteq S$. A left ideal L of a semigroup S will be called a *minimal left ideal* with respect to a subset B of a semigroup S (or in S), if $L \cap B \neq \emptyset$ and there is no left ideal L' in S such that $L' \subset L$ and $L' \cap B \neq \emptyset$.

Remark 1. a) If we put $B = S$ ($B = S \setminus \{0\}$) in Definition 1, then we have for each $\emptyset \neq L \subseteq S$:

L is a minimal left (0-minimal left) ideal with respect to a subset B of the semigroup S (of the semigroup S with 0) if and only if L is a minimal left ideal of the semigroup S (of the semigroup with 0).

b) Let S be a semigroup with the kernel K and let $K \neq S$. A left ideal L of the semigroup S is called a simple left ideal of the semigroup S , if $K \subset L$ and there is no left ideal L' in S such that $K \subset L' \subset L$ (see [10]). Put $B = S \setminus K$. In the paper it is shown how to get theorems on minimal left ideals with respect to the subset B of the semigroup S using theorems on simple left ideals of the semigroup with the kernel K .

Example 1. Let S_1 be the set of all real numbers $x \in \mathbb{R}$ such that $0 < x < 1$. A binary operation on S_1 will be defined in the following way: $xy = \min\{x, y\}$ for each two elements $x, y \in S_1$. Then S_1 is a semigroup.

Let $S_2 = \{a, b, c\}$ and let a binary operation on S_2 be defined in the following way:

	a	b	c
a	a	b	c
b	a	b	c
c	a	b	c

Then S_2 is a semigroup. Let $S_3 = S_1 \times S_2$ be the direct product of semigroups S_1, S_2 . For each $\alpha \in (0, 1)$ put $M^\alpha = \{y \mid y \in \mathbb{R} \text{ and } \alpha \leq y < 1\}$ and $B^\alpha = M^\alpha \times S_2$. Then for each $\alpha \in (0, 1)$ the set $\{L(\alpha, u) \mid u \in S_2\}$ is the set of all minimal left ideals with respect to the set B^α of the semigroup S_3 . It is easy to prove that the semigroup S_3 contains no minimal two-sided ideal. In this example instead of the set S_1 take a set S_{10} of all real numbers $x \in \mathbb{R}$ such that $0 \leq x < 1$. Define the binary operation on S_{10} analogously as on S_1 . Then S_{10} is a semigroup. Let $S_{30} = S_{10} \times S_2$ be the direct product of semigroups S_{10}, S_2 . Then we can easily prove that the semigroup S_{30} has the following properties:

a) S_{30} contains at least one minimal left and one minimal right ideal and hence S_{30} has the kernel,

b) S_{30} does not contain any simple left (two-sided) ideal,

c) S_{30} contains infinitely many mutually different subsets $(B^\alpha, \alpha \in (0, 1))$ such that with respect to each of them S_{30} has minimal left ideals (none of them is a minimal ideal of S).

For each $\beta \in (0, 1)$ put $N^\beta = \{y \mid y \in \mathbb{R} \text{ and } 0 < \beta < y < 1\}$ and $B^\beta = N^\beta \times S_2$. Then for each $\beta \in (0, 1)$ the set of all minimal left (right, two-sided) ideals with respect to the set B^β of the semigroup S_3 is empty.

Remark 2. By means of an example it can be shown that there exists a semigroup having a kernel and containing no minimal left (right), simple left ideal, while containing infinitely many mutually different subsets such that with respect to each of them it has both a minimal left ideal and the kernel.

Theorem 1. *Let S be a semigroup and let $\emptyset \neq B \subseteq S$. Then for each $\emptyset \neq L \subseteq S$ the following holds:*

(a) *L is a minimal left ideal with respect to the subset B of the semigroup S if and only if there exists an element $b \in B$ such that $L = L(b)$ and L_b is a minimal element of $\overline{NL(B)}/\mathcal{L}$.*

(b) *For each $b \in B$: $L(b)$ is a minimal left ideal with respect to the subset B of the semigroup S if and only if $L(b) \cap \overline{NL(B)} = L_b$.*

Proof. (a) I. Suppose that L is a minimal left ideal with respect to the subset B of the semigroup S . Let $b \in L \cap B$. Then $L(b) \subseteq L$ and $L(b) \cap B \neq \emptyset$. It follows from the assumption that $L = L(b)$. Let $a \in \overline{NL(B)}$ and let $L_a \leq L_b$. Then there exists an element $c \in B$ such that $L_c \leq L_a$. This implies that $L(c) = L(b)$, hence $L_a = L_b$. Therefore, L_b is a minimal element of $\overline{NL(B)}/\mathcal{L}$.

II. Let $b \in B$, $L = L(b)$ and let L_b be a minimal element of $\overline{NL(B)}/\mathcal{L}$. Let L' be a left ideal of the semigroup S such that $L' \subset L$ and $L' \cap B \neq \emptyset$. Let $c \in L' \cap B$. Hence $L(c) \subseteq L'(c) \subset L(b)$. Therefore $L_c < L_b$ and $L_b, L_c \in \overline{NL(B)}/\mathcal{L}$. This is a contradiction with the fact that L_b is a minimal element of $\overline{NL(B)}/\mathcal{L}$. Therefore $L(b)$ is a minimal ideal with respect to the subset B of the semigroup S .

(b) Let $b \in B$.

I. Suppose that $L(b)$ is a minimal left ideal with respect to the subset B of the semigroup S . Using (a) we get that $L_b \subseteq L(b) \cap \overline{NL(B)}$. Suppose that there is an element $d \in L(b) \cap \overline{NL(B)}$ such that $d \notin L_b$. Hence $L_d \subseteq \overline{NL(B)}$ and $L_d < L_b$. This is a contradiction with the fact that L_b is a minimal element of $\overline{NL(B)}/\mathcal{L}$. Therefore $L(b) \cap \overline{NL(B)} \subseteq L_b$.

II. Suppose that $L(b) \cap \overline{NL(B)} = L_b$. Further suppose that there exists a left ideal L of the semigroup S such that $L \subset L(b)$ and $L \cap B \neq \emptyset$. Then $L \cap L_b \neq \emptyset$. Hence

$L(b) \subset L$, which contradicts $L \subset L(b)$. Therefore $L(b)$ is a minimal left ideal with respect to the subset B of the semigroup S . \square

Corollary 1. *Let S be a semigroup. Then for each $\emptyset \neq L \subseteq S$ the following holds:*

(a) *L is a minimal left ideal in S if and only if there exists an element $b \in S$ such that $L = L(b)$ and L_b is a minimal element in S/\mathcal{L} .*

(b) *For each $b \in S$: $L(b)$ is a minimal left ideal in S if and only if $L(b) = L_b$.*

Proof. Put $B = S$. Then $\overline{NL(B)} = S$. Using Theorem 1 we get Corollary 1. \square

Corollary 2. *Let S be a semigroup S with zero 0 . Put $B = S \setminus \{0\}$. Then for each $\emptyset \neq L \subseteq S$ the following holds:*

(a) *L is a 0-minimal left ideal of the semigroup S if and only if there exists an element $b \in B$ such that $L = L(b)$ and L_b is a minimal element in B/\mathcal{L} .*

(b) *For each $b \in B$, $L(b)$ is a 0-minimal left ideal of the semigroup S if and only if $L(b) = \{0\} \cup L_b$.*

Proof. From the assumption we have that $B = S \setminus \{0\}$. Then $\overline{NL(B)} = S \setminus \{0\}$. Using Theorem 1 we get Corollary 2. \square

Corollary 3. *Let S be a semigroup with the kernel K and let S be not simple. Put $B = S \setminus K$. Then for each $L \subseteq S$ the following holds:*

L is a simple left ideal in S if and only if there exists an element $b \in B$ such that $L = K \cup L(b)$ and $L(b)$ is a minimal left ideal with respect to the subset B of the semigroup S .

Proof. I. Let L be a simple left ideal in S . Let $b \in L \setminus K$. Then $K \cup L(b) \subseteq L$ and $K \cup L(b)$ is a left ideal containing the kernel K . Then the assumption implies that $L = K \cup L(b)$. Suppose that L_b is not a minimal element of B/\mathcal{L} . There exists an element $c \in B$ such that $L_c < L_b$. Then $L(b) \setminus L_b \neq \emptyset$ and $(L(b) \setminus L_b) \cap B \neq \emptyset$. Then $L_1 = K \cup (L(b) \setminus L_b)$ is a left ideal of the semigroup S and $K \subset L_1 \subset L$. This is a contradiction with the fact that L is a simple left ideal of S . It follows that L_b is a minimal element in B/\mathcal{L} . Using Theorem 1 we get that $L = K \cup L(b)$ and $L(b)$ is a minimal ideal with respect to the subset B of the semigroup S .

II. Let $L = K \cup L(b)$ and let $L(b)$ be a minimal left ideal with respect to the subset B of the semigroup S . Suppose that there exists a left ideal L' in S such that $K \subset L' \subseteq L$. Let $d \in L' \cap L_b$. Then $L_b = L_d \subseteq L(d) \subseteq L'$. We get $L \subseteq L'$. Hence $L' = L$. Therefore L is a simple left ideal of S . \square

Definition 2. We will say that a semigroup S satisfies the condition m_{LB} (m_B) if $\emptyset \neq B \subseteq S$ and the set of all minimal left (two-sided) ideals with respect to the subset B in S is nonempty.

Let S be a semigroup and let $\emptyset \neq B \subseteq S$. A minimal left ideal L with respect to the subset B of the semigroup S will be called a left mB -ideal of the semigroup S if L has the following property: for each left ideal L' of S the following holds: If $L' \subset L$ and $c \in S$ then $L'c \cap \overline{NL(B)} = \emptyset$.

Lemma 1. Let a semigroup S satisfy the condition m_{LB} . Let either $NL(B) = \emptyset$, or let $NL(B)$ be a two-sided ideal of S . Then its every minimal left ideal with respect to the subset B of the semigroup S is a left mB -ideal of the semigroup S .

The proof is clear.

Let S be a semigroup without zero (with zero 0). Put $B = S$ ($B = S \setminus \{0\}$). Let S satisfy the condition m_{LB} . Then each minimal (0 -minimal) left ideal with respect to the set $B = S$ ($B = S \setminus \{0\}$) of the semigroup is a left mB -ideal of S .

It can be shown by means of an example that there is a semigroup S and its nonempty subset $B \subseteq S$ with the following properties:

- a) $\overline{NL(B)} \neq S$ and $NL(B)$ is not a two-sided ideal of S ,
- b) S satisfies the condition m_{LB} ,
- c) S contains a minimal left ideal with respect to the subset B of S that is its left mB -ideal and contains a minimal left ideal with respect to the subset B of S that is left mB -ideal of S .

Example 2. Let $S = \{0, \alpha, \beta, u, v, e\}$. Define on S a binary operation as follows:

	α	β	u	v	e
α	α	0	0	v	e
β	0	β	u	0	0
u	u	0	0	β	u
v	0	v	e	0	0
e	e	0	0	v	e

Then S is a semigroup. Put $B = \{\alpha, \beta\}$. Then $\overline{NL(B)} \neq \emptyset$, $\overline{NL(B)}$ is not a two-sided ideal of S . S satisfies the condition m_{LB} and contains a minimal left ideal with respect to the subset of S that is not its left mB -ideal of S and contains a minimal left ideal that is its left mB -ideal of S .

Lemma 2. Let a semigroup S satisfy the condition m_{LB} . Let L be a left mB -ideal of the semigroup S . Then for each $c \in \overline{NL(B)}$ the following holds: If $Lc \cap \overline{NL(B)} \neq \emptyset$, then Lc is a minimal left ideal with respect to the subset B of the semigroup S .

Proof. Let $c \in \overline{NL(B)}$, and let $Lc \cap B \neq \emptyset$. Suppose there exists a left ideal L^* of S such that $L^* \subset Lc$ and $L^* \cap B \neq \emptyset$. By L_1 we will denote the set of all elements $a \in L$ such that $ac \in L^*$. Then by the assumption we get that $L_1 \neq \emptyset$ and $L_1 \cap \overline{NL(B)} \neq \emptyset$. If $s \in S$ and $a \in L_1$, then $(sa)c = s(ac) \in sL^* \subseteq L^*$. Hence L_1 is a left ideal of S . Due to the assumption we have $L_1 = L$. Hence $Lc = L_1c \subseteq L^*$. This is a contradiction with $L^* \subset Lc$. Therefore Lc is a minimal left ideal with respect to the subset B of the semigroup S . \square

Corollary 4. (See [3].) *Let L be a minimal left ideal of a semigroup S and let $c \in S$. Then Lc be a minimal left ideal of the semigroup S .*

Proof. Put $B = S$. Then L is a left mB -ideal of S . By Lemmas 1 and 2 we get Corollary 4. \square

Corollary 5. (See [4].) *Let S be a semigroup with zero 0 . Let L be a 0 -minimal left ideal of S , and let $c \in S$. Then either $Lc = \{0\}$ or Lc is a 0 -minimal left ideal of S .*

Proof. Put $B = S \setminus \{0\}$. Then $\overline{NL(B)} = S \setminus \{0\}$ and $NL(B) = \{0\}$. Due to Lemmas 1 and 2 we get Corollary 5. \square

Corollary 6. (See [10].) *Let S be a semigroup with the kernel K and let L be a simple left ideal of S . Let $c \in S$. Then the set $K \cup Lc$ is either a simple left ideal of S or $K = K \cup Lc$.*

Proof. Put $B = S \setminus K$. Then using Corollary 3 and Lemma 2 we get Corollary 6. \square

Let a semigroup S satisfy the condition m_{LB} . By $*B$ we will denote the set of all elements of B such that for each minimal left ideal with respect to the subset B there exists exactly one element $b \in *B$ such that $L = L(b)$ (see Theorem 1) and $L(b)$ is a minimal left ideal with respect to the subset B of S for each $b \in *B$. The set $*B$ will be called the left lower basic (minimal) set of the subset B of the semigroup S . Clearly $*B$ is such a minimal subset of the set B that the sets of all minimal ideals with respect to B and of those with respect to $*B$ coincide.

Definition 3. We will say that a semigroup S satisfies the condition m_{LB}^* if S satisfies the condition m_{LB} and the left lower basic set $*B$ of the set B has the following properties:

- i) If $b \in *B$, $c \in S$ and $L(b)c \cap \overline{NL(B)} = \emptyset$, then there exists an element $d \in *B$ such that $\overline{L(b)c} \subseteq \overline{L(d)}$.
- ii) $\overline{NL(B)} = \overline{N(B)}$.

Remark 4. It is easy to prove that the following assertion holds:

(a) Let a semigroup S contain at least one minimal left ideal. Put $B = S$. Then the semigroup S satisfies the condition m_{LB}^* .

(b) Let a semigroup S with 0 contain at least one 0-minimal left ideal. Put $B = S \setminus \{0\}$. Then the semigroup S satisfies the condition m_{LB}^* .

Lemma 3. *Let a semigroup S satisfy the condition m_{LB}^* . Then the set union of all minimal left ideals with respect to the subset B of S is a two-sided ideal of S .*

Proof. Put $M = \cup\{L(b) \mid b \in {}_*B\}$. Let $a \in M$ and $c \in S$. There exists an element $d \in {}_*B$ such that $a \in L(d)$. Then either $\alpha) L(d)c \cap \overline{NL(B)} = \emptyset$, or $\beta) L(d)c \cap \overline{NL(B)} \neq \emptyset$. First suppose that $\alpha)$ holds. Then by the assumption, there exists an element $d' \in {}_*B$ such that $L(d)c \subseteq L(d')$. It follows that $ac \in M$. In the case $\beta)$, due to Lemma 2 we get that there exists $h \in {}_*B$ such that $L(b)c = L(h)$. It follows that M is a right ideal of S . Clearly M is a left ideal of S . Hence M is a two-sided ideal of S . \square

Definition 4. We will say that a semigroup S satisfies the condition m_{LB}^{**} if S satisfies the condition m_{LB}^* and for each $b, c \in {}_*B$ there exists an element $d \in \overline{NL(B)}$ such that $L(b)d = L(c)$.

Example 3. Let a semigroup S contain at least one minimal left ideal. Put $B = S$. Then $\overline{NL(B)} = S$. Let ${}_*B$ be the left lower basic set of the subset of the set $B (\subseteq S)$. Then it is easy to prove that the semigroup S satisfies the condition m_{LB}^{**} .

Theorem 2. *Let a semigroup S satisfy the condition m_{LB}^{**} . Then:*

(a) *For each two-sided ideal M of the semigroup S the following holds: If $M \cap {}_*B \neq \emptyset$, then $L({}_*B) \subseteq M$.*

(b) *The set $L({}_*B) = \cup\{L(b) \mid b \in {}_*B\}$ is a minimal two-sided ideal with respect to the subset ${}_*B$ of the semigroup S .*

Proof. (a) Let $b \in M \cap {}_*B$. Suppose that $c \in {}_*B$ and $c \notin M$. By the assumption there exists an element $d \in \overline{NL(B)}$ such that $L(b)d = L(c)$. This is a contradiction with $L(b) \subseteq M$ and $c \notin M$. Hence $L({}_*B) \subseteq M$.

(b) By the assumption and Lemma 3, $L({}_*B)$ is a two-sided ideal of the semigroup S . Suppose that there exists a two-sided ideal M' of the semigroup S such that $M' \subset L({}_*B)$ and $M' \cap {}_*B \neq \emptyset$. Using (a) we get $L({}_*B) \subseteq M'$. This contradicts the assumption. \square

Corollary 7. *Let a semigroup S contain at least one minimal left ideal. Then the set union of all minimal left ideals of the semigroup S is its minimal two-sided ideal (for the kernel of the semigroup S see e.g. [3], [9]).*

Remark 5. Let S be a semigroup in Example 2 and $B = \{\alpha, \beta\}$. Then

a) S satisfies the condition m_{LB}^* and does not satisfy the condition m_{LB}^{**} .

b) The set union of all minimal ideals with respect to the set B of a semigroup S is not a minimal two-sided ideal of S and $L_B \neq R_B$.

Definition 5. Let S be a semigroup and let $\emptyset \neq B \subseteq S$. Denote by K_B the intersection of all two-sided ideals N of the semigroup S such that $N \cap B \neq \emptyset$. If $K_B \neq \emptyset$ then the two-sided ideal K_B of S will be called the kernel with respect to the subset B of the semigroup S .

Clearly the following holds: If $B = S$ and $K_B \neq \emptyset$, then K_B is the kernel of the semigroup S .

Corollary 8. Let a semigroup S satisfy the condition m_{LB}^{**} . Then $L(*B)$ is the kernel with respect to the subset $*B$ of the semigroup S .

We get Corollary 8 using Theorem 2.

Example 4. Let $S_1, S_2, S_3, S_{10}, S_{30}$ be semigroups from Example 1. Let for each $\alpha \in (0, 1)$, M^α and B^α be the sets from Example 1. It is easy to show that each semigroup S_3 (S_{30}) satisfies the condition m_{LB}^{**} for each $\alpha \in (0, 1)$ ($\alpha \in \langle 0, 1 \rangle$). The semigroup S_3 (S_{30}) has the kernel with respect to its every subset B^α , $\alpha \in (0, 1)$ ($\alpha \in \langle 0, 1 \rangle$), contains the kernel and does not contain any simple left (right, two-sided) ideal.

Definition 6. Let S be a semigroup and let $\emptyset \neq B \subseteq S$. We will say that the semigroup S satisfies the condition mu_{LB}^{**} (mu_{RB}^{**}) if it satisfies the condition m_{LB}^{**} (m_{RB}^{**}) and for each $a, b \in *B$ ($a, b \in B_*$) we have $L_a b = L_b$ ($b R_a = R_b$).

Further, we denote by $D_l(B)$ ($D_r(B)$) the set of all elements $b \in B$ such that $bB = B$ ($Bb = B$).

Definition 7. A semigroup S will be called a partial group if and only if $D_r(S) \neq \emptyset$ and $D_r(S) = D_l(S)$ (see [2]).

Further, we will use the following lemma (its proof see e.g. [1], [2]).

Lemma 4. Let S be a partial group. Then

(a) $D_r(S) = S$ if and only if S is a group.

(b) If $D_r(S) \neq S$, then $S \setminus D_r(S)$ is a two-sided ideal of S and $D_r(S)$ is a group.

(c) The unit of the group $D_r(S)$ is a unit of the semigroup S .

A nonempty subset H of the semigroup S will be called a filter of the semigroup S if for each two elements $a, b \in S$ the following holds: $ab \in H$ ($a, b \in S$) if and only

if $a \in H, b \in H$. If H is filter of the semigroup S and $S \setminus H \neq \emptyset$, then $S \setminus H$ is a two-sided ideal in S .

Lemma 5. *Let a semigroup S satisfy the conditions $mu_{LB}^{**}, mu_{RB}^{**}$. Let $L_{*B} = R_{B*}$ and let c, d be arbitrary elements of L_{*B} . Put $G = R(c)L(d)$ and $D = G \cap L_B$. Then*

- (a) L_{*B} is a filter in $L(*B)$,
- (b) $D \neq \emptyset$,
- (c) $D \subseteq R_c \cap L_d$,
- (d) $D = D_r(G) = D_l(G)$.

Proof. By the assumption and Theorem 2 we get that $L(*B)$ is a two-sided ideal of S and $R(B_*) \subseteq L(*B), L(*B) \subseteq R(B_*)$. Therefore $L(*B) = R(B_*)$. By the assumption, we get that $L(*B) \setminus L_{*B} = R(B_*) \setminus R_{B*}$. Put $K = L(*B) \setminus L_{*B}$. Then $K = \cup\{L_b \cup L(b) \setminus L_b \mid b \in *B\} \setminus \cup\{L_b \mid b \in *B\} = \cup\{L(b) \setminus L_b \mid b \in *B\}$. Hence either (i) $L(b) \setminus L_b = \emptyset$ for all $b \in *B$, or (ii) there exists an element $b \in B$ such that $L(b) \setminus L_b \neq \emptyset$. Suppose that (ii) holds. Then $K \neq \emptyset$ and K is a two-sided ideal of S . Let a and b be elements of L_{*B} . Then by the assumption, $L_a b = L_b \subseteq L_{*B}$. It follows that L_{*B} is a filter in $L(*B)$ (in the case (i) we have $L(*B) = L_{*B}$).

b) Let c, d be elements of L_B . Then $cd \in R(c)L(d) = G$ and by (a) we get $cd \in L_{*B}$. Hence $D \neq \emptyset$.

c) Since $G \cap L_{*B} = [R(c)L(d)] \cap L_{*B} \subseteq [R(c) \cap L(d)] \cap L_{*B} = [R(c) \cap L_{*B}] \cap [L(d) \cap L_{*B}]$, the assumption and Theorem 1 yield that $D \subseteq R_c \cap L_d$.

d) Let g be an element of D . By (c) we get $g \in R_c$ and $g \in L_d$. By the assumption we get that $L_d = L_g = L_d g \subseteq L(d)g \subseteq L(d)L(d) \subseteq L(d)$. Then $L(d) = L(d)g$. Analogously $gR(c) = R(c)$. Hence $gG = gR(c)L(d) = G$ and $Gg = R(c)L(d)g = G$.

Let g be an element of G such that $g \notin D$. Then $g \in L(d)$ and $g \notin L_{*B}$. Therefore $g \in K$. By (a), L_{*B} is a filter in $L(*B)$ and $K \neq \emptyset$, hence K is a two-sided ideal in $L(*B)$. It follows that $Gg \cap L_{*B} = \emptyset$ and $gG \cap L_{*B} = \emptyset$. According to (b) we get $Gg \neq G$ and $gG \neq G$. The above considerations imply that the assertion (d) of Lemma 5 holds. \square

Theorem 3. *Let the assumptions of Lemma 5 hold. Then:*

- (a) G is a partial group.
- (b) $L(d) = Se, R(c) = eS$ and $G = R(c) \cap L(d) = eSe$ where e is the unit of the partial group G .
- (c) $D = R_c \cap L_d$.

Proof. (a) Since $L(d)$ is a left ideal of the semigroup S , we get that $GG = R(c)L(d)R(c)L(d) \subseteq R(c)L(d) = G$. According to (b) and (d) of Lemma 5 we get that G is a partial group.

(b) Let e be the unit of the partial group G . Then by Lemmas 4 and 5 we have $e \in R_c$ and $e \in L_d$. It means that $R(c) = eS$ and $L(d) = Se$. Then $eSe \subseteq eL(d) \subseteq L(d)$ and $eSe \subseteq R(c)e \subseteq R(c)$. It follows that $eSe \subseteq R(c) \cap L(d)$. Let x be an arbitrary element of $R(c) \cap L(d)$. Then there exists elements $u, v \in S$ such that $x = eu = ve$. Then $x = eu = e(eu) = e(ev) = eve$, i.e. $x \in eSe$. Hence $R(c) \cap L(d) \subseteq eSe$.

Clearly $R(c)L(d) \subseteq L(d)$ and $R(c)L(d) \subseteq R(c)$. Therefore $G \subseteq R(c) \cap L(d)$. Let x be an element of $R(c) \cap L(d)$. By (b) there exists an element $u \in S$ such that $x = ue$. Then $xe = (ue)e = ue = x$. Therefore $x \in R(c)L(e) = R(c)L(d)$. Hence $G = R(c) \cap L(d) = eSe$, where e is the unit element of the partial group G .

(c) By (b), $R_c \cap L_d \subseteq R(c) \cap L(d) = G$. Since $R_c \subseteq L_{*B}$ and $L_d \subseteq L_{*B}$, we get that $R_c \cap L_d \subseteq G \cap L_{*B}$. Lemma 7 implies that $D = R_c \cap L_d$. \square

Corollary 9. (See [3].) *Let $L(d)$ be a minimal left ideal and $R(c)$ a minimal right ideal of a semigroup S ($c, d \in S$). Put $B = S$, $G = R(c)L(d)$ and $D = G \cap L_{*B}$. Then:*

(a) G is a group.

(b) $R(c) = eS$, $L(d) = Se$ and $G = R(c) \cap L(d) = eSe$, where e is the unit of the group G .

(c) $G = R_c \cap L_d$.

Proof. By the assumption, the semigroup S satisfies the conditions m_{LB} , m_{RB} , where $B = S$. Let ${}_*B$ (B_*) be the left (right) lower basic set of the subset B of the semigroup S . Then it is easy to prove that the semigroup S satisfies the assumptions of Theorem 3. Because by the assumption, $L(d)$ is a minimal ideal of the semigroup S , using Theorem 1 ($B = S$) we have $L(d) = L_d$. It follows that $G = R(c)L(d) \subseteq L(d) = L_d \cap L_{*B}$. Therefore $D = G$. Using (d) of Lemma 6 we conclude that G is a group. \square

Example 5. Let $S_1 = \{0, 1, 2, 4, 5, 7, 8, 10, 11\}$ be a semigroup of the semigroup $S_{12} = \{0, 1, 2, \dots, 11\} \bmod 12$. S_2 is the semigroup from Example 1. Let $S_3 = S_1 \times S_2$ be the direct product of S_1, S_2 . Put $B_1 = \{2\} \times S_2$ and $B_2 = \{1\} \times S_2$. Then:

a) If $B = B_1$ then the semigroup S_3 satisfies the condition m_{LB}^{**} and does not satisfy the condition mu_{LB}^{**} .

b) If $B = B_2$ then ${}_*B = \{1\} \times S_2$, $B_* = \{(1, a)\}$, $L_{*B} = R_{*B}$ and the semigroup S satisfies the condition $mu_{LB}^{**}, mu_{RB}^{**}$.

References

- [1] *A. H. Clifford, G. B. Preston*: The algebraic theory of semigroups I, II. Amer. Math. Soc., Providence, 1961, 1967.
- [2] *E. S. Ljapin*: Semigroups. FIZMATGIZ, Moskva, 1960. (In Russian.)
- [3] *A. H. Clifford*: Semigroups containing minimal ideals. Amer. J. Math. *70* (1948), 521–526.
- [4] *A. H. Clifford*: Semigroups without nilpotent ideals. Amer. J. Math. *71* (1949), 834–844.
- [5] *L. Fuchs*: On semigroups admitting relative inverses and having minimal ideals. Publ. Math. Debrecen *1* (1950), 227–231.
- [6] *G. B. Preston*: Inverse semigroups with minimal right ideals. J. London Math. Soc. *29* (1954), 404–411.
- [7] *D. Rees*: Note on semigroups. Proc. Cambridge Philos. Soc. *37* (1941), 334–435.
- [8] *Š. Schwarz*: Theory of semigroups. Sborník prác Prír. fak. Slov. univerzity, IV. Bratislava, 1943. (In Slovak.)
- [9] *Š. Schwarz*: On the structure of simple semigroups without zero. Czechoslovak Math. J. *1* (1951), 41–58. (In Russian.)
- [10] *Š. Schwarz*: On semigroups having a kernel. Czechoslovak Math. J. *1* (1951), 259–301. (In Russian.)
- [11] *A. Suschkiewitsch*: Über die endlichen Gruppen ohne Gesetz der eindeutigen Umkehrbarkeit. Math. Ann. *99* (1928), 30–50.
- [12] *O. Steinfeld*: Über die Quasiideale von Halbgruppen. Publ. Math. Debrecen *4* (1956), 262–275.
- [13] *I. Abrhan*: On \mathcal{J} -subalgebras in unary algebras, on simple ideals and \mathcal{J} -ideals in groupoids and semigroups. Math. Slovaca *28* (1978), no. 1, 61–80. (In Russian.)

Author's address: Imrich Abrhan, Department of Mathematics, Faculty of Mechanical Engineering, Slovak Technical University, Nám. slobody 17, 812 31 Bratislava, Slovakia.