

# Computing the constant in Friedrichs' inequality

Tomáš Vejchodský

[vejchod@math.cas.cz](mailto:vejchod@math.cas.cz)

Institute of Mathematics, Academy of Sciences  
Žitná 25, 115 67 Praha 1  
Czech Republic



February 8, 2012, SIGA 2012, Prague

# Motivation

Classical formulation:

$$-\Delta u = f \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega$$

Weak formulation:  $V = H_0^1(\Omega)$

$$u \in V : \quad a(u, v) = \mathcal{F}(v) \quad \forall v \in V$$

Error bound:  $u_h \in V$

$$\|u - u_h\| \leq \|\mathbf{y} - \nabla u_h\|_0 + C_F \|f + \operatorname{div} \mathbf{y}\|_0 \quad \forall \mathbf{y} \in \mathbf{H}(\operatorname{div}, \Omega)$$

Notation:

- ▶  $a(u, v) = (\nabla u, \nabla v)$
- ▶  $\mathcal{F}(v) = (f, v)$
- ▶  $(\varphi, \psi) = \int_{\Omega} \varphi \psi \, dx$
- ▶ Energy norm:  $\|e\|^2 = a(e, e) = (\nabla e, \nabla e) = \|\nabla e\|_0^2$

# Friedrichs' inequality

Standard:

$$\|v\|_0 \leq C_F \|\nabla v\|_0 \quad \forall v \in H_0^1(\Omega)$$

# Friedrichs' inequality

Standard:

$$\|v\|_0 \leq C_F \|\nabla v\|_0 \quad \forall v \in H_0^1(\Omega)$$

Generalization:

$$\|v\|_0 \leq C_F \|v\| \quad \forall v \in V$$

# Friedrichs' inequality

Standard:

$$\|v\|_0 \leq C_F \|\nabla v\|_0 \quad \forall v \in H_0^1(\Omega)$$

Generalization:

$$\|v\|_0 \leq C_F \|v\| \quad \forall v \in V$$

Variants:

►  $\|v\|^2 = \|\nabla u\|_{\mathcal{A}} = (\mathcal{A}\nabla u, \nabla u)$

# Friedrichs' inequality

Standard:

$$\|v\|_0 \leq C_F \|\nabla v\|_0 \quad \forall v \in H_0^1(\Omega)$$

Generalization:

$$\|v\|_0 \leq C_F \|v\| \quad \forall v \in V$$

Variants:

- ▶  $\|v\|^2 = \|\nabla u\|_{\mathcal{A}}^2 = (\mathcal{A}\nabla u, \nabla u)$
- ▶  $\|v\|^2 = \|\nabla u\|_{\mathcal{A}}^2 + \|u\|_c^2 = (\mathcal{A}\nabla u, \nabla u) + (cu, u)$

# Friedrichs' inequality

Standard:

$$\|v\|_0 \leq C_F \|\nabla v\|_0 \quad \forall v \in H_0^1(\Omega)$$

Generalization:

$$\|v\|_0 \leq C_F \|v\| \quad \forall v \in V$$


Variants:

- ▶  $\|v\|^2 = \|\nabla u\|_{\mathcal{A}}^2 = (\mathcal{A}\nabla u, \nabla u)$
- ▶  $\|v\|^2 = \|\nabla u\|_{\mathcal{A}}^2 + \|u\|_c^2 = (\mathcal{A}\nabla u, \nabla u) + (cu, u)$

▶  $V = H_0^1(\Omega)$



▶  $V = \{v \in H^1(\Omega) : v = 0 \text{ on } \Gamma\}$



# Relation with eigenvalues

Friedrichs' inequality:

$$\|v\|_0 \leq C_F \|\nabla v\|_0 \quad \forall v \in V \quad \Rightarrow \quad C_F = \sup_{v \in V} \frac{\|v\|_0}{\|\nabla v\|_0}$$

Laplace eigenvalue problem

$$-\Delta u_i = \lambda_i u_i \quad \text{in } \Omega, \quad u_i = 0 \text{ on } \partial\Omega$$

Theorem:  $C_F^2 = \frac{1}{\lambda_1}$  where  $\lambda_1 = \min_i \lambda_i$ .

# Relation with eigenvalues

Friedrichs' inequality:

$$\|v\|_0 \leq C_F \|\nabla v\|_0 \quad \forall v \in V \quad \Rightarrow \quad C_F = \sup_{v \in V} \frac{\|v\|_0}{\|\nabla v\|_0}$$

Laplace eigenvalue problem

$$-\Delta u_i = \lambda_i u_i \quad \text{in } \Omega, \quad u_i = 0 \text{ on } \partial\Omega$$

Theorem:  $C_F^2 = \frac{1}{\lambda_1}$  where  $\lambda_1 = \min_i \lambda_i$ .

Proof:

Weak formulation:  $u_i \in V : (\nabla u_i, \nabla v) = \lambda_i(u_i, v) \quad \forall v \in V$

$$\lambda_1 = \inf_{v \in V} \frac{\|\nabla v\|_0^2}{\|v\|_0^2} \quad \Leftrightarrow \quad \frac{1}{\lambda_1} = \sup_{v \in V} \frac{\|v\|_0^2}{\|\nabla v\|_0^2}$$

# Rayleigh–Ritz approximation of $\lambda_1$

Weak formulation:

$$u_i \in V : \quad (\nabla u_i, \nabla v) = \lambda_i(u_i, v) \quad \forall v \in V$$

Rayleigh–Ritz method:  $V^h \subset V$ ,  $\dim V^h < \infty$

$$u_i^h \in V^h : \quad (\nabla u_i^h, \nabla v^h) = \lambda_i^h(u_i^h, v^h) \quad \forall v^h \in V^h$$

Theorem:  $\lambda_1 \leq \lambda_1^h$

# Rayleigh–Ritz approximation of $\lambda_1$

Weak formulation:

$$u_i \in V : \quad (\nabla u_i, \nabla v) = \lambda_i(u_i, v) \quad \forall v \in V$$

Rayleigh–Ritz method:  $V^h \subset V$ ,  $\dim V^h < \infty$

$$u_i^h \in V^h : \quad (\nabla u_i^h, \nabla v^h) = \lambda_i^h(u_i^h, v^h) \quad \forall v^h \in V^h$$

Theorem:  $\lambda_1 \leq \lambda_1^h$

Proof:

$$\lambda_1 = \inf_{v \in V} \frac{\|\nabla v\|_0^2}{\|v\|_0^2} \leq \inf_{v^h \in V^h} \frac{\|\nabla v^h\|_0^2}{\|v^h\|_0^2} = \lambda_1^h$$



# Rayleigh–Ritz approximation of $\lambda_1$

Weak formulation:

$$u_i \in V : \quad (\nabla u_i, \nabla v) = \lambda_i(u_i, v) \quad \forall v \in V$$

Rayleigh–Ritz method:  $V^h \subset V$ ,  $\dim V^h < \infty$

$$u_i^h \in V^h : \quad (\nabla u_i^h, \nabla v^h) = \lambda_i^h(u_i^h, v^h) \quad \forall v^h \in V^h$$

Theorem:  $\lambda_1 \leq \lambda_1^h$

Proof:

$$\lambda_1 = \inf_{v \in V} \frac{\|\nabla v\|_0^2}{\|v\|_0^2} \leq \inf_{v^h \in V^h} \frac{\|\nabla v^h\|_0^2}{\|v^h\|_0^2} = \lambda_1^h$$

□

Corollary:  $C_F^h \leq C_F$

# Lower bound on $\lambda_1$

Method of *a priori-a posteriori inequalities*.

Theorem (Kuttler and Sigillito, 1978):

- ▶ Let  $H$  be a separable Hilbert space.
- ▶ Let  $A : H \mapsto H$  be a symmetric operator with dense domain  $D(A)$ .
- ▶ Other technical assumptions on  $A$ .
- ▶ Let  $\lambda_*$  and  $u_* \in D(A)$  be arbitrary.
- ▶ Consider  $w \in D(A)$  such that  $Aw = Au_* - \lambda_* u_*$ .

Then

$$\min_i \left| \frac{\lambda_i - \lambda_*}{\lambda_i} \right| \leq \frac{\|w\|_H}{\|u_*\|_H}.$$

Usage:  $H = L^2(\Omega)$ ,  $A = -\Delta \Rightarrow$

$$\min_i \left| \frac{\lambda_i - \lambda_*}{\lambda_i} \right| \leq \frac{\|w\|_0}{\|u_*\|_0} \leq C_F \frac{\|\nabla w\|_0}{\|u_*\|_0}$$

## Algorithm

$$\min_i \left| \frac{\lambda_i - \lambda_*}{\lambda_i} \right| \leq C_F \frac{\|\nabla w\|_0}{\|u_*\|_0}$$

Theorem:

$$\|\nabla w\|_0 \leq \|\nabla u_* - \mathbf{q}\|_0 + C_F \|\lambda_* u_* + \operatorname{div} \mathbf{q}\|_0 \quad \forall \mathbf{q} \in \mathbf{H}(\operatorname{div}, \Omega).$$

# Algorithm

$$\min_i \left| \frac{\lambda_i - \lambda_*}{\lambda_i} \right| \leq C_F \left( \frac{\|\nabla u_* - \mathbf{q}\|_0}{\|u_*\|_0} + C_F \frac{\|\lambda_* u_* + \operatorname{div} \mathbf{q}\|_0}{\|u_*\|_0} \right)$$

Theorem:

$$\|\nabla w\|_0 \leq \|\nabla u_* - \mathbf{q}\|_0 + C_F \|\lambda_* u_* + \operatorname{div} \mathbf{q}\|_0 \quad \forall \mathbf{q} \in \mathbf{H}(\operatorname{div}, \Omega).$$

## Algorithm

$$\min_i \left| \frac{\lambda_i - \lambda_*}{\lambda_i} \right| \leq C_F \left( \frac{\|\nabla u_* - \mathbf{q}\|_0}{\|u_*\|_0} + C_F \frac{\|\lambda_* u_* + \operatorname{div} \mathbf{q}\|_0}{\|u_*\|_0} \right)$$

Theorem:

$$\|\nabla w\|_0 \leq \|\nabla u_* - \mathbf{q}\|_0 + C_F \|\lambda_* u_* + \operatorname{div} \mathbf{q}\|_0 \quad \forall \mathbf{q} \in \mathbf{H}(\operatorname{div}, \Omega).$$

- ▶ Compute Rayleigh–Ritz approximations  $\lambda_1^h$  and  $u_1^h$

# Algorithm

$$\min_i \left| \frac{\lambda_i - \lambda_*}{\lambda_i} \right| \leq C_F \left( \frac{\|\nabla u_* - \mathbf{q}\|_0}{\|u_*\|_0} + C_F \frac{\|\lambda_* u_* + \operatorname{div} \mathbf{q}\|_0}{\|u_*\|_0} \right)$$

Theorem:

$$\|\nabla w\|_0 \leq \|\nabla u_* - \mathbf{q}\|_0 + C_F \|\lambda_* u_* + \operatorname{div} \mathbf{q}\|_0 \quad \forall \mathbf{q} \in \mathbf{H}(\operatorname{div}, \Omega).$$

- ▶ Compute Rayleigh–Ritz approximations  $\lambda_1^h$  and  $u_1^h$
- ▶ Set  $\lambda_* = \lambda_1^h$  and  $u_* = u_1^h$

# Algorithm

$$\min_i \left| \frac{\lambda_i - \lambda_1^h}{\lambda_i} \right| \leq C_F \left( \frac{\|\nabla u_1^h - \mathbf{q}\|_0}{\|u_1^h\|_0} + C_F \frac{\|\lambda_1^h u_1^h + \operatorname{div} \mathbf{q}\|_0}{\|u_1^h\|_0} \right)$$

Theorem:

$$\|\nabla w\|_0 \leq \|\nabla u_* - \mathbf{q}\|_0 + C_F \|\lambda_* u_* + \operatorname{div} \mathbf{q}\|_0 \quad \forall \mathbf{q} \in \mathbf{H}(\operatorname{div}, \Omega).$$

- ▶ Compute Rayleigh–Ritz approximations  $\lambda_1^h$  and  $u_1^h$
- ▶ Set  $\lambda_* = \lambda_1^h$  and  $u_* = u_1^h$

## Algorithm

$$\min_i \left| \frac{\lambda_i - \lambda_1^h}{\lambda_i} \right| \leq C_F \left( \frac{\|\nabla u_1^h - \mathbf{q}\|_0}{\|u_1^h\|_0} + C_F \frac{\|\lambda_1^h u_1^h + \operatorname{div} \mathbf{q}\|_0}{\|u_1^h\|_0} \right)$$

Theorem:

$$\|\nabla w\|_0 \leq \|\nabla u_* - \mathbf{q}\|_0 + C_F \|\lambda_* u_* + \operatorname{div} \mathbf{q}\|_0 \quad \forall \mathbf{q} \in \mathbf{H}(\operatorname{div}, \Omega).$$

- ▶ Compute Rayleigh–Ritz approximations  $\lambda_1^h$  and  $u_1^h$
- ▶ Set  $\lambda_* = \lambda_1^h$  and  $u_* = u_1^h$
- ▶ Find approximate minimizer  $\mathbf{q}_h \in \mathbf{H}(\operatorname{div}, \Omega)$

## Algorithm

$$\min_i \left| \frac{\lambda_i - \lambda_1^h}{\lambda_i} \right| \leq C_F \left( \frac{\|\nabla u_1^h - \mathbf{q}_h\|_0}{\|u_1^h\|_0} + C_F \frac{\|\lambda_1^h u_1^h + \operatorname{div} \mathbf{q}_h\|_0}{\|u_1^h\|_0} \right)$$

Theorem:

$$\|\nabla w\|_0 \leq \|\nabla u_* - \mathbf{q}\|_0 + C_F \|\lambda_* u_* + \operatorname{div} \mathbf{q}\|_0 \quad \forall \mathbf{q} \in \mathbf{H}(\operatorname{div}, \Omega).$$

- ▶ Compute Rayleigh–Ritz approximations  $\lambda_1^h$  and  $u_1^h$
- ▶ Set  $\lambda_* = \lambda_1^h$  and  $u_* = u_1^h$
- ▶ Find approximate minimizer  $\mathbf{q}_h \in \mathbf{H}(\operatorname{div}, \Omega)$

## Algorithm

$$\min_i \left| \frac{\lambda_i - \lambda_1^h}{\lambda_i} \right| \leq C_F \left( \frac{\|\nabla u_1^h - \mathbf{q}_h\|_0}{\|u_1^h\|_0} + C_F \frac{\|\lambda_1^h u_1^h + \operatorname{div} \mathbf{q}_h\|_0}{\|u_1^h\|_0} \right)$$

Theorem:

$$\|\nabla w\|_0 \leq \|\nabla u_* - \mathbf{q}\|_0 + C_F \|\lambda_* u_* + \operatorname{div} \mathbf{q}\|_0 \quad \forall \mathbf{q} \in \mathbf{H}(\operatorname{div}, \Omega).$$

- ▶ Compute Rayleigh–Ritz approximations  $\lambda_1^h$  and  $u_1^h$
- ▶ Set  $\lambda_* = \lambda_1^h$  and  $u_* = u_1^h$
- ▶ Find approximate minimizer  $\mathbf{q}_h \in \mathbf{H}(\operatorname{div}, \Omega)$
- ▶  $\alpha = \frac{\|\nabla u_1^h - \mathbf{q}_h\|_0}{\|u_1^h\|_0}$ ,    $\beta = \frac{\|\lambda_1^h u_1^h + \operatorname{div} \mathbf{q}_h\|_0}{\|u_1^h\|_0}$ ,    $C_F = \frac{1}{\sqrt{\lambda_1}}$

# Algorithm

$$\min_i \left| \frac{\lambda_i - \lambda_1^h}{\lambda_i} \right| \leq C_F \left( \frac{\|\nabla u_1^h - \mathbf{q}_h\|_0}{\|u_1^h\|_0} + C_F \frac{\|\lambda_1^h u_1^h + \operatorname{div} \mathbf{q}_h\|_0}{\|u_1^h\|_0} \right)$$

Theorem:

$$\|\nabla w\|_0 \leq \|\nabla u_* - \mathbf{q}\|_0 + C_F \|\lambda_* u_* + \operatorname{div} \mathbf{q}\|_0 \quad \forall \mathbf{q} \in \mathbf{H}(\operatorname{div}, \Omega).$$

- ▶ Compute Rayleigh–Ritz approximations  $\lambda_1^h$  and  $u_1^h$
- ▶ Set  $\lambda_* = \lambda_1^h$  and  $u_* = u_1^h$
- ▶ Find approximate minimizer  $\mathbf{q}_h \in \mathbf{H}(\operatorname{div}, \Omega)$
- ▶  $\alpha = \frac{\|\nabla u_1^h - \mathbf{q}_h\|_0}{\|u_1^h\|_0}, \quad \beta = \frac{\|\lambda_1^h u_1^h + \operatorname{div} \mathbf{q}_h\|_0}{\|u_1^h\|_0}, \quad C_F = \frac{1}{\sqrt{\lambda_1}}$
- ▶  $\Rightarrow \frac{\lambda_1^h - \lambda_1}{\lambda_1} \leq \frac{1}{\sqrt{\lambda_1}} \left( \alpha + \frac{1}{\sqrt{\lambda_1}} \beta \right)$

## Algorithm

$$\min_i \left| \frac{\lambda_i - \lambda_1^h}{\lambda_i} \right| \leq C_F \left( \frac{\|\nabla u_1^h - \mathbf{q}_h\|_0}{\|u_1^h\|_0} + C_F \frac{\|\lambda_1^h u_1^h + \operatorname{div} \mathbf{q}_h\|_0}{\|u_1^h\|_0} \right)$$

Theorem:

$$\|\nabla w\|_0 \leq \|\nabla u_* - \mathbf{q}\|_0 + C_F \|\lambda_* u_* + \operatorname{div} \mathbf{q}\|_0 \quad \forall \mathbf{q} \in \mathbf{H}(\operatorname{div}, \Omega).$$

- ▶ Compute Rayleigh–Ritz approximations  $\lambda_1^h$  and  $u_1^h$
- ▶ Set  $\lambda_* = \lambda_1^h$  and  $u_* = u_1^h$
- ▶ Find approximate minimizer  $\mathbf{q}_h \in \mathbf{H}(\operatorname{div}, \Omega)$
- ▶  $\alpha = \frac{\|\nabla u_1^h - \mathbf{q}_h\|_0}{\|u_1^h\|_0}$ ,  $\beta = \frac{\|\lambda_1^h u_1^h + \operatorname{div} \mathbf{q}_h\|_0}{\|u_1^h\|_0}$ ,  $C_F = \frac{1}{\sqrt{\lambda_1}}$
- ▶  $\Rightarrow \frac{\lambda_1^h - \lambda_1}{\lambda_1} \leq \frac{1}{\sqrt{\lambda_1}} \left( \alpha + \frac{1}{\sqrt{\lambda_1}} \beta \right)$
- ▶  $\Leftrightarrow 0 \leq X^2 + \alpha X + \beta - \lambda_1^h$ , where  $X = \sqrt{\lambda_1}$

## Algorithm

$$\min_i \left| \frac{\lambda_i - \lambda_1^h}{\lambda_i} \right| \leq C_F \left( \frac{\|\nabla u_1^h - \mathbf{q}_h\|_0}{\|u_1^h\|_0} + C_F \frac{\|\lambda_1^h u_1^h + \operatorname{div} \mathbf{q}_h\|_0}{\|u_1^h\|_0} \right)$$

Theorem:

$$\|\nabla w\|_0 \leq \|\nabla u_* - \mathbf{q}\|_0 + C_F \|\lambda_* u_* + \operatorname{div} \mathbf{q}\|_0 \quad \forall \mathbf{q} \in \mathbf{H}(\operatorname{div}, \Omega).$$

- ▶ Compute Rayleigh–Ritz approximations  $\lambda_1^h$  and  $u_1^h$
- ▶ Set  $\lambda_* = \lambda_1^h$  and  $u_* = u_1^h$
- ▶ Find approximate minimizer  $\mathbf{q}_h \in \mathbf{H}(\operatorname{div}, \Omega)$
- ▶  $\alpha = \frac{\|\nabla u_1^h - \mathbf{q}_h\|_0}{\|u_1^h\|_0}$ ,  $\beta = \frac{\|\lambda_1^h u_1^h + \operatorname{div} \mathbf{q}_h\|_0}{\|u_1^h\|_0}$ ,  $C_F = \frac{1}{\sqrt{\lambda_1}}$
- ▶  $\Rightarrow \frac{\lambda_1^h - \lambda_1}{\lambda_1} \leq \frac{1}{\sqrt{\lambda_1}} \left( \alpha + \frac{1}{\sqrt{\lambda_1}} \beta \right)$
- ▶  $\Leftrightarrow 0 \leq X^2 + \alpha X + \beta - \lambda_1^h$ , where  $X = \sqrt{\lambda_1}$
- ▶  $\Rightarrow X_2^2 \leq \lambda_1$ , where  $X_2 = \left( \sqrt{\alpha^2 + 4(\lambda_1^h - \beta)} - \alpha \right) / 2$

## Algorithm

$$\min_i \left| \frac{\lambda_i - \lambda_1^h}{\lambda_i} \right| \leq C_F \left( \frac{\|\nabla u_1^h - \mathbf{q}_h\|_0}{\|u_1^h\|_0} + C_F \frac{\|\lambda_1^h u_1^h + \operatorname{div} \mathbf{q}_h\|_0}{\|u_1^h\|_0} \right)$$

Theorem:

$$\|\nabla w\|_0 \leq \|\nabla u_* - \mathbf{q}\|_0 + C_F \|\lambda_* u_* + \operatorname{div} \mathbf{q}\|_0 \quad \forall \mathbf{q} \in \mathbf{H}(\operatorname{div}, \Omega).$$

- ▶ Compute Rayleigh–Ritz approximations  $\lambda_1^h$  and  $u_1^h$
- ▶ Set  $\lambda_* = \lambda_1^h$  and  $u_* = u_1^h$
- ▶ Find approximate minimizer  $\mathbf{q}_h \in \mathbf{H}(\operatorname{div}, \Omega)$
- ▶  $\alpha = \frac{\|\nabla u_1^h - \mathbf{q}_h\|_0}{\|u_1^h\|_0}$ ,  $\beta = \frac{\|\lambda_1^h u_1^h + \operatorname{div} \mathbf{q}_h\|_0}{\|u_1^h\|_0}$ ,  $C_F = \frac{1}{\sqrt{\lambda_1}}$
- ▶  $\Rightarrow \frac{\lambda_1^h - \lambda_1}{\lambda_1} \leq \frac{1}{\sqrt{\lambda_1}} \left( \alpha + \frac{1}{\sqrt{\lambda_1}} \beta \right)$
- ▶  $\Leftrightarrow 0 \leq X^2 + \alpha X + \beta - \lambda_1^h$ , where  $X = \sqrt{\lambda_1}$
- ▶  $\Rightarrow X_2^2 \leq \lambda_1$ , where  $X_2 = \left( \sqrt{\alpha^2 + 4(\lambda_1^h - \beta)} - \alpha \right) / 2$
- ▶  $\Rightarrow C_F \leq 1/X_2$

# Computing $\mathbf{q}_h \in \mathbf{H}(\text{div}, \Omega)$

$$\begin{aligned}
& \left( \|\nabla u_* - \mathbf{q}\|_0 + C_F \|\lambda_* u_* + \operatorname{div} \mathbf{q}\|_0 \right)^2 \\
& \approx \left( \|\nabla u_1^h - \mathbf{q}\|_0 + (\lambda_1^h)^{-1/2} \|\lambda_1^h u_1^h + \operatorname{div} \mathbf{q}\|_0 \right)^2 \\
& \leq \frac{1+\varrho}{\varrho} \|\nabla u_1^h - \mathbf{q}\|_0^2 + \frac{1+\varrho}{\lambda_1^h} \|\lambda_1^h u_1^h - \operatorname{div} \mathbf{q}\|_0^2, \quad \forall \varrho > 0
\end{aligned}$$

Minimize over  $W_h \subset \mathbf{H}(\text{div}, \Omega)$ :

Find  $\mathbf{q}_h \in W_h$ :

$$(\operatorname{div} \mathbf{q}_h, \operatorname{div} \psi_h) + \frac{\lambda_1^h}{\varrho} (\mathbf{q}_h, \psi_h) = \frac{\lambda_1^h}{\varrho} (\nabla u_1^h, \psi_h) - (\lambda_1^h u_1^h, \operatorname{div} \psi_h)$$

$$\forall \psi_h \in W_h$$


Solve by standard Raviart-Thomas finite elements.

# Example 1

$$-\Delta u = f \text{ in } (0, 2) \times (0, 1)$$

$$u = 0 \text{ on } \Gamma_D$$

$$\mathbf{n}^\top \nabla u = 0 \text{ on } \Gamma_N$$



$$f = \frac{5\pi^2}{16} u$$

$$u = \sin \frac{\pi x_1}{4} \sin \frac{\pi x_2}{2}$$

## Example 1

$$-\Delta u = f \text{ in } (0, 2) \times (0, 1)$$

$$u = 0 \text{ on } \Gamma_D$$

$$\mathbf{n}^\top \nabla u = 0 \text{ on } \Gamma_N$$


$$f = \frac{5\pi^2}{16} u$$

$$u = \sin \frac{\pi x_1}{4} \sin \frac{\pi x_2}{2}$$


$$C_F = \frac{4}{\sqrt{5}\pi} \doteq 0.5694$$

$$C_F^{\text{low}} = 0.5693$$

$$C_F^{\text{up}} = 0.6004$$



Friedrichs' constant – Example 1




# Example 1

$$-\Delta u = f \text{ in } (0, 2) \times (0, 1)$$


$$u = 0 \text{ on } \Gamma_D$$

$$\mathbf{n}^\top \nabla u = 0 \text{ on } \Gamma_N$$



Lower bound:  
reference solution

Upper bound:  
error majorant




## Example 2

$$-\Delta u = f \text{ in } (0, 2) \times (0, 1)$$

$$u = 0 \text{ on } \Gamma_D$$

$$\mathbf{n}^\top \nabla u = 0 \text{ on } \Gamma_N$$




$$f = \frac{5\pi^2}{16} \sin \frac{\pi x_1}{4} \sin \frac{\pi x_2}{2}$$

## Example 2

$$-\Delta u = f \text{ in } (0, 2) \times (0, 1)$$

$$u = 0 \text{ on } \Gamma_D$$

$$\mathbf{n}^\top \nabla u = 0 \text{ on } \Gamma_N$$




$$f = \frac{5\pi^2}{16} \sin \frac{\pi x_1}{4} \sin \frac{\pi x_2}{2}$$

$$C_F = ?$$

$$C_F^{\text{low}} = 0.7750$$

$$C_F^{\text{up}} = 0.8712$$




## Example 2

$$-\Delta u = f \text{ in } (0, 2) \times (0, 1)$$

$$u = 0 \text{ on } \Gamma_D$$


$$\mathbf{n}^\top \nabla u = 0 \text{ on } \Gamma_N$$



Lower bound:  
reference solution

Upper bound:  
error majorant

Error bounds – Example 2



# Conclusions



- ▶ Practical method
- ▶ Guaranteed upper bound on Friedrichs' constant
- ▶ Easy to generalize to similar inequalities
- ▶ Computationally demanding
- ▶ Exact representation of the domain  $\Omega$
- ▶ ⇒ curved elements
- ▶ ⇒ **Splines!**

Thank you for your attention

Tomáš Vejchodský

[vejchod@math.cas.cz](mailto:vejchod@math.cas.cz)

Institute of Mathematics, Academy of Sciences  
Žitná 25, 115 67 Praha 1  
Czech Republic



February 8, 2012, SIGA 2012, Prague