

CLOSED CONVEX ℓ -SUBGROUPS AND RADICAL CLASSES
OF CONVERGENCE ℓ -GROUPS

JÁN JAKUBÍK, Košice

(Received March 20, 1996)

Abstract. In this paper we prove that the system of all closed convex ℓ -subgroups of a convergence ℓ -group is a Brouwer lattice and that a similar result is valid for radical classes of convergence ℓ -groups.

Keywords: convergence ℓ -group, closed convex ℓ -subgroup, radical class of convergence ℓ -groups

MSC 1991: 06F20, 22C05

All ℓ -groups considered in the present paper are assumed to be abelian. For convergence ℓ -groups (shorter: cl-groups) we apply the same notation and definitions as in [6].

Let (G, α) be a cl-group (where G is an ℓ -group and α is a convergence in G). For the definition of a closed ℓ -subgroup (shorter: cl-subgroup) of (G, α) cf. Section 1 below. The system of all convex cl-subgroups of (G, α) will be denoted by $c(G, \alpha)$; this system is partially ordered by the set-theoretical inclusion.

In the present paper we prove that $c(G, \alpha)$ is a Brouwer lattice. The lattice operations in $c(G, \alpha)$ are constructively described.

For $X \subseteq G$ the meaning of $\lim_{\alpha} X$ is defined in a natural way. We show that if X is an ℓ -subgroup of G such that $\lim_{\alpha} X$ can be represented as a direct product of a finite number of linearly ordered groups, then

$$\lim_{\alpha} \lim_{\alpha} X = \lim_{\alpha} X.$$

A nonempty class A of cl-groups is called a radical class of cl-groups if it is closed with respect to isomorphisms, convex cl-subgroups and joins of convex cl-subgroups. For radical classes A_1 and A_2 we put $A_1 \leq A_2$ if A_1 is a subclass of A_2 .

We prove that a certain form of distributive law (analogous to the condition applied when defining a Brouwer lattice) is valid for radical classes of cl-groups.

The analogous notion of a radical class of ℓ -groups was introduced in [5] and studied in several papers (cf., e.g., [1], [2], [7], [8]).

1. PRELIMINARIES

For an ℓ -group G we denote by $c(G)$ the system of all convex ℓ -subgroups of G ; this system is partially ordered by the set theoretical inclusion. Then $c(G)$ is a complete lattice. The lattice operations in $c(G)$ will be denoted by \vee and \wedge .

Let \mathcal{G} be the class of all ℓ -groups. A nonempty subclass X of \mathcal{G} is said to be a radical class of ℓ -groups if it satisfies the following conditions:

- (i) X is closed with respect to isomorphisms.
- (ii) Whenever $G \in X$ and $G_1 \in c(G)$, then $G_1 \in X$.
- (iii) Whenever $G \in \mathcal{G}$ and $\{G_i\}_{i \in I}$ is a nonempty subset of $X \cap c(G)$, then $\bigvee_{i \in I} G_i$ belongs to X .

We suppose that the reader is acquainted with the definitions from Section 1 of [6].

Let (G, α) and (G_1, α_1) be cl-groups.

1.1. Definition. (G_1, α_1) is said to be a cl-subgroup of (G, α) if

- (i) G_1 is an ℓ -subgroup of G ;
- (ii) whenever (x_n) is a sequence in G_1 , $x \in G$ and $x_n \rightarrow_\alpha x$, then $x \in G_1$ and $x_n \rightarrow_{\alpha_1} x$;
- (iii) whenever (x_n) is a sequence in G_1 , $x \in G_1$ and $x_n \rightarrow_{\alpha_1} x$, then $x_n \rightarrow_\alpha x$.

If (G_1, α_1) is a cl-subgroup of (G, α) , then we often write (G_1, α) instead of (G_1, α_1) .

The meaning of a convex cl-subgroup of (G, α) is obvious. The system of all convex cl-subgroups of (G, α) will be denoted by $c(G, \alpha)$. If (G_1, α_1) and (G_2, α_2) belong to $c(G, \alpha)$ and $G_1 \subseteq G_2$, then we put $(G_1, \alpha_1) \leq (G_2, \alpha_2)$. It is easy to verify that under the relation \leq , the system $c(G, \alpha)$ is a complete lattice. The lattice operations in $c(G, \alpha)$ will be denoted by \vee^c and \wedge^c .

1.2. Definition. A mapping φ of G into G_1 is called a cl-homomorphism if

- (i) φ is a homomorphism of the ℓ -group G into the ℓ -group G_1 ;
- (ii) whenever (x_n) is a sequence in G , $x \in G$ and $x_n \rightarrow_\alpha x$, then $\varphi(x_n) \rightarrow_{\alpha_1} \varphi(x)$.

If there exists a cl-homomorphism of (G, α) onto (G_1, α_1) , then (G_1, α_1) is said to be a homomorphic image of (G, α) .

1.3. Definition. Let φ be a cl-homomorphism of (G, α) onto (G_1, α_1) such that

- (i) φ is a monomorphism;
- (ii) the inverse mapping φ^{-1} is a cl-homomorphism of (G_1, α_1) onto (G, α) .

Then φ is an isomorphism of (G, α) onto (G_1, α_1) ; if such φ does exist, then (G_1, α_1) is said to be cl-isomorphic to (G, α) .

Let \mathcal{G}_c be the class of all cl-groups.

1.4. Definition. A nonempty subclass Y of \mathcal{G}_c is said to be a radical class of a cl-group if the following conditions are satisfied:

- (i) Y is closed with respect to cl-isomorphisms;
- (ii) whenever $(G, \alpha) \in Y$ and $(G_1, \alpha_1) \in c(G, \alpha)$, then $(G_1, \alpha_1) \in Y$;
- (iii) whenever $(G, \alpha) \in \mathcal{G}_c$ and $\{(G_i, \alpha_i)\}_{i \in I}$ is a nonempty subset of $Y \cap c(G, \alpha)$, then $\bigvee_{i \in I}^c (G_i, \alpha_i) \in Y$.

We shall often apply without quotation the following facts:

- (a₁) If $a_n \rightarrow_\alpha a$ and $a_n \leq a$ for each $n \in \mathbb{N}$, then $\bigvee_{n \in \mathbb{N}} a_n = a$ (and dually).
- (a₂) If G is linearly ordered, $a_n \rightarrow_\alpha a$, $c_1 < a < c_2$, then there is $m \in \mathbb{N}$ such that for each $n > m$ the relation $c_1 < a_n < c_2$ is valid.

(The assertion (a₁) is easy to verify; (a₂) is a consequence of (a₁).

2. THE SYSTEM $c(G, \alpha)$

Again, let $(G, \alpha) \in \mathcal{G}_c$.

A subset S of G is said to be closed with respect to (G, α) if, whenever (x_n) is a sequence in S , $x \in G$ and $x_n \rightarrow_\alpha x$, then $x \in S$.

2.1. Lemma. Let H be an ℓ -subgroup of G such that it is closed with respect to (G, α) . For a sequence (x_n) in H and $x \in H$ we put $x_n \rightarrow_{\alpha(H)} x$ if $x_n \rightarrow_\alpha x$. Then

- (i) $(H, \alpha(H))$ is a cl-group.
- (ii) $(H, \alpha(H))$ is a cl-subgroup of (G, α) .

Proof. The first assertion is an immediate consequence of the definition of the cl-group. Since H is closed with respect to (G, α) and in view of (i), the assertion (ii) holds as well. □

In view of the above remark concerning the notation (cf. Section 1) we will write (H, α) instead of $(H, \alpha(H))$.

Let X be a nonempty subset of G . We denote by $\lim_{\alpha} X$ the set of all $y \in G$ such that there exists a sequence (x_n) in X with $x_n \rightarrow_{\alpha} y$.

2.2. Lemma. *Let H be an ℓ -subgroup of G . Then $\lim_{\alpha} H$ is an ℓ -subgroup of G . If, moreover, H is convex in G , then $\lim_{\alpha} H$ is convex in $\overset{\alpha}{G}$ as well.*

Proof. Let $y_1, y_2 \in \lim_{\alpha} H$. Hence there are sequences (x_n^i) in H such that $x_n^i \rightarrow_{\alpha} y_i$ ($i = 1, 2$). Thus $x_n^1 + x_n^2 \rightarrow_{\alpha} y_1 + y_2$, and analogously for the operations \wedge and \vee . Also, $-x_n^1 \rightarrow -y_1$. Hence $\lim_{\alpha} H$ is an ℓ -subgroup of G .

Now suppose that H is convex in $\overset{\alpha}{G}$ and that $z \in G$, $y_1 \leq z \leq y_2$. Then

$$x_n^1 \wedge x_n^2 \rightarrow_{\alpha} y_1, \quad x_n^1 \vee x_n^2 \rightarrow_{\alpha} y_2.$$

Put

$$z_n = ((x_n^1 \wedge x_n^2) \vee z) \wedge (x_n^1 \vee x_n^2).$$

Hence $z_n \in H$ and $z_n \rightarrow_{\alpha} (y_1 \vee z) \wedge y_2 = z$. Thus $z \in \lim_{\alpha} H$. \square

Let H be as in 2.2. We put $H_0 = H$ and for each ordinal $t > 0$ we construct H_t by transfinite induction as follows. Suppose that for $t_1 < t$ all H_{t_1} are already defined and that they are ℓ -subgroups of G such that, whenever $t_1 < t_2 < t$, then $H_{t_1} \subseteq H_{t_2}$. If t is a limit ordinal, then we put

$$H_t = \bigcup_{t_1 < t} H_{t_1}.$$

If t is non-limit, then there exists t_1 with $t = t_1 + 1$. In this case we set

$$H_t = \lim_{\alpha} H_{t_1}.$$

There exists an ordinal t such that $H_t = H_{t_2}$ whenever $t_2 > t$. We denote

$$\lim_{\alpha} H = H_t.$$

From 2.1, 2.2 and from the construction of $\lim_{\alpha} H$ we immediately obtain

2.3. Lemma. *Let H be an ℓ -subgroup of G . Put $\lim_{\alpha} H = H^*$. Then*

- (i) (H^*, α) is a cl-subgroup of (G, α) ;
- (ii) if (K, α) is a cl-subgroup of (G, α) and $H \subseteq K$, then $H^* \subseteq K$;
- (iii) if, moreover, H is convex in G , then H^* is convex in G as well.

2.4. Lemma. Let $\{(H_i, \alpha)\}_{i \in I}$ be a nonempty subset of $c(G, \alpha)$. Put $H_0 = \bigcap_{i \in I} H_i$, $H^0 = \bigvee_{i \in I} H_i$. Then

- (i) $\bigwedge_{i \in I}^c (H_i, \alpha) = (H_0, \alpha)$;
- (ii) $\bigvee_{i \in I}^c (H_i, \alpha) = (\lim_{\alpha} H^0, \alpha)$.

Proof. The first assertion is obvious; the second is a consequence of 2.3. \square

2.5. Lemma. Let H be an ℓ -subgroup of G . Then the following conditions are equivalent:

- (i) H is closed with respect to (G, α) ;
- (ii) H^+ is closed with respect to (G, α) .

Proof. Let (i) be valid and let (x_n) be a sequence in H^+ , $x \in G$, $x_n \rightarrow_{\alpha} x$. Then $x_n = x_n \vee 0 \rightarrow_{\alpha} x \vee 0$, whence $x \vee 0 = x$ and thus (ii) holds. Conversely, suppose that (ii) is satisfied. Let (x_n) be a sequence in H , $x \in G$, $x_n \rightarrow_{\alpha} x$. Then $x_n^+ \rightarrow_{\alpha} x^+$ and $x_n^- \rightarrow_{\alpha} x^-$. We have $x_n^+, x_n^- \in H^+$ for each $n \in \mathbb{N}$ and thus, in view of (ii), both x^+ and x^- belong to H^+ . Hence $x = x^+ - x^-$ is an element of H . \square

For subsets X and Y of G we denote

$$X - Y = \{x - y : x \in X \text{ and } y \in Y\}.$$

2.6. Lemma. Let X be a subset of G^+ such that

- (i) X is a sublattice and a subsemigroup of G^+ ;
- (ii) $0 \in X$.

Then $X - X$ is an ℓ -subgroup of G and $(X - X)^+ = X$. If, moreover, X is a convex subset of G^+ , then $X - X$ is a convex ℓ -subgroup of G .

The proof is routine, it will be omitted.

For each nonempty subset X of G we can perform an analogous construction as we did above for H ; in this way we obtain a subset of G which will be denoted by $\lim_{\alpha} X$ or by X^* .

From the construction of X^* we immediately obtain

2.7. Lemma. Let X be as in 2.6. Then

- (i) X^* is a subset of G^+ and it satisfies the conditions (i), (ii) from 2.6;
- (ii) X^* is closed with respect to (G, α) ;
- (iii) if, moreover, X is convex in G , then X^* is convex in G as well.

2.8. Lemma. *Let H be an ℓ -subgroup of G ; put $X = H^+$. Then $H^* = X^* - X^*$.*

Proof. In view of the constructions of H^* and X^* we have $X^* \subseteq H^*$. Then according to 2.3 (i), $X^* - X^* \subseteq H^*$. Further, 2.7 and 2.5 yield that $X^* - X^*$ is closed with respect to (G, α) . Moreover, $H = H^+ - H^+ \subseteq X^* - X^*$. Hence according to 2.3 (ii) we obtain the relation $H^* \subseteq X^* - X^*$, which completes the proof. \square

2.9. Lemma. *Let $\{(H_i, \alpha)\}_{i \in I}$ and H^0 be as in 2.4. Put $(H^0)^+ = X$. Then*

$$\bigvee_{i \in I}^c (H_i, \alpha) = X^* - X^*.$$

Proof. This is a consequence of 2.4 and 2.8. \square

Now, let (A, α) and (B_i, α) ($i \in I$) be elements of $c(G, \alpha)$. Put

$$X = X_0 = \left(\bigvee_{i \in I} B_i \right)^+,$$

and let X^* be as above. For each ordinal t we define X_t analogously as when defining H_t .

Further, we put

$$Y = Y_0 = \left(\bigvee_{i \in I} (A \wedge B_i) \right)^+;$$

the symbols Y^* and Y_t are defined analogously as X^* and X_t .

It is well-known that the relation

$$A \wedge \left(\bigvee_{i \in I} B_i \right) = \bigvee_{i \in I} (A \wedge B_i)$$

is valid (cf., e.g., [5]). From this relation we immediately obtain that

$$A \wedge X_0 = Y_0$$

holds. Let t be an ordinal with $t > 0$ and assume that for each ordinal $t_1 < t$ the relation

$$A \wedge X_{t_1} = Y_{t_1}$$

is valid.

a) Suppose that t is a limit ordinal. Then we have

$$\begin{aligned} Y_t &= \bigcup_{t_1 < t} Y_{t_1} = \bigcup_{t_1 < t} (A \wedge X_{t_1}) = \bigcup_{t_1 < t} (A \cap X_{t_1}) \\ &= A \cap \left(\bigcup_{t_1 < t} X_{t_1} \right) = A \cap X_t = A \wedge X_t. \end{aligned}$$

b) Further, suppose that t is a non-limit ordinal. Hence there is an ordinal t_1 with $t = t_1 + 1$. Then

$$X_t = \lim_{\alpha} X_{t_1}, \quad Y_t = \lim_{\alpha} Y_{t_1} = \lim_{\alpha} (A \cap X_{t_1}).$$

Let $z \in A \wedge X_t$. Hence $z \in A$ and $z \in X_t$. Also, $z \geq 0$. There exists a sequence (z_n) in X_{t_1} such that $z_n \rightarrow_{\alpha} z$. Clearly $z_n \geq 0$. Then $0 \leq z_n \wedge z \leq z$, whence $z_n \wedge z \in A \cap X_{t_1} = Y_{t_1}$ and $z_n \wedge z \rightarrow_{\alpha} z$. Thus $z \in Y_t$ and therefore $A \wedge X_t \subseteq Y_t$.

Assume that $v \in Y_t$. There exists a sequence (v_n) in Y_{t_1} with $v_n \rightarrow_{\alpha} v$. We have $v_n \in A$ for each $n \in \mathbb{N}$. Since A is closed with respect to (G, α) we obtain that $v \in A$. Further, $v_n \in X_{t_1}$ for each $n \in \mathbb{N}$ and thus $v \in X_t$. Therefore $v \in A \wedge X_t$.

By summarizing, we obtain the relation

$$A \wedge X_t = Y_t$$

for each ordinal t . Thus

$$(*) \quad A \wedge X^* = Y^*.$$

2.10. Theorem. Let (A, α) and (B_i, α) , $i \in I$, be elements of (G, α) . Then

$$(A, \alpha) \wedge^c \left(\bigvee_{i \in I}^c (B_i, \alpha) \right) = \bigvee_{i \in I}^c ((A, \alpha) \wedge^c (B_i, \alpha)).$$

Proof. This is a consequence of 2.8, 2.9 and of the relation (*). □

2.11. Corollary. The system $c(G, \alpha)$ is a Brouwer lattice.

Let the symbol ω_1 have the usual meaning. It is easy to verify that if X is a nonempty subset of G and if t is an ordinal with $X^* = X_t$, then $t \leq \omega_1$.

If t is the first ordinal with $X^* = X_t$, then t will be said to be the degree of X in (G, α) .

Further, let t' be the first ordinal such that, whenever X is a nonempty subset of G , then the degree of X in (G, α) is less or equal to t' . We denote $d(G, \alpha) = t'$.

The following questions remain open:

a) For which ordinals t there exist $(G, \alpha) \in \mathcal{G}_c$ and $X \subseteq G$ such that t is the degree of X in (G, α) ?

b) For which ordinals t there exists $(G, \alpha) \in \mathcal{G}_c$ such that $d(G, \alpha) = t$?

For a related open question concerning convergence groups cf. [3].

3. THE CONDITION $\lim_{\alpha}^2 X = \lim_{\alpha} X$

Let (G, α) be as above. For $X \subseteq G$ we denote $\lim_{\alpha} \lim_{\alpha} X = \lim_{\alpha}^2 X$. In this section we prove that if X is an ℓ -subgroup of G such that X is a direct product of a finite number of linearly ordered groups, then the relation

$$(1) \quad \lim_{\alpha}^2 X = \lim_{\alpha} X$$

is valid. In other words, the degree of X is either 0 or 1.

3.1. Lemma. *Let X be a linearly ordered ℓ -subgroup of G and $g \in \lim X$. Then the set $X \cup \{g\}$ is linearly ordered and there are $x^1, x^2 \in X$ such that $x^1 \leq g \leq x^2$.*

Proof. In the case $X = \{0\}$ we have $g = 0$. Assume that $X \neq \{0\}$. Then there exists $x_0^1 \in X$ with $x_0^1 > 0$. First we prove that the element g cannot be an upper bound of the set X . By way of contradiction, suppose that $g > x$ for each $x \in X$.

Since $g \in \lim_{\alpha} X$, there is a sequence (x_n) in X such that $x_n \rightarrow_{\alpha} g$. Because $x_n \leq g$ for each $n \in \mathbb{N}$, we obtain that

$$\sup\{x_n\}_{n \in \mathbb{N}} = g$$

and this yields that $\sup X = g$. For each $x \in X$ we have $x + x_0^1 \in X$, thus $x + x_0^1 \leq g$, hence $x \leq g - x_0^1 < g$. This is a contradiction with the relation $\sup X = g$. Hence there is $x^2 \in X$ such that $x^2 \not\leq g$.

If x^2 is any element of X with this property, then there is a positive integer $m(x^2)$ such that for each $n \in \mathbb{N}$ with $n \geq m(x^2)$ we have $x_n \leq x^2$ (otherwise the relation $g \geq x^2$ would be valid). Then $g \leq x^2$. By a dual argument we prove that there is $x^1 \in X$ with $x^1 \leq g$. Moreover, if x^3 is any element of X with $x^3 \not\leq g$, then $g \geq x^3$. \square

3.2. Lemma. *Let X be a linearly ordered ℓ -subgroup of G . Then $\lim_{\alpha} X$ is also a linearly ordered ℓ -subgroup of G .*

Proof. In view of 2.2, $\lim_{\alpha} X$ is an ℓ -subgroup of G . Hence it suffices to verify that whenever g_1 and g_2 are distinct elements of $\lim_{\alpha} X$, then g_1 and g_2 are comparable. In view of 3.1 there are ideals X_1 and X_2 of the linearly ordered set X such that

- (i) $X_1 \neq X \neq X_2$;
- (ii) $x \leq g_1$ if $x \in X_1$, and $x > g_1$ if $x \in X \setminus X_1$;
- (iii) $x \leq g_2$ if $x \in X_2$, and $x > g_2$ if $x \in X \setminus X_2$.

The ideals X_1 and X_2 are comparable. Since $g_1 \neq g_2$, we obtain that $X_1 \neq X_2$. Thus without loss of generality we can suppose that $X_1 \subset X_2$. Hence there is $z \in X_2 \setminus X_1$. Then in view of (ii), $z > g_1$. Further, according to (iii), $z \leq g_2$. Therefore $g_1 \leq g_2$. \square

3.3. Lemma. *Let X be a linearly ordered ℓ -subgroup of G . Then (1) holds.*

Proof. Let (y_n) be a sequence in $\lim_{\alpha} X$, $g \in G$ and $y_n \rightarrow_{\alpha} g$. Then in view of 3.1 and 3.2, g is comparable with all elements of $\lim_{\alpha} X$. Hence there exists a subsequence (y_n^1) of (y_n) such that either (i) $y_n^1 \geq g$ for each $n \in \mathbb{N}$, or (ii) $y_n^1 \leq g$ for each $n \in \mathbb{N}$. Suppose that (i) holds (in the case of (ii) the method is similar). If $y_n^1 = g$ for some $n \in \mathbb{N}$, then $g \in \lim_{\alpha} X$. Thus it suffices to suppose that $y_n^1 < g$ for each $n \in \mathbb{N}$, and in this case we can assume without loss of generality that $y_n^1 < y_{n+1}^1$ for each $n \in \mathbb{N}$.

Let $n \in \mathbb{N}$. There exists a sequence $(x_k^n)_{k \in \mathbb{N}}$ in X such that $x_k^n \rightarrow_{\alpha} y_n^1$ (as $k \rightarrow \infty$). Hence there is $m(n) \in \mathbb{N}$ such that

$$y_{n-1}^1 < x_k^n < y_{n+1}^1$$

whenever $k \geq m(n)$. Since $y_{n-1}^1 \rightarrow_{\alpha} g$ and $y_{n+1}^1 \rightarrow_{\alpha} g$ we obtain that

$$x_{m(n)}^n \rightarrow_{\alpha} g$$

and thus $g \in \lim_{\alpha} X$. Hence (1) is valid. \square

3.4. Lemma. *Let L be a distributive lattice with the least element 0. Let A and B be sublattices of L such that*

- (i) $0 \in A \cap B$;
- (ii) $a \wedge b = 0$ for each $a \in A$ and each $b \in B$;
- (iii) for each $g \in L$ there are $a \in A$ and $b \in B$ with $g = a \vee b$.

Then the elements a, b from (iii) are uniquely determined and the mapping $g \rightarrow (a, b)$ gives an isomorphism of L onto the direct product $A \times B$.

The proof is routine, it will be omitted.

3.5. Lemma. *Let X be an ℓ -subgroup of G such that X is a direct product of linearly ordered groups X_1, X_2, \dots, X_k . Then the ℓ -group $\lim_{\alpha} X$ is a direct product of linearly ordered groups $\lim_{\alpha} X_1, \lim_{\alpha} X_2, \dots, \lim_{\alpha} X_k$.*

Proof. We proceed by induction with respect to k . The case $k = 1$ is trivial. Suppose that $k > 1$ and that the assertion is valid for $k - 1$.

Without loss of generality we can assume that $X_i \neq \{0\}$ for $i = 1, 2, \dots, k$. Put $Y_i = \lim_{\alpha} X_i$ ($i = 1, 2, \dots, k$). According to 3.2, all Y_i are linearly ordered ℓ -subgroups of G . Also, $\lim_{\alpha} X = Y$ is an ℓ -subgroup of G . In view of Theorem 2.3, [4] it suffices to verify that the lattice Y^+ is a direct product of lattices Y_1^+, \dots, Y_k^+ .

Let $g \in Y^+$. In the same way as in the proof of 3.1 we can verify that g fails to be an upper bound of the set X^+ . For each $x \in X^+$ we have

$$x = x(X_1) \vee \dots \vee x(X_k), \quad x(X_i) \geq 0 \quad (i = 1, 2, \dots, k),$$

where $x(X_i)$ is the component of x in X_i . Hence g fails to be an upper bound of the set $X_1^+ \cup X_2^+ \cup \dots \cup X_k^+$. Thus we can suppose that g is not an upper bound of the set X_k^+ . Therefore there is $x_0 \in X_k^+$ such that $x_0 \not\leq g$.

There is a sequence (z_n) in X such that $z_n \rightarrow_{\alpha} g$. Put $z'_n = z_n \vee 0$. Then we have $z'_n \rightarrow_{\alpha} g$ as well. Further,

$$\begin{aligned} z'_n \wedge x_0 &= (z'_n(X_1) \vee z'_n(X_2) \vee \dots \vee z'_n(X_k)) \wedge x_0 = \\ &= z'_n(X_k) \wedge x_0 \in X_k \end{aligned}$$

and $z'_n(X_k) \wedge x_0 \rightarrow_{\alpha} g \wedge x_0$, whence $g \wedge x_0 \in \lim_{\alpha} X_k \subseteq \lim_{\alpha} X$.

Put $N_1 = \{n \in \mathbb{N} : z'_n(X_k) \geq x_0\}$. If the set N_1 is infinite, then there exists a subsequence (z''_n) of (z'_n) such that $z''_n \geq x_0$ for each $n \in \mathbb{N}$ and then we would have $g \geq x_0$, which is a contradiction. Hence the set N_1 is finite; thus there is a subsequence (z''_n) of (z'_n) such that $z''_n(X_k) < x_0$ for each $n \in \mathbb{N}$, whence

$$z''_n(X_k) \wedge x_0 = z''_n(X_k)$$

and then $z''_n(X_k) \rightarrow_{\alpha} g \wedge x_0$. Therefore

$$z''_n - z''_n(X_k) = z''_n(X_1) + z''_n(X_2) + \dots + z''_n(X_{k-1}) \rightarrow_{\alpha} g - (g \wedge x_0).$$

Therefore by the induction hypothesis (since $z_n''(X_1) + \dots + z_n''(X_{k-1})$ belongs to $X_1 \times \dots \times X_{k-1}$) the element $g - (g \wedge x_0)$ belongs to the direct product $Y_1 \times Y_2 \times \dots \times Y_{k-1}$. Since $g - (g \wedge x_0) \geq 0$ we obtain, moreover, that this element belongs to the direct product of lattices Y_1^+, \dots, Y_{k-1}^+ .

Let $t \in Y_1^+ \times Y_2^+ \times \dots \times Y_{k-1}^+$. Then by the induction hypothesis, there is a sequence (t_n) in $X_1^+ \times \dots \times X_{k-1}^+$ such that $t_n \rightarrow_\alpha t$. We have

$$t_n \wedge z_n''(X_k) = 0 \quad \text{for each } n \in \mathbb{N},$$

thus

$$\begin{aligned} t \wedge (g \wedge x_0) &= 0, \\ t + (g \wedge x_0) &= t \vee (g \wedge x_0). \end{aligned}$$

In particular,

$$g = (g - (g \wedge x_0)) + (g \wedge x_0) = (g - (g \wedge x_0)) \vee (g \wedge x_0)$$

with $g - (g \wedge x_0) \in Y_1^+ \times \dots \times Y_{k-1}^+$ and $g \wedge x_0 \in Y_k^+$.

Hence in view of 3.4 we obtain that for the lattice Y^+ there exists a direct product decomposition

$$Y^+ = Y_1^+ \times Y_2^+ \times \dots \times Y_k^+.$$

Now we apply again Theorem 2.3 of [4] concluding that the ℓ -group Y has a direct product decomposition

$$(2) \quad Y = Y_1 \times Y_2 \times \dots \times Y_k.$$

□

3.6. Theorem. *Let X be an ℓ -subgroup of G such that X is a direct product of a finite number of linearly ordered groups. Then (1) holds.*

Proof. We apply the notation as in the proof of 3.5 and similarly as in 3.5 we proceed by induction with respect to k . The case $k = 1$ was dealt with in 3.3; let $k > 1$.

Since all Y_i are linearly ordered we can apply 3.5 to the relation (2) obtaining

$$\lim_\alpha Y = \lim_\alpha Y_1 \times \lim_\alpha Y_2 \times \dots \times \lim_\alpha Y_k.$$

Since $\lim_\alpha Y = \lim_\alpha^2 X$ and $\lim_\alpha Y_i = \lim_\alpha^2 X_i$ ($i = 1, 2, \dots, k$), by applying 3.3 we infer

$$\lim_\alpha^2 X = Y_1 \times \dots \times Y_k = \lim_\alpha X.$$

□

4. THE RELATION OF PARTIAL ORDER BETWEEN RADICAL CLASSES

For a class X of cl-groups we denote by

$\text{Sub}_0 X$ —the class of all cl-groups (G, α) having the property that there exist (H, β) in X and $(H_1, \beta) \in c(H, \beta)$ such that (G, α) and (H_1, β) are cl-isomorphic;

$\text{Join } X$ —the class of all cl-groups (G, α) having the property that there exist (H_i, β_i) in X and $(G_i, \alpha) \in c(G, \alpha)$ ($i \in I$) such that

- a) for each $i \in I$, (H_i, β_i) and (G_i, α) are cl-isomorphic, and
- b) $(G, \alpha) = \bigvee_{i \in I}^c (G_i, \alpha)$.

4.1. Proposition. *Let X be a nonempty class of cl-groups. Then*

- a) $\text{Join Sub}_c X$ is a radical class of cl-groups.
- b) If Y is a radical class of cl-groups and $X \subseteq Y$, then $\text{Join Sub}_c X \subseteq Y$.

Proof. Put $\text{Join Sub}_c X = Z$. We have to verify that Z satisfies the conditions (i), (ii) and (iii) from 1.4. It is obvious that Z is closed with respect to cl-isomorphisms. For each nonempty class Z_1 of cl-groups we have $\text{Join Join } Z_1 = \text{Join } Z_1$, whence Z satisfies the condition (iii) from 1.4.

Let $(G, \alpha) \in Z$ and $(G_1, \alpha) \in c(G, \alpha)$. Hence there exist (H_i, α_i) ($i \in I$) belonging to $\text{Sub}_c X \cap c(G, \alpha)$ such that

$$(G, \alpha) = \bigvee_{i \in I}^c (H_i, \alpha).$$

Then by applying 2.10

$$\begin{aligned} (G_1, \alpha) &= (G_1, \alpha) \wedge^c (G, \alpha) = (G_1, \alpha) \wedge^c \left(\bigvee_{i \in I}^c (H_i, \alpha) \right) \\ &= \bigvee_{i \in I}^c ((G_1, \alpha) \wedge^c (H_i, \alpha)). \end{aligned}$$

For each $i \in I$, the cl-group $(G_1, \alpha) \wedge^c (H_i, \alpha)$ belongs to $\text{Sub}_c \text{Sub}_c X = \text{Sub}_c X$ and therefore (G_1, α) belongs to Z . Hence the condition (ii) from 1.3 is valid, which completes the proof of a).

Let Y be a radical class of cl-groups and $X \subseteq Y$. Then $\text{Sub}_c X \subseteq \text{Sub}_c Y = Y$ and $\text{Join Sub}_c X \subseteq \text{Join } Y = Y$. Thus b) is valid. \square

Let Y_1 and Y_2 be radical classes of cl-groups. We put $Y_1 \leq Y_2$ if Y_1 is a subclass of Y_2 .

We denote by Y_0 the class of all cl-groups (G, α) such that G is a one-element set. Then Y_0 is a radical class of cl-groups and for each radical class Y of cl-groups we have $Y_0 \leq Y \leq \mathcal{G}_c$.

Let G be an ℓ -group. For a sequence (x_n) in G and for $x \in G$ we put $x_n \rightarrow_{\alpha(G)} x$ if there exists $m \in \mathbb{N}$ such that $x_n = x$ for each positive integer n with $n \geq m$. Then $(G, \alpha(G))$ is a cl-group; $\alpha(G)$ is the discrete convergence on G .

If X is a class of ℓ -groups, then we put

$$\varphi(X) = \{(G; \alpha(G)) : G \in X\}.$$

Then we obviously have

4.2. Lemma. *If X is a radical class of ℓ -groups, then $\varphi(X)$ is a radical class of cl-groups. Moreover, if X_1 and X_2 are distinct radical classes of ℓ -groups, then $\varphi(X_1) \neq \varphi(X_2)$.*

Let \mathcal{R}_a and \mathcal{R}_c be the collection of all radical classes of ℓ -groups or the collection of all radical classes of cl-groups, respectively. (Let us remark that in [5] the symbol \mathcal{R} was used, but in [5] it was not assumed that the ℓ -groups under consideration were abelian.)

There exists an injective mapping of the class of all infinite cardinals into \mathcal{R}_a (this follows from the construction in [5], Section 3). Hence in view of 4.2, there exists an injective mapping of the class of all infinite cardinals into \mathcal{R}_c .

Suppose that I is a nonempty class and that for each $i \in I$, Y_i is a radical class of cl-groups. Put

$$Z_1 = \bigcap_{i \in I} Y_i.$$

Then in view of 1.4, Z_1 is a radical class of cl-groups. We obviously have

$$Z_1 = \inf\{Y_i\}_{i \in I}.$$

We express this fact by writing

$$Z_1 = \bigwedge_{i \in I} Y_i.$$

Further, we put

$$Z_2 = \text{Join Sub}_c \bigcup_{i \in I} Y_i.$$

Then 4.1 yields that the relation

$$Z_2 = \sup\{Y_i\}_{i \in I}$$

is valid in \mathcal{R}_c . We express this fact by writing

$$Z_2 = \bigvee_{i \in I} Y_i.$$

We clearly have

$$\text{Sub}_c \bigcup_{i \in I} Y_i = \bigcup_{i \in I} \text{Sub}_c Y_i.$$

Since each Y_i is a radical class of cl-groups we obtain $\text{Sub}_c Y_i = Y_i$. Hence

$$\bigvee_{i \in I} Y_i = \text{Join} \bigcup_{i \in I} Y_i.$$

4.3. Theorem. *Let $\{Y_i\}_{i \in I}$ be as above and let Y be a radical class of cl-groups. Then*

$$Y \wedge \left(\bigvee_{i \in I} Y_i \right) = \bigvee_{i \in I} (Y \wedge Y_i).$$

Proof. We have

$$\bigvee_{i \in I} (Y \wedge Y_i) \leq Y \wedge \left(\bigvee_{i \in I} Y_i \right).$$

Let $(G, \alpha) \in Y \wedge \left(\bigvee_{i \in I} Y_i \right)$. Thus $(G, \alpha) \in Y$ and

$$(G, \alpha) \in \text{Join} \bigcup_{i \in I} Y_i.$$

Then there exist cl-groups (G_k, α) ($k \in K$) such that, for each $k \in K$,

$$(G_k, \alpha) \in c(G, \alpha) \cap \left(\bigcup_{i \in I} Y_i \right)$$

and

$$(G, \alpha) = \bigvee_{k \in K}^c (G_k, \alpha).$$

Hence for each $k \in K$ there exists $i(k) \in I$ with $(G_k, \alpha) \in Y_{i(k)}$. Denote

$$I_1 = \{i(k) : k \in K\}.$$

Thus $(G_k, \alpha) \in Y \wedge Y_{i(k)}$ and

$$(G, \alpha) \in \text{Join} \bigcup_{i \in I_1} (Y \wedge Y_{i(k)}) \leq \text{Join} \bigcup_{i \in I} (Y \wedge Y_i) = \bigvee_{i \in I} (Y \wedge Y_i).$$

□

References

- [1] *P. Conrad*: K -radical classes of lattice ordered groups. Algebra, Proc. Conf. Carbondale (1980), Lecture Notes Math. 848 (1981), 186–207.
- [2] *M. Darnel*: Closure operations on radicals of lattice ordered groups. Czechoslovak Math. J. 37 (1987), 51–64.
- [3] *R. Frič, V. Koutník*: Sequential convergence spaces: Iteration, extension, completion, enlargement. Recent Progress in General Topology. Elsevier Sci. Publ., Amsterdam, 1992, pp. 201–213.
- [4] *J. Jakubík*: Direct decompositions of partially ordered groups, II. Czechoslovak Math. J. 11 (1961), 490–515. (In Russian.)
- [5] *J. Jakubík*: Radical mappings and radical classes of lattice ordered groups. Symposia Math. 21. Academic Press, New York, 1977, pp. 451–477.
- [6] *J. Jakubík*: Sequential convergences in ℓ -groups without Urysohn's axiom. Czechoslovak Math. J. 42 (1992), 101–116.
- [7] *N. Ya. Medvedev*: On the lattice of radicals of a finitely generated ℓ -group. Math. Slovaca 33 (1983), 185–188. (In Russian.)
- [8] *Dao-Rong Ton*: Product radical classes of ℓ -groups. Czechoslovak Math. J. 42 (1992), 129–142.

Author's address: Ján Jakubík, Matematický ústav SAV, Grešákova 6, 040 01 Košice, Slovakia.