125 (2000) MATHEMATICA BOHEMICA No. 1, 39–54

A GENERALIZED MAXIMUM PRINCIPLE FOR BOUNDARY VALUE PROBLEMS FOR DEGENERATE PARABOLIC OPERATORS WITH DISCONTINUOUS COEFFICIENTS

Salvatore Bonafede, Palermo, Francesco Nicolosi, Catania

(Received March 8, 1999)

Dedicated to Professor Alois Kufner on the occasion of his 65th birthday

Abstract. We prove a generalized maximum principle for subsolutions of boundary value problems, with mixed type unilateral conditions, associated to a degenerate parabolic second-order operator in divergence form.

Keywords: weak subsolution, generalized maximum principle, comparison theorem, degenerate equation

MSC 2000: 35B50, 35K10, 35K65, 35K85

1. Introduction

In [14] M. G. Platone Garroni has extended the classical generalized maximum principle (see, for instance, [15]), when the coefficients of the operator are discontinuous, to subsolutions of elliptic linear second order equations with mixed type boundary unilateral conditions, that is, on a portion of the boundary $\partial\Omega$ of Ω, the values of the solution are assigned, while on the other part a unilateral condition on the solution and its conormal derivative is given. In the present paper we will establish a similar result (see Theorem 5.1) for degenerate parabolic equations, using a technique different from that of [14]. As a corollary, we obtain a comparison theorem (see Theorem 6.1). Our procedure, rather similar to that followed in [12] and in [13] allows us to obtain more general results. Other sufficient conditions for the boundedness of weak subsolutions of Cauchy-Dirichlet problem, in the non degenerate case, may be obtained from [6] and [17], while in the degenerate case some results are announced in [3] and in [4].

2. Functional spaces

Let $\mathbb{R}^m$ be the Euclidean space ($m > 2$) with a generic point $x = (x_1, x_2, \dots, x_m)$, Ω a bounded open subset of $\mathbb{R}^m$ whose boundary satisfies locally a Lipschitz condition, T a real positive number. Let us denote by $Q(\tau_1, \tau_2)$ ($0 \leqslant \tau_1 < \tau_2 \leqslant T$) the cylinder $\Omega\times]\tau_1, \tau_2[$ and let $Q = Q(0, T)$; Γ is the parabolic boundary of Q, that is $\Gamma = \big(\Omega \times \{t = 0\}\big) \cup \big(\partial\Omega\times]0, T[\big)$.

Let $\partial\Omega_2$ be a closed subset of $\partial\Omega$, $\Gamma_2 = \partial\Omega_2 \times [0, T]$, and let us set $\partial\Omega_1 = \partial\Omega \setminus \partial\Omega_2$, $\Gamma_1 = \partial\Omega_1 \times [0, T]$.

The symbol meas_x will henceforth denote the m-dimensional measure.

If $u(x)$ is a measurable function defined in Ω, we will denote by $|u|_p$ ($1 \leqslant p \leqslant \infty$) the usual norm in the space $L^p(\Omega)$.

Hypothesis 2.1. Let $\nu(x)$ be a positive function defined in Ω such that

$$\nu(x) \in L^{\frac{q(m-1)}{q-m}}(\Omega), \quad \nu^{-1}(x) \in L^q(\Omega), \ \ q > m.$$

The symbol $\widetilde{H}^1(\nu, \Omega)$ stands for the completion of $C^1\big(\overline{\Omega}\big)$ with respect to the norm

$$\|u\|_1 = \left(|u|_2^2 + \sum_{i=1}^m \nu \Big|\frac{\partial u}{\partial x_i}\Big|_2^2\right)^{\frac{1}{2}};$$

$C^*(\Omega)$ denotes the following linear subspace of $C^\infty\big(\overline{\Omega}\big)$:

$$C^*(\Omega) = \big\{u \in C^\infty\big(\overline{\Omega}\big) \colon \ u = 0 \text{ on } \partial\Omega_2\big\}.$$

$H^*(\nu, \Omega)$ denotes the closure of $C^*(\Omega)$ in $\widetilde{H}^1(\nu, \Omega)$.

If $u(x, t)$ is a measurable real function in Q, we will denote by $|u|_{p,q}$ ($1 \leqslant p$, $q \leqslant +\infty$) the usual norm in the space $L^{p,q}(Q)$, with the obvious modification if p or q are $+\infty$.

Hypothesis 2.2. Let $\psi(t)$ be a positive monotone nondecreasing function defined in $]0, T[$ such that

$$\psi(t) \in L^1(0, T).$$

The symbol $\widetilde{H}^{1,0}\big(\nu\psi, Q(\tau_1, \tau_2)\big)$ ($0 \leqslant \tau_1 < \tau_2 \leqslant T$) stands for the completion of $C^1\big(\overline{Q(\tau_1, \tau_2)}\big)$ with respect to the norm

$$\|u\|_{1,0,(\tau_1,\tau_2)} = \left(\int_{Q(\tau_1,\tau_2)} \left(|u|^2 + \sum_{i=1}^m \nu\psi \Big|\frac{\partial u}{\partial x_i}\Big|^2\right) \mathrm{d}x\,\mathrm{d}t\right)^{\frac{1}{2}};$$

$$\|u\|_{1,0} = \|u\|_{1,0,(0,T)}.$$

– $\widetilde{H}^{1,0}\big(\nu\psi, Q(\tau_1,\tau_2)\big)$ is a Hilbert space with respect to the norm $\|u\|_{1,0,(\tau_1,\tau_2)}$.
– $C^*\big(Q(\tau_1,\tau_2)\big)$ $(0 \leqslant \tau_1 < \tau_2 \leqslant T)$ denotes the following linear subspace of $C^\infty\big(Q(\tau_1,\tau_2)\big) \cap C^0\big(\overline{Q(\tau_1,\tau_2)}\big)$:
– $C^*\big(Q(\tau_1,\tau_2)\big) = \big\{u \in C^\infty\big(Q(\tau_1,\tau_2)\big) \cap C^0\big(\overline{Q(\tau_1,\tau_2)}\big)\colon\ u = 0 \text{ on } \partial\Omega_2 \times [\tau_1,\tau_2]\big\}$.
– $\widetilde{H}^{1,0}_*\big(\nu\psi, Q(\tau_1,\tau_2)\big)$ $(0 \leqslant \tau_1 < \tau_2 \leqslant T)$ is the closure of $C^*\big(Q(\tau_1,\tau_2)\big)$ in $\widetilde{H}^{1,0}\big(\nu\psi, Q(\tau_1,\tau_2)\big)$.

Finally, we will denote by $V^{1,0}(\nu\psi, Q)$ the space of functions $u(x,t)$ belonging to $\widetilde{H}^{1,0}(\nu\psi, Q)$, continuous in $[0,T]$ with values in $L^2(\Omega)$.[1]

Definition 1. Given a real number h, if $u \in \widetilde{H}^{1,0}\big(\nu\psi, Q(\tau_1,\tau_2)\big)$ $(0 \leqslant \tau_1 < \tau_2 \leqslant T)$, we will say that $u(x,t) \leqslant h$ $(\geqslant h)$ on $\partial\Omega_i \times [\tau_1,\tau_2]$ $(i = 1, 2)$ if there exists a sequence $\{u_n\}$ of functions from $C^1\big(\overline{Q(\tau_1,\tau_2)}\big)$ such that

$$u_n(x,t) \leqslant h \ (\geqslant h) \text{ on } \partial\Omega_i \times [\tau_1,\tau_2]$$

and

$$(*) \qquad \lim_{n\to\infty} \big\|u_n - u\big\|_{1,0,(\tau_1,\tau_2)} = 0.$$

If k is such that $u(x,t) \leqslant k$ on $\partial\Omega_i \times [\tau_1,\tau_2]$, we will say that $u(x,t)$ is bounded from above on $\partial\Omega_i \times [\tau_1,\tau_2]$.

Definition 2. If $u(x,t)$, $w(x,t)$ belong to $\widetilde{H}^{1,0}\big(\nu\psi, Q(\tau_1,\tau_2)\big)$ $(0 \leqslant \tau_1 < \tau_2 \leqslant T)$ and $w(x,t) \geqslant 0$ on $\partial\Omega_i \times [\tau_1,\tau_2]$ $(i = 1, 2)$, let us denote[2]

$$\sup{}^* \frac{u}{w} = \inf\big\{h \in \mathbb{R}\colon\ u(x,t) - hw(x,t) \leqslant 0 \text{ on } \partial\Omega_i \times [\tau_1,\tau_2]\big\}.$$

We will consider the following generalized problem:

$$(2.1) \qquad \begin{cases} -\sum_{i=1}^m \frac{\partial}{\partial x_i}\Big(\sum_{j=1}^m a_{ij}\frac{\partial u}{\partial x_j} + d_i u\Big) + \Big(\sum_{i=1}^m b_i \frac{\partial u}{\partial x_i} + cu\Big) + \frac{\partial u}{\partial t} = 0 & \text{in } Q \\ \frac{\partial u}{\partial \nu} + \alpha\, u + \sum_{i=1}^m d_i u \cos nx_i \geqslant 0 & \text{on } \Gamma_1, \end{cases}$$

where

$$\frac{\partial u}{\partial \nu} = \sum_{i,j=1}^m a_{ij} \cos nx_j \frac{\partial u}{\partial x_i},$$

and $\cos nx_j$ is the j-th directional cosine of x, normal to Γ_1 and external to Q.

[1] For more details concerning hypotheses (2.1), (2.2) see also [5], [7], [8] and [9].
[2] We suppose that $\inf \emptyset = +\infty$.

By an L_{Γ_1}-subsolution (L_{Γ_1}-supersolution) of problem (2.1) we mean any function $u \in V^{1,0}(\nu\psi, Q)$ satisfying the following conditions:

$$\tilde{a}(u,\varphi) = \int_Q \Bigg(\sum_{i,j=1}^m a_{ij} \frac{\partial u}{\partial x_j} \frac{\partial \varphi}{\partial x_i} + \sum_{i=1}^m b_i \frac{\partial u}{\partial x_i} \varphi + cu\varphi + \sum_{i=1}^m d_i u \frac{\partial \varphi}{\partial x_i} - u \frac{\partial \varphi}{\partial t} \Bigg) \mathrm{d}x\, \mathrm{d}t + \int_{\Gamma_1} \alpha u \varphi \, \mathrm{d}\sigma \, \mathrm{d}t \leqslant 0 \quad (\geqslant 0) \tag{2.2}$$

for any $\varphi \in C^*(Q)$ such that $\varphi(x,t) \geqslant 0$ a.e. in Q, $\varphi(x,0) = \varphi(x,T) = 0$ for a.e. $x \in \Omega$.

Of particular interest are L_{Γ_1}-subsolutions (L_{Γ_1}-supersolutions) such that

$$\int_Q \Bigg(\sum_{i,j=1}^m a_{ij} \frac{\partial u}{\partial x_j} \frac{\partial \varphi}{\partial x_i} + \sum_{i=1}^m b_i \frac{\partial u}{\partial x_i} \varphi + cu\varphi + \sum_{i=1}^m d_i u \frac{\partial \varphi}{\partial x_i} - u \frac{\partial \varphi}{\partial t} \Bigg) \mathrm{d}x\, \mathrm{d}t + \int_{\Gamma_1} \alpha u \varphi \, \mathrm{d}\sigma \, \mathrm{d}t \leqslant 0 \quad (\geqslant 0) \tag{2.3}$$

for any $\varphi \in C^*(Q)$, $\varphi(x,t) \geqslant 0$ on Γ_1, $\varphi(x,0) = \varphi(x,T) = 0$ for a.e. $x \in \Omega$.

In fact, problem (2.3) is equivalent, at least "formally," to the problem

$$\begin{cases} -\sum\limits_{i=1}^m \dfrac{\partial}{\partial x_i} \Bigg(\sum\limits_{j=1}^m a_{ij} \dfrac{\partial u}{\partial x_j} + d_i u \Bigg) + \Bigg(\sum\limits_{i=1}^m b_i \dfrac{\partial u}{\partial x_i} + cu \Bigg) + \dfrac{\partial u}{\partial t} = 0 & \text{in } Q \\ \dfrac{\partial u}{\partial \nu} + \alpha u + \sum\limits_{i=1}^m d_i u \cos n x_i \leqslant 0 \quad (\geqslant 0) & \text{on } \Gamma_1. \end{cases}$$

Let us consider the problem

$$\begin{cases} \displaystyle\int_Q \Bigg(\sum_{i,j=1}^m a_{ij} \frac{\partial u}{\partial x_j} \frac{\partial \varphi}{\partial x_i} + \sum_{i=1}^m b_i \tfrac{\partial u}{\partial x_i} \varphi + cu\varphi + \sum_{i=1}^m d_i u \frac{\partial \varphi}{\partial x_i} - u \frac{\partial \varphi}{\partial t} \Bigg) \mathrm{d}x\, \mathrm{d}t \\ \qquad + \displaystyle\int_{\Gamma_1} \alpha u \varphi \, \mathrm{d}\sigma \, \mathrm{d}t = \int_Q f\varphi \, \mathrm{d}x \, \mathrm{d}t + \int_{\Gamma_1} g_1 \varphi \, \mathrm{d}\sigma \, \mathrm{d}t \\ \qquad\qquad\qquad\qquad\qquad\qquad \text{for any } \varphi \in C^*(Q),\ \varphi(x,T) = 0 \text{ in } \Omega \\ u(x,t) - g_2(x,t) \in \widetilde{H}^{1,0}_*(\nu\psi, Q) \\ u(x,0) = 0 \text{ in } \Omega. \end{cases} \tag{2.4}$$

The problem (2.4) is formally equivalent to the problem

$$\begin{cases} -\sum\limits_{i=1}^m \dfrac{\partial}{\partial x_i} \Bigg(\sum\limits_{j=1}^m a_{ij} \dfrac{\partial u}{\partial x_j} + d_i u \Bigg) + \Bigg(\sum\limits_{i=1}^m b_i \dfrac{\partial u}{\partial x_i} + cu \Bigg) + \dfrac{\partial u}{\partial t} = f & \text{in } Q \\ \dfrac{\partial u}{\partial \nu} + \alpha u + \sum\limits_{i=1}^m d_i u \cos n x_i = g_1 & \text{on } \Gamma_1 \\ u = g_2 & \text{on } \Gamma_2 \\ u(x,0) = 0 & \text{in } \Omega. \end{cases}$$

3. Hypotheses on coefficients

Let us denote by A the set of pairs (α^*, α) with $2 \leqslant \alpha^*, \alpha \leqslant +\infty$, such that there exists a positive constant β for which

$$\|u\|_{\alpha^*,\alpha} \leqslant \beta\big(\|u\|_{2,\infty} + \|u\|_{1,0}\big) \tag{3.1}$$

for any $u \in L^{2,\infty}(Q) \cap \widetilde{H}^{1,0}(\nu\psi, Q)$. The set A obviously contains[3] the pair $(2, +\infty)$.

Let us observe that, under the hypotheses on Ω, we have[4]

$$|u|_{\frac{2(m-1)}{m-2+\frac{m}{g}},\partial\Omega_1} \leqslant \gamma\|u\|_1 \quad \text{for any } u \in \widetilde{H}^1(\nu, \Omega). \tag{3.2}$$

Consequently, we obtain:

$$\left(\int_0^T \psi(t)\,|u|^2_{\frac{2(m-1)}{m-2+\frac{m}{g}},\partial\Omega_1}\,\mathrm{d}t\right)^{\frac{1}{2}} \leqslant \gamma\big(\|u\|_{2,\infty} + ||u||_{1,0}\big) \tag{3.3}$$

for any $u \in L^{2,\infty}(Q) \cap \widetilde{H}^{1,0}(\nu\psi, Q)$.

The constant in (3.2) and (3.3) depends on Ω_1.

H y p o t h e s i s 3.1. The functions $a_{ij}(x,t)$, $b_i(x,t)$, $c(x,t)$, $d_i(x,t)$ $(1 \leqslant i, j \leqslant m)$ are defined and measurable in Q;

$$\frac{a_{ij}}{\nu\psi} \in L^\infty(Q), \quad \frac{b_i}{\sqrt{\nu\psi}} \in L^{p^*,p}(Q), \quad c \in L^{q^*,q}(Q), \quad \frac{d_i}{\sqrt{\nu\psi}} \in L^{r^*,r}(Q),$$

where

$$\frac{1}{p^*} + \frac{1}{\alpha_1^*} = \frac{1}{2}, \qquad \frac{1}{p} + \frac{1}{\alpha_1} = \frac{1}{2}, \qquad \frac{1}{q^*} + \frac{2}{\alpha_2^*} = 1,$$
$$\frac{1}{q} + \frac{2}{\alpha_2} = 1, \qquad \frac{1}{r^*} + \frac{1}{\alpha_3^*} = \frac{1}{2}, \qquad \frac{1}{r} + \frac{1}{\alpha_3} = \frac{1}{2}$$

with (α_1^*, α_1), (α_2^*, α_2) and (α_3^*, α_3) belonging to A.

Moreover, if $p = +\infty$ $[q = +\infty,\ r = +\infty]$ and $p^* < +\infty$ $[q^* < +\infty,\ r^* < +\infty]$, then there exists a function $\eta_1(\sigma)$ $[\eta_2(\sigma),\ \eta_3(\sigma)]$, defined for $\sigma \geqslant 0$, non decreasing,

[3] If $\frac{1}{\psi(t)} \in L^t(0,T)$ $(0 < t \leqslant +\infty)$, the set A contains the pair $\big(\frac{2mg}{mg+m\theta-2\theta g}, \frac{2t}{\theta(t+1)}\big)$ for any $\theta \in \big[0, \frac{t}{t+1}\big[$, see, for instance, [13].

[4] See, for instance, [11] Theorem 3.9.

vanishing for σ approaching zero and satisfying for almost all t in the interval $]0,T[$ the inequalities

$$\sum_{i=1}^{m}\Big(\int_E\Big(\frac{|b_i(x,t)|}{\sqrt{\nu(x)}}\Big)^{p^*}\,\mathrm{d}x\Big)^{\frac{1}{p}}\leqslant\eta_1(\sigma)\sqrt{\psi(t)},$$
$$\Big[\Big(\int_E\big(|c(x,t)|-c(x,t)\big)^{q^*}\,\mathrm{d}x\Big)^{\frac{1}{q}}\leqslant\eta_2(\sigma),$$
$$\sum_{i=1}^{m}\Big(\int_E\Big(\frac{|d_i(x,t)|}{\sqrt{\nu(x)}}\Big)^{r^*}\,\mathrm{d}x\Big)^{\frac{1}{r}}\leqslant\eta_3(\sigma)\sqrt{\psi(t)}\Big]$$

for all measurable subsets E of Ω such that $\mathrm{meas}_x\,E\leqslant\sigma$.

The function $\alpha(x,t)$ is defined and measurable on Γ_1 and

$$\frac{\alpha}{\psi}\in L^\infty\Big(0,T;L^{\frac{m-1}{1-\frac{m}{g}}}\big(\partial\Omega_1\big)\Big).$$

H y p o t h e s i s 3.2. The functions $f(x,t)$, $g(x,t)$ are defined and measurable respectively in Q and in Γ_1, moreover

$$f\in L^2(Q),\quad \frac{g_1}{\sqrt{\psi}}\in L^2\Big(0,T;L^{\frac{2(m-1)}{m-\frac{m}{g}}}\big(\partial\Omega_1\big)\Big).$$

The function $g_2(x,t)$ is defined and measurable in Q and

$$g_2\in\widetilde{H}^{1,0}(\nu\psi,Q),\quad \frac{\partial g_2}{\partial t}\in L^2(Q),\quad g_2(x,t)\leqslant 0\text{ on }\Gamma_1.$$

Finally, the functions $f^*(x,t)$, $g_1^*(x,t)$ are defined and measurable respectively in Q and in Γ_1, moreover

$$f^*\in L^2(Q),\quad \frac{g_1^*}{\sqrt{\psi}}\in L^2\Big(0,T;L^{\frac{2(m-1)}{m-\frac{m}{g}}}\big(\partial\Omega_1\big)\Big).$$

H y p o t h e s i s 3.3. The following inequality holds for a.e. (x,t) in Q and for all real numbers $\chi_1,\chi_2,\ldots,\chi_m$:

$$\sum_{i,j=1}^{m}a_{ij}(x,t)\chi_i\chi_j\geqslant\nu(x)\psi(t)\sum_{i=1}^{m}\chi_i^2.$$

4. Preliminary lemmas

Lemma 4.1. *Let us assume that hypotheses* (2.1), (2.2), (3.1) *hold and let* $u(x,t)$ *be an* L_{Γ_1}*-subsolution of the problem* (2.1) *bounded from above on* $\partial\Omega_2\times[0,T]$. *Then if* $0 \leqslant \tilde{\tau}_1 < \tau < T$ *and* $k > \sup^* u$, *we get*

$$\int_{Q(\tilde{\tau}_1,\tau)} \Big(\sum_{i,j=1}^{m} a_{ij}\frac{\partial v}{\partial x_j}\frac{\partial v}{\partial x_i} + \sum_{i=1}^{m} b_i \frac{\partial v}{\partial x_i} v + cuv + \sum_{i=1}^{m} d_i u \frac{\partial v}{\partial x_i} \Big)\,\mathrm{d}x\,\mathrm{d}t + \frac{1}{2}\int_\Omega v^2(x,\tau)\,\mathrm{d}x + \int_{\tilde{\tau}_1}^{\tau}\int_{\partial\Omega_1} \alpha uv\,\mathrm{d}\sigma\,\mathrm{d}t \leqslant \frac{1}{2}\int_\Omega v^2(x,\tilde{\tau}_1)\,\mathrm{d}x,$$

where $v = u - \min(u,k)$ *in* Q; *moreover,* $v \in \widetilde{H}^{1,0}_*(\nu\psi, Q)$.[5]

P r o o f. Let $\tilde{\tau}_1$, τ be such that $0 < \tilde{\tau}_1 < \tau < T$; setting $\tau_1 = \frac{\tau+T}{2}$, $\tau_2 = T - \tau_1$, we denote by $C^\infty_\tau(Q)$ the set of nonnegative functions from $C^*(Q)$ equal to zero for $t \geqslant \tau_1$. Let φ be a function from $C^\infty_\tau(Q)$. We extend u, φ and the coefficients of (2.1) to $\Omega\times(-\infty,+\infty)$, assuming that these functions are equal at zero in those points where they are not defined.

We define in $\Omega\times\,]-\infty,+\infty[$ and for any integer ϱ:

$$\Phi_\varrho(x,t) = \frac{\varrho}{\tau_2}\int_{t-\frac{\tau_2}{\varrho}}^{t} \varphi(x,\lambda)\,\mathrm{d}\lambda,$$

$$U_\varrho(x,t) = \frac{\varrho}{\tau_2}\int_{t}^{t+\frac{\tau_2}{\varrho}} u(x,\lambda)\,\mathrm{d}\lambda,$$

$$B_\varrho(x,t) = \frac{\varrho}{\tau_2}\int_{t}^{t+\frac{\tau_2}{\varrho}} \sum_{i=1}^{m} b_i(x,\lambda)\frac{\partial u(x,\lambda)}{\partial x_i}\,\mathrm{d}\lambda,$$

$$C_\varrho(x,t) = \frac{\varrho}{\tau_2}\int_{t}^{t+\frac{\tau_2}{\varrho}} c(x,\lambda)u(x,\lambda)\,\mathrm{d}\lambda,$$

$$A_{i,\varrho}(x,t) = \frac{\varrho}{\tau_2}\int_{t}^{t+\frac{\tau_2}{\varrho}} \sum_{j=1}^{m} a_{ij}(x,\lambda)\frac{\partial u(x,\lambda)}{\partial x_j}\,\mathrm{d}\lambda,$$

$$D_{i,\varrho}(x,t) = \frac{\varrho}{\tau_2}\int_{t}^{t+\frac{\tau_2}{\varrho}} d_i(x,\lambda)u(x,\lambda)\,\mathrm{d}\lambda,$$

$$\alpha_\varrho(x,t) = \frac{\varrho}{\tau_2}\int_{t}^{t+\frac{\tau_2}{\varrho}} \alpha(x,\lambda)u(x,\lambda)\,\mathrm{d}\lambda.$$

[5] Let us observe that, for t a.e. in $]0,T[$, $v(x,t) \in H^*(\nu,\Omega)$.

From (2.2), in virtue of $\varphi = \Phi_\varrho(x,t)$, via an exchange of the order of integration with respect to t and λ,[6] we get

$$\int_Q \Big(\sum_{i=1}^m A_{i,\varrho} \frac{\partial \varphi}{\partial x_i} + B_\varrho \varphi + C_\varrho \varphi + \sum_{i=1}^m D_{i,\varrho} \frac{\partial \varphi}{\partial x_i} + \frac{\partial U_\varrho}{\partial t} \varphi \Big) \,\mathrm{d}x \,\mathrm{d}t + \int_{\Gamma_1} \alpha_\varrho \varphi \,\mathrm{d}\sigma \,\mathrm{d}t \leqslant 0 \tag{4.1}$$

for all φ belonging to the functional class $C^\infty_\tau(Q)$.

Let $\{u_n\}$ be a sequence of functions of $C^1\big(\overline{Q}\big)$ such that $u_n < \sup^* u$ on Γ_2 and satisfying $(*)$. For all pairs of positive integers ν and n, we define

$$U_{\varrho,n}(x,t) = \frac{\varrho}{\tau_2} \int_t^{t+\frac{\tau_2}{\varrho}} u_n(x,\lambda) \,\mathrm{d}\lambda;$$

the function $U_{\varrho,n}(x,t)$ belongs to $C^1\big(\overline{Q(0,\tau_1)}\big)$.

Let us now introduce the function[7]

$$V_{\varrho,n} = \begin{cases} U_{\varrho,n} - \min(U_{\varrho,n}, k) & \text{in } Q(\tilde\tau_1, \tau), \\ 0 & \text{in } Q \setminus Q(\tilde\tau_1, \tau). \end{cases}$$

Let $\{\Phi_\mu\}_{\mu\in\mathbb{N}}$ be a sequence of nonnegative equibounded functions from $C^\infty_\tau(Q)$ converging to $V_{\varrho,n}$ in $\widetilde{H}^{1,0}\big(\nu\psi, Q(\tilde\tau_1,\tau)\big)$;[8] moreover, let also the functions in the sequence $\big\{\frac{\partial \Phi_\mu}{\partial x_i}\big\}_{\mu\in\mathbb{N}}$ be equibounded.

From (4.1), in virtue of $\varphi = \Phi_\mu(x,t)$, as μ diverges to $+\infty$, we obtain the following relation:

$$\int_{Q(\tilde\tau_1,\tau)} \Big(\sum_{i=1}^m A_{i,\varrho} \frac{\partial V_{\varrho,n}}{\partial x_i} + B_\varrho V_{\varrho,n} + C_\varrho V_{\varrho,n} + \sum_{i=1}^m D_{i,\varrho} \frac{\partial V_{\varrho,n}}{\partial x_i} + \frac{\partial U_\varrho}{\partial t} V_{\varrho,n} \Big) \,\mathrm{d}x \,\mathrm{d}t + \int_{\tilde\tau_1}^{\tau} \int_{\partial\Omega_1} \alpha_\nu V_{\varrho,n} \,\mathrm{d}\sigma \,\mathrm{d}t \leqslant 0. \tag{4.2}$$

Setting now in Q:

$$V_\varrho = \begin{cases} U_\varrho - \min(U_\varrho, k) & \text{in } Q(\tilde\tau_1, \tau), \\ 0 & \text{in } Q \setminus Q(\tilde\tau_1, \tau); \end{cases}$$

[6] See [10], p. 141.

[7] Since $k > \sup^* u$ there exists a neighbourhood of $\partial\Omega_2$ such that $V_{\varrho,n}(x,t) = 0$ for any $t \in]0,T[$ (see Lemma 4.2 of [3]).

[8] See remark 4.1 of [3].

the sequence $\{V_{\varrho,n}\}$ converges to V_ϱ in $\widetilde{H}^{1,0}\big(\nu\psi, Q(\tilde\tau_1,\tau)\big)\cap L^{2,\infty}\big(Q(\tilde\tau_1,\tau)\big)$ and satisfies the relation

$$\lim_{n\to\infty}\big\|(V_{\varrho,n}-V_\varrho)\big\|_{\frac{2(m-1)}{m-2+\frac{m}{g}},2,\Gamma_1}=0.$$

On the other hand, the functions of the sequence $\{V_{\varrho,n}\}$ belong to $\widetilde{H}^{1,0}_*\big(\nu\psi, Q(\tilde\tau_1,\tau)\big)$ and so also V_ϱ belongs to this space.

From (3.1)–(3.6) we deduce, as n goes to $+\infty$, the following inequality:[9]

$$\begin{aligned}(4.3)\qquad &\int_Q\Big(\sum_{i=1}^m A_{i,\varrho}\frac{\partial V_\varrho}{\partial x_i}+B_\varrho V_\varrho+C_\varrho V_\varrho+\sum_{i=1}^m D_{i,\varrho}\frac{\partial V_\varrho}{\partial x_i}+\frac{\partial U_\varrho}{\partial t}V_{\varrho,n}\Big)\,\mathrm{d}x\,\mathrm{d}t\\ &\quad+\frac12\int_{\Omega_\varrho(\tau,k)}\big|U_\varrho(x,\tau)-k\big|^2\,\mathrm{d}x\\ &\quad-\frac12\int_{\Omega_\varrho(0,k)}\big|U_\varrho(x,0)-k\big|^2\,\mathrm{d}x+\int_{\tilde\tau_1}^\tau\int_{\partial\Omega_1}\alpha_\varrho V_\varrho\,\mathrm{d}\sigma\,\mathrm{d}t\leqslant 0.\end{aligned}$$

Let us remark that the sequence $\{V_\varrho\}$ converges in $\widetilde{H}^{1,0}(\nu\psi,Q)\cap L^{2,\infty}(Q)$ to the function equal to v in $Q(\tilde\tau_1,\tau)$ and equal to zero in $Q\setminus Q(\tilde\tau_1,\tau)$.

From (4.3), the conclusion follows via another passage to the limit.

For example, we prove that

$$\lim_{n\to\infty}\int_{\tilde\tau_1}^\tau\int_{\partial\Omega_1}\alpha_\varrho V_\varrho\,\mathrm{d}\sigma\,\mathrm{d}t=\int_{\tilde\tau_1}^\tau\int_{\partial\Omega_1}\alpha uv\,\mathrm{d}\sigma\,\mathrm{d}t.$$

We get

$$\begin{aligned}&\Big|\int_{\tilde\tau_1}^\tau\int_{\partial\Omega_1}\alpha_\varrho V_\varrho-\alpha uv\,\mathrm{d}\sigma\,\mathrm{d}t\Big|\\ &\qquad\leqslant\gamma\,\frac{\psi(\tau)}{\psi(\tilde\tau_1)}\Big\|\frac{\alpha}{\psi}\Big\|_{\frac{m-1}{1-\frac{m}{g}},\infty,\Gamma_1}\big(\|u\|_{2,\infty}+\|u\|_{1,0}\big)\\ &\qquad\quad\times\big(\|V_\varrho-v\|_{2,\infty,(\tilde\tau_1,\tau)}+\|V_\varrho-v\|_{1,0,(\tilde\tau_1,\tau)}\big)+\Big(\frac{1}{\psi(\tilde\tau_1)}\Big)^{\frac12}\\ &\qquad\quad\times\big(\|v\|_{2,\infty,(\tilde\tau_1,\tau)}+\|v\|_{1,0,(\tilde\tau_1,\tau)}\big)\Big(\int_{\tilde\tau_1}^\tau\Big(\big|\alpha_\varrho-\alpha u\big|^{\frac{2(m-1)}{m-\frac{m}{g}}}\,\mathrm{d}\sigma\Big)^{\frac{m-\frac{m}{g}}{m-1}}\,\mathrm{d}t\Big)^{\frac12}\end{aligned}$$

for any $\varrho\in\mathbb{N}$.[10]

[9] For a fixed $t\in]0,T[$, we set $\Omega_\varrho(t,k)=\{x\in\Omega\colon\ U_\varrho(x,t)>k\}$.

[10] Let us remark that, by the properties of Steklov averages, it follows that α_ϱ converges to αu in $L^2\Big(\tilde\tau_1,\tau;L^{\frac{2(m-1)}{m-\frac{m}{g}}}(\partial\Omega_1)\Big)$.

Next, it is easy to verify that the restriction of the function v to $Q(\tilde\tau_1,\tau)$ belongs to $\widetilde H^{1,0}_*(\nu\psi,Q)$ for any $0<\tilde\tau_1<\tau<T$ and, therefore, since v by definition belongs to $\widetilde H^{1,0}\big(\nu\psi,Q(\tilde\tau_1,\tau)\big)$, it belongs to $\widetilde H^{1,0}_*(\nu\psi,Q)$, too.

Finally, if $\tilde\tau_1=0$, as $\tau>0$ is assumed, it suffices to consider $\tau_n=\frac{\tau}{n+1}$ for $n\in\mathbb{N}$ recalling that the function $v(x,t)$ is continuous in $[0,T]$ with values in $L^2(\Omega)$. □

Lemma 4.2. *Let us assume the hypotheses* (2.1), (2.2), (3.1), (3.3) *hold and let* $u(x,t)$ *be an* L_{Γ_1}*-subsolution of the problem* (2.1) *satisfying the conditions*

$$\operatorname*{ess\,sup}_{\Omega} u(x,0)\leqslant 0,\quad \sup{}^{*}\,u\leqslant 0.$$

Then, we have:

$$\operatorname*{ess\,sup}_{Q} u(x,t)\leqslant 0.$$

P r o o f. For any integer n, we consider the functions

$$v=u-\min(u,0),\quad v_n=u-\min\Big(u,\frac{1}{n}\Big).$$

From Lemma 4.1 we deduce that v_n belongs to $\widetilde H^{1,0}_*(\nu\psi,Q)$ and that, provided $\tau\in]0,T[$, we have

$$\int_{Q(\tau)}\Big(\sum_{i,j=1}^m a_{ij}\frac{\partial v_n}{\partial x_j}\frac{\partial v_n}{\partial x_i}+\sum_{i=1}^m b_i\frac{\partial v_n}{\partial x_i}v_n+cuv_n+\sum_{i=1}^m d_iu\frac{\partial v_n}{\partial x_i}\Big)\,\mathrm{d}x\,\mathrm{d}t + \frac12\int_\Omega v_n^2(x,\tau)\,\mathrm{d}x+\int_0^\tau\int_{\partial\Omega_1}\alpha uv_n\,\mathrm{d}\sigma\,\mathrm{d}t\leqslant 0. \tag{4.4}$$

On the other hand, we obtain

$$\lim_{n\to\infty}v_n(x,t)=v(x,t),\quad |v_n(x,t)|\leqslant|u(x,t)|\ \text{ in } Q$$

and

$$\lim_{n\to\infty}\frac{\partial v_n}{\partial x_i}=\frac{\partial v}{\partial x_i},\quad \Big|\frac{\partial v_n}{\partial x_i}\Big|\leqslant\Big|\frac{\partial u}{\partial x_i}\Big|\ \text{a.e. in } Q.$$

Furthermore, also v belongs to $\widetilde H^{1,0}_*(\nu\psi,Q)$ and so, as n goes to $+\infty$ in (4.4), we get

$$\int_{Q(\tau)}\Big(\sum_{i,j=1}^m a_{ij}\frac{\partial v}{\partial x_j}\frac{\partial v}{\partial x_i}+\sum_{i=1}^m b_i\frac{\partial v}{\partial x_i}v+cv^2+\sum_{i=1}^m d_iv\frac{\partial v}{\partial x_i}\Big)\,\mathrm{d}x\,\mathrm{d}t + \frac12\int_\Omega v^2(x,\tau)\,\mathrm{d}x+\int_0^\tau\int_{\partial\Omega_1}\alpha v^2\,\mathrm{d}\sigma\,\mathrm{d}t\leqslant 0. \tag{4.5}$$

From (4.5) we deduce that $|v|_{2,\infty}=0$ and the conclusion easily follows.

The proof is similar to that given in Lemma 4.1 of [13]; let us remark that since $v \in \widetilde{H}^{1,0}(\nu\psi, Q) \cap L^{2,\infty}(Q)$ we can apply the relations (3.1) and (3.3) instead of the hypothesis A) of [13]. □

5. A generalized maximum principle

We will prove

Theorem 5.1. *Let us assume the hypotheses* (2.1), (2.2), (3.1), (3.3) *hold and let* $w(x,t)$ *be an* L_{Γ_1}*-supersolution of the problem* (2.1) *satisfying the conditions*

$$w(x,t) > 0 \quad \text{a.e. in } Q,$$
$$w(x,0) > 0 \quad \text{a.e. in } \Omega, \quad w(x,t) \geqslant 0 \quad \text{on } \Gamma_2.$$

Then

$$\operatorname*{ess\,sup}_{Q} \frac{u(x,t)}{w(x,t)} \leqslant \max\Big(0, \operatorname*{ess\,sup}_{\Omega} \frac{u(x,0)}{w(x,0)}, \sup{}^{*} \frac{u}{w}\Big) \tag{5.1}$$

for any L_{Γ_1}*-subsolution* $u(x,t)$ *of the problem* (2.1).

Proof. The conclusion is obvious if the second term of (5.1) is equal to $+\infty$. Let us suppose, now, that this term is finite and let us denote by h some real number greater than its value. Consequently, the function $u(x,t) - hw(x,t)$ is an L_{Γ_1}-subsolution of the problem (2.1) such that $\operatorname{ess\,sup}_{\Omega}\big[u(x,0) - hw(x,0)\big] \leqslant 0$, $\sup^{*}(u - hw) \leqslant 0$. From Lemma 4.2 we can see that $u(x,t) - hw(x,t) \leqslant 0$ a.e. in Q. So, we obtain $\operatorname{ess\,sup}_{Q} \frac{u(x,t)}{w(x,t)} \leqslant h$ and the conclusion easily follows. □

6. A comparison theorem

Let us define the following closed convex sets:

$$K^{*} = \big\{z \in \widetilde{H}^{1,0}(\nu\psi, Q),\ z \in C\big([0,T]; L^2(\Omega)\big),\ z(x,0) = 0,\ z \geqslant g_2 \text{ on } \Gamma_2\big\},$$
$$\Phi^{*} = \Big\{\varphi \in \widetilde{H}^{1,0}(\nu\psi, Q),\ \frac{\partial\varphi}{\partial t} \in L^2(Q),\ \varphi(x,T) = 0,\ \varphi \geqslant g_2 \text{ on } \Gamma_2\Big\}$$

and let us suppose that there exists a solution $z \in K^{*}$ of the variational inequality

$$\begin{aligned} \int_Q \Big(\sum_{i,j=1}^{m} a_{ij} \frac{\partial z}{\partial x_j} \frac{\partial(\varphi - z)}{\partial x_i} + \sum_{i=1}^{m} b_i \frac{\partial z}{\partial x_i}(\varphi - z) + cz(\varphi - z) \\ + \sum_{i=1}^{m} d_i z \frac{\partial(\varphi - z)}{\partial x_i} - z\frac{\partial\varphi}{\partial t}\Big)\,\mathrm{d}x\,\mathrm{d}t + \int_{\Gamma_1} \alpha z(\varphi - z)\,\mathrm{d}\sigma\,\mathrm{d}t \\ \geqslant \int_Q f^{*}(\varphi - z)\,\mathrm{d}x\,\mathrm{d}t + \int_{\Gamma_1} g_1^{*}(\varphi - z)\,\mathrm{d}\sigma\,\mathrm{d}t \end{aligned} \tag{6.1}$$

for any $\varphi \in \Phi^*$. The problem (6.1) is formally equivalent to the problem

$$\begin{cases} -\sum_{i=1}^{m} \frac{\partial}{\partial x_i}\Big(\sum_{j=1}^{m} a_{ij}\frac{\partial z}{\partial x_j} + d_i z\Big) + \Big(\sum_{i=1}^{m} b_i \frac{\partial z}{\partial x_i} + cz\Big) + \frac{\partial z}{\partial t} = f^* & \text{in } Q \\ \frac{\partial z}{\partial \nu} + \alpha z + \sum_{i=1}^{m} d_i z \cos nx_i = g_1^* & \text{on } \Gamma_1 \\ z \geqslant g_2,\ \frac{\partial z}{\partial \nu} + \sum_{i=1}^{m} d_i z \cos nx_i \geqslant 0,\ (z-g_2)\Big(\frac{\partial z}{\partial \nu} + \sum_{i=1}^{m} d_i z \cos nx_i\Big) = 0 & \text{on } \Gamma_2 \\ z(x,0) = 0 & \text{in } \Omega. \end{cases}$$

We will prove

Theorem 6.1. *Let us assume the hypotheses* (2.1), (2.2), (3.1), (3.2), (3.3) *hold and let* $w(x,t)$ *be an* L_{Γ_1}*-supersolution of the problem* (2.1) *satisfying the conditions*

$$w(x,t) > 0 \quad \text{a.e. in } Q,$$
$$w(x,0) > 0 \quad \text{a.e. in } \Omega, \quad w(x,t) \geqslant 0 \text{ on } \Gamma_2.$$

Let $z(x,t)$ *be a solution of the problem* (6.1) *with* $f^* \geqslant f$ *in* Q, $g_1^* \geqslant g_1$ *on* Γ_1.

Then, we have the inequality

$$u(x,t) \leqslant z(x,t) \quad \text{a.e. in } Q$$

for any solution $u(x,t)$ *of the problem* (2.4).

P r o o f. Let us extend $z(x,t)$ to $\mathbb{R}^{m+1}$ assuming that it vanishes at points not belonging to Q; for a fixed $\tau \in]0,T[$ and for any pairs of integers ϱ, n we introduce the functions

$$\theta_n(t) = \begin{cases} 0 & \text{if } t < \tau - \frac{2}{n}, \\ n\big(t + \frac{2}{n} - \tau\big) & \text{if } \tau - \frac{2}{n} \leqslant t \leqslant \tau - \frac{1}{n}, \\ 1 & \text{if } t > \tau - \frac{1}{n}; \end{cases}$$
$$z_{n,\varrho}(x,t) = \varrho\theta_n(t) \int_{t-\frac{1}{2\varrho}}^{t+\frac{1}{2\varrho}} z(x,y)\theta_n(y)\,\mathrm{d}y.$$

We have

$$\frac{\partial z_{n,\varrho}}{\partial t} = \varrho\theta_n'(t)\int_{t-\frac{1}{2\varrho}}^{t+\frac{1}{2\varrho}} z(x,y)\theta_n(y)\,\mathrm{d}y + \varrho\theta_n(t)\Big(z\Big(x,t+\frac{1}{2\varrho}\Big)\theta_n\Big(t+\frac{1}{2\varrho}\Big) - z\Big(x,t-\frac{1}{2\varrho}\Big)\theta_n\Big(t-\frac{1}{2\varrho}\Big)\Big).$$

Choosing $\varphi = z_{n,\varrho} + \beta$, $0 \leqslant \beta \in C^*(Q)$, $\beta(x,t) = 0$ a.e. in Q, we get from (6.1)

$$\begin{aligned}(6.2)\qquad &\int_Q \Big(\sum_{i,j=1}^m a_{ij}\frac{\partial z}{\partial x_j}\frac{\partial(z_{n,\varrho}+\beta-z)}{\partial x_i} + \sum_{i=1}^m b_i\frac{\partial z}{\partial x_i}(z_{n,\varrho}+\beta-z) \\ &\qquad + cz(z_{n,\varrho}+\beta-z) + \sum_{i=1}^m d_i z\frac{\partial(z_{n,\varrho}+\beta-z)}{\partial x_i}\Big)\,\mathrm{d}x\,\mathrm{d}t \\ &\quad - \int_Q z\frac{\partial z_{n,\varrho}}{\partial t}\,\mathrm{d}x\,\mathrm{d}t - \int_Q z\frac{\partial\beta}{\partial t}\,\mathrm{d}x\,\mathrm{d}t + \int_{\Gamma_1}\alpha z(z_{n,\varrho}+\beta-z)\,\mathrm{d}\sigma\,\mathrm{d}t \\ &\geqslant \int_Q f^*(z_{n,\varrho}+\beta-z)\,\mathrm{d}x\,\mathrm{d}t + \int_{\Gamma_1} g_1^*(z_{n,\varrho}+\beta-z)\,\mathrm{d}\sigma\,\mathrm{d}t;\end{aligned}$$

now, taking into account the relation

$$\begin{aligned}&\int_Q z(x,t)\varrho\theta_n(t)\theta_n\Big(t+\frac{1}{2\varrho}\Big)z\Big(x,t+\frac{1}{2\varrho}\Big)\,\mathrm{d}x\,\mathrm{d}t \\ &\qquad = \int_\Omega \mathrm{d}x\int_{-\infty}^{+\infty} z(x,t)\varrho\theta_n(t)\theta_n\Big(t+\frac{1}{2\varrho}\Big)z\Big(x,t+\frac{1}{2\varrho}\Big)\,\mathrm{d}t \\ &\qquad = \int_\Omega \mathrm{d}x\int_{-\infty}^{+\infty} z\Big(x,t-\frac{1}{2\varrho}\Big)\varrho\theta_n\Big(t-\frac{1}{2\varrho}\Big)\theta_n(t)z(x,t)\,\mathrm{d}t \\ &\qquad = \int_Q z(x,t)\varrho\theta_n(t)\theta_n\Big(t-\frac{1}{2\varrho}\Big)z\Big(x,t-\frac{1}{2\varrho}\Big)\,\mathrm{d}x\,\mathrm{d}t,\end{aligned}$$

we obtain from (6.2)

$$\begin{aligned}&\int_Q \Big(\sum_{i,j=1}^m a_{ij}\frac{\partial z}{\partial x_j}\frac{\partial(z_{n,\varrho}+\beta-z)}{\partial x_i} + \sum_{i=1}^m b_i\frac{\partial z}{\partial x_i}(z_{n,\varrho}+\beta-z) \\ &\qquad + cz(z_{n,\varrho}+\beta-z) + \sum_{i=1}^m d_i z\frac{\partial(z_{n,\varrho}+\beta-z)}{\partial x_i}\Big)\,\mathrm{d}x\,\mathrm{d}t \\ &\quad - \int_Q \Big(z\varrho\theta_n'(t)\int_{t-\frac{1}{2\varrho}}^{t+\frac{1}{2\varrho}} z(x,y)\theta_n(y)\,\mathrm{d}y\Big)\,\mathrm{d}x\,\mathrm{d}t - \int_Q z\frac{\partial\beta}{\partial t}\,\mathrm{d}x\,\mathrm{d}t \\ &\quad + \int_{\Gamma_1}\alpha z(z_{n,\varrho}+\beta-z)\,\mathrm{d}\sigma\,\mathrm{d}t \\ &\geqslant \int_Q f^*(z_{n,\varrho}+\beta-z)\,\mathrm{d}x\,\mathrm{d}t + \int_{\Gamma_1} g_1^*(z_{n,\varrho}+\beta-z)\,\mathrm{d}\sigma\,\mathrm{d}t,\end{aligned}$$

and therefore, letting ϱ tend to $+\infty$, we find that

$$
\begin{aligned}
(6.3)\qquad &\int_Q \Big(\sum_{i,j=1}^m a_{ij} \frac{\partial z}{\partial x_j} \frac{\partial(\theta_n^2(t)z + \beta - z)}{\partial x_i} + \sum_{i=1}^m b_i \frac{\partial z}{\partial x_i} \big(\theta_n^2(t)z + \beta - z\big) \\
&\qquad + cz\big(\theta_n^2(t)z + \beta - z\big) + \sum_{i=1}^m d_i z \frac{\partial(\theta_n^2(t)z + \beta - z)}{\partial x_i} \Big) \,\mathrm{d}x\,\mathrm{d}t \\
&\quad - \int_Q z^2 \theta_n(t)\theta_n'(t)\,\mathrm{d}x\,\mathrm{d}t - \int_Q z \frac{\partial \beta}{\partial t}\,\mathrm{d}x\,\mathrm{d}t + \int_{\Gamma_1} \alpha z\big(\theta_n^2(t)z + \beta - z\big)\,\mathrm{d}\sigma\,\mathrm{d}t \\
&\geqslant \int_Q f^*\big(\theta_n^2(t)z + \beta - z\big)\,\mathrm{d}x\,\mathrm{d}t + \int_{\Gamma_1} g_1^*\big(\theta_n^2(t)z + \beta - z\big)\,\mathrm{d}\sigma\,\mathrm{d}t.
\end{aligned}
$$

Let us observe that $\theta_n(t)\theta_n'(t) \geqslant 0$ a.e. in $]0,T[$ and $\theta_n(t)\theta_n'(t) > \frac{n}{2}$ if $\tau - \frac{3}{2n} < t < \tau - \frac{1}{n}$. Then, from (6.3) we have

$$
\begin{aligned}
&\int_Q \Big(\sum_{i,j=1}^m a_{ij} \frac{\partial z}{\partial x_j} \frac{\partial(\theta_n^2(t)z + \beta - z)}{\partial x_i} + \sum_{i=1}^m b_i \frac{\partial z}{\partial x_i} \big(\theta_n^2(t)z + \beta - z\big) \\
&\qquad + cz\big(\theta_n^2(t)z + \beta - z\big) + \sum_{i=1}^m d_i z \frac{\partial(\theta_n^2(t)z + \beta - z)}{\partial x_i} \Big) \,\mathrm{d}x\,\mathrm{d}t \\
&\quad - \int_Q z \frac{\partial \beta}{\partial t}\,\mathrm{d}x\,\mathrm{d}t + \int_{\Gamma_1} \alpha z\big(\theta_n^2(t)z + \beta - z\big)\,\mathrm{d}\sigma\,\mathrm{d}t \\
&\geqslant \int_Q z^2 \theta_n(t)\theta_n'(t)\,\mathrm{d}x\,\mathrm{d}t + \int_Q f^*\big(\theta_n^2(t)z + \beta - z\big)\,\mathrm{d}x\,\mathrm{d}t \\
&\quad + \int_{\Gamma_1} g_1^*\big(\theta_n^2(t)z + \beta - z\big)\,\mathrm{d}\sigma\,\mathrm{d}t \\
&\geqslant \frac{n}{2} \int_{\tau - \frac{3}{2n}}^{\tau - \frac{1}{n}} \mathrm{d}t \int_\Omega z^2(x,t)\,\mathrm{d}x + \int_Q f^*\big(\theta_n^2(t)z + \beta - z\big)\,\mathrm{d}x\,\mathrm{d}t \\
&\quad + \int_{\Gamma_1} g_1^*\big(\theta_n^2(t)z + \beta - z\big)\,\mathrm{d}\sigma\,\mathrm{d}t.
\end{aligned}
$$

Finally, as $n \to \infty$ and $\tau \to 0$, we get

$$
\begin{aligned}
(6.4)\qquad &\int_Q \Big(\sum_{i,j=1}^m a_{ij} \frac{\partial z}{\partial x_j} \frac{\partial \beta}{\partial x_i} + \sum_{i=1}^m b_i \frac{\partial z}{\partial x_i}\beta + cz\beta + \sum_{i=1}^m d_i z \frac{\partial \beta}{\partial x_i} - z\frac{\partial \beta}{\partial t} \Big) \,\mathrm{d}x\,\mathrm{d}t \\
&\qquad + \int_{\Gamma_1} \alpha z\beta\,\mathrm{d}\sigma\,\mathrm{d}t \geqslant \int_Q f^*\beta\,\mathrm{d}x\,\mathrm{d}t + \int_{\Gamma_1} g_1^*\beta\,\mathrm{d}\sigma\,\mathrm{d}t
\end{aligned}
$$

for any $0 \leqslant \beta \in C^*(Q)$, $\beta(x,T) = 0$ a.e. in Q.

By virtue of (6.4) and (2.4) we conclude that

$$\tilde{a}(u-z,\varphi) \leqslant \int_Q (f-f^*)\beta \,\mathrm{d}x\,\mathrm{d}t + \int_{\Gamma_1} (g_1 - g_1^*)\beta \,\mathrm{d}\sigma\,\mathrm{d}t \leqslant 0$$

for any $0 \leqslant \beta \in C^*(Q)$, $\beta(x,T) = 0$ a.e. in Q.

Applying the above maximum principle to the L_{Γ_1}-subsolution $(u-z)$ and to the L_{Γ_1}-supersolution w, we obtain

$$\operatorname*{ess\,sup}_Q \frac{u(x,t)-z(x,t)}{w(x,t)} \leqslant \max\Big(0, \sup{}^* \frac{u-z}{w}\Big) = 0.$$

This completes the proof. □

References

[1] *Adams, R. A.*: Sobolev Spaces. Academic Press, New York, 1975.

[2] *Baiocchi, C.*: Soluzioni deboli dei problemi ai limiti per le equazioni paraboliche del tipo del calore. Ist. Lombardo Accad. Sci. Lett. Rend. Sez. A *103* (1969), 704–726.

[3] *Bonafede, S.*: On maximum principle for weak subsolutions of degenerate parabolic linear equations. Comment. Math. Univ. Carolin. *35* (1994), 417–430.

[4] *Bonafede, S.*: A generalized maximum principle for weak subsolutions of degenerate parabolic equations. Rend. Circ. Mat. Palermo (2) *41* (1992), 81–95.

[5] *Drábek, P., Nicolosi, F.*: Existence of bounded solutions for some degenerated quasilinear elliptic equations. Ann. Mat. Pura Appl. (4) *165* (1993), 217–238.

[6] *Eklund, N. A.*: Generalized super-solution of parabolic operators. Trans. Amer. Math. Soc. *220* (1976), 235–242.

[7] *Guglielmino, F., Nicolosi, F.*: W-solutions of boundary value problems for degenerate elliptic operators. Ricerche Mat. Suppl. *36* (1987), 59–72.

[8] *Guglielmino, F., Nicolosi, F.*: Existence theorems for boundary value problems associated with quasilinear elliptic equations. Ricerche Mat. *37* (1988), 157–176.

[9] *Guglielmino, F., Nicolosi, F.*: Existence results for boundary value problems for a class of quasilinear parabolic equations. Actual problems in analysis and mathematical physics, Proceeding of the international symposium in Taormina, Italy, 1992. Univ. di Roma, Roma, 1993, pp. 95–117.

[10] *Ladyzhenskaja, O. A., Solonnikov, V. A., Ural'tseva, N. N.*: Linear and Quasi-Linear Equations of Parabolic Type. Translation of mathematical monographs, vol. 23, A. M. S. Providence, 1968.

[11] *Murthy, M. K. V., Stampacchia, G.*: Boundary value problems for some degenerate-elliptic operators. Ann. Mat. Pura Appl. (4) *80* (1968), 1–122.

[12] *Nicolosi, F.*: Sottosoluzioni deboli delle equazioni paraboliche lineari del secondo ordine superiormente limitate. Matematiche *28* (1973), 361–378.

[13] *Nicolosi, F.*: Soluzioni dei problemi al contorno per operatori parabolici che possono degenerare. Ann. Mat. Pura Appl., IV. Ser. *125* (1980), 135–155.

[14] *Platone Garroni, M. G.*: A generalized maximum principle for boundary value problems for elliptic operators with discontinuos coefficients. Boll. U. M. I. (4) *9* (1974), 1–9.

[15] *Protter, M. H., Weinberger H. F.*: Regular points for elliptic equations with discontinuous coefficients. Ann. Scuola Sup. Pisa *17* (1963), 45–79.

[16] *Stampacchia, G.*: Le probleme de Dirichlet pour les equations elliptiques du second ordre a coefficients discontinus. Annal. Inst. Fourier *15* (1965), 187–257.
[17] *Troianello, G. M.*: On weak subsolutions for parabolic second-order operators. Comm. Partial Differential Equations *3*, 933–948.

Authors' addresses: *Salvatore Bonafede*, Dipartimento di Economia dei Sistemi Agro-Forestali, University of Palermo, Viale delle Scienze, 90128 Palermo, Italy; *Francesco Nicolosi*, Department of Mathematics, University of Catania, Viale A. Doria 6, 95125 Catania, Italy.