

ON THE FROBENIUS NUMBER OF A MODULAR
DIOPHANTINE INEQUALITY

J. C. ROSALES, Granada, P. VASCO, Vila Real

(Received April 25, 2007)

Abstract. We present an algorithm for computing the greatest integer that is not a solution of the modular Diophantine inequality $ax \bmod b \leq x$, with complexity similar to the complexity of the Euclid algorithm for computing the greatest common divisor of two integers.

Keywords: numerical semigroup, Diophantine inequality, Frobenius number, multiplicity

MSC 2000: 11D75, 20M14

1. INTRODUCTION

Given two integers m and n with $n \neq 0$, we denote by $m \bmod n$ the remainder of the division of m by n . Following the terminology used in [6], a proportionally modular Diophantine inequality is an expression of the form $ax \bmod b \leq cx$, where a , b and c are positive integers. The set $S(a, b, c)$ of integer solutions of this inequality is a numerical semigroup, that is, it is a subset of \mathbb{N} (here \mathbb{N} denotes the set of nonnegative integers) that is closed under addition, contains the zero element and its complement in \mathbb{N} is finite. We say that a numerical semigroup is proportionally modular if it is the set of integer solutions of a proportionally modular Diophantine inequality.

The integers a , b and c in the inequality $ax \bmod b \leq cx$ are, respectively, the factor, the modulus and the proportion of the inequality. Following the terminology used in [7], proportionally modular Diophantine inequalities with proportion 1, that

The first author is supported by the projects MTM2004-01446 and MTM2007-62346, and FEDER funds. This paper has been supported by the Luso-Espanhola action HP2004-0056. The authors want to thank P. A. García-Sánchez and the referee for their comments and suggestions.

is, such that $c = 1$, are simply called modular Diophantine inequalities. A numerical semigroup is modular if it is the set of integer solutions of a modular Diophantine inequality.

If S is a numerical semigroup, then the greatest integer that does not belong to S is an important invariant of S , called the Frobenius number of S (see [3]) and denoted here by $g(S)$. Giving a formula for the Frobenius number of $S(a, b, 1)$, as a function of a and b , is still an open problem. Some progress was made in [7] and [4]. In [2] an algorithm to determine the Frobenius number of $S(a, b, c)$ is described. The aim of the present paper is to give an algorithm that computes the Frobenius number of $S(a, b, 1)$, with complexity similar to the complexity of the Euclid algorithm for computing the greatest common divisor of two integers. This algorithm has considerably smaller complexity than the one presented in [2] in most of the cases.

2. PRELIMINARIES

Given a nonempty subset A of \mathbb{Q}_0^+ (here \mathbb{Q}_0^+ is the set of nonnegative rational numbers), we will denote by $\langle A \rangle$ the submonoid of $(\mathbb{Q}_0^+, +)$ generated by A , that is, $\langle A \rangle = \{\lambda_1 a_1 + \dots + \lambda_n a_n; n \in \mathbb{N} \setminus \{0\}, \lambda_1, \dots, \lambda_n \in \mathbb{N} \text{ and } a_1, \dots, a_n \in A\}$. Clearly, $\langle A \rangle \cap \mathbb{N}$ is a submonoid of $(\mathbb{N}, +)$, represented here by $S(A)$. We will refer to $S(A)$ as the submonoid of \mathbb{N} associated to A .

Let p and q be two positive rational numbers with $p < q$. We use the notation

$$[p, q] = \{x \in \mathbb{Q}; p \leq x \leq q\} \quad \text{and} \quad]p, q[= \{x \in \mathbb{Q}; p < x < q\}.$$

The following result is a reformulation of [6, Corollary 9].

Proposition 1.

- (1) *Let a, b and c be positive integers such that $c < a < b$. Then $S([\frac{b}{a}, \frac{b}{a-c}]) = S(a, b, c)$.*
- (2) *Conversely, if a_1, b_1, a_2 and b_2 are positive integers such that $\frac{b_1}{a_1} < \frac{b_2}{a_2}$, then $S([\frac{b_1}{a_1}, \frac{b_2}{a_2}]) = S(a_1 b_2, b_1 b_2, a_1 b_2 - a_2 b_1)$.*

Since the inequality $ax \bmod b \leq cx$ has the same solutions as the inequality $(a \bmod b)x \bmod b \leq cx$, we can assume that $a < b$. Moreover, if $c \geq a$, then $S(a, b, c) = \mathbb{N}$. Therefore, we can suppose that a, b and c are positive integers such that $c < a < b$. Consequently, the condition imposed in (1) of the above proposition is not restrictive.

The next proposition is [8, Proposition 5].

Proposition 2. *If I is an interval of positive rational numbers (not necessarily closed), then $S(I)$ is a proportionally modular numerical semigroup.*

As an immediate consequence of Propositions 1 and 2 we have the following result.

Proposition 3. *Let I be an interval of rational numbers greater than one. Then $S(I)$ is a proportionally modular numerical semigroup. Moreover, every proportionally modular numerical semigroup not equal to \mathbb{N} is of this form.*

The following lemma can be easily deduced from [8, Lemma 2] and will be used several times in this paper.

Lemma 4. *Let I be an interval of positive rational numbers and let x be a positive integer. Then $x \in S(I)$ if and only if there exists a positive integer y such that $x/y \in I$.*

If S is a numerical semigroup, then the smallest positive integer that belongs to S is the multiplicity of S (see [1]) and it is denoted by $m(S)$. If a_1, b_1, a_2 and b_2 are positive integers such that $\frac{b_1}{a_1} < \frac{b_2}{a_2}$, then [9, Algorithm 12] allows us to compute the multiplicity of $S(\left[\frac{b_1}{a_1}, \frac{b_2}{a_2}\right])$. In essence, this algorithm follows the steps of the Euclid algorithm for computing the greatest common divisor of two integers.

Note that, by Proposition 1, we have $S(a, b, 1) = S(\left[\frac{b}{a}, \frac{b}{a-1}\right])$. In Theorem 18 we will see that

$$g\left(S\left(\left[\frac{b}{a}, \frac{b}{a-1}\right]\right)\right) = b - m\left(S\left(\left[\frac{b}{a}, \frac{b}{a-1}\right]\right)\right)$$

and in Theorem 9 that

$$S\left(\left[\frac{b}{a}, \frac{b}{a-1}\right]\right) = S\left(\left[\frac{2b^2+1}{2ab}, \frac{2b^2-1}{2b(a-1)}\right]\right).$$

Therefore

$$g\left(S\left(\left[\frac{b}{a}, \frac{b}{a-1}\right]\right)\right) = b - m\left(S\left(\left[\frac{2b^2+1}{2ab}, \frac{2b^2-1}{2b(a-1)}\right]\right)\right)$$

and $m\left(S\left(\left[\frac{2b^2+1}{2ab}, \frac{2b^2-1}{2b(a-1)}\right]\right)\right)$ can be computed by applying [9, Algorithm 12].

3. A PROPORTIONALLY MODULAR REPRESENTATION FOR AN OPEN MODULAR NUMERICAL SEMIGROUP

If $x_1 < x_2 < \dots < x_k$ are integers, then we use $\{x_1, x_2, \dots, x_k, \rightarrow\}$ to denote the set $\{x_1, x_2, \dots, x_k\} \cup \{z \in \mathbb{Z}; z > x_k\}$. Following the terminology used in [8], a numerical semigroup S is a half-line if $S = \{0\} \cup \{m(S), \rightarrow\}$, and it is open modular if either S is a half-line or there exist integers a and b such that $2 \leq a < b$ and $S = S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$.

If $S = S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$, then by Proposition 2 we know that S is a proportionally modular numerical semigroup and therefore it admits a proportionally modular representation, that is, there exist positive integers x, y and z such that $S = S(x, y, z)$. Observe that, in view of Proposition 1, it suffices to find positive integers a_1, b_1, a_2 and b_2 such that $S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor) = S(\lfloor \frac{b_1}{a_1}, \frac{b_2}{a_2} \rfloor)$. Finding these positive integers is the fundamental aim of this section. To this end, we need some preliminary results and concepts.

If S is a numerical semigroup, then $\mathbb{N} \setminus S$ is finite. The elements of $\mathbb{N} \setminus S$ are the so called gaps of S . The cardinality of $\mathbb{N} \setminus S$ is known as the singularity degree of S (see [1]).

The Frobenius number and the singularity degree of an open modular numerical semigroup can be easily computed by using the following result.

Lemma 5 [8, Theorem 11]. *Let $2 \leq a < b$ be integers, $\alpha = \gcd\{a, b\}$ and $\beta = \gcd\{a - 1, b\}$. Then $S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$ is a proportionally modular numerical semigroup with Frobenius number b and singularity degree $\frac{1}{2}(b - 1 + \alpha + \beta)$.*

The next lemma is straightforward to prove.

Lemma 6. *Let $2 \leq a < b$ be integers. Then $b - 1 \in S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$.*

Proof. A simple check shows that $\frac{b}{a} < \frac{b-1}{a-1} < \frac{b}{a-1}$. By applying Lemma 4 we have that $b - 1 \in S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$. □

It is well-known (see for instance [5]) that every numerical semigroup S is finitely generated and therefore there exists a finite subset A of \mathbb{N} such that $S = \langle A \rangle$. We say that A is a minimal system of generators of S if no proper subset of A generates S . It is also well-known (see [5]) that $S^* \setminus (S^* + S^*)$ is the unique minimal system of generators of S , with $S^* = S \setminus \{0\}$. The cardinality of the minimal system of generators of S is also an important invariant of S called the embedding dimension of S (see [1]).

From Lemmas 5 and 6 we deduce the following result which gives an upper bound to the minimal generators of $S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$.

Lemma 7. *Let $2 \leq a < b$ be integers. Then every minimal generator of $S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$ is smaller than $2b$.*

Proof. From Lemmas 5 and 6 we know that $\{b-1, b+1, \rightarrow\} \subseteq S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$. We conclude the proof by pointing out that every positive integer greater than or equal to $2b$ belongs to $\{b-1, b+1, \rightarrow\} + \{b-1, b+1, \rightarrow\}$. \square

A simple check proves the next result.

Lemma 8. *Let $2 \leq a < b$ be integers. Then $\frac{b}{a} < \frac{2b^2+1}{2ab} < \frac{2b^2-1}{2b(a-1)} < \frac{b}{a-1}$.*

We are now ready to state the principal result of this section.

Theorem 9. *Let $2 \leq a < b$ be integers. Then $S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor) = S(\lfloor \frac{2b^2+1}{2ab}, \frac{2b^2-1}{2b(a-1)} \rfloor)$.*

Proof. From Lemma 8 we have that $\lfloor \frac{2b^2+1}{2ab}, \frac{2b^2-1}{2b(a-1)} \rfloor \subseteq \lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor$ and so $S(\lfloor \frac{2b^2+1}{2ab}, \frac{2b^2-1}{2b(a-1)} \rfloor) \subseteq S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$. To prove the other inclusion we only need to show that every minimal generator of $S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$ belongs to $S(\lfloor \frac{2b^2+1}{2ab}, \frac{2b^2-1}{2b(a-1)} \rfloor)$.

Let x be a minimal generator of $S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$. Then by Lemma 4 there exists a positive integer y such that $\frac{b}{a} < \frac{x}{y} < \frac{b}{a-1}$. Moreover, by applying Lemma 7 we have that $x \leq 2b-1$ and, since $1 < \frac{b}{a} < \frac{x}{y}$, we deduce that $y < 2b-1$. Let us show that $\frac{2b^2+1}{2ab} \leq \frac{x}{y} \leq \frac{2b^2-1}{2b(a-1)}$. As $\frac{b}{a} < \frac{x}{y}$, we have $by < ax$ and so $ax - by \geq 1$. Hence $2abx - 2b^2y \geq 2b$. Since $y < 2b-1$, we infer that $2abx - 2b^2y \geq y$, and consequently $\frac{2b^2+1}{2ab} \leq \frac{x}{y}$. Arguing in a similar way with $\frac{x}{y} < \frac{b}{a-1}$, we get $2b^2y - 2b(a-1)x \geq y$, which is equivalent to $\frac{x}{y} \leq \frac{2b^2-1}{2b(a-1)}$. Finally, by applying Lemma 4 we obtain that $x \in S(\lfloor \frac{2b^2+1}{2ab}, \frac{2b^2-1}{2b(a-1)} \rfloor)$. \square

As an immediate consequence of the previous theorem we have the following result.

Corollary 10. *Let $2 \leq a < b$ be integers and let α and β be rational numbers such that $\frac{b}{a} < \alpha \leq \frac{2b^2+1}{2ab} < \frac{2b^2-1}{2b(a-1)} \leq \beta < \frac{b}{a-1}$. Then $S([\alpha, \beta]) = S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$.*

From this we deduce the following.

Corollary 11. *Let $2 \leq a < b$ be integers. If k is an integer greater than or equal to $2b^2$, then $S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor) = \{x \in \mathbb{N}; (ka-1)x \bmod kb \leq (k-2)x\}$.*

Proof. A simple check shows that

$$\frac{b}{a} < \frac{kb}{ka-1} \leq \frac{2b^2+1}{2ab} < \frac{2b^2-1}{2b(a-1)} \leq \frac{kb}{k(a-1)+1} < \frac{b}{a-1}.$$

By applying Corollary 10, we have that $S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor) = S(\lfloor \frac{kb}{ka-1}, \frac{kb}{k(a-1)+1} \rfloor)$. We conclude the proof by using Proposition 1. \square

The next result is an immediate consequence of Lemma 5 and Corollary 11.

Corollary 12. *Let $2 \leq a < b$ be integers. Set $\alpha = \gcd\{a, b\}$, $\beta = \gcd\{a - 1, b\}$, and let k be an integer greater than or equal to $2b^2$. Then the numerical semigroup $S(ka - 1, kb, k - 2)$ has Frobenius number b and singularity degree $\frac{1}{2}(b - 1 + \alpha + \beta)$.*

4. AN ALGORITHM FOR COMPUTING THE FROBENIUS NUMBER OF A MODULAR NUMERICAL SEMIGROUP

In this section, our first goal will be to prove Theorem 18, which establishes a relationship between the Frobenius number of $S([\frac{b}{a}, \frac{b}{a-1}])$ and the multiplicity of $S([\frac{b}{a}, \frac{b}{a-1}])$, for a and b integers such that $2 \leq a < b$. Before that, we need to recall and establish some results.

The next result is deduced from Proposition 1 and [7, Corollary 6].

Lemma 13. *Let $2 \leq a < b$ be integers. If $x \in \mathbb{N} \setminus S([\frac{b}{a}, \frac{b}{a-1}])$, then $b - x \in S([\frac{b}{a}, \frac{b}{a-1}])$.*

The following lemma follows from [7, Lemma 11] and describes the integers x for which both x and $b - x$ belong to $S([\frac{b}{a}, \frac{b}{a-1}])$.

Lemma 14. *Let $2 \leq a < b$ be integers. Then $\{x, b - x\} \subseteq S([\frac{b}{a}, \frac{b}{a-1}])$ if and only if*

$$x \in \left\{ 0, \frac{b}{\alpha}, 2\frac{b}{\alpha}, \dots, (\alpha - 1)\frac{b}{\alpha}, \frac{b}{\beta}, 2\frac{b}{\beta}, \dots, (\beta - 1)\frac{b}{\beta}, b \right\},$$

where $\alpha = \gcd\{a, b\}$ and $\beta = \gcd\{a - 1, b\}$.

The next result gives an upper bound for the Frobenius number of $S([\frac{b}{a}, \frac{b}{a-c}])$.

Lemma 15. *Let $1 \leq c < a < b$ be integers. Then the Frobenius number of $S([\frac{b}{a}, \frac{b}{a-c}])$ is smaller than $b - 1$.*

Proof. By Proposition 1 we know that $S([\frac{b}{a}, \frac{b}{a-c}]) = \{x \in \mathbb{N}; ax \bmod b \leq cx\}$. Note that, if $x \geq b - 1$, then $ax \bmod b \leq b - 1 \leq c(b - 1) \leq cx$ and therefore $x \in S([\frac{b}{a}, \frac{b}{a-c}])$. □

Next we discard some values for the multiplicity of $S([\frac{b}{a}, \frac{b}{a-1}])$.

Lemma 16. Let $2 \leq a < b$ be integers, $\alpha = \gcd\{a, b\}$, $\beta = \gcd\{a - 1, b\}$ and $S' = S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$. Then $m(S') \notin \{0, \frac{b}{\alpha}, 2\frac{b}{\alpha}, \dots, (\alpha - 1)\frac{b}{\alpha}, \frac{b}{\beta}, 2\frac{b}{\beta}, \dots, (\beta - 1)\frac{b}{\beta}, b\}$.

Proof. By Lemma 4 there exists a positive integer y such that $\frac{b}{a} < \frac{m(S')}{y} < \frac{b}{a-1}$. Let us assume that $m(S') = k\frac{b}{\alpha}$ with $k \in \{1, \dots, \alpha\}$. Then $\frac{b}{a} = \frac{kb/\alpha}{ka/\alpha} = \frac{m(S')}{ka/\alpha} < \frac{m(S')}{y} < \frac{b}{a-1}$. Hence $S(\lfloor \frac{m(S')}{ka/\alpha}, \frac{m(S')}{ka/\alpha-1} \rfloor) \subseteq S'$. In view of Lemma 5 we have that $g(S(\lfloor \frac{m(S')}{ka/\alpha}, \frac{m(S')}{ka/\alpha-1} \rfloor)) = m(S')$ and also $g(S') = b$. So we deduce that $b \leq m(S')$. From Lemma 6 we know that $m(S') \leq b - 1$, which is not possible. Similarly we can prove that $m(S') \neq k\frac{b}{\beta}$ for $k \in \{1, \dots, \beta\}$. \square

Now we study which elements of $S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$ belong to $S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$.

Lemma 17. Let $2 \leq a < b$ be integers, $\alpha = \gcd\{a, b\}$ and $\beta = \gcd\{a - 1, b\}$. If $x \in S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor) \setminus \{0, \frac{b}{\alpha}, 2\frac{b}{\alpha}, \dots, (\alpha - 1)\frac{b}{\alpha}, \frac{b}{\beta}, 2\frac{b}{\beta}, \dots, (\beta - 1)\frac{b}{\beta}, b\}$, then $x \in S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$.

Proof. Since $x \in S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$, by Lemma 4 there exists a positive integer y such that $\frac{b}{a} \leq \frac{x}{y} \leq \frac{b}{a-1}$. If $\frac{x}{y} = \frac{b}{a}$, then $x = k\frac{b}{\alpha}$ for some positive integer k . Suppose that $k \geq \alpha + 1$. Let us prove that $k\frac{b}{\alpha} \in S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$. To this end, in view of Lemma 4, it suffices to see that $\frac{b}{a} < \frac{kb/\alpha}{ka/\alpha-1} < \frac{b}{a-1}$. But a simple check shows that these inequalities hold. The case $\frac{x}{y} = \frac{b}{a-1}$ is analogous to the previous one. \square

We are now ready to state the theorem announced at the beginning of this section.

Theorem 18. Let $2 \leq a < b$ be integers. Define $S = S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$ and $S' = S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$. Then $g(S) = b - m(S')$.

Proof. From Lemmas 14 and 16 we deduce that $b - m(S') \notin S$. By Lemma 17 we obtain that, if $x \in \{1, \dots, m(S') - 1\}$, then either $x \notin S$ or $x \in \{0, \frac{b}{\alpha}, 2\frac{b}{\alpha}, \dots, (\alpha - 1)\frac{b}{\alpha}, \frac{b}{\beta}, 2\frac{b}{\beta}, \dots, (\beta - 1)\frac{b}{\beta}, b\}$. Hence, by Lemmas 13 and 14 we have that $\{b - 1, b - 2, \dots, b - (m(S') - 1)\} \subseteq S$. Moreover, Lemma 15 asserts that $\{b - 1, \rightarrow\} \subseteq S$. Therefore $g(S) = b - m(S')$. \square

Now, we present an algorithm that allows us to compute the Frobenius number of $S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$ for a and b integers such that $2 \leq a < b$. In view of Proposition 1, we have $S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor) = S(a, b, 1)$. Therefore, this algorithm computes the Frobenius number of a modular numerical semigroup.

In [9] we gave an algorithm for computing the multiplicity of a proportionally modular numerical semigroup defined by a closed interval. Thus the idea is to combine this algorithm with Theorems 9 and 18.

Algorithm 19. Input: a and b integers such that $2 \leq a < b$.

Output: The Frobenius number of $S(\lfloor \frac{b}{a}, \frac{b}{a-1} \rfloor)$.

- (1) Compute the multiplicity m of $S(\left[\frac{2b^2+1}{2ab}, \frac{2b^2-1}{2b(a-1)}\right])$ by using [9, Algorithm 12].
- (2) Return $b - m$.

Next we briefly recall [9, Algorithm 12]. In order to do this, we need to introduce some concepts.

Let a_1, b_1, a_2 and b_2 be positive integers. Define

$$R\left(\left[\frac{b_1}{a_1}, \frac{b_2}{a_2}\right]\right) = \left[\frac{a_2}{b_2 \bmod a_2}, \frac{a_1}{b_1 \bmod a_1}\right].$$

Given a closed interval I of positive rational numbers we can construct recursively the following sequence of closed intervals:

$$I_1 = I,$$

$$I_{n+1} = R(I_n), \text{ if } I_n \text{ contains no integers, and } I_{n+1} = I_n, \text{ otherwise.}$$

We will refer to $\{I_n\}_{n \in \mathbb{N} \setminus \{0\}}$ as the sequence of intervals associated with I .

Given a rational number q we denote by $\lfloor q \rfloor$ the integer $\max\{z \in \mathbb{Z}; z \leq q\}$ and by $\lceil q \rceil$ the integer $\min\{z \in \mathbb{Z}; q \leq z\}$. Let I be a closed interval. If I does not contain an integer, then $\lfloor x \rfloor = \lfloor y \rfloor$ for every $x, y \in I$. This integer is denoted by $\lfloor I \rfloor$. We are now ready to recall [9, Algorithm 12].

Algorithm 20. Input: I a closed interval of positive rational numbers such that $S(I) \neq \mathbb{N}$.

Output: The multiplicity of the semigroup $S(I)$.

- (1) Compute the sequence of intervals associated to I until we find the first interval of the sequence that contains an integer. Let us denote such intervals by I_1, I_2, \dots, I_l .
- (2) If $I_l = [\alpha, \beta]$, then $P(I_l) = \lceil \alpha \rceil / 1$.
- (3) Calculate $P(I_1)$ by applying successively $P(I_{n-1}) = P(I_n)^{-1} + \lfloor I_{n-1} \rfloor$.
- (4) The multiplicity of $S(I)$ is the numerator of $P(I_1)$.

We end this section with an example that illustrates Algorithm 19.

Example 21. Let us compute the Frobenius number of the modular numerical semigroup $S(17, 108, 1)$. By Proposition 1, we have $S(17, 108, 1) = S(\left[\frac{108}{17}, \frac{108}{16}\right])$.

- (1) (a)

$$I_1 = \left[\frac{23329}{3672}, \frac{23327}{3456}\right], \quad I_2 = \left[\frac{3456}{2591}, \frac{3672}{1297}\right].$$

Note that $2 \in I_2$.

- (b) $P(I_2) = \frac{2}{1}$.
- (c) $P(I_1) = \frac{1}{2} + 6 = \frac{13}{2}$.
- (d) The multiplicity of $S(\left[\frac{23329}{3672}, \frac{23327}{3456}\right])$ is 13.
- (2) The Frobenius number of $S(\left[\frac{108}{17}, \frac{108}{16}\right])$ is $108 - 13 = 95$.

References

- [1] *V. Barucci, D. E. Dobbs, M. Fontana*: Maximality Properties in Numerical Semigroups and Applications to One-Dimensional Analytically Irreducible Local Domains. *Memoirs of the Amer. Math. Soc.* *598* (1997). zbl
- [2] *M. Delgado, J. C. Rosales*: On the Frobenius number of a proportionally modular Diophantine inequality. *Portugaliae Mathematica* *63* (2006), 415–425. zbl
- [3] *J. L. Ramírez Alfonsín*: *The Diophantine Frobenius Problem*. Oxford Univ. Press, 2005. zbl
- [4] *J. C. Rosales*: Modular Diophantine inequalities and some of their invariants. *Indian J. Pure App. Math.* *36* (2005), 417–429. zbl
- [5] *J. C. Rosales, P. A. García-Sánchez*: *Finitely Generated Commutative Monoids*. Nova Science Publishers, New York, 1999. zbl
- [6] *J. C. Rosales, P. A. García-Sánchez, J. I. García-García, J. M. Urbano-Blanco*: Proportionally modular Diophantine inequalities. *J. Number Theory* *103* (2003), 281–294. zbl
- [7] *J. C. Rosales, P. A. García-Sánchez, J. M. Urbano-Blanco*: Modular Diophantine inequalities and numerical semigroups. *Pacific J. Math.* *218* (2005), 379–398. zbl
- [8] *J. C. Rosales, J. M. Urbano-Blanco*: Opened modular numerical semigroups. *J. Algebra* *306* (2006), 368–377. zbl
- [9] *J. C. Rosales, P. Vasco*: The smallest positive integer that is solution of a proportionally modular Diophantine inequality. *Math. Inequal. Appl.* *11* (2008), 203–212.

Authors' addresses: *J. C. Rosales*, Departamento de Álgebra, Universidad de Granada, E-18071 Granada, Spain, e-mail: jrosales@ugr.es; *P. Vasco*, Departamento de Matemática, Universidade de Trás-os-Montes e Alto Douro, 5001-801 Vila Real, Portugal, e-mail: pvasco@utad.pt.