

CONTINUUM SPECTRUM FOR THE LINEARIZED EXTREMAL EIGENVALUE PROBLEM WITH BOUNDARY REACTIONS

FUTOSHI TAKAHASHI, Osaka

(Received August 6, 2013)

Abstract. We study the semilinear problem with the boundary reaction

$$-\Delta u + u = 0 \quad \text{in } \Omega, \quad \frac{\partial u}{\partial \nu} = \lambda f(u) \quad \text{on } \partial\Omega,$$

where $\Omega \subset \mathbb{R}^N$, $N \geq 2$, is a smooth bounded domain, $f: [0, \infty) \rightarrow (0, \infty)$ is a smooth, strictly positive, convex, increasing function which is superlinear at ∞ , and $\lambda > 0$ is a parameter. It is known that there exists an extremal parameter $\lambda^* > 0$ such that a classical minimal solution exists for $\lambda < \lambda^*$, and there is no solution for $\lambda > \lambda^*$. Moreover, there is a unique weak solution u^* corresponding to the parameter $\lambda = \lambda^*$. In this paper, we continue to study the spectral properties of u^* and show a phenomenon of continuum spectrum for the corresponding linearized eigenvalue problem.

Keywords: continuum spectrum; extremal solution; boundary reaction

MSC 2010: 35J25, 35J20

1. INTRODUCTION

In this paper, we consider the boundary value problem with the boundary reaction

$$(1.1) \quad -\Delta u + u = 0 \quad \text{in } \Omega, \quad \frac{\partial u}{\partial \nu} = \lambda f(u) \quad \text{on } \partial\Omega$$

where $\lambda > 0$ and $\Omega \subset \mathbb{R}^N$, $N \geq 2$ is a smooth bounded domain. Throughout the paper, we assume

$$(1.2) \quad f: [0, \infty) \rightarrow (0, \infty) \text{ is smooth, convex, increasing, } f(0) > 0,$$

Part of this work was supported by JSPS Grant-in-Aid for Challenging Exploratory Research, No. 24654043, and JSPS Grant-in-Aid for Scientific Research (B), No. 23340038.

and superlinear at ∞ in the sense that

$$(1.3) \quad \lim_{t \rightarrow \infty} \frac{f(t)}{t} = \infty.$$

Then the maximum principle implies that solutions are positive on $\overline{\Omega}$.

It is known that there exists an extremal parameter $\lambda^* \in (0, \infty)$ such that

- (i) for every $\lambda \in (0, \lambda^*)$, $(1.1)_\lambda$ has a positive, classical, minimal solution $u_\lambda \in C^2(\overline{\Omega})$ which is strictly stable in the sense that

$$(1.4) \quad \int_{\Omega} (|\nabla \varphi|^2 + \varphi^2) dx > \lambda \int_{\partial\Omega} f'(u_\lambda) \varphi^2 ds_x$$

for every $\varphi \in C^1(\overline{\Omega})$, $\varphi \not\equiv 0$,

- (ii) for $\lambda = \lambda^*$, the pointwise limit

$$(1.5) \quad u^*(x) = \lim_{\lambda \uparrow \lambda^*} u_\lambda(x), \quad x \in \overline{\Omega},$$

becomes a weak solution of $(1.1)_{\lambda^*}$,

- (iii) for $\lambda > \lambda^*$, there exists no solution of $(1.1)_\lambda$, not even in the weak sense.

Here, we call a function $u = (u_1, u_2) \in L^1(\Omega) \times L^1(\partial\Omega)$ a *weak solution* to $(1.1)_\lambda$ if $f(u_2) \in L^1(\partial\Omega)$ and

$$(1.6) \quad \int_{\Omega} (-\Delta \zeta + \zeta) u_1 dx = \int_{\partial\Omega} \left(\lambda f(u_2) \zeta - \frac{\partial \zeta}{\partial \nu} u_2 \right) ds_x$$

holds for any $\zeta \in C^2(\overline{\Omega})$. The statement (ii) says that, under the assumption (1.3), $u^* = (u^*|_{\Omega}, u^*|_{\partial\Omega})$ is a weak solution in the above sense. We proved in [10] Theorem 11 that $u^* \in W^{1,\gamma}(\Omega)$ for any $\gamma \in [1, N/(N-2))$ when $N \geq 3$ (for any $\gamma \in [1, \infty)$ when $N = 2$), so $u^*|_{\partial\Omega} \in W^{1-1/\gamma,\gamma}(\partial\Omega) \subset L^{(N-1)\gamma/(N-\gamma)}(\partial\Omega)$ is the usual trace of the $W^{1,\gamma}$ function u^* on $\partial\Omega$. For the facts (ii), (iii), we refer the reader to [10]. In the following, we call u^* the *extremal solution* of (1.1). In [10], the author obtained several properties such as regularity and uniqueness of the extremal solution u^* . This paper is a sequel to [10]. For related elliptic problems with boundary reaction terms, see, e.g., [4], [6], [9]. For a well-studied problem

$$-\Delta u = \lambda f(u) \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega$$

where f satisfies (1.2), (1.3), see [1], [2], [3], [5], [7], [8], and the references therein.

For $\lambda \in (0, \lambda^*)$, we denote by $\mu_1(\lambda f'(u_\lambda))$ the first eigenvalue of the eigenvalue problem

$$-\Delta \varphi + \varphi = 0 \quad \text{in } \Omega, \quad \frac{\partial \varphi}{\partial \nu} = \lambda f'(u_\lambda) \varphi + \mu \varphi \quad \text{on } \partial\Omega.$$

By the variational characterization, we have

$$\mu_1(\lambda f'(u_\lambda)) = \inf_{\varphi \in C^1(\overline{\Omega}), \varphi \neq 0} \frac{\int_{\Omega} (|\nabla \varphi|^2 + \varphi^2) dx - \int_{\partial\Omega} \lambda f'(u_\lambda) \varphi^2 ds_x}{\int_{\partial\Omega} \varphi^2 ds_x}.$$

Note that $\mu_1(\lambda f'(u_\lambda)) > 0$ since the minimal solution u_λ is strictly stable, and decreases as $\lambda \uparrow \lambda^*$. Denote

$$(1.7) \quad \mu_1^* = \lim_{\lambda \uparrow \lambda^*} \mu_1(\lambda f'(u_\lambda)).$$

If u^* is classical, it must hold that $\mu_1^* = 0$ by considering (iii) above. However, if $u^* = (u^*|_{\Omega}, u^*|_{\partial\Omega}) \notin L^\infty(\Omega) \times L^\infty(\partial\Omega)$, it could happen that μ_1^* is positive. In [10], we proved that even when $\mu_1^* > 0$, there exists a nonnegative weak solution of

$$(1.8) \quad -\Delta \varphi + \varphi = 0 \quad \text{in } \Omega, \quad \frac{\partial \varphi}{\partial \nu} = \lambda^* f'(u^*) \varphi + \mu \varphi \quad \text{on } \partial\Omega$$

for $\mu = 0$. This is a phenomenon of the existence of (L^1) -zero eigenvalue for the eigenvalue problem (1.8). The main purpose of this paper is to prove the following result, which is a generalization of the result by Cabré and Martel [3] to our setting, and may be seen as a phenomenon of the existence of (L^1) -continuum spectrum for the eigenvalue problem (1.8).

Theorem 1.1. *Let μ_1^* be defined by (1.7). Then for any $\mu \in [0, \mu_1^*]$ there exists a weak solution φ to (1.8), $\varphi \in W^{1,q}(\Omega)$ ($1 \leq q < N/(N-1)$), $\varphi \geq 0$, in the sense that $f'(u^*)\varphi|_{\partial\Omega} \in L^1(\partial\Omega)$ and*

$$\int_{\Omega} (-\Delta \zeta + \zeta) \varphi dx = \int_{\partial\Omega} \left\{ (\lambda^* f'(u^*) \varphi|_{\partial\Omega} + \mu \varphi|_{\partial\Omega}) \zeta - \frac{\partial \zeta}{\partial \nu} \varphi|_{\partial\Omega} \right\} ds_x$$

for all $\zeta \in C^2(\overline{\Omega})$. Here $\varphi|_{\partial\Omega}$ is the usual trace of $\varphi \in W^{1,q}(\Omega)$.

2. PROOF OF THEOREM 1.1

In this section we prove Theorem 1.1. We need the uniqueness theorem from [10], which is an analogue of the result by Y. Martel [8].

Theorem 2.1 ([10], Theorem 14). Assume $(1.1)_{\lambda^*}$ has a weak supersolution $w = (w_1, w_2) \in L^1(\Omega) \times L^1(\partial\Omega)$, in the sense that $f(w_2) \in L^1(\partial\Omega)$ and

$$\int_{\Omega} (-\Delta\zeta + \zeta)w_1 \, dx \geq \int_{\partial\Omega} \left\{ \lambda^* f(w_2)\zeta - \frac{\partial\zeta}{\partial\nu} w_2 \right\} ds_x$$

for any $\zeta \in C^2(\overline{\Omega})$, $\zeta \geq 0$ on $\overline{\Omega}$. Then $(w_1, w_2) = (u^*|_{\Omega}, u^*|_{\partial\Omega})$, where u^* is defined by (1.5).

The following is Lemma 17 in [10].

Lemma 2.2. Let $\{u_n\} \subset C^2(\overline{\Omega})$ be a sequence of functions such that

$$-\Delta u_n + u_n = 0 \quad \text{in } \Omega, \quad \frac{\partial u_n}{\partial\nu} \geq 0 \quad \text{on } \partial\Omega.$$

Assume $\|u_n\|_{L^1(\partial\Omega)} \leq C$ for some $C > 0$ independent of n . Then there exists a subsequence (denoted again by u_n) and $u \in W^{1,q}(\Omega)$ such that

$$u_n \rightharpoonup u \quad \text{weakly in } W^{1,q}(\Omega), \quad 1 < q < \frac{N}{N-1}.$$

Moreover, for any $1 \leq p < (N-1)/(N-2)$ there exists a constant $C_p > 0$ depending only on p such that

$$\|u_n\|_{L^p(\partial\Omega)} \leq C_p \|u_n\|_{L^1(\partial\Omega)} \quad \text{for any } n \in \mathbb{N}.$$

Now, we prove Theorem 1.1.

Proof. We follow the argument by X. Cabré and Y. Martel [3].

Step 1. For $n \in \mathbb{N}$, define a sequence of functions f_n as

$$f_n(s) = \begin{cases} f(s) & \text{if } s \leq n, \\ f(n) + f'(n)(s-n) & \text{if } s > n, \end{cases}$$

and consider the approximated problem

$$(2.1) \quad -\Delta u + u = 0 \quad \text{in } \Omega, \quad \frac{\partial u}{\partial\nu} = \lambda f_n(u) \quad \text{on } \partial\Omega.$$

Denote $\lambda_n^* = \sup\{\lambda > 0: (2.1)_{\lambda} \text{ admits a minimal solution} \in C^2(\overline{\Omega})\}$, and let $u_{n,\lambda} \in C^2(\overline{\Omega})$ be the classical minimal solution to $(2.1)_{\lambda}$ for $\lambda < \lambda_n^*$. Since $f_n \leq f_{n+1} \leq f$, we have $u_{n,\lambda} \leq u_{n+1,\lambda} \leq u_{\lambda}$ and $\lambda^* \leq \lambda_{n+1}^* \leq \lambda_n^*$ for any $n \in \mathbb{N}$. Define

$$(2.2) \quad \mu_1(\lambda f'_n(u_{n,\lambda})) = \inf_{\varphi \in C^1(\overline{\Omega}), \varphi \neq 0} \frac{\int_{\Omega} (|\nabla\varphi|^2 + \varphi^2) \, dx - \int_{\partial\Omega} \lambda f'_n(u_{n,\lambda}) \varphi^2 \, ds_x}{\int_{\partial\Omega} \varphi^2 \, ds_x}.$$

Note that $\mu_1(\lambda f'_n(u_{n,\lambda}))$ is continuous with respect to λ by (2.2). Take $0 \leq \mu \leq \mu_1^*$ where μ_1^* is defined by (1.7). Since u_{n,λ_n^*} is classical (which is because f_n is asymptotically linear) and there is no classical solution of (2.1) $_\lambda$ for $\lambda > \lambda_n^*$, the linearized problem around $(\lambda_n^*, u_{n,\lambda_n^*})$ must have zero eigenvalue. Thus

$$\mu_1(\lambda_n^* f'_n(u_{n,\lambda_n^*})) = 0 \leq \mu \leq \mu_1^* \leq \mu_1(\lambda^* f'_n(u_{n,\lambda^*}));$$

here we have used the fact that $f'_n \leq f'$ and $u_{n,\lambda} \leq u_\lambda$, which implies $\mu_1(\lambda f'_n(u_\lambda)) \leq \mu_1(\lambda f'_n(u_{n,\lambda}))$. By the Intermediate Value Theorem, there exists $\lambda_n \in [\lambda^*, \lambda_n^*]$ such that

$$\mu_1(\lambda_n f'_n(u_{n,\lambda_n})) = \mu,$$

which in turn implies there exists $\varphi_n > 0$ with $\int_{\partial\Omega} \varphi_n \, ds_x = 1$ such that

$$(2.3) \quad -\Delta \varphi_n + \varphi_n = 0 \quad \text{in } \Omega, \quad \frac{\partial \varphi_n}{\partial \nu} = \lambda_n f'_n(u_{n,\lambda_n}) \varphi_n + \mu \varphi_n \quad \text{on } \partial\Omega.$$

Recall also that u_{n,λ_n} satisfies

$$(2.4) \quad -\Delta u_{n,\lambda_n} + u_{n,\lambda_n} = 0 \quad \text{in } \Omega, \quad \frac{\partial u_{n,\lambda_n}}{\partial \nu} = \lambda_n f_n(u_{n,\lambda_n}) \quad \text{on } \partial\Omega.$$

We claim there exists $n_0 \in \mathbb{N}$ such that

$$(2.5) \quad \|u_{n,\lambda_n}\|_{L^1(\partial\Omega)} \leq C \quad \text{for any } n \geq n_0.$$

Indeed, let ψ_1 be the first eigenfunction of the Steklov type eigenvalue problem

$$(2.6) \quad -\Delta \psi_1 + \psi_1 = 0 \quad \text{in } \Omega, \quad \frac{\partial \psi_1}{\partial \nu} = \kappa_1 \psi_1 \quad \text{on } \partial\Omega$$

with the first eigenvalue κ_1 , which is normalized as $\int_{\partial\Omega} \psi_1 \, ds_x = 1$. Multiplying (2.4) by ψ_1 and using Jensen's inequality for f_n , we obtain

$$\begin{aligned} \kappa_1 \int_{\partial\Omega} \psi_1 u_{n,\lambda_n} \, ds_x &= \lambda_n \int_{\partial\Omega} f_n(u_{n,\lambda_n}) \psi_1 \, ds_x \\ &\geq \lambda_n f_n \left(\int_{\partial\Omega} \psi_1 u_{n,\lambda_n} \, ds_x \right) \geq \lambda^* f_n \left(\int_{\partial\Omega} \psi_1 u_{n,\lambda_n} \, ds_x \right). \end{aligned}$$

Put $a_n = \int_{\partial\Omega} \psi_1 u_{n,\lambda_n} \, ds_x$. Then we have

$$(2.7) \quad a_n \geq \frac{\lambda^*}{\kappa_1} f_n(a_n).$$

Assume by contradiction that $f_n(a_n) = f'(n)(a_n - n) + f(n)$ for some $n \in \mathbb{N}$ sufficiently large. Then, since $a_n > n$ and $f(n) > (\kappa_1/\lambda^*)n$, $f'(n) > (\kappa_1/\lambda^*)$ for n sufficiently large by (1.2) and (1.3), we have, by (2.7),

$$\begin{aligned} a_n &\geq \frac{\lambda^*}{\kappa_1} f_n(a_n) = \frac{\lambda^*}{\kappa_1} \{f'(n)(a_n - n) + f(n)\} \\ &> a_n - n + n = a_n, \end{aligned}$$

which is a contradiction. Thus we conclude there exists $n_0 \in \mathbb{N}$ such that $f_n(a_n) = f(a_n)$ for any $n \geq n_0$. Again, this and (2.7) imply $a_n \geq (\lambda^*/\kappa_1)f(a_n)$ for any $n \geq n_0$. Now, by the assumption on f , we have $C > 0$ such that $f(s) \geq (2\kappa_1/\lambda^*)s - C$ holds for any $s > 0$. From this and the former estimate, we have $a_n \leq (\lambda^*/\kappa_1)C$ for $n \geq n_0$. This implies the claim (2.5).

Step 2. By (2.5), we have $\|u_{n,\lambda_n}\|_{L^1(\partial\Omega)} \leq C$ for some C independent of n . Also recall that $\|\varphi_n\|_{L^1(\partial\Omega)} = 1$ for a solution φ_n of (2.3). Thus we can apply Lemma 2.2 and the trace Sobolev embedding to obtain $w, \varphi \in L^1(\Omega)$, $\varphi \geq 0$ a.e. satisfying

$$\begin{aligned} (2.8) \quad u_{n,\lambda_n} &\rightharpoonup w, \quad \varphi_n \rightharpoonup \varphi \quad \text{weakly in } W^{1,q}(\Omega), \\ u_{n,\lambda_n} &\rightarrow w, \quad \varphi_n \rightarrow \varphi \quad \text{strongly in } L^p(\partial\Omega) \text{ and a.e. on } \partial\Omega \end{aligned}$$

for any $1 < q < N/(N-1)$ and $1 \leq p < (N-1)/(N-2)$. Since $\int_{\partial\Omega} \varphi \, ds_x = 1$, we see $\varphi \not\equiv 0$ on $\partial\Omega$.

In the following, we prove that $\lambda_n \downarrow \lambda^*$ as $n \rightarrow \infty$ and $w = u^*$. We will show that $w \in W^{1,q}(\Omega)$ is a weak supersolution in the sense of Theorem 2.1. Then the conclusion is obtained by Theorem 2.1. To prove that w is a weak supersolution, put $\bar{\lambda} = \inf_{n \in \mathbb{N}} \lambda_n$. Since $\lambda_n \geq \lambda^*$, we have $\bar{\lambda} \geq \lambda^*$. We observe that

$$\begin{aligned} \int_{\Omega} (-\Delta\zeta + \zeta)u_{n,\lambda_n} \, dx &= \lambda_n \int_{\partial\Omega} f_n(u_{n,\lambda_n})\zeta \, ds_x - \int_{\partial\Omega} \frac{\partial\zeta}{\partial\nu} u_{n,\lambda_n} \, ds_x \\ &\geq \bar{\lambda} \int_{\partial\Omega} f_n(u_{n,\lambda_n})\zeta \, ds_x - \int_{\partial\Omega} \frac{\partial\zeta}{\partial\nu} u_{n,\lambda_n} \, ds_x \end{aligned}$$

holds for all $\zeta \in C^2(\bar{\Omega})$, $\zeta \geq 0$. Using the fact that $u_{n,\lambda_n} \rightarrow w$ in $L^1(\Omega)$ or $L^1(\partial\Omega)$, respectively, and Fatou's lemma, we have

$$\begin{aligned} \int_{\Omega} (-\Delta\zeta + \zeta)w \, dx &\geq \bar{\lambda} \int_{\partial\Omega} f(w)\zeta \, ds_x - \int_{\partial\Omega} \frac{\partial\zeta}{\partial\nu} w \, ds_x \\ &\geq \lambda^* \int_{\partial\Omega} f(w)\zeta \, ds_x - \int_{\partial\Omega} \frac{\partial\zeta}{\partial\nu} w \, ds_x, \quad \forall \zeta \in C^2(\bar{\Omega}), \zeta \geq 0. \end{aligned}$$

This implies also $f(w) \in L^1(\partial\Omega)$ if we take $\zeta \equiv 1$. Thus, we conclude that w is a weak supersolution to (1.1) $_{\lambda^*}$.

Step 3. Let φ_n, φ be as in Step 2. We claim that

$$(2.9) \quad \lambda_n f'_n(u_{n,\lambda_n})\varphi_n \rightarrow \lambda^* f'(u^*)\varphi \quad \text{strongly in } L^1(\partial\Omega)$$

as $n \rightarrow \infty$. For the proof, we invoke Vitali's Convergence Theorem. First, by (2.8), we see

$$\lambda_n f'_n(u_{n,\lambda_n}(x))\varphi_n(x) \rightarrow \lambda^* f'(u^*(x))\varphi(x) \quad \text{a.e. } x \in \partial\Omega$$

for a subsequence. Next, we prove the uniformly absolute continuity property of the sequence $\{\lambda_n f'_n(u_{n,\lambda_n})\varphi_n\}_{n \in \mathbb{N}}$. For that purpose, let $A \subset \partial\Omega$ be measurable and $\varepsilon > 0$ be given arbitrary. Since f_n is convex, we have

$$(2.10) \quad f_n\left(\frac{\chi_A(x)}{\varepsilon}\right) \geq f_n(u_{n,\lambda_n}(x)) + f'_n(u_{n,\lambda_n}(x))\left(\frac{\chi_A(x)}{\varepsilon} - u_{n,\lambda_n}(x)\right)$$

a.e. $x \in \partial\Omega$; here χ_A is the characteristic function of A . By (2.3) and (2.4), we have

$$(2.11) \quad \begin{aligned} \lambda_n \int_{\partial\Omega} f_n(u_{n,\lambda_n})\varphi_n \, ds_x &= \lambda_n \int_{\partial\Omega} f'_n(u_{n,\lambda_n})u_{n,\lambda_n}\varphi_n \, ds_x + \mu \int_{\partial\Omega} u_{n,\lambda_n}\varphi_n \, ds_x \\ &\geq \lambda_n \int_{\partial\Omega} f'_n(u_{n,\lambda_n})u_{n,\lambda_n}\varphi_n \, ds_x. \end{aligned}$$

Also an easy consideration shows that

$$(2.12) \quad \left\{f_n\left(\frac{\chi_A(x)}{\varepsilon}\right) - f(0)\right\}\varphi_n(x) \leq f\left(\frac{1}{\varepsilon}\right)\varphi_n(x)\chi_A(x) \quad \text{a.e. on } \partial\Omega.$$

Thus by (2.10), (2.11) and (2.12), we have

$$(2.13) \quad \begin{aligned} &\int_{\partial\Omega} f'_n(u_{n,\lambda_n})\frac{\chi_A}{\varepsilon}\varphi_n \, ds_x \\ &\leq \int_{\partial\Omega} f_n\left(\frac{\chi_A}{\varepsilon}\right)\varphi_n \, ds_x + \int_{\partial\Omega} f'_n(u_{n,\lambda_n})u_{n,\lambda_n}\varphi_n \, ds_x - \int_{\partial\Omega} f_n(u_{n,\lambda_n})\varphi_n \, ds_x \\ &\leq \int_{\partial\Omega} f_n\left(\frac{\chi_A}{\varepsilon}\right)\varphi_n \, ds_x = \int_{\partial\Omega} \left\{f_n\left(\frac{\chi_A}{\varepsilon}\right) - f(0)\right\}\varphi_n \, ds_x + \int_{\partial\Omega} f(0)\varphi_n \, ds_x \\ &\leq \int_{\partial\Omega} f\left(\frac{1}{\varepsilon}\right)\varphi_n\chi_A \, ds_x + f(0) \leq f\left(\frac{1}{\varepsilon}\right)|A|^{1/p'}\|\varphi_n\|_{L^p(\partial\Omega)} + f(0) \\ &\leq Cf\left(\frac{1}{\varepsilon}\right)|A|^{1/p'} + f(0) \end{aligned}$$

for any $1 \leq p < (N-1)/(N-2)$, where $|A|$ denotes the $(N-1)$ dimensional Hausdorff measure of $A \subset \partial\Omega$ and $p' = p/(p-1)$. In (2.13) we have used $\|\varphi_n\|_{L^p(\partial\Omega)} \leq C$ for some $C > 0$ independent of n by (2.8). Define

$$\delta(\varepsilon) = \left(\frac{f(0)}{f(1/\varepsilon)C}\right)^{p'}.$$

Then for any $\varepsilon > 0$ we obtain $\int_A f'_n(u_{n,\lambda_n})\varphi_n \, ds_x \leq 2f(0)\varepsilon$ if $A \subset \partial\Omega$ satisfies that $|A| < \delta(\varepsilon)$ by (2.13). This implies the uniform absolute continuity of the sequence $\{\lambda_n f'_n(u_{n,\lambda_n})\varphi_n\}_{n \in \mathbb{N}}$. Also for any $\varepsilon > 0$, if we take $E \subset \partial\Omega$ such that $|\partial\Omega \setminus E| < \delta(\varepsilon)$ where $\delta(\varepsilon)$ is as above, we obtain that $\int_{\partial\Omega \setminus E} \lambda_n f'_n(u_{n,\lambda_n})\varphi_n \, ds_x \leq C\varepsilon$. This implies the uniform integrability of $\{\lambda_n f'_n(u_{n,\lambda_n})\varphi_n\}_{n \in \mathbb{N}}$. Therefore, Vitali's Convergence Theorem ensures the claim (2.9).

By (2.9), we pass to the limit $n \rightarrow \infty$ in the weak formulation of (2.3):

$$\int_{\Omega} (-\Delta\zeta + \zeta)\varphi_n \, dx = \int_{\partial\Omega} (\lambda_n f'_n(u_{n,\lambda_n}) + \mu)\varphi_n \zeta - \frac{\partial\zeta}{\partial\nu} \varphi_n \, ds_x, \quad \forall \zeta \in C^2(\overline{\Omega}),$$

and conclude that φ is a weak solution of

$$-\Delta\varphi + \varphi = 0 \quad \text{in } \Omega, \quad \frac{\partial\varphi}{\partial\nu} = \lambda^* f'(u^*)\varphi + \mu\varphi \quad \text{on } \partial\Omega.$$

Recall $\varphi \in W^{1,q}(\Omega)$ for any $1 \leq q < N/(N-1)$. The proof of Theorem 1.1 is completed. \square

References

- [1] *H. Brezis, T. Cazenave, Y. Martel, A. Ramiandrisoa*: Blow up for $u_t - \Delta u = g(u)$ revisited. *Adv. Differ. Equ.* **1** (1996), 73–90. [zbl](#) [MR](#)
- [2] *H. Brezis, J. L. Vázquez*: Blow-up solutions of some nonlinear elliptic problems. *Rev. Mat. Univ. Complutense Madr.* **10** (1997), 443–469. [zbl](#) [MR](#)
- [3] *X. Cabré, Y. Martel*: Weak eigenfunctions for the linearization of extremal elliptic problems. *J. Funct. Anal.* **156** (1998), 30–56. [zbl](#) [MR](#)
- [4] *M. Chipot, I. Shafrir, M. Fila*: On the solutions to some elliptic equations with nonlinear Neumann boundary conditions. *Adv. Differ. Equ.* **1** (1996), 91–110. [zbl](#) [MR](#)
- [5] *J. Dávila*: Singular solutions of semi-linear elliptic problems. *Handbook of Differential Equations: Stationary Partial Differential Equations* (M. Chipot, ed.). Elsevier, Amsterdam, 2008, pp. 83–176. [zbl](#) [MR](#)
- [6] *J. Dávila, L. Dupaigne, M. Montenegro*: The extremal solution of a boundary reaction problem. *Commun. Pure Appl. Anal.* **7** (2008), 795–817. [zbl](#) [MR](#)
- [7] *L. Dupaigne*: *Stable Solutions of Elliptic Partial Differential Equations*. Chapman & Hall Monographs and Surveys in Pure and Applied Mathematics 143, CRC Press, Boca Raton, 2011. [zbl](#) [MR](#)
- [8] *Y. Martel*: Uniqueness of weak extremal solutions of nonlinear elliptic problems. *Houston J. Math.* **23** (1997), 161–168. [zbl](#) [MR](#)
- [9] *P. Quittner, W. Reichel*: Very weak solutions to elliptic equations with nonlinear Neumann boundary conditions. *Calc. Var. Partial Differ. Equ.* **32** (2008), 429–452. [zbl](#) [MR](#)
- [10] *F. Takahashi*: Extremal solutions to Liouville-Gelfand type elliptic problems with nonlinear Neumann boundary conditions. *Commun. Contemp. Math.* (2014), 27 pages, DOI:10.1142/S0219199714500163.

Author's address: Futoshi Takahashi, Department of Mathematics, Osaka City University, Osaka, Japan, e-mail: futoshi@sci.osaka-cu.ac.jp.