

# **Savage-Hutter model of the motion of a gravity driven avalanche flow**

Eduard Feireisl

joint work with P. Gwiazda and A. Świerczewska-Gwiazda

Institute of Mathematics, Academy of Sciences of the Czech Republic, Prague  
MFO Oberwolfach, 9 August - 15 August 2015

The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ ERC Grant Agreement 320078

# Savage-Hutter model for avalanches

## Unknowns

- flow height .....  $h = h(t, x)$   
depth-averaged velocity .....  $\mathbf{u} = \mathbf{u}(t, x)$

$$\partial_t h + \operatorname{div}_x(h\mathbf{u}) = 0$$

$$\partial_t(h\mathbf{u}) + \operatorname{div}_x(h\mathbf{u} \otimes \mathbf{u}) + \nabla_x(ah^2) = h \left( -\gamma \frac{\mathbf{u}}{|\mathbf{u}|} + \mathbf{f} \right)$$

## Periodic boundary conditions

$$\Omega = ([0, 1]|_{\{0,1\}})^2$$

# Transformation - Step I

## Helmholtz decomposition

$$h\mathbf{u} = \mathbf{v} + \mathbf{V} + \nabla_x \Psi$$

where

$$\operatorname{div}_x \mathbf{v} = 0, \quad \int_{\Omega} \Psi \, dx = 0, \quad \int_{\Omega} \mathbf{v} \, dx = 0, \quad \mathbf{V} \in R^2$$

## Fixing $h$ and the potential $\Psi$

$$\partial_t h + \Delta \Psi = 0$$

$$h(0, \cdot) = h_0, \quad -\partial_t h(0, \cdot) = \Delta \Psi_0$$

# Problem I

## Equation

$$\begin{aligned} \partial_t \mathbf{v} + \operatorname{div}_x \left( \frac{(\mathbf{v} + \mathbf{V} + \nabla_x \Psi) \otimes (\mathbf{v} + \mathbf{V} + \nabla_x \Psi)}{h} + (ah^2 + \partial_t \Psi) \mathbb{I} \right) \\ + \partial_t \mathbf{V} \\ = h \left( -\gamma \frac{\mathbf{v} + \mathbf{V} + \nabla_x \Psi}{|\mathbf{v} + \mathbf{V} + \nabla_x \Psi|} + \mathbf{f} \right), \end{aligned}$$

## Constraints and initial conditions

$$\operatorname{div}_x \mathbf{v} = 0, \quad \int_{\Omega} \mathbf{v}(t, \cdot) \, dx = 0$$

$$\mathbf{v}(0, \cdot) = \mathbf{v}_0, \quad \mathbf{V}(0) = \mathbf{V}_0$$

# Transformation - Step II

Prescribing the kinetic energy

$$\frac{1}{2} \frac{|\mathbf{v} + \mathbf{V} + \nabla_x \Psi|^2}{h} = E \equiv \Lambda(t) - ah^2 - \partial_t \Psi$$

Problem II

$$\begin{aligned} & \partial_t \mathbf{v} + \partial_t \mathbf{V} \\ & + \operatorname{div}_x \left( \frac{(\mathbf{v} + \mathbf{V} + \nabla_x \Psi) \otimes (\mathbf{v} + \mathbf{V} + \nabla_x \Psi)}{h} - \frac{1}{2} \frac{|\mathbf{v} + \mathbf{V} + \nabla_x \Psi|^2}{h} \mathbb{I} \right) \\ & = -\gamma \left( \frac{h}{2E} \right)^{1/2} (\mathbf{v} + \mathbf{V} + \nabla_x \Psi) + h \mathbf{f} \end{aligned}$$

# Transformation - Step III

Determining function  $\mathbf{V}$

$$\begin{aligned} & \partial_t \mathbf{v} - \left[ \frac{1}{|\Omega|} \int_{\Omega} \gamma \left( \frac{h}{2E} \right)^{1/2} dx \right] \mathbf{v} \\ &= \frac{1}{|\Omega|} \int_{\Omega} \left[ \gamma \left( \frac{h}{2E} \right)^{1/2} (\mathbf{v} + \nabla_x \Psi) + h \mathbf{f} \right] dx, \quad \mathbf{v}(0) = \mathbf{v}_0 \end{aligned}$$

# Problem III

## Equation

$$\begin{aligned} \partial_t \mathbf{v} + \operatorname{div}_x \left( \frac{(\mathbf{v} + \mathbf{V}[\mathbf{v}] + \nabla_x \psi) \odot (\mathbf{v} + \mathbf{V}[\mathbf{v}] + \nabla_x \psi)}{h} \right) \\ = -\gamma \left( \frac{h}{2E} \right)^{1/2} (\mathbf{v} + \mathbf{V}[\mathbf{v}] + \nabla_x \psi) \\ + \frac{1}{|\Omega|} \int_{\Omega} \gamma \left( \frac{h}{2E} \right)^{1/2} (\mathbf{v} + \mathbf{V}[\mathbf{v}] + \nabla_x \psi) \, dx + h \mathbf{f} - \frac{1}{|\Omega|} \int_{\Omega} h \mathbf{f} \, dx \end{aligned}$$

$$\mathbf{v} \odot \mathbf{w} = \mathbf{v} \otimes \mathbf{w} - \frac{1}{2} \mathbf{v} \cdot \mathbf{w} \mathbb{I}$$

# Transformation - Step IV

## Solving elliptic problem

$$\operatorname{div}_x \mathbb{M} \equiv \operatorname{div}_x (\nabla_x \mathbf{m} + \nabla_x^t \mathbf{m} - \operatorname{div}_x \mathbf{m} \mathbb{I})$$

$$= -\gamma \left( \frac{h}{2E} \right)^{1/2} (\mathbf{v} + \mathbf{V}[\mathbf{v}] + \nabla_x \Psi)$$

$$+ \frac{1}{|\Omega|} \int_{\Omega} \gamma \left( \frac{h}{2E} \right)^{1/2} (\mathbf{v} + \mathbf{V}[\mathbf{v}] + \nabla_x \Psi) \, dx + h \mathbf{f} - \frac{1}{|\Omega|} \int_{\Omega} h \mathbf{f} \, dx,$$

$$\int_{\Omega} \mathbb{M}(t, \cdot) \, dx = 0 \text{ for any } t \in [0, T].$$

# Abstract formulation

Variable coefficients “Euler system”

$$\partial_t \mathbf{v} + \operatorname{div}_x \left( \frac{(\mathbf{v} + \mathbf{H}[\mathbf{v}]) \odot (\mathbf{v} + \mathbf{H}[\mathbf{v}])}{h[\mathbf{v}]} + \mathbb{M}[\mathbf{v}] \right) = 0$$
$$\operatorname{div}_x \mathbf{v} = 0,$$

Kinetic energy

$$\frac{1}{2} \frac{|\mathbf{v} + \mathbf{H}[\mathbf{v}]|^2}{h[\mathbf{v}]} = E[\mathbf{v}]$$

Data

$$\mathbf{v}(0, \cdot) = \mathbf{v}_0, \quad \mathbf{v}(T, \cdot) = \mathbf{v}_T$$

# Abstract operators

## Boundedness

$b$  maps bounded sets in  $L^\infty((0, T) \times \Omega; R^N)$  on bounded sets in  $C_b(Q, R^M)$

## Continuity

$b[\mathbf{v}_n] \rightarrow b[\mathbf{v}]$  in  $C_b(Q; R^M)$  (uniformly for  $(t, x) \in Q$ )

whenever

$\mathbf{v}_n \rightarrow \mathbf{v}$  in  $C_{\text{weak}}([0, T]; L^2(\Omega; R^N))$

## Causality

$\mathbf{v}(t, \cdot) = \mathbf{w}(t, \cdot)$  for  $0 \leq t \leq \tau \leq T$  implies  $b[\mathbf{v}] = b[\mathbf{w}]$  in  $[(0, \tau] \times \Omega]$

# Subsolutions

## Field equations, differential constraints

$$\partial_t \mathbf{v} + \operatorname{div}_x \mathbb{F} = 0, \quad \operatorname{div}_x \mathbf{v} = 0$$

$$\mathbf{v}(0, \cdot) = \mathbf{v}_0, \quad \mathbf{v}(T, \cdot) = \mathbf{v}_T$$

## Non-linear constraint

$$\mathbf{v} \in C(Q; \mathbb{R}^N), \quad \mathbb{F} \in C(Q; \mathbb{R}_{\text{sym},0}^{N \times N}),$$

$$\frac{N}{2} \lambda_{\max} \left[ \frac{(\mathbf{v} + \mathbf{H}[\mathbf{v}]) \otimes (\mathbf{v} + \mathbf{H}[\mathbf{v}])}{h[\mathbf{v}]} - \mathbb{F} + \mathbb{M}[\mathbf{v}] \right] < E[\mathbf{v}]$$

# Subsolution relaxation

## Algebraic inequality

$$\frac{1}{2} \frac{|\mathbf{v} + \mathbf{H}[\mathbf{v}]|^2}{h[\mathbf{v}]} \leq \frac{N}{2} \lambda_{\max} \left[ \frac{(\mathbf{v} + \mathbf{H}[\mathbf{v}]) \otimes (\mathbf{v} + \mathbf{H}[\mathbf{v}])}{h[\mathbf{v}]} - \mathbb{F} + \mathbb{M}[\mathbf{v}] \right] \\ < E[\mathbf{v}]$$

## Solutions

$$\frac{1}{2} \frac{|\mathbf{v} + \mathbf{H}[\mathbf{v}]|^2}{h[\mathbf{v}]} = E[\mathbf{v}]$$

$\Rightarrow$

$$\mathbb{F} = \frac{(\mathbf{v} + \mathbf{H}[\mathbf{v}]) \odot (\mathbf{v} + \mathbf{H}[\mathbf{v}])}{h[\mathbf{v}]} + \mathbb{M}[\mathbf{v}]$$

# Oscillatory lemma

Hypotheses:

$U \subset R \times R^N$ ,  $N = 2, 3$  bounded open set

$\tilde{\mathbf{h}} \in C(U; R^N)$ ,  $\tilde{\mathbb{H}} \in C(U; R_{\text{sym},0}^{N \times N})$ ,  $\tilde{e}$ ,  $\tilde{r} \in C(U)$ ,  $\tilde{r} > 0$ ,  $\tilde{e} \leq \bar{e}$  in  $U$

$$\frac{N}{2} \lambda_{\max} \left[ \frac{\tilde{\mathbf{h}} \otimes \tilde{\mathbf{h}}}{\tilde{r}} - \tilde{\mathbb{H}} \right] < \tilde{e} \text{ in } U.$$

## Conclusion:

$$\mathbf{w}_n \in C_c^\infty(U; \mathbb{R}^N), \quad \mathbb{G}_n \in C_c^\infty(U; \mathbb{R}_{\text{sym},0}^{N \times N}), \quad n = 0, 1, \dots$$

$$\partial_t \mathbf{w}_n + \operatorname{div}_x \mathbb{G}_n = 0, \quad \operatorname{div}_x \mathbf{w}_n = 0 \text{ in } \mathbb{R} \times \mathbb{R}^N,$$

$$\frac{N}{2} \lambda_{\max} \left[ \frac{(\tilde{\mathbf{h}} + \mathbf{w}_n) \otimes (\tilde{\mathbf{h}} + \mathbf{w}_n)}{\tilde{r}} - (\tilde{\mathbb{H}} + \mathbb{G}_n) \right] < \tilde{e} \text{ in } U,$$

$$\mathbf{w}_n \rightarrow 0 \text{ weakly in } L^2(U; \mathbb{R}^N)$$

$$\liminf_{n \rightarrow \infty} \int_U \frac{|\mathbf{w}_n|^2}{\tilde{r}} \, dxdt \geq \Lambda(\bar{e}) \int_U \left( \tilde{e} - \frac{1}{2} \frac{|\tilde{\mathbf{h}}|^2}{\tilde{r}} \right)^2 \, dxdt$$

# Basic ideas of proof

## Localization

Localizing the result of DeLellis and Széhelyhidi to “small” cubes by means of scaling arguments

## Linearization

Replacing all continuous functions by their means on any of the “small” cubes

## Eliminating singular sets

Applying Whitney's decomposition lemma to the non-singular sets (e.g. out of the vacuum  $\{h = 0\}$ )

## Energy and other coefficients depending on solutions

Applying compactness of the abstract operators in  $C$

# Results

## Result (A)

The set of subsolutions is non-empty  $\Rightarrow$  there exists infinitely many weak solutions of the problem with the same initial data

## Initial energy jump

$$\frac{1}{2} \frac{|\mathbf{v}_0 + \mathbf{H}[\mathbf{v}_0]|^2}{h[\mathbf{v}_0]} \quad \boxed{<} \quad \liminf_{t \rightarrow 0} \frac{1}{2} \frac{|\mathbf{v} + \mathbf{H}[\mathbf{v}]|^2}{h[\mathbf{v}]}$$

## Result (B)

The set of subsolutions is non-empty  $\Rightarrow$  there exists a dense set of times such that the values  $\mathbf{v}(t)$  give rise to non-empty subsolution set with

$$\frac{1}{2} \frac{|\mathbf{v}_0 + \mathbf{H}[\mathbf{v}_0]|^2}{h[\mathbf{v}_0]} \quad \boxed{=} \quad \liminf_{t \rightarrow 0} \frac{1}{2} \frac{|\mathbf{v} + \mathbf{H}[\mathbf{v}]|^2}{h[\mathbf{v}]}$$

# Application to Savage-Hutter model

## Theorem

(i) Let the initial data

$$h_0 \in C^2(\Omega), \mathbf{u}_0 \in C^2(\Omega; \mathbb{R}^2), h_0 > 0 \text{ in } \Omega$$

be given, and let  $\mathbf{f}$  and  $a$  be smooth.

Then the Savage-Hutter system admits infinitely many weak solutions in  $(0, T) \times \Omega$ .

(ii) Let  $T > 0$  and

$$h_0 \in C^2(\Omega), h_0 > 0$$

be given.

Then there exists

$$\mathbf{u}_0 \in L^\infty(\Omega; \mathbb{R}^2)$$

such that the Savage-Hutter system admits infinitely many weak solutions in  $(0, T) \times \Omega$  satisfying the energy inequality.



# Example II, Euler-Fourier system

**Mass conservation**

$$\partial_t \varrho + \operatorname{div}_x(\varrho \mathbf{u}) = 0$$

**Momentum balance**

$$\partial_t(\varrho \mathbf{u}) + \operatorname{div}_x(\varrho \mathbf{u} \otimes \mathbf{u}) + \nabla_x(\varrho \vartheta) = 0$$

**Internal energy balance**

$$\frac{3}{2} \left[ \partial_t(\varrho \vartheta) + \operatorname{div}_x(\varrho \vartheta \mathbf{u}) \right] - \Delta \vartheta = -\varrho \vartheta \operatorname{div}_x \mathbf{u}$$

# Example III, Euler-Korteweg-Poisson system

**Mass conservation - equation of continuity**

$$\partial_t \varrho + \operatorname{div}_x(\varrho \mathbf{u}) = 0$$

**Momentum equations - Newton's second law**

$$\begin{aligned} & \partial_t(\varrho \mathbf{u}) + \operatorname{div}_x(\varrho \mathbf{u} \otimes \mathbf{u}) + \nabla_x p(\varrho) \\ &= \boxed{\varrho \nabla_x \left( K(\varrho) \Delta_x \varrho + \frac{1}{2} K'(\varrho) |\nabla_x \varrho|^2 \right) - \varrho \mathbf{u} + \varrho \nabla_x V} \end{aligned}$$

**Poisson equation**

$$\Delta_x V = \varrho - \bar{\varrho}$$

# Example IV, Euler-Cahn-Hilliard system

Model by Lowengrub and Truskinovsky

## Mass conservation

$$\partial_t \varrho + \operatorname{div}_x(\varrho \mathbf{u}) = 0$$

## Momentum balance

$$\partial_t(\varrho \mathbf{u}) + \operatorname{div}_x(\varrho \mathbf{u} \otimes \mathbf{u}) + \nabla_x p_0(\varrho, c) = \operatorname{div}_x \left( \varrho \nabla_x c \otimes \nabla_x c - \frac{\varrho}{2} |\nabla_x c|^2 \mathbb{I} \right)$$

## Cahn-Hilliard equation

$$\partial_t(\varrho c) + \operatorname{div}_x(\varrho c \mathbf{u}) = \Delta \left( \mu_0(\varrho, c) - \frac{1}{\varrho} \operatorname{div}_x(\varrho \nabla_x c) \right)$$