

Endothelial glycocalyx integrity is preserved in young, healthy men during a single bout of strenuous physical exercise

Joanna Majerczak¹, Krzysztof Duda², Stefan Chlopicki^{3,4}, Grzegorz Bartosz⁵, Agnieszka Zakrzewska³, Aneta Balcerczyk⁵, Ryszard T. Smoleński⁶, Jerzy A. Zoladz¹

¹Department of Muscle Physiology, Chair of Physiology and Biochemistry, Faculty of Rehabilitation, University School of Physical Education, Kraków, Poland, ²Department of Oncological Surgery, Center of Oncology, M. Skłodowska-Curie Memorial Institute, Kraków, Poland, ³Jagiellonian Center for Experimental Therapeutics (JCET), Jagiellonian University, Kraków, Poland, ⁴Department of Experimental Pharmacology, Chair of Pharmacology, Jagiellonian University Medical College, Kraków, Poland, ⁵Department of Molecular Biophysics, University of Łódź, Łódź, Poland, ⁶Department of Biochemistry, Medical University of Gdańsk, Gdańsk, Poland

Short title: Endothelial glycocalyx integrity during exercise in humans

Corresponding author: Jerzy A. Zoladz, DSc, PhD

Department of Muscle Physiology, Chair of Physiology and Biochemistry, Faculty of Rehabilitation, University School of Physical Education, Al. Jana Pawła II 78, 31-571 Kraków, Poland

tel/fax.: +48 12 683 1316, E-mail.: jerzy.zoladz@awf.krakow.pl

Summary

In the present study we aimed to evaluate whether oxidative stress and inflammation induced by strenuous exercise affect glycocalyx integrity and endothelial function. Twenty one young, untrained healthy men performed a maximal incremental cycling exercise - until exhaustion. Markers of glycocalyx shedding (syndecan-1, heparan sulfate and hyaluronic acid), endothelial status (nitric oxide and prostacyclin metabolites - nitrate, nitrite, 6-keto-prostaglandin $F_{1\alpha}$), oxidative stress (8-oxo-2'-deoxyguanosine) and antioxidant capacity (uric acid, non-enzymatic antioxidant capacity) as well as markers of inflammation (sVCAM-1 and sICAM-1) were analysed in venous blood samples taken at rest and at the end of exercise. The applied strenuous exercise caused a 5-fold increase in plasma lactate and hypoxanthine concentrations ($p < 0.001$), a fall in plasma uric acid concentration and non-enzymatic antioxidant capacity ($p < 10^{-4}$), accompanied by an increase ($p = 0.003$) in sVCAM-1 concentration. Plasma 6-keto-prostaglandin $F_{1\alpha}$ concentration increased ($p = 0.006$) at exhaustion, while nitrate and nitrite concentrations were not affected. Surprisingly, no significant changes in serum syndecan-1 and heparan sulfate concentrations were observed. We have concluded, that a single bout of severe-intensity exercise is well accommodated by endothelium in young, healthy men as it neither results in evident glycocalyx disruption nor in the impairment of nitric oxide and prostacyclin production.

Key words: Endothelium·Exercise·Glycocalyx·Inflammation·Oxidative stress

Introduction

It is generally accepted that regular physical exercise of moderate intensity enhances health status in humans (Matheson *et al.* 2011, Walker *et al.* 2014). In the recent years special attention has been given to the effect of regular physical activity on the cardiac and vascular protection (Seals *et al.* 2009, Walker *et al.* 2014). An important role in this process seems to play the exercise-induced increase in shear stress *i.e.* the frictional force along endothelial cell membranes exerted by blood flow (Laughlin *et al.* 2008, Seals *et al.* 2009).

Endothelial nitric oxide synthase (eNOS)-derived nitric oxide (NO) is a key factor mediating acute and sustained cardiovascular protective effects of exercise (Laughlin *et al.* 2008). Namely, it has been demonstrated that NO is responsible not only for vasodilation effect, but also inhibits activity of pro-inflammatory factors (cytokines, adhesion molecules) and participates in redox signalling (Hare and Stamler 2005). Other studies indicated that a regular physical activity, leading to an increase in laminar shear stress during exercise, increases the expression of eNOS and its phosphorylation (Hambrecht *et al.* 2003) resulting in an enhancement in systemic NO bioavailability (Green *et al.* 2004). Moreover, nitrate supplementation showed to be effective in enhancement of exercise tolerance in healthy people and in athletes (for review see Jones 2014). The training-induced changes in NO bioavailability common with an increase in antioxidant capacity (Gomez-Cabrera *et al.* 2008), which result in an attenuation of NO degradation by free radicals, leads to an anti-atherogenic phenotype of the endothelium and vascular wall (Laughlin *et al.* 2008). Accordingly, a number of recent studies showed that regular moderate intensity training enhances cardiovascular health status in humans (Walker *et al.* 2014, Seals *et al.* 2009). Moreover, there is growing body of evidence that exercise-based rehabilitation, individually prescribed, provide considerable protection in the primary and secondary prevention of coronary heart disease and improves quality of life of patient with cardiovascular disease (Swift *et al.* 2013).

On the other hand, a single bout of strenuous exercise especially when performed by previously untrained subjects enhances cardiovascular risk, as reported in the cross-sectional and retrospective study (Albert *et al.* 2000). Accordingly, it could well be that despite the fact that regular low and moderate intensity physical activity is indeed associated with reduced long-term morbidity and mortality, an acute challenging exercise can transiently increase the risk of fatal or nonfatal cardiovascular events (Dangardt *et al.* 2013) possibly *via* maladaptive endothelial response. In fact, an impairment of flow-mediated dilation response after heavy exercise intensity (>80% $\text{VO}_{2\text{max}}$) has been clearly demonstrated in healthy humans (Dawson *et al.* 2013). Still relatively little is known concerning the mechanism(s) by which single bout of strenuous exercise develops harmful effects on endothelial cells (Dawson *et al.* 2013).

Even less is known concerning the effect of a single bout of a strenuous exercise on the glycocalyx - a key structure for adequate endothelial function (Florian *et al.* 2003, Reitsma *et al.* 2007). Glycocalyx is a thin carbohydrate-rich gel-like layer (~0.1 μm in capillaries) negatively charged, covering endothelial cells, which constitutes a part of blood-to-tissue permeability barrier and plays various other functions. In fact, the endothelial glycocalyx not only prevents transvascular protein leakage, but also reduces leukocyte adhesion and participates in the mechanotransduction *i.e.* transmits mechanical forces induced by blood flow into biochemical signals (Reitsma *et al.* 2007).

Number of reports suggested that glycocalyx damage might be the earliest stage of the alterations of endothelial function that not only precedes but also promotes the development of endothelial dysfunction (Reitsma *et al.* 2007, Annecke *et al.* 2011). Interestingly, in cultured endothelial cells, degradation of glycocalyx by heparinase, which specifically break down heparan sulfate glycosaminoglycan, impairs NO production by endothelial cells suggesting a close link between glycocalyx integrity and NO bioavailability (Florian *et al.* 2003). Furthermore, the glycocalyx might be damaged by the well-known pathological

condition inducing endothelial dysfunction such as reactive oxygen species (ROS), inflammation, oxidized lipoproteins (Reitsma *et al.* 2007), and some of them are known to constitute the physiological response to exercise.

To our best knowledge, no data has been reported so far concerning the effect of single bout of strenuous exercise undertaken by untrained, healthy people on the endothelial glycocalyx integrity. Therefore, we have hypothesized that a single bout of strenuous exercise, which requires to reach maximal oxygen uptake (VO_{2max}), in young, healthy, but previously untrained men will be accompanied by glycocalyx degradation and this damage will be related to the exercise-induced increase in oxidative stress and inflammation. The question regarding the effect of a single bout of strenuous exercise on glycocalyx seems to be especially relevant, since currently a maximal exercise tests are routinely performed in untrained subjects in order to assess their VO_{2max} (Wasserman *et al.* 2005).

Methods

Subjects

Twenty one young, healthy, non-smoking men (mean \pm SE: age 22.7 ± 0.3 years; body mass 74.9 ± 2.9 kg; height 179 ± 1 cm; BMI 23.3 ± 0.7 kg \cdot m⁻²) with moderate level of physical fitness (VO_{2max} amounting to about 43.9 ± 1.0 mL \cdot min⁻¹ \cdot kg⁻¹) participated in this study. The subjects had no previous experience with maximal incremental exercise tests. Subjects were asked to avoid heavy exercise the day before and on the day of the test as well as to avoid caffeinated and alcoholic beverages 24 hours before exercise. The subjects provided written informed consent to participate in the study. The study was conducted with permission of the Ethical Committee (Commission for Bioethics at the Regional Medical Chamber in Krakow, opinion No 48/KBL/OIL/2009) according to principles established in the Declaration of Helsinki for research on human subjects.

Exercise protocol

An incremental exercise (IT) was performed on the cycloergometer Ergo-Line GmbH & Co KG 800s (Bitz, Germany) following a 6 minute resting period allowed to establish basic cardio-respiratory parameters. The exercise test was performed at pedalling rates of 60 rev · min⁻¹ starting from power output 30 W and followed by gradual increase by 30 W every 3 minutes until exhaustion (Zoladz *et al.* 1995). Based on our previous experience (Zoladz *et al.* 1995, Majerczak *et al.* 2008, Majerczak *et al.* 2014) reaching maximal power output (PO_{max}) during this exercise protocol requires to reach VO_{2max}.

Exercise test was performed about 2h after an intake of light meal (~500 kcal) composed of ~55% of carbohydrates, ~25% of fat and ~20% of protein in order to avoid hypoglycemic events.

Heart rate (HR)

The heart rate in this study was monitored using a heart rate monitor (Polar S810, Polar Electro Oy, Kempele, Finland).

Blood sampling

Blood samples were taken from an antecubital vein at two time points: (a) at rest in the fasting state and (b) in the final stage of the maximal incremental exercise test lasting about 25 minutes, which started at about 2 h after a light meal *i.e.* the second blood sample was taken at about 2.5 hours after a light meal. Blood for parameters analysed in plasma (except lactate, see below) was collected in plain tubes containing EDTA and then centrifuged at 653×g for 15 min in 4°C. Plasma samples were stored at -80°C until analysis. Blood for parameters analysed in serum was collected into plain tubes with clotting activator and left to clot for a minimum of 30 minutes at room temperature and then centrifuged at 1469×g for 10 min in 4°C. Serum was stored at -80°C until analysis.

Plasma lactate concentration

Blood samples were transferred to 1.8 mL Eppendorf tubes containing 1mg ammonium oxalate and 5mg sodium fluoride, mixed for about 20 s and centrifuged at $1055\times g$ for 4 min. The supernatants containing blood plasma (about 0.2 mL) were stored at -40°C until further analysis of plasma lactate concentration $[\text{La}^{-}]_{\text{pl}}$ (Vitros 250 Dry Chemistry System; Kodak, Rochester, NY, USA).

Plasma hypoxanthine and uric acid concentrations

Plasma hypoxanthine $[\text{Hx}]_{\text{pl}}$ and uric acid concentrations $[\text{UA}]_{\text{pl}}$ were evaluated according to previously described reversed-phase high performance liquid chromatography (HPLC) procedure (Smolenski and Yacoub 1993).

Plasma 6-keto-PGF_{1 α} concentration

The concentration of 6-keto-PGF_{1 α} in plasma was assayed using commercially available colorimetric competitive enzyme immunoassay kits (ADI-900-004, Assay Design / Enzo Life Sciences). Multi-Mode Microplate Reader (BioTek, Winooski, VT, USA) was used to measure absorbance at 405 nm.

Plasma nitrite (NO₂⁻) and nitrate (NO₃⁻) concentration

The nitrate and nitrite concentration in plasma was measured by sensitive high-pressure liquid chromatography techniques (ENO-20 NO_x Analyser; EiCom, Kyoto, Japan).

Serum heparan sulfate and syndecan-1 concentrations

Heparan sulfate was measured after pretreatment of serum with actinase E (Sigma-Aldrich, St. Louis, MO, USA). Serum heparan sulfate concentration $[\text{HS}]_{\text{s}}$ was evaluated with special ELISA kit (Seikagaku Corp., Tokyo, Japan). Syndecan-1 concentration $[\text{Syndecan-1}]_{\text{s}}$ was determined in serum using human Syndecan-1 ELISA Kit (Diacclone CD138, No 950.640.096, Diacclone Research, Besancon, France). Both tests were conducted according to manufacturer's instructions. Intensity of the colorful reaction was estimated in optical density

(OD) using the microplate reader BioTek ELx808 (BioTek Instruments, Winooski, VT, USA) at 450 nm.

Serum hyaluronic acid concentration

Hyaluronic acid concentration [HA]_s was determined using Corgenix, Inc. Broomfield, Colorado, USA according to manufacturer's instructions. Multi-Mode Microplate Reader (BioTek, Winooski, VT, USA) was used to measure OD at 450 nm.

Serum VCAM-1 and serum ICAM-1 concentrations

Serum soluble Vascular Cell Adhesion Molecule [sVCAM-1]_s and soluble Intercellular Adhesion Molecule-1 [sICAM-1]_s concentrations were evaluated with special ELISA kit (Human sVCAM-1, #KHT0602/KHT0601, ELISA Kit, Invitrogen and Human sICAM-1, #KHS5411, ELISA Kit, Invitrogen) according to manufacturer's instructions. Intensity of the colorful reaction was estimated by optical density using a spectrophotometer MINI-BOS; Biomedica, Italy (at 450 nm).

Plasma 8-oxo-2'-deoxyguanosine concentration

Plasma 8-oxo-2'-deoxyguanosine concentration [8-oxo-dG]_{pl} was estimated using Cayman Chemical, ELISA Kit (N^o589320) developed for the measurement of DNA/RNA oxidative damage, according to the manufacturer's instruction. The absorbance was measured at a wavelength 405 nm using microplate reader (BioTek®, Winooski, VT, USA). Before the final estimation of the 8-oxo-dG level in the plasma, we have tested the dilution of the plasma samples. Two samples of each group were taken and diluted 25-, 50-, and 100-times, based on the manufacturer's instruction. For the final assay, 50-times dilution was chosen. The 8-oxo-dG level was calculated based on the standard curve, performed every time on the each tested plate, according to the protocol attached to the kit.

Estimation of Non-Enzymatic Antioxidant Capacity

Non-Enzymatic Antioxidant Capacity (NEAC) of blood plasma was estimated by ABTS^{•+} decolorization method (Re *et al.* 1999) modified for a microplate reader.

Data analysis

Estimation of maximal heart rate (HR_{max})

Maximal heart rate (HR_{max}) was predicted for each subject from the formula developed by Tanaka *et al.* (2001).

$$HR_{max} = 208 - 0.7 \cdot \text{age} \quad (r = -0.90) \quad (\text{equation 1})$$

Estimation of maximal oxygen uptake (VO_{2max})

The estimation of the VO_{2max} in this study was performed based on a linear relationship between maximal power output (PO_{max}) reached during maximal incremental exercise test and the measured VO_{2max} , established during the same exercise protocol for the group of young, healthy subjects (n=38) (Majerczak *et al.* 2014). Namely, the VO_{2max} ($\text{mL} \cdot \text{min}^{-1}$) in the present study was estimated based on the PO_{max} (expressed in Watts), reached during the maximal incremental exercise test, using the following formula:

$$VO_{2max} = 1307 + 8.43 \cdot PO_{max} \quad (r = 0.84, p < 10^{-4}) \quad (\text{equation 2})$$

Plasma volume change during exercise (ΔPV)

Plasma volume change during exercise was calculated for each subject based on the hematocrit level using Van Beaumont' equation (Van Beaumont 1972). Then the concentration of blood parameters measured at exhaustion were recalculated using the formula presented by Berthoin *et al.* (2002). In the Table 1 we have presented raw data (value measured at PO_{max}) and also data after recalculation for the changes in plasma volume during incremental exercise (value corrected).

Statistics

The results presented in tables and in figures are expressed as means and standard error (SE). The statistical significance of differences for paired samples was tested using the non-parametric Wilcoxon test and non-asymptotic, exact, two-sided p-values are presented. Significance was set at $p < 0.05$. The statistics was done using the statistical packet STATISTICA 10.0 (StatSoft, Tulsa, OK, USA) and StatXact 9.0 (Cytel Software Corporation, Cambridge, MA, USA).

Results

Power output reached at exhaustion, estimated maximal oxygen uptake and estimated maximal heart rate

The incremental cycling exercise performed by our subjects lasted for 23 ± 0.7 min. The heart rate at the exhaustion (HR_{\max}) has amounted to 188 ± 2 beats per minute and it constituted 98% of age-predicted HR_{\max} (see equation 1) for this group of subjects.

The power output reached at the end of maximal incremental cycling exercise (PO_{\max}) has amounted to 229 ± 7 W. The estimated $VO_{2\max}$ (at PO_{\max} , see equation 2) for the entire group of subjects ($n = 21$) has amounted to 3237 ± 60 mL \cdot min⁻¹ *i.e.* 43.9 ± 1.0 mL \cdot min⁻¹ \cdot kg⁻¹.

Plasma lactate and plasma hypoxanthine concentration

Single bout of strenuous exercise incremental exercise resulted in a decrease in plasma volume (PV) amounting to $14.2 \pm 1.4\%$. The incremental exercise performed in the studied group of subjects was accompanied by a ~5-fold increase ($p < 10^{-4}$) in plasma lactate and in plasma hypoxanthine concentrations (Table 1).

Serum syndecan-1, heparan sulfate and hyaluronic acid concentrations

No significant changes in serum syndecan-1 (Fig.1A, $p = 0.23$) and serum heparan sulfate concentrations after strenuous exercise were observed (Fig. 1B, $p = 0.76$). A significantly

lower serum hyaluronic acid concentration at exhaustion was found when compared to its basal value (by ~34%, Fig. 1C, $p = 0.0003$).

Plasma nitrite, nitrate and 6-keto-Prostaglandin $F_{1\alpha}$ concentrations

No significant changes ($p > 0.05$) in plasma nitrite $[\text{NO}_2^-]_{\text{pl}}$, nitrate $[\text{NO}_3^-]_{\text{pl}}$, and sum of plasma nitrite and nitrate concentrations after exercise were found when compared to their levels at rest (Table 1). In contrast, a significant increase (by ~70%) in plasma 6-keto-PGF- $_{1\alpha}$ concentration after strenuous exercise has been presented ($p = 0.006$, Table 1).

Serum soluble VCAM-1 and serum soluble ICAM-1 concentrations

A significant but small increase in $[\text{sVCAM-1}]_s$ by ~17% has been observed ($p = 0.003$, Fig. 2A) after exercise. No significant changes in $[\text{sICAM-1}]_s$ during exercise was noticed ($p = 0.65$, Fig. 2B).

Plasma 8-oxo-2'-deoxyguanosine concentration, non-enzymatic antioxidant capacity and uric acid concentration

No significant changes in $[\text{8-oxo-dG}]_{\text{pl}}$ ($p = 0.75$) during strenuous incremental cycling exercise was noticed (Table 1), however, a significant decrease (by ~14%, Fig. 3A, $p < 10^{-4}$) in plasma NEAC accompanied by a significant decrease in plasma uric acid concentration (by ~12%, Fig. 3B, $p < 10^{-4}$) after this strenuous exercise was found.

Discussion

The main and original finding of the present study is that a single bout of metabolically highly demanding exercise leading to a mild endothelial inflammatory response (Fig. 2A) and attenuation of antioxidant defense (Fig. 3), appears to have no harmful effect on the endothelial glycocalyx integrity (Fig. 1) and did not impair endothelial function in young, healthy men (Table 1) with normal $\text{VO}_{2\text{max}}$ (amounting to about $43.9 \pm 1.0 \text{ mL} \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$).

Exercise-induced metabolic stress

In the present study we have used a maximal incremental cycling exercise lasting about 25 – 30 min, as an exercise model. Such exercise protocols are frequently used to assess human physical capacity, including $\text{VO}_{2\text{max}}$ considered as the main physiological indicator of aerobic capacity (Åstrand and Rodahl 1986, Wasserman *et al.* 2005). In its final stage this fatiguing exercise requires to reach HR_{max} , $\text{VO}_{2\text{max}}$ (Zoladz *et al.* 1995, Majerczak *et al.* 2008, Majerczak *et al.* 2014) and presumably the maximal cardiac output (Åstrand and Rodahl 1986), therefore it generates the highest challenge to cardiovascular system (Åstrand and Rodahl 1986, Wasserman *et al.* 2005). In the present study, heart rate reached by subjects at the end of incremental exercise has amounted to $188 \pm 2 \text{ beats} \cdot \text{min}^{-1}$ (~98% of age predicted HR_{max}). Moreover, we have found a pronounced increase in plasma lactate concentration from $1.5 \pm 0.2 \text{ mmol} \cdot \text{L}^{-1}$ at rest to $7.0 \pm 0.4 \text{ mmol} \cdot \text{L}^{-1}$ at fatigue, indicating a high contribution of anaerobic glycolysis to ATP resynthesis in the final stage of this exercise protocol. We have also found a 5-fold increase in plasma hypoxanthine concentration at fatigue, considered as an important marker of energetic state of muscle, showing the magnitude of adenine nucleotide degradation during exercise (Sahlin *et al.* 1991). In general, in the present study we have used rather demanding mode of exercise, that requires to reach $\text{VO}_{2\text{max}}$, therefore generating a maximal challenge to the circulatory system in humans.

Strenuous exercise and the markers of glycocalyx damage

It has been reported that an acute high intensity exercise is accompanied by a transient endothelial dysfunction (a decrease in flow-mediated vasodilation response) (Dawson *et al.* 2013), however the knowledge concerning the effect of strenuous exercise on glycocalyx in humans is limited. Nevertheless, it has been reported in an animal model that chronic electrical stimulation of skeletal muscles, which mimics muscle contraction, is accompanied by an disruption of glycocalyx, which as suggested by authors is needed to induce muscle

capillarisation (Brown *et al.* 1996). Moreover, it is known that high concentration of catecholamines in trauma patients, becomes maladaptive and contributes to the glycocalyx damage as well as to endothelial dysfunction (Johansson *et al.* 2011). Therefore, one would expect that a single bout of high intensity exercise, which requires to reach ~100% of VO_{2max} , leading to a marked increase in plasma catecholamine concentration (Galbo 1983) and to an increase in oxidative stress (Powers and Jackson 2008) as well as inflammatory response (Ostrowski *et al.* 1999) might have a harmful impact on the glycocalyx layer in young, untrained men. However, in the present study, we have found no harmful effect of this stressful exercise on the glycocalyx, as judged by no changes in serum syndecan-1 and serum heparan sulfate concentrations (Fig. 1). Surprisingly, in the present study, instead of an expected increase in the plasma hyaluronic acid concentration (a marker of glycocalyx damage), we have observed a significant decrease in hyaluronic acid plasma concentration during this strenuous exercise (Fig. 1C). Although we have no direct explanation for this phenomena there are number of possible reasons for this unexpected response. One might speculate that decrease in hyaluronic acid concentration was due to an exercise-induced oxidative stress (Hrabarova *et al.* 2011) or exercise-induced shear stress enhancing hyaluronic acid incorporation into glycocalyx (Gouverneur *et al.* 2006). Unfortunately, no data concerning hyaluronan incorporation to glycocalyx during exercise has been presented so far. One should also take into account that a decrease in hyaluronic acid concentration during exhaustive exercise might be related to the activation of scavenger activity of liver sinusoidal endothelial cells (LSEC) known to uptake hyaluronic acid (Smedstød 1991). On the other hand, it has been reported that during heavy intensity exercise liver blood flow is diminished by ~50% when compared to its resting value (Bergeron *et al.* 2001) hence the observed decrease in HA concentration at exhaustion cannot be simply explained by a greater clearance of HA by sinusoidal liver endothelial cells and therefore the concept of the exercise-induced

activation of HA uptake by LSEC requires further studies. Another option could be increased exchange of water between intravascular and interstitial compartment during exercise that may help to selectively export hyaluronan into the extravascular space. Nevertheless, our results indicate that a single bout of strenuous exercise seem to be not harmful to the glycocalyx layer in young, healthy men, as judged by no changes in plasma serum syndecan-1 and serum heparan sulfate concentrations.

Exercise-induced endothelial inflammation

In the present study we have found a significant increase ($p = 0.003$) in soluble VCAM-1 concentration (Fig. 2A) during exercise when compared to its level at rest, suggesting an activation of endothelial inflammatory response, which is in agreement with the earlier reports by Ostrowski *et al.* (1999). Inflammatory stimulation during exercise leads to an increase in soluble adhesion molecules (VCAM and ICAM), which regulate attachment and transendothelial migration of leukocytes (Bartzeliotou *et al.* 2007). Our results suggest that a strenuous exercise undertaken occasionally even in case of a young, healthy people with normal VO_{2max} , may enhance endothelial inflammation, which is not accompanied however by a glycocalyx degradation.

Exercise-induced oxidative stress and glycocalyx status

It is well known, that free radicals might induce glycocalyx damage (Marechal *et al.* 2008) and endothelial dysfunction leading to an attenuation of NO bioavailability (Florian *et al.* 2003, Green *et al.* 2004). It is suggested, that high intensity exercise or exercise of long duration is accompanied by an appropriate increase in reactive oxygen species (ROS) and reactive nitrogen species (RNS), overwhelming the antioxidant defence (Lovlin *et al.* 1987). In our study, no significant difference in the plasma 8-oxo-2'-deoxyguanosine measured at exhaustion when compared to its value at rest has been found (Table 1). Despite no changes in plasma 8-oxo-dG, which is a marker of oxidative damage to nucleotides- we have found a

significant exercise-induced decrease in plasma non-enzymatic antioxidant capacity (Fig. 3A), which in this study represents the slow-reacting antioxidants such as tyrosine and tryptophan (Lewinska *et al.* 2007). Moreover, we have observed an exercise-induced attenuation in uric acid concentration (Fig. 3B), a low-molecular mass, fast reacting-antioxidant, which constitutes one half of the antioxidant capacity of serum (Ames *et al.* 1981). Uric acid/urate is a product of purine metabolism and acts as a scavenger of peroxy radicals, hydroxyl radicals and single oxygen. In skeletal muscle during intense exercise urate in reaction with free radicals is converted to allantoin (Hellsten *et al.* 2001, Kandár *et al.* 2014). Additionally, urate can be taken from blood by skeletal muscle, heart and pulmonary bed and it can act there as a scavenger (Becker 1993, Hellsten *et al.* 2001). A decrease in plasma urate concentration, as found in the present study (Fig. 3B), may indicate that uric acid acts indeed as a scavenger during exercise in blood but, this result might suggest that it can also be taken up by skeletal muscle during exercise. The attenuation of antioxidant defence (Fig. 3), suggesting an increase in oxidative stress, was not however accompanied by a glyocalyx damage as reflected by no changes in serum syndecan-1 and heparan sulfate concentrations (Fig. 1).

Nitric oxide and prostacyclin metabolites during exercise

It is well known, that endothelium produces and secretes number of vasoactive substances including vasodilators such as nitric oxide (NO) and prostacyclin (PGI₂) (Clough 2015). In the present study, similarly to an early reports by our group (Zoladz *et al.* 2009, Zoladz *et al.* 2010) we have observed a pronounced exercise-induced increase in plasma 6-keto-PGF_{1 α} (Table 1). Surprisingly, the strenuous exercise, applied in this study, which required to reach maximal oxygen uptake (Zoladz *et al.* 1995, Majerczak *et al.* 2014) and most likely maximal cardiac output (Åstrand and Rodahl 1986) was not accompanied by changes in plasma nitrite (NO₂⁻) and nitrate (NO₃⁻) concentrations (Table 1). Our results are in agreement with previous study showing no effect of moderate as well as high intensity

exercise on the plasma nitrate concentration in humans (St Croix *et al.* 1999). Lack of the increase in NO metabolites in plasma during fatiguing maximal exercise is rather surprising in view of postulated a major role of NO in vasodilatation during exercise in humans (Joyner and Green 2009). However, lack of increase in nitrite/nitrate concentration after high-intensity exercise may be explained by quick NO inactivation in reaction with superoxide and in consequence peroxynitrite formation (Pacher *et al.* 2007) in the oxidative stress condition of high intensity exercise.

Altogether, our results indicate that the endothelial glycocalyx layer in young, healthy men is relatively resistant to the metabolic stress induced by strenuous exercise as judged by an increase in plasma lactate and hypoxanthine concentrations (Table 1) despite the presence of endothelial inflammatory response (Fig. 2A) and an exercise-induced decrease in antioxidant defence (Fig. 3). It should be added that no effect of oxidative stress and inflammation on glycocalyx layer might be due to the fact that in our experiment the exposition to oxidative stress and inflammation was rather short, since this exercise test lasted about 25 minutes and its fatiguing part (above the lactate threshold, *i.e.* in heavy intensity exercise domain) about 10 minutes. Perhaps upon prolonged exposure to oxidative stress and inflammation or in susceptible individuals in whom endothelium represent *locus minoris resistentiae* glycocalyx damage could be observed. Therefore, our results cannot be simply extrapolated to the older subject or patients with various diseases accompanied by endothelial dysfunction such as hypertension, atherosclerosis, diabetes or heart failure. Nevertheless, the present study suggests that a single bout of metabolically highly demanding exercise appears to be well accommodated by endothelium as it neither results in evident glycocalyx disruption nor in the impairment of NO and prostacyclin production in young, healthy men. These results have an important health message, since measurements of VO_{2max} during maximal

exercise tests are routinely performed in humans when determining their exercise physical capacity (Åstrand and Rodahl 1986, Wasserman *et al.* 2005, Majerczak *et al.* 2014).

Conclusions

The present study showed that a single bout of metabolically highly demanding exercise leading to a mild endothelial inflammatory response and attenuation of antioxidant defense, appears to have no harmful effect on the endothelial glycocalyx and endothelial function in young, healthy men with normal physical capacity. This shows that the endothelial glycocalyx in young, healthy men is relatively well protected against challenges induced by a single bout of exercise which requires reaching maximal oxygen uptake.

Conflict of interest

All authors declare no conflict of interest to disclose.

Acknowledgments

We would like to thank the subjects who volunteered for this study.

This work was supported by the European Union from the resources of the European Regional Development Fund under the Innovative Economy Programme (grant coordinated by JCET-UJ, No. POIG.01.01.02-00-069/09).

References

ALBERT CM, MITTLEMAN MA, CHAE CU, LEE IM, HENNEKENS CH, MANSON JE: Triggering of sudden death from cardiac causes by vigorous exertion. *N Engl J Med* **343**: 1355-1361, 2000.

AMES BN, CATHCART R, SCHWIERS E, HOCHSTEIN P: Uric acid provides an antioxidant defense in humans against oxidant- and radical-caused aging and cancer: a hypothesis. *Proc Natl Acad Sci USA* **78**: 6858-6862, 1981.

ANNECKE T, FISCHER J, HARTMANN H, TSCHOEP J, REHM M, CONZEN P, SOMMERHOFF CP, BECKER BF: Shedding of the coronary endothelial glycocalyx: effects of hypoxia/reoxygenation vs ischaemia/reperfusion. *Br J Anaesth* **107**: 679-686, 2011.

ÅSTRAND PO, RODAHL K: Evaluation of physical performance on the basis of tests. In: Textbook of work physiology, Physiological Bases of Exercise. PO ÅSTRAND, K RODAHL (eds), McGraw-Hill Book Company, 1986, pp. 354-390.

BARTZELIOTOU AI, MARGELI AP, TSIRONI M, SKENDERI K, BACOULA C, CHROUSOS GP, PAPASSOTIRIOU I: Circulating levels of adhesion molecules and markers of endothelial activation in acute inflammation induced by prolonged brisk exercise. *Clin Biochem* **40**: 765-770, 2007.

BECKER BF: Towards the physiological function of urate. *Free Radic Biol Med* **14**: 615-631, 1993.

BERGERON R, KJAER M, SIMONSEN L, BÜLOW J, SKOVGAARD D, GALBO H: Splanchnic blood flow and hepatic glucose production in exercising humans: role of renin-angiotensin system. *Am J Physiol Regul Integr Comp Physiol* **281**: R1854-R1861, 2001.

BERTHOIN S, PELAYO P, BAQUET G, MARAIS G, ALLENDER H, ROBIN H: Plasma lactate recovery from maximal exercise with correction for variations in plasma volume. *J Sports Med Phys Fitness* **42**: 26-30, 2002.

BROWN MD, EGGINTON S, HUDLICKA O, ZHOU AL: Appearance of the capillary endothelial glycocalyx in chronically stimulated rat skeletal muscles in relation to angiogenesis. *Exp Physiol* **81**: 1043-1046, 1996.

CLOUGH GF: Developmental Conditioning of the Vasculature. *Compr Physiol* **5**: 397-438, 2015.

DANGARDT FJ, MCKENNA WJ, LÜSCHER TF, DEANFIELD JE: Exercise: friend or foe? *Nat Rev Cardiol* **10**: 495-507, 2013.

DAWSON EA, GREEN DJ, CABLE NT, THIJSSSEN DH: Effects of acute exercise on flow-mediated dilatation in healthy humans. *J Appl Physiol* **115**: 1589-1598, 2013.

FLORIAN JA, KOSKY JR, AINSLIE K, PANG Z, DULL RO, TARBELL JM: Heparan sulfate proteoglycan is a mechanosensor on endothelial cells. *Circ Res* **93**: e136-e142, 2003.

GALBO H: Hormonal and Metabolic Adaptation to Exercise. Thieme-Stratton Inc., New York, 1983.

GOMEZ-CABRERA MC, DOMENECH E, VIÑA J: Moderate exercise is an antioxidant: upregulation of antioxidant genes by training. *Free Radic Biol Med* **44**: 126-131, 2008.

GOUVERNEUR M, SPAAN JA, PANNEKOEK H, FONTIJN RD, VINK H: Fluid shear stress stimulates incorporation of hyaluronan into endothelial cell glycocalyx. *Am J Physiol Heart Circ Physiol* **290**: H458-H462, 2006.

GREEN DJ, MAIORANA A, O'DRISCOLL G, TAYLOR R: Effect of exercise training on endothelium-derived nitric oxide function in humans. *J Physiol* **561**: 1-25, 2004.

HAMBRECHT R, ADAMS V, ERBS S, LINKE A, KRÄNKEL N, SHU Y, BAITHER Y, GIELEN S, THIELE H, GUMMERT JF, MOHR FW, SCHULER G: Regular physical activity improves endothelial function in patients with coronary artery disease by increasing phosphorylation of endothelial nitric oxide synthase. *Circulation* **107**: 3152-3158, 2003.

HARE JM, STAMLER JS: NO/redox disequilibrium in the failing heart and cardiovascular system. *J Clin Invest* **115**: 509-517, 2005.

HELLSTEN Y, SVENSSON M, SJÖDIN B, SMITH S, CHRISTENSEN A, RICHTER EA, BANGSBO J: Allantoin formation and urate and glutathione exchange in human muscle during submaximal exercise. *Free Radic Biol Med* **31**: 1313-1322, 2001.

HRABAROVA E, JURANEK I, SOLTES L: Pro-oxidative effect of peroxynitrite regarding biological systems: a special focus on high-molar-mass hyaluronan degradation. *Gen Physiol Biophys* **30**: 223-238, 2011.

JOHANSSON PI, STENSBALLE J, RASMUSSEN LS, OSTROWSKI SR: A high admission syndecan-1 level, a marker of endothelial glycocalyx degradation, is associated with inflammation, protein C depletion, fibrinolysis, and increased mortality in trauma patients. *Ann Surg* **254**: 194-200, 2011.

JONES AM: Influence of dietary nitrate on the physiological determinants of exercise performance: a critical review. *Appl Physiol Nutr Metab* **39**: 1019-1028, 2014.

JOYNER MJ, GREEN DJ: Exercise protects the cardiovascular system: effects beyond traditional risk factors. *J Physiol* **587**: 5551-5558, 2009.

KANDÁR R, ŠTRAMOVÁ X, DRÁBKOVÁ P, KŘENKOVÁ J: A monitoring of allantoin, uric acid, and malondialdehyde levels in plasma and erythrocytes after ten minutes of running activity. *Physiol Res* **63**: 753-762, 2014.

LAUGHLIN MH, NEWCOMER SC, BENDER SB: Importance of hemodynamic forces as signals for exercise-induced changes in endothelial cell phenotype. *J Appl Physiol* **104**: 588-600, 2008.

LEWINSKA A, WNUK M, SLOTA E, BARTOSZ G: Total anti-oxidant capacity of cell culture media. *Clin Exp Pharmacol Physiol* **34**: 781-786, 2007.

LOVLIN R, COTTLE W, PYKE I, KAVANAGH M, BELCASTRO AN: Are indices of free radical damage related to exercise intensity. *Eur J Appl Physiol Occup Physiol* **56**: 313-316, 1987.

MAJERCZAK J, NIECKARZ Z, KARASINSKI J, ZOLADZ JA: Myosin heavy chain composition in the vastus lateralis muscle in relation to oxygen uptake and heart rate during cycling in humans. *J Physiol Pharmacol* **65**: 217-227, 2014.

MAJERCZAK J, SZKUTNIK Z, DUDA K, KOMOROWSKA M, KOLODZIEJSKI L, KARASINSKI J, ZOLADZ JA: Effect of pedaling rates and myosin heavy chain composition

in the vastus lateralis muscle on the power generating capability during incremental cycling in humans. *Physiol Res* **57**: 873-884, 2008.

MARECHAL X, FAVORY R, JOULIN O, MONTAIGNE D, HASSOUN S, DECOSTER B, ZERIMECH F, NEVIERE R: Endothelial glycocalyx damage during endotoxemia coincides with microcirculatory dysfunction and vascular oxidative stress. *Shock* **29**: 572-576, 2008.

MATHESON GO, KLÜGL M, DVORAK J, ENGBRETSSEN L, MEEUWISSE WH, SCHWELLNUS M, BLAIR SN, VAN MECHELEN W, DERMAN W, BÖRJESSON M, BENDIKSEN F, WEILER R: Responsibility of sport and exercise medicine in preventing and managing chronic disease: applying our knowledge and skill is overdue. *Br J Sports Med* **45**: 1272-1282, 2011.

OSTROWSKI K, ROHDE T, ASP S, SCHJERLING P, PEDERSEN BK: Pro- and anti-inflammatory cytokine balance in strenuous exercise in humans. *J Physiol* **515**: 287-291, 1999.

PACHER P, BECKMAN JS, LIAUDET L: Nitric oxide and peroxynitrite in health and disease. *Physiol Rev* **87**: 315-424, 2007.

POWERS SK, JACKSON MJ: Exercise-induced oxidative stress: cellular mechanisms and impact on muscle force production. *Physiol Rev* **88**: 1243-1276, 2008.

RE R, PELLEGRINI N, PROTEGGENTE A, PANNALA A, YANG M, RICE-EVANS C: Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med* **26**: 1231-1237, 1999.

REITSMA S, SLAAF DW, VINK H, VAN ZANDVOORT MA, OUDE EGBRINK MG: The endothelial glycocalyx: composition, functions, and visualization. *Pflügers Arch* **454**: 345-359, 2007.

SAHLIN K, EKBERG K, CIZINSKY S: Changes in plasma hypoxanthine and free radical markers during exercise in man. *Acta Physiol Scand* **142**: 275-281, 1991.

SEALS DR, WALKER AE, PIERCE GL, LESNIEWSKI LA: Habitual exercise and vascular ageing. *J Physiol* **587**: 5541-5549, 2009.

SMEDSTØD B: Cellular events in the uptake and degradation of hyaluronan. *Adv Drug Deliv Rev* **7**: 265-278, 1991.

SMOLENSKI RT, YACOUB MH: Liquid chromatographic evaluation of purine production in the donor human heart during transplantation. *Biomed Chromatogr* **7**: 189-195, 1993.

ST CROIX CM, WETTER TJ, PEGELOW DF, MEYER KC, DEMPSEY JA: Assessment of nitric oxide formation during exercise. *Am J Respir Crit Care Med* **159**: 1125-1133, 1999.

SWIFT DL, LAVIE CJ, JOHANNSEN NM, ARENA R, EARNEST CP, O'KEEFE JH, MILANI RV, BLAIR SN, CHURCH TS: Physical activity, cardiorespiratory fitness, and exercise training in primary and secondary coronary prevention. *Circ J* **77**: 281-292, 2013.

TANAKA H, MONAHAN K, SEALS D: Aged-predicted maximal heart rate revisited. *JACC* **37**: 153-156, 2001.

VAN BEAUMONT W: Evaluation of hemoconcentration from hematocrit measurements. *J Appl Physiol* **32**: 712-713, 1972.

WALKER AE, KAPLON RE, PIERCE GL, NOWLAN MJ, SEALS DR: Prevention of age-related endothelial dysfunction by habitual aerobic exercise in healthy humans: possible role of nuclear factor κ B. *Clin Sci (Lond)* **127**: 645-654, 2014.

WASSERMAN K, HANSEN JE, SUE DY, STRINGER WW, WHIPP BJ: Principles of Exercise Testing and Interpretation, 4th edn. Lippincott, Williams and Wilkins, Philadelphia, 2005.

ZOLADZ JA, MAJERCZAK J, DUDA K, CHLOPICKI S: Endurance training increases exercise-induced prostacyclin release in young, healthy men-relationship with VO_{2max} . *Pharmacol Rep* **62**: 494-502, 2010.

ZOLADZ JA, MAJERCZAK J, DUDA K, CHLOPICKI S: Exercise-induced prostacyclin release positively correlates with VO_{2max} in young healthy men. *Physiol Res* **58**: 229-238, 2009.

ZOLADZ JA, RADEMAKER AC, SARGEANT AJ: Non-linear relationship between O_2 uptake and power output at high intensities of exercise in humans. *J Physiol* **488**: 211-217, 1995.

Table 1. Selected biochemical blood variables measured at rest in fasted state (basal) and at the end of the maximal incremental test (at PO_{max}).

	Basal Value	Value at PO _{max} measured	Value at PO _{max} corrected
[Syndecan-1] _s (ng · mL ⁻¹)	39.5 ± 3.4	47.2 ± 3.6	41.3 ± 3.7
[HS] _s (ng · mL ⁻¹)	4.91 ± 1.09	5.38 ± 1.27	4.73 ± 1.13
[HA] _s (ng · mL ⁻¹)	40.4 ± 3.5	25.2 ± 2.4**	22.2 ± 2.1***
[6-keto-PGF _{1α}] _{pl} (pg · mL ⁻¹)	179 ± 26	256 ± 33**	223 ± 28**
[NO ₂ ⁻] _{pl} (μmol · L ⁻¹)	0.56 ± 0.09	0.61 ± 0.09	0.53 ± 0.08
[NO ₃ ⁻] _{pl} (μmol · L ⁻¹)	26.5 ± 1.9	31.1 ± 2.2	27.5 ± 2.0
[NO ₂ ⁻] _{pl} + [NO ₃ ⁻] _{pl} (μmol · L ⁻¹)	27.04 ± 1.8	31.7 ± 2.3	28.1 ± 2.0
[sVCAM-1] _s (ng · mL ⁻¹)	702 ± 46	913 ± 70***	802 ± 63**
[sICAM-1] _s (ng · mL ⁻¹)	217 ± 10	243 ± 9***	213 ± 8
[8-oxo-dG] _{pl} (pg · mL ⁻¹)	5427 ± 389	6190 ± 311**	5433 ± 294
NEAC (Trolox Equivalents)	1467 ± 25	1437 ± 18**	1258 ± 18***
[UA] _{pl} (μmol · L ⁻¹)	204 ± 7	201 ± 7	180 ± 7.4***
[Hx] _{pl} (μmol · L ⁻¹)	0.006 ± 0.001	0.032 ± 0.003***	0.028 ± 0.003***
[La ⁻] _{pl} (mmol · L ⁻¹)	1.5 ± 0.2	8.0 ± 0.5***	7.0 ± 0.4***

Values presented as means ± SE. Non asymptotic, exact, 2-sided p-values are presented. *p < 0.05, **p < 0.01, ***p < 0.001. Values at PO_{max} are presented as measured and as corrected for exercise-induced plasma volume change (ΔPV).

[Syndecan-1]_s (serum syndecan-1 concentration), [HS]_s (serum heparan sulfate concentration), [HA]_s (serum hyaluronic acid concentration), [6-keto-PGF_{1α}]_{pl} (plasma 6-keto-prostaglandin F_{1α} concentration), [NO₂⁻]_{pl} (plasma nitrite concentration), [NO₃⁻]_{pl} (plasma nitrate concentration), [NO₂⁻]_{pl} + [NO₃⁻]_{pl} (sum of plasma nitrite and nitrate), [sVCAM-1]_s (serum soluble vascular cell adhesion molecule-1), [sICAM-1]_s (serum soluble intercellular adhesion molecule-1), [8-oxo-dG]_{pl} (plasma 8-oxo-2'-deoxyguanosine concentration), NEAC (non-enzymatic antioxidant capacity), [UA]_{pl} (plasma uric acid concentration), [Hx]_{pl} (plasma hypoxanthine concentration), [La⁻]_{pl} (plasma lactate concentration).




Fig. 1. Markers of glyocalyx shedding during strenuous exercise. Serum syndecan-1 concentration at rest (basal) and at exhaustion (panel A), serum heparan sulfate concentration ([HS]_s) at rest (basal) and at exhaustion (panel B) as well as serum hyaluronic acid concentration ([HA]_s) at rest (basal) and at exhaustion (panel C). Values presented as means ± SE.




Fig. 2. Markers of inflammation during strenuous exercise. Serum soluble vascular adhesion molecule-1 concentration ([sVCAM-1]_s) at rest (basal) and at exhaustion (panel A) and serum soluble intracellular adhesion molecule-1 concentration ([sICAM-1]_s) at rest (basal) and at exhaustion (panel B). Values presented as means ± SE.




Fig. 3. Antioxidant capacity during strenuous exercise. Plasma non-enzymatic antioxidant capacity ($[NEAC]_{pl}$) at rest (basal) and at exhaustion (panel A) and plasma uric acid concentration ($[UA]_{pl}$) at rest (basal) and at exhaustion (panel B). Values presented as means \pm SE.