

Fluids in motion


Eduard Feireisl

Institute of Mathematics of the Academy of Sciences of the Czech Republic, Praha

CMSE 2015 in honour of 70-th anniversary of Z.Dostál, 25–27 May 2016

The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ ERC Grant Agreement 320078

Do we need mathematics?



However beautiful the strategy, you should occasionally look at the results...

Sir Winston Churchill
[1874-1965]

Some naive questions concerning the model

Well posedness

- Does the problem admit a solution for given data?
- In which sense the solution is understood?
- Is solution determined uniquely by the data?
- Are solutions stable under data perturbation?

Typical answers

We don't know yet. But in certain cases we do know...



Euler system (compressible inviscid)

- $\mathbf{u} = \mathbf{u}(t, \mathbf{x})$ fluid velocity
- $\rho = \rho(t, \mathbf{x})$ density



Leonhard Paul Euler
[1707-1783]

Mass conservation

$$\partial_t \rho + \operatorname{div}_x(\rho \mathbf{u}) = 0$$

Balance of momentum

$$\partial_t(\rho \mathbf{u}) + \operatorname{div}_x(\rho \mathbf{u} \otimes \mathbf{u}) + \nabla_x p(\rho) = 0$$


Oscillations vs. nonlinearity

Oscillatory solutions - velocity

$$U(x) \approx \sin(nx), \quad U \rightarrow 0 \text{ in the sense of averages (weakly)}$$

Oscillatory solutions - kinetic energy

$$\frac{1}{2}|U|^2(x) \approx \frac{1}{2}\sin^2(nx) \rightarrow \frac{1}{4} \neq \frac{1}{2}0^2 \text{ in the sense of averages (weakly)}$$



Do some solutions lose/produce energy?



Rudolph Clausius,
[1822–1888]

First and Second law of thermodynamics

Die Energie der Welt ist constant; Die Entropie der Welt strebt einem Maximum zu

Mechanical energy balance for compressible fluid

classical: $\frac{d}{dt} \int \frac{1}{2} \varrho |\mathbf{u}|^2 + P(\varrho) dx = 0, P(\varrho) = \varrho \int_1^\varrho \frac{p(z)}{z^2} dz$

weak: $\frac{d}{dt} \int \frac{1}{2} \varrho |\mathbf{u}|^2 + P(\varrho) dx \boxed{\leq} 0$

Existence

Global-in-time solutions (in general) do not exist. Weak solutions may exist but may not be uniquely determined by the initial data.

Mechanical energy

$$E = \frac{1}{2} \varrho |\mathbf{u}|^2 + P(\varrho)$$

Admissibility criteria - mechanical energy dissipation

$$\partial_t E + \operatorname{div}_x (E \mathbf{u} + p(\varrho) \mathbf{u}) \leq 0$$

Bad or good news for compressible Euler?



Camillo DeLellis [*1976]

Existence

Good news: There exists a global-in-time weak solution of compressible Euler system for “any” initial data

Bad news: There are infinitely many...

Admissible solutions?

Good news: Most of these “wild” solutions produce energy.

Bad news: There is a vast class of data for which there exist infinitely many admissible solutions



László Székelyhidi
[*1977]

Known facts about compressible Euler...

Many admissible solutions [E.Chiodaroli, EF]

For any (or a large class) of initial densities, there exists an initial bounded velocity field such that the compressible Euler system admits infinitely many *physically admissible* solutions

Many admissible solutions [E.Chiodaroli, C.DeLellis, O.Kreml]

There exists Lipschitz (regular) initial data for which the compressible Euler system admits infinitely many admissible solutions

Many admissible solutions [E.Chiodaroli, O.Kreml]

There exist initial data for the Riemann problem (in 2D) and a weak solution of the compressible Euler system that dissipates “globally” more energy than the Riemann solution

Do we compute the right object?

Young measures

$$U(t, x) \approx \nu_{t,x}[U]$$

$\nu(B)$, $B \subset \mathbb{R}^3$ probability that \mathbf{U} belongs to the set B



Laurence Chisholm Young [1905-2000]



Siddhartha Mishra

Numerical results

Certain numerical solutions of “inviscid” problems exhibit scheme independent oscillatory behavior

The way out?

Possible solutions...

- Forget Euler and similar (inviscid) systems
- Take into account only viscosity solutions to Euler but what kind?
- Formulate a local condition of maximality of energy dissipation
- Add more equations...