

CUT-VERTICES AND DOMINATION IN GRAPHS

PREBEN DAHL VESTERGAARD, Aalborg, BOHDAN ZELINKA, Liberec

(Received October 12, 1993)

Summary. The paper studies the domatic numbers and the total domatic numbers of graphs having cut-vertices.

Keywords: domatic number, total domatic number, cut-vertex, bridge

MSC 1991: 05035, 05040

We shall study the domatic number $d(G)$ and the total domatic number $d_t(G)$ of a graph G . A survey of the related theory is given in [3]. We consider finite, undirected graphs without loops or multiple edges.

A subset D of the vertex set $V(G)$ of a graph G is called dominating (total dominating), if for each $x \in V(G) - D$ (for each $x \in V(G)$, respectively) there exists a vertex $y \in D$ adjacent to x . A partition \mathcal{D} of $V(G)$ is called a domatic (total domatic) partition of G , if each class of \mathcal{D} is a dominating (total dominating, respectively) set.

The maximum number of classes of a domatic (total domatic) partition of $V(G)$ was in [1] ([2]) named the domatic (total domatic, respectively) number of G , and it is denoted by $d(G)$ ($d_t(G)$, respectively). Note that $d(G)$ is well-defined for every finite, undirected graph, while $d_t(G)$ is defined only for graphs without isolated vertices.

Consider in G a vertex v of minimum valency $\delta(G)$. Then a dominating set must contain v or a neighbour of v , thus it is obvious that $d(G) \leq \delta(G) + 1$. A total dominating set must contain a neighbour of v , thus $d_t(G) \leq \delta(G)$.

We shall consider the case when a graph G is the union of two graphs G_1, G_2 having exactly one common vertex a ; this vertex a is a cut-vertex of G . The graphs obtained from G_1 and G_2 by deleting a will be denoted respectively by G'_1, G'_2 .

Theorem 1. *With the above notation, for every graph G the domatic numbers satisfy*

$$(1) \quad \min\{d(G_1), d(G_2)\} \leq d(G) \leq 1 + \min\{d(G'_1), d(G'_2)\}.$$

The gaps in the inequalities can be arbitrarily large:

(2) *For any positive integer q there exists a graph G such that*

$$d(G) = \min\{d(G_1), d(G_2)\} + q.$$

(3) *For any positive integer q there exists a graph G such that*

$$d(G) = \min\{d(G'_1), d(G'_2)\} - q.$$

Proof. (1): Let $d_1 = d(G_1)$, $d_2 = d(G_2)$. Let $\{D_1^1, \dots, D_{d_1}^1\}$ be a domatic partition of G_1 with d_1 classes, let $\{D_1^2, \dots, D_{d_2}^2\}$ be a domatic partition of G_2 with d_2 classes. Without loss of generality assume $d_1 \leq d_2$. For $i = 1, \dots, d_1 - 1$ define $D_i = D_i^1 \cup D_i^2$ and let $D_{d_1} = D_{d_1}^1 \cup \bigcup_{i=d_1}^{d_2} D_i^2$. The sets D_1, \dots, D_{d_1} evidently form a domatic partition of G and thus $d(G) \geq d_1 = \min\{d(G_1), d(G_2)\}$.

For the right side inequality in (1) let $d = d(G)$ and consider a domatic partition D_1, \dots, D_d of G ; without loss of generality let $a \in D_d$. For $i = 1, \dots, d$ let $D_i^1 = D_i \cap V(G_1)$, $D_i^2 = D_i \cap V(G_2)$. Consider D_i^1 for $1 \leq i \leq d - 1$. Any vertex $x \in V(G'_1) - D_i^1$ must be adjacent to a vertex of D_i ; as x cannot be adjacent to any vertex of $V(G_2)$, x necessarily is adjacent to a vertex of D_i^1 and thus D_i^1 is a dominating set in G'_1 . Therefore $\{D_1^1, \dots, D_{d-2}^1, D_{d-1}^1 \cup D_d^1\}$ is a domatic partition of G'_1 and $d(G'_1) \geq d(G) - 1$. Analogously $d(G'_2) \geq d(G) - 1$. This proves (1).

Next, we shall construct graphs demonstrating (2) and (3).

(2): Let the vertex set of G_1 be $V(G_1) = \{a, u_1^1, \dots, u_{q+1}^1, v_1^1, \dots, v_{q+1}^1\}$. The set $V(G_1) - \{a\}$ induces the complete subgraph G'_1 with $2q + 2$ vertices. The vertex a is adjacent to the vertices u_1^1, \dots, u_{q+1}^1 . The graph G_2 is isomorphic to G_1 and has the vertex a in common with it. There exists an isomorphism φ of G_1 onto G_2 such that $\varphi(a) = a$. For $i = 1, \dots, q + 1$ denote $u_i^2 = \varphi(u_i^1)$, $v_i^2 = \varphi(v_i^1)$. The vertex a has degree $q + 1$ in G_1 , therefore $d(G_1) \leq q + 2$. Consider the partition of $V(G_1)$ formed by the sets $\{u_i^1\}$ for $i = 1, \dots, q + 1$ and by the set $\{a, v_1^1, \dots, v_{q+1}^1\}$. This is evidently a domatic partition of G_1 with $q + 2$ classes and thus $d(G_1) = q + 2$. As $G_2 \cong G_1$, also $d(G_2) = q + 2 = \min\{d(G_1), d(G_2)\}$. The vertex v_1^1 has degree $2q + 1$ in G and therefore $d(G) \leq 2q + 2$. Consider the partition of $V(G)$ formed by the set $\{a, u_1^1, v_1^2\}$, the sets $\{u_i^1, v_i^2\}$ for $i = 2, \dots, q + 1$ and the sets $\{u_i^2, v_i^1\}$ for $i = 1, \dots, q + 1$. This

is a domatic partition of G with $2q + 2$ classes and thus $d(G) = 2q + 2$. This implies assertion (2).

(3): Let both G'_1, G'_2 be complete graphs with $q + 2$ vertices. Let G_1 be obtained from G'_1 by adding the vertex a and joining it by an edge to exactly one vertex of G'_1 ; analogously let G_2 be constructed. Then $d(G_1) = d(G_2) = \min\{d(G'_1), d(G'_2)\} = q + 2$. For G we have $d(G) \leq 3$, because the vertex a has degree 2. We can easily construct a domatic partition of G with three classes and thus $d(G) = 3$. This implies assertion (3) and Theorem 1 is proven. \square

We shall now express analogous assertions for the total domatic number.

Theorem 2. *With the above notation, for every graph G without isolated vertices the total domatic numbers satisfy*

$$(1) \quad \min\{d_t(G_1), d_t(G_2)\} \leq d_t(G) \leq 1 + \min\{d_t(G'_1), d_t(G'_2)\}.$$

(2) *For any positive integer q there exists a graph G such that*

$$d_t(G) = \min\{d_t(G_1), d_t(G_2)\} + q.$$

(3) *For any positive integer q there exists a graph G such that*

$$d_t(G) = \min\{d_t(G'_1), d_t(G'_2)\} - q.$$

Proof. (1): The proof is analogous to the proof of Theorem 1.

(2): The vertex set of G_1 is

$$V(G_1) = \{a, u_1^1, \dots, u_{q+2}^1, v_1^1, \dots, v_{q+1}^1, w_1^1, \dots, w_{q+1}^1, x_1^1, \dots, x_{q+1}^1\}.$$

The set $V(G_1) - \{a\}$ induces a complete bipartite graph G'_1 on the bipartition classes $\{u_1^1, \dots, u_{q+2}^1, w_1^1, \dots, w_{q+1}^1\}, \{v_1^1, \dots, v_{q+1}^1, x_1^1, \dots, x_{q+1}^1\}$. The vertex a is adjacent to the vertices u_1^1, \dots, u_{q+2}^1 . The graph G_2 is isomorphic to G_1 and has the vertex a in common with it. There exists an isomorphism φ of G_1 onto G_2 such that $\varphi(a) = a$. For $i = 1, \dots, q + 1$ denote $u_i^2 = \varphi(u_i^1), v_i^2 = \varphi(v_i^1), w_i^2 = \varphi(w_i^1), x_i^2 = \varphi(x_i^1)$ and $u_{q+2}^2 = \varphi(u_{q+2}^1)$. The vertex a has degree $q + 2$ in G_1 , therefore $d_t(G_1) \leq q + 2$. Consider the partition of $V(G_1)$ formed by the sets $\{u_i^1, v_i^1\}$ for $i = 1, \dots, q + 1$ and by the set $\{a, u_{q+2}^1, w_1^1, \dots, w_{q+1}^1, x_1^1, \dots, x_{q+1}^1\}$. It is evident that this is a total domatic partition of G_1 with $q + 2$ classes and thus $d_t(G_1) = q + 2$. As $G_2 \cong G_1$, also $d_t(G_2) = q + 2 = \min\{d_t(G_1), d_t(G_2)\}$. The vertex w_1^1 has degree $2q + 2$ in G , therefore $d_t(G) \leq 2q + 2$.

Consider the partition of $V(G)$ formed by the set $\{a, u_1^1, v_1^1, w_1^2, x_1^2, u_{q+2}^1, u_{q+2}^2\}$, the sets $\{u_i^1, v_i^1, w_1^2, x_1^2\}$ for $i = 2, \dots, q + 1$ and the sets $\{u_i^2, v_i^2, w_i^1, x_i^1\}$ for $i = 1, \dots, q + 1$.

This is a total domatic partition of G with $2q + 2$ classes and thus $d_t(G) = 2q + 2$. This implies assertion (2).

(3): The proof is analogous to the proof of Theorem 1(3); the graphs G'_1, G'_2 are complete bipartite graphs in which each bipartition class has $q + 2$ vertices. This proves Theorem 2. \square

Now we shall consider the case when a graph H is obtained from two disjoint graphs H_1, H_2 by joining a vertex a_1 of H_1 with a vertex a_2 of H_2 by a bridge b . By H'_1 we denote the graph obtained from H_1 by deleting a_1 , by H'_2 the graph obtained from H_2 by deleting a_2 .

Theorem 3. *For the domatic numbers of H, H_1, H_2 the following inequalities hold:*

$$\min\{d(H_1), d(H_2)\} \leq d(H) \leq 1 + \min\{d(H_1), d(H_2)\}.$$

Proof. The proof of the first inequality is analogous to the proof of Theorem 1. We shall prove the second inequality. Let $d(H) = d$ and let $\{D_1, \dots, D_d\}$ be a domatic partition of H with d classes. For $i = 1, \dots, d$ let $D_i^1 = D_i \cap V(H_1), D_i^2 = D_i \cap V(H_2)$. Without loss of generality let $a_1 \in D_1$. Consider the case when $a_2 \in D_1$, too. For $1 \leq i \leq d$ each vertex x of H_1 not belonging to D_i^1 is adjacent to some vertex y of D_i . If $x \neq a_1$, then x is adjacent to no vertex of H_2 and $y \in D_i^1$. If $x = a_1$ then $i \neq 1$ and x is adjacent to exactly one vertex a_2 of H_2 and $a_2 \in D_1^2$, i.e. $a_2 \notin D_i^2$; the vertex x must be again adjacent to $y \in D_i^1$. The partition D_1^1, \dots, D_d^1 is a domatic partition of H_1 and $d(H_1) \geq d(H)$. Now let $a_2 \notin D_1$; without loss of generality let $a_2 \in D_d$. Analogously to the preceding case we prove that D_1^1, \dots, D_{d-1}^1 are dominating sets in H_1 ; the set D_d^1 need not be, because a_1 may be adjacent to only one vertex of D_d , namely a_2 , and to no vertex of D_d^1 . The partition $\{D_1^1, \dots, D_{d-2}^1, D_{d-1}^1 \cup D_d^1\}$ is a domatic partition of H_1 and $d(H_1) \geq d(H) - 1$. Analogously $d(H_2) \geq d(H) - 1$ and thus the assertion is proved. \square

Theorem 4. *For the graphs H, H_1, H_2 in the above notation the equality*

$$d(H) = 1 + \min\{d(H_1), d(H_2)\}$$

holds if and only if the following condition is fulfilled: For each $i \in \{1, 2\}$ such that $d(H_i) = \min\{d(H_1), d(H_2)\}$ there exists a partition $\{D_1^i, \dots, D_{d+1}^i\}$ (where $d = d(H_i)$) of the vertex set of H_i such that D_1^i, \dots, D_d^i are dominating sets in H_i and D_{d+1}^i is a dominating set in H'_i but not in H_i .

Proof. Suppose that $d(H) = 1 + \min\{d(H_1), d(H_2)\}$. Let i and d have the described meaning. Consider a domatic partition $\{D_1, \dots, D_{d+1}\}$ of H . For each

$j = 1, \dots, d + 1$ let $D_j^i = D_j \cap V(H_i)$. Without loss of generality let the end vertex of b not belonging to H_i be in D_{d+1} . Let $1 \leq j \leq d$. For each vertex $x \in V(H_i) \setminus D_j^i$ there exists a vertex $y \in D_j$ adjacent to it. A vertex of H_i can be adjacent to no vertex outside of H_i except that end vertex of b which belongs to D_{d+1} and thus not to D_j ; therefore $y \in D_j^i$ and all the sets D_1^i, \dots, D_d^i are dominating in H_i . For each vertex $x \in V(H_i) \setminus D_{d+1}^i$ there also exists a vertex $y \in D_{d+1}$ adjacent to it. No vertex of H_i can be adjacent to a vertex outside of H_i and thus $y \in D_{d+1}^i$; the set D_{d+1}^i is dominating in H_i . It cannot be dominating in H_1 , because then the domatic number of H_i would be $d + 1$.

Now suppose that the condition is fulfilled. Without loss of generality let $d(H_1) = \min\{d(H_1), d(H_2)\}$. Then in H_1 there exists a partition $\{D_1^1, \dots, D_{d+1}^1\}$ with the described property. Choose the subscripts in such a way that $a_1 \in D_1^1$. If $d(H_2) = d(H_1)$, then such a partition $\{D_1^2, \dots, D_{d+1}^2\}$ by assumption exists also in H_2 . If $d(H_2) > d(H_1)$, then there exists a domatic partition $\{D_1^2, \dots, D_{d+1}^2\}$ of H_2 . In both cases choose the subscripts in such a way that $a_2 \in D_1^2$. Now define $D_1 = D_1^1 \cup D_{d+1}^2, D_{d+1} = D_{d+1}^1 \cup D_1^2, D_j = D_j^1 \cup D_j^2$ for $j = 2, \dots, d$. Then the partition $\{D_1, \dots, D_{d+1}\}$ is a domatic partition of H and $d(H) = d + 1 = 1 + \min\{d(H_1), d(H_2)\}$. \square

Theorem 5. *Let for the graphs H, H_1, H_2 in the above notation the equality $d(H) = 1 + d(H_1)$ hold. Then there exists a vertex of H_1 non-adjacent to a_1 with the property that by joining it by an edge to a_1 a graph \hat{H}_1 with domatic number $d(\hat{H}_1) = d(H_1) + 1$ is obtained from H_1 .*

Proof. Consider the partition $\{D_1^1, \dots, D_{d+1}^1\}$ introduced above. Let $u \in D_{d+1}^1$. As D_{d+1}^1 is a dominating set in H_1' but not in H_1 , the vertex a_1 is not adjacent to u . If we join a_1 and u by an edge, then a_1 is adjacent to a vertex of D_{d+1}^1 and D_{d+1}^1 is dominating in the resulting graph H_1 . Then $\{D_1^1, \dots, D_{d+1}^1\}$ is a domatic partition in \hat{H}_1 and $d(\hat{H}_1) = d(H_1) + 1$. (As we have added only one edge, it cannot be greater.) \square

Note that the inverse assertion is not true. An example is a circuit C_4 of length 4. Its domatic number is 2, after adding one chord it is 3, but no graph having a circuit C_4 as a terminal block has domatic number greater than 2.

Theorem 6. *For the total domatic numbers of H, H_1, H_2 the following inequalities hold:*

$$\min\{d_t(H_1), d_t(H_2)\} \leq d_t(H) \leq 1 + \min\{d_t(H_1), d_t(H_2)\}.$$

The proof is analogous to the proof of Theorem 3. \square

Before stating the next theorem, we shall express a slight modification of the definition of a total dominating set.

Let G be a graph, and let G_0 be a subgraph of G . We say that a subset D of $V(G)$ is total dominating for G_0 , if for each vertex $x \in V(G_0)$ there exists a vertex $y \in D$ adjacent to x .

Note that in this definition we do not suppose that $D \subseteq V(G_0)$ but only $D \subseteq V(G)$.

Theorem 7. *If for the graphs H, H_1, H_2 in the above notation the equality*

$$d_t(H) = 1 + \min\{d_t(H_1), d_t(H_2)\}$$

holds, then for each $i \in \{1, 2\}$ such that $d_t(H_i) = \min\{d_t(H_1), d_t(H_2)\}$ there exists a partition $\{D_1^i, \dots, D_{d+1}^i\}$ (where $d = d_t(H_i)$) of the vertex set of H_i such that D_1^i, \dots, D_d^i are total dominating sets in H_i and D_{d+1}^i is a total dominating set for H_i' but not for H_i .

The proof is analogous to the first part of the proof of Theorem 4.

Note that Theorem 7 differs from Theorem 4 by the fact that it is only an implication, not an equivalence. Before investigating the inverse assertion, we introduce some notation.

If a graph H_i with a vertex a_i has the property that $d_t(H_i) = d$ and there exists a partition as described in Theorem 7, we say that the pair (H_i, a_i) is in the class $\kappa(d)$. If $(H_i, a_i) \in \kappa(d)$ and the described partition has the property that $a_i \in D_{d+1}^i$ (or $a_i \notin D_{d+1}^i$), we write $(H_i, a_i) \in \kappa_1(d)$ (or $(H_i, a_i) \in \kappa_0(d)$, respectively). Obviously $\kappa_0(d) \cup \kappa_1(d) = \kappa(d)$, note that $\kappa_0(d) \cap \kappa_1(d) \neq \emptyset$ may occur.

Theorem 8. *Let H, H_1, H_2 be graphs in the above notation. The equality*

$$d_t(H) = 1 + \min\{d_t(H_1), d_t(H_2)\}$$

holds if and only if at least one of the following three cases occurs:

- (i) *exactly one of the pairs $(H_1, a_1), (H_2, a_2)$ is in $\kappa(d)$ and the graph from the other pair has total domatic number greater than d ;*
- (ii) *both the pairs $(H_1, a_1), (H_2, a_2)$ are in $\kappa_0(d)$;*
- (iii) *both the pairs $(H_1, a_1), (H_2, a_2)$ are in $\kappa_1(d)$.*

Proof. Suppose that the above mentioned equality holds, say $d_t(H_1) \leq d_t(H_2)$ and $d_t(H) = 1 + d_t(H_1)$. Then by Theorem 7 $(H_1, a_1) \in \kappa(d)$. With the same notation as in Theorem 7 we let $\mathcal{D} = \{D_1, \dots, D_{d+1}\}$ be a total domatic partition of H and let $D_j^1 = D_j \cap V(H_1)$, $D_j^2 = D_j \cap V(H_2)$ for $j = 1, \dots, d+1$. The notation is chosen such that D_{d+1}^1 is a total dominating set for H_1' but not for H_1 . Then a_1 is

adjacent to no vertex of D_{d+1}^1 and necessarily $a_2 \in D_{d+1}$. Hence if $a_1 \in D_{d+1}^1$, then a_1, a_2 belong to the same class of \mathcal{D} ; otherwise they belong to different classes.

If also $(H_2, a_2) \in \kappa(d)$, then one of the classes D_1^2, \dots, D_{d+1}^2 is total dominating for H_2' but not for H_2 ; let this class be D_k^2 for some $k, 1 \leq k \leq d+1$. Then a_1 must be in D_k . If $a_1 \in D_{d+1}^1$, then $k = d+1$ and both $(H_1, a_1), (H_2, a_2)$ are in $\kappa_1(d)$. If $a_1 \notin D_{d+1}^1$, then $k \neq d+1$ and both $(H_1, a_1), (H_2, a_2)$ are in $\kappa_0(d)$. If $(H_2, a_2) \notin \kappa(d)$ and hence by Theorem 7 $d_t(H_2) > d$ then (i) is satisfied. We have proved that one of the cases (i), (ii), (iii) occurs.

Conversely, assume that $(H_1, a_1) \in \kappa_0(d)$. Construct the described partition $\{D_1^1, \dots, D_{d+1}^1\}$ such that $a_1 \notin D_{d+1}^1$; choose the notation so that $a_1 \in D_1^1$. If $d_t(H_2) > d$, choose a total domatic partition $\{D_1^2, \dots, D_{d+1}^2\}$ of H_2 ; choose the notation so that $a_2 \in D_{d+1}^2$. If we define $D_j = D_j^1 \cup D_j^2$ for $j = 1, \dots, d+1$, then $\{D_1, \dots, D_{d+1}\}$ is a total domatic partition of H and $d_t(H) = d+1$. If $(H_2, a_2) \in \kappa_0(d)$, then construct the described partition $\{D_1^2, \dots, D_{d+1}^2\}$ for H_2 such that $a_2 \in D_1^2$. If we put $D_1 = D_1^1 \cup D_{d+1}^2, D_{d+1} = D_{d+1}^1 \cup D_1^2, D_j = D_j^1 \cup D_j^2$ for $j = 2, \dots, d$, then $\{D_1, \dots, D_{d+1}\}$ is a total domatic partition of H and $d_t(H) = d+1$.

Suppose $(H_1, a_1) \in \kappa_1(d)$. Construct the described partition $\{D_1^1, \dots, D_{d+1}^1\}$ such that $a_1 \in D_{d+1}^1$; if $d_t(H_2) > d$, choose a total domatic partition $\{D_1^2, \dots, D_{d+1}^2\}$ of H_2 ; again choose the notation so that $a_2 \in D_{d+1}^2$. If we define $D_j = D_j^1 \cup D_j^2$ for $j = 1, \dots, d+1$, then $\{D_1, \dots, D_{d+1}\}$ is a total domatic partition of H and $d_t(H) = d+1$. If $(H_2, a_2) \in \kappa_1(d)$, then construct the described partition $\{D_1^2, \dots, D_{d+1}^2\}$ for H_2 such that $a_2 \in D_{d+1}^2$. Now we define again $D_j = D_j^1 \cup D_j^2$, and $\{D_1, \dots, D_{d+1}\}$ is a total domatic partition of H and $d_t(H) = d+1$. This proves Theorem 8. \square

A vertex x of the graph G is called saturated, if it is adjacent to all other vertices of G .

Theorem 9. *Let for the graphs H, H_1, H_2 in the above notation the equality $d_t(H) = 1 + d_t(H_1)$ hold. If a_1 is not saturated in H_1 , then there exists a vertex of H_1 non-adjacent to a_1 with the property that by joining it by an edge to a_1 a graph \hat{H}_1 with total domatic number $d_t(\hat{H}_1) = d_t(H_1) + 1$ is obtained from H_1 .*

The proof is analogous to the proof of Theorem 5. If a_1 is saturated in H_1 , then the unique subset of $V(H_1)$ which is total dominating for H_1' but not for H_1 can be only the set $\{a_1\}$ and thus $D_{d+1}^1 = \{a_1\}$ and $D_{d+1}^1 \cap V(H_1') = \emptyset$.

At the end of the paper we shall prove a theorem on circuits. Let C_n be the circuit of length n . Its vertices will be denoted by u_1, \dots, u_n so that the edges of C_n are (u_i, u_{i+1}) for $i = 1, \dots, n-1$ and (u_n, u_1) . It is known (cf. [2]) that $d_t(C_n) = 2$ if and only if $n \equiv 0 \pmod{4}$; otherwise $d_t(C_n) = 1$. \square

In the following theorem the circuit C_n will be considered as a graph H_1 or H_2 in the notation introduced above; in this sense we shall write the pair (C_n, a) and the classes $\kappa(1)$, $\kappa_0(1)$ and $\kappa_1(1)$.

Theorem 10. *Let C_n be a circuit of length $n \not\equiv 0 \pmod{4}$, let a be an arbitrary vertex of C_n . Then*

- (1) $(C_n, a) \in \kappa_1(1) \setminus \kappa_0(1)$ for $n \equiv 3 \pmod{4}$;
- (2) $(C_n, a) \in \kappa_0(1) \setminus \kappa_1(1)$ for $n \equiv 1 \pmod{4}$;
- (3) $(C_n, a) \notin \kappa(1)$ for $n \equiv 2 \pmod{4}$.

Proof. Without loss of generality put $a = u_n$. Suppose that $(C_n, a) \in \kappa(1)$. Then there exists a partition $\{D_1, D_2\}$ of $V(C_n)$ such that D_1 is a total dominating set in C_n and D_2 is total dominating for the path obtained from C_n by deletion of u_n , but not for C_n . None of the vertices adjacent to u_n belongs to D_2 , therefore $u_1 \in D_1$, $u_{n-1} \in D_1$. Suppose that $(C_n, a) \in \kappa_0(1)$, i.e. $u_n \in D_1$. Each vertex of C_n distinct from u_n must be adjacent to a vertex of D_1 and to a vertex of D_2 . As $u_n \in D_1$, $u_1 \in D_1$, we have $u_i \in D_2$ for $i \equiv 2 \pmod{4}$ or $i \equiv 3 \pmod{4}$ and $u_i \in D_1$ for $i \equiv 0 \pmod{4}$ or $i \equiv 1 \pmod{4}$; in all cases $i \neq n$. But as was mentioned above, $u_{n-1} \in D_1$. This is possible only if $n-1 \equiv 0 \pmod{4}$ or $n-1 \equiv 1 \pmod{4}$, i.e., if $n \equiv 1 \pmod{4}$ or $n \equiv 2 \pmod{4}$. If $n \equiv 2 \pmod{4}$, then also $u_{n-2} \in D_1$ and u_{n-1} is adjacent to two vertices u_{n-2} and u_n of D_1 ; this is a contradiction. Therefore $(C_n, a) \in \kappa_0(1)$ implies $n \equiv 1 \pmod{4}$, and conversely for $n \equiv 1 \pmod{4}$ the described partition exists so that $(C_n, a) \in \kappa_0(1)$.

Next, assume that $(C_n, a) \in \kappa_1(1)$, i.e. $u_n \in D_2$. Then $u_i \in D_1$ for $i \equiv 1 \pmod{4}$ or $i \equiv 2 \pmod{4}$ and $u_i \in D_2$ for $i \equiv 0 \pmod{4}$ or $i \equiv 3 \pmod{4}$ again for all $i \neq n$. We have $u_{n-1} \in D_1$ and thus $n-1 \equiv 1 \pmod{4}$ or $n-1 \equiv 2 \pmod{4}$, i.e. $n \equiv 2 \pmod{4}$ or $n \equiv 3 \pmod{4}$. If $n \equiv 2 \pmod{4}$, then $u_{n-2} \in D_2$ and u_{n-1} is adjacent to two vertices u_{n-2} and u_n of D_2 ; this is a contradiction. Therefore $(c_n, a) \in \kappa_1(1)$ implies that $n \equiv 3 \pmod{4}$, and conversely for $n \equiv 3 \pmod{4}$ the described partition exists and $(C_n, a) \in \kappa_1(1)$. We have proved that $(C_n, a) \in \kappa_0(1)$ if and only if $n \equiv 1 \pmod{4}$ and $(C_n, a) \in \kappa_1(1)$ if and only if $n \equiv 3 \pmod{4}$. This proves Theorem 10. \square

We are now able to illustrate Theorem 8 by Figures 1–5 below.

In Fig. 1 we see a graph H with $H_1 \cong H_2 \cong C_5$, in Fig. 2 with $H_1 \cong H_2 \cong C_7$. The set D_1 (or D_2) is the set of all vertices labelled by 1 (or 2, respectively). From Theorems 8 and 10 we see that $d_t(H) = 2$ in both cases. In Fig. 3 there is a graph H and $H_1 \cong C_5$, $H_2 \cong C_7$; its total domatic number is 1. Figures 4 and 5 demonstrate (ii) and (iii) in Theorem 8 for a graph H with $(H_1, a_1) \in \kappa_0(1) \cap \kappa_1(1)$. Here H_1 is a C_4 , one vertex of which is joined to a new vertex, a_1 .




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5

References

- [1] *E. J. Cockayne, S. T. Hedetniemi*: Towards a theory of domination on graphs. *Networks* 7 (1977), 247–261.
- [2] *E. J. Cockayne, R. M. Daves, S. T. Hedetniemi*: Total domination in graphs. *Networks* 10 (1980), 211–219.
- [3] *S. T. Hedetniemi, R. C. Laskar eds.*: Topics on domination. A special volume of *Discrete Mathematics* 86 (1990), no. 5, 1–13.

Authors' addresses: *Preben Dahl Vestergaard*, Department of Mathematics and Computer Science, Aalborg University, Frederik Bajers Vej 7, DK 9220 Aalborg 0, Denmark; *Bohdan Zelinka*, Department of Discrete Mathematics and Statistics, Technical University of Liberec, Hálkova 6, 461 17 Liberec, Czech Republic.