

## A NOTE ON RADIO ANTIPODAL COLOURINGS OF PATHS

RIADH KHENNOUFA, OLIVIER TOGNI, Dijon

(Received November 12, 2004)

*Abstract.* The radio antipodal number of a graph  $G$  is the smallest integer  $c$  such that there exists an assignment  $f: V(G) \rightarrow \{1, 2, \dots, c\}$  satisfying  $|f(u) - f(v)| \geq D - d(u, v)$  for every two distinct vertices  $u$  and  $v$  of  $G$ , where  $D$  is the diameter of  $G$ . In this note we determine the exact value of the antipodal number of the path, thus answering the conjecture given in [G. Chartrand, D. Erwin and P. Zhang, *Math. Bohem.* 127 (2002), 57–69]. We also show the connections between this colouring and radio labelings.

*Keywords:* radio antipodal colouring, radio number, distance labeling

*MSC 2000:* 05C78, 05C12, 05C15

## 1. INTRODUCTION

Let  $G$  be a connected graph and let  $k$  be an integer,  $k \geq 1$ . The distance between two vertices  $u$  and  $v$  of  $G$  is denoted by  $d(u, v)$  and the diameter of  $G$  by  $D(G)$  or simply  $D$ . A *radio  $k$ -colouring*  $f$  of  $G$  is an assignment of positive integers to the vertices of  $G$  such that

$$|f(u) - f(v)| \geq 1 + k - d(u, v)$$

for every two distinct vertices  $u$  and  $v$  of  $G$ .

Following the notation of [1], [3], we define the *radio  $k$ -colouring number*  $rc_k(f)$  of a radio  $k$ -colouring  $f$  of  $G$  to be the maximum colour assigned to a vertex of  $G$  and the *radio  $k$ -chromatic number*  $rc_k(G)$  to be  $\min\{rc_k(f)\}$  taken over all radio  $k$ -colourings  $f$  of  $G$ .

Radio  $k$ -colourings generalize many graph colourings. For  $k = 1$ ,  $rc_1(G) = \chi(G)$ , the chromatic number of  $G$ . For  $k = 2$ , the radio 2-colouring problem corresponds to the well studied  $L(2, 1)$ -colouring problem and  $rc_2(G) = \lambda(G)$  (see [5] and references therein). For  $k = D(G) - 1$ , the radio  $(D - 1)$ -colouring is referred to as the *radio*

*antipodal colouring*, because only antipodal vertices can have the same colour. In that case,  $rc_k(G)$  is called the *radio antipodal number*, also denoted by  $ac(G)$ . Finally, for the case  $k = D(G)$ ,  $rc_k(G)$  is called the *radio number* and is studied in [1], [6].

In [2] the antipodal number for cycles was discussed and bounds were given. In [3], Chartrand et al. gave general bounds for the antipodal number of a graph. The authors proved the following result for the radio antipodal number of the path:

**Theorem 1** ([3]). *For every positive integer  $n$ ,*

$$ac(P_n) \leq \binom{n-1}{2} + 1.$$

Moreover, they conjectured that the above upper bound is the value of the antipodal number of the path. In [4], the authors found a sharper bound for the antipodal number of an odd path (thus showing that the conjecture was false):

**Theorem 2** ([4]). *For the path  $P_n$  of odd order  $n \geq 7$ ,*

$$ac(P_n) \leq \binom{n-1}{2} - \frac{n-1}{2} + 4.$$

In this note we completely determine the antipodal number of the path:

**Theorem 3.** *For any  $n \geq 5$ ,*

$$ac(P_n) = \begin{cases} 2p^2 - 2p + 3 & \text{if } n = 2p + 1, \\ 2p^2 - 4p + 5 & \text{if } n = 2p. \end{cases}$$

Notice that for  $n = 2p + 1$  we have  $\binom{n-1}{2} - \frac{n-1}{2} + 4 = p(2p-1) - p + 4 = 2p^2 - 2p + 4$ , thus the bound of Theorem 2 is one from the optimal.

Examples of minimal antipodal colourings of  $P_7$  and  $P_8$  are given in Figure 1.




Figure 1. Antipodal colouring of  $P_7$  and  $P_8$ .

In order to prove Theorem 3, we shall use a result of Liu and Zhu [6] about the radio number of the path. Notice that Liu and Zhu allow 0 to be used as a colour but we do not. Then, when presenting their result, we will make the necessary adjustment (adding “one”) to be consistent with the rest of the paper.

**Theorem 4** ([6]). For any  $n \geq 3$

$$\text{rc}_{n-1}(P_n) = \begin{cases} 2p^2 + 3 & \text{if } n = 2p + 1, \\ 2p^2 - 2p + 2 & \text{if } n = 2p. \end{cases}$$

## 2. RADIO $k$ -COLOURINGS

**Lemma 1.** Let  $G$  be a graph of order  $n$  and let  $k$  be an integer. If  $f$  is a radio  $k$ -colouring of  $G$  then, for any integer  $k' > k$ , there exists a radio  $k'$ -colouring  $f'$  of  $G$  with  $\text{rc}_{k'}(f') \leq \text{rc}_k(f) + (n-1)(k' - k)$ .

*Proof.* We construct a radio  $k'$ -colouring  $f'$  of  $G$  with  $\text{rc}_{k'}(f') = c + (n-1)(k' - k)$  from a radio  $k$ -colouring  $f$  with  $\text{rc}_k(f) = c$  in the following way: Let  $x_1, x_2, \dots, x_n$  be an ordering of the vertices of  $G$  such that  $f(x_i) \leq f(x_{i+1})$ ,  $1 \leq i \leq n-1$ , and set

$$f'(x_i) = f(x_i) + (i-1)(k' - k).$$

For any two integers  $i$  and  $j$ ,  $1 \leq i < j \leq n$ , we have  $|f'(x_j) - f'(x_i)| = |f(x_j) - f(x_i)| + (j-i)(k' - k)$ .

As  $|f(x_j) - f(x_i)| \geq 1 + k - d(x_j, x_i)$  and  $j - i \geq 1$ , we obtain  $|f'(x_j) - f'(x_i)| \geq 1 + k + (j-i)(k' - k) - d(x_j, x_i) \geq 1 + k' - d(x_j, x_i)$ . Thus  $f'$  is a radio  $k'$ -colouring of  $G$  and  $\text{rc}_{k'}(f') = c + (n-1)(k' - k)$ .  $\square$

The above result can be strengthened a little in some cases:

**Lemma 2.** Let  $G$  be a graph of order  $n$  and let  $k, k'$  be integers,  $k' > k$ . Given a radio  $k$ -colouring  $f$  of  $G$ , let  $x_1, x_2, \dots, x_n$  be an ordering of the vertices of  $G$  such that  $f(x_i) \leq f(x_{i+1})$ ,  $1 \leq i \leq n-1$  and let  $\varepsilon_i = |f(x_i) - f(x_{i-1})| - (1 + k - d(x_i, x_{i-1}))$ ,  $2 \leq i \leq n$ . Consider a set  $I = \{i_1, i_2, \dots, i_s\} \subset \{2, \dots, n\}$ , where  $1 \leq s \leq n-1$ , such that  $i_{j+1} > i_j + 1$  for all  $j$ ,  $1 \leq j \leq s-1$ . Then there exists a radio  $k'$ -colouring  $f'$  of  $G$  with  $\text{rc}_{k'}(f') \leq \text{rc}_k(f) + (n-1)(k' - k) - \sum_{i \in I} \min(k' - k, \varepsilon_i)$ .

*Proof.* A radio  $k'$ -colouring  $f'$  of  $G$  is obtained simply by setting for all  $j$  with  $1 \leq j \leq n-1$ :

$$f'(x_j) = f(x_j) + (j-1)(k' - k) - \sum_{i \in I, i \leq j} \min(k' - k, \varepsilon_i).$$

The vertex  $x_n$  has the maximum colour:  $f'(x_n) = f(x_n) + (n-1)(k' - k) - \sum_{i \in I} \min(k' - k, \varepsilon_i) = \text{rc}_k(f) + (n-1)(k' - k) - \sum_{i \in I} \min(k' - k, \varepsilon_i)$ .

Then, for any two integers  $j_1$  and  $j_2$ ,  $1 \leq j_1 < j_2 \leq n$ , let us show that the condition

$$|f'(x_{j_2}) - f'(x_{j_1})| \geq 1 + k' - d(x_{j_2}, x_{j_1})$$

is verified, i.e. that

$$|f(x_{j_2}) - f(x_{j_1})| + (j_2 - j_1)(k' - k) - \sum_{i \in I, j_1 < i \leq j_2} \min(k' - k, \varepsilon_i) \geq 1 + k' - d(x_{j_2}, x_{j_1}).$$

If  $j_2 = j_1 + 1$ , then  $|f(x_{j_2}) - f(x_{j_1})| = 1 + k - d(x_{j_2}, x_{j_1}) + \varepsilon_{j_2}$ . Thus  $|f'(x_{j_2}) - f'(x_{j_1})| \geq 1 + k - d(x_{j_2}, x_{j_1}) + \varepsilon_{j_2} + (k' - k) - \min(k' - k, \varepsilon_{j_2}) \geq 1 + k' - d(x_{j_2}, x_{j_1})$ .

If  $j_2 > j_1 + 1$ , then  $\sum_{i \in I, j_1 < i \leq j_2} \min(k' - k, \varepsilon_i) \leq (j_2 - j_1 - 1)(k' - k)$  since by the hypothesis there are no two consecutive integers in the set  $I$ . Thus  $|f'(x_{j_2}) - f'(x_{j_1})| \geq 1 + k - d(x_{j_2}, x_{j_1}) + (j_2 - j_1)(k' - k) - (j_2 - j_1 - 1)(k' - k) = 1 + k' - d(x_{j_2}, x_{j_1})$ .

Therefore,  $f'$  is a radio  $k'$ -colouring of  $G$  and  $\text{rc}_{k'}(f') = \text{rc}_k(f) + (n - 1)(k' - k) - \sum_{i \in I} \min(k' - k, \varepsilon_i)$ .  $\square$

### 3. ANTIPODAL COLOURINGS OF PATHS

Theorem 3 derives from the next two theorems.

**Theorem 5.** For any  $n \geq 5$ ,

$$\text{ac}(P_n) \leq \begin{cases} 2p^2 - 2p + 3 & \text{if } n = 2p + 1, \\ 2p^2 - 4p + 5 & \text{if } n = 2p. \end{cases}$$

*Proof.* The fact that  $\text{ac}(P_5) = 7$  is easily checked (see [3]). Thus take  $n \geq 6$  and let  $P_n = (u_1, u_2, \dots, u_n)$ . We consider two cases depending on whether  $n$  is even or odd.

*Case 1.*  $n = 2p + 1$  is odd for an integer  $p \geq 3$ . Define a colouring  $f$  of  $P_{2p+1}$  by

$$\begin{cases} f(u_1) = 3p + 2, \\ f(u_2) = p + 1, \\ f(u_i) = i(2p - 1) - p + 3, & 3 \leq i \leq p, \\ f(u_{p+1}) = 2p + 2, \\ f(u_{p+2}) = 1, \\ f(u_{p+i}) = i(2p - 1) - 2p + 3, & 3 \leq i \leq p, \\ f(u_{2p+1}) = p + 2. \end{cases}$$

Then the vertex  $u_p$  has the maximum colour:  $f(u_p) = p(2p-1)-p+3 = 2p^2-2p+3$ . We only have to show that the distance condition is verified for two vertices  $u_i$  and  $u_{p+j}$ ,  $3 \leq i, j \leq p$  (the other cases can be easily checked). We want

$$\begin{aligned} |f(u_{p+j}) - f(u_i)| &\geq 1 + (D-1) - d(u_{p+j}, u_i) \Leftrightarrow \\ |j(2p-1) - 2p + 3 - (i(2p-1) - p + 3)| &\geq 2p - (p+j-i) \Leftrightarrow \\ |(j-i)(2p-1) - p| &\geq p - j + i. \end{aligned}$$

If  $j-i \geq 1$  then  $|(j-i)(2p-1) - p| = (j-i)(2p-1) - p \geq 2p-1-p = p-1 \geq p-j+i$ .  
If  $j-i < 1$  then  $|(j-i)(2p-1) - p| = -(j-i)(2p-1) + p = (i-j)(2p-1) + p \geq p-j+i$  for  $p \geq 1$ .

C a s e 2.  $n = 2p$  is even for an integer  $p \geq 3$ . Define a colouring  $f$  of  $P_{2p}$  by

$$\begin{cases} f(u_1) = p, \\ f(u_i) = (p-i)(2p-1) + 2, & 2 \leq i \leq p-1, \\ f(u_p) = 2p^2 - 4p + 5, \\ f(u_{p+i}) = 2p^2 - 4p + 6 - f(u_{p-i+1}), & 1 \leq i \leq p. \end{cases}$$

Then the vertex  $u_p$  has the maximum colour:  $f(u_p) = 2p^2 - 4p + 5$ . We only have to show that the distance condition is verified for two vertices  $u_i$  and  $u_{p+j}$ ,  $2 \leq i \leq p-1$ ,  $1 \leq j \leq p$  (the other cases can be easily checked). We want

$$\begin{aligned} |f(u_{p+j}) - f(u_i)| &\geq 1 + (D-1) - d(u_{p+j}, u_i) \Leftrightarrow \\ |(p-j)(2p-1) + 3 - ((p-i)(2p-1) - p + 2)| &\geq 2p-1 - (p+j-i) \Leftrightarrow \\ |(i-j)(2p-1) + p + 1| &\geq p - j + i - 1. \end{aligned}$$

If  $i-j \geq 0$  then  $|(i-j)(2p-1) + p + 1| = (i-j)(2p-1) + p + 1 \geq p - j + i - 1$  since  $(i-j)(2p-2) \geq -1$  for  $p \geq 1$ .

If  $i-j < 0$ , i.e. if  $j-i \geq 1$  then  $|(i-j)(2p-1) + p + 1| = (j-i)(2p-1) - p - 1 \geq p - j + i - 1$  since  $2p(j-i) \geq 2p$ .  $\square$

**Theorem 6.** For any  $n \geq 5$ ,

$$\text{ac}(P_n) \geq \begin{cases} 2p^2 - 2p + 3 & \text{if } n = 2p + 1, \\ 2p^2 - 4p + 5 & \text{if } n = 2p. \end{cases}$$

**P r o o f.** For  $n = 2p + 1$ , by Lemma 1 we have  $\text{rc}_{n-1}(P_n) \leq \text{ac}(P_n) + (n-1)$ . This together with Theorem 4 gives  $\text{ac}(P_{2p+1}) \geq 2p^2 + 3 - 2p$ .

For  $n = 2p$ , let  $D = D(P_{2p}) = 2p - 1$ . We will use Lemma 2 with the radio  $(D - 1)$ -colouring  $f$  of  $P_{2p}$  described in the proof of Theorem 5 and with  $k = D - 1 = 2p - 1$  and  $k' = D = 2p$ . Keeping the notation of Lemma 2, one can see that  $f$  is such that  $x_1 = u_{p+1}$ ,  $x_2 = u_1$ ,  $x_3 = u_{2p-1}$ ,  $x_4 = u_{p-1}$ ,  $\dots$ ,  $x_{2j+1} = u_{2p-j+1}$ ,  $x_{2j} = u_{p-j+1}$ ,  $\dots$ ,  $x_{2p-1} = u_{2p}$ ,  $x_{2p} = u_p$ . Thus  $\varepsilon_3$  verifies

$$\begin{aligned} \varepsilon_3 &= |f(x_3) - f(x_2)| - (1 + k - d(x_3, x_2)) \\ &= |f(u_{2p-1}) - f(u_1)| - (1 + 2p - 2 - (2p - 2)) \\ &= |2p^2 - 4p + 6 - f(u_2) - f(u_1)| - 1 \\ &= |2p^2 - 4p + 6 - (p - 2)(2p - 1) - 2 - p| - 1 = 1. \end{aligned}$$

A similar calculus gives  $\varepsilon_{2p-1} = 1$  and  $\varepsilon_i = 0$  for all other indices.

Thus, as  $k' - k = 1$  and  $p \geq 3$ , applying Lemma 2 with  $I = \{3, 2p - 1\}$  gives

$$\text{rc}_{2p-1}(P_{2p}) \leq \text{ac}(P_{2p}) + (2p - 1) - \varepsilon_3 - \varepsilon_{2p-1},$$

that is

$$\text{ac}(P_{2p}) \geq \text{rc}_{2p-1}(P_{2p}) - (2p - 1) + \varepsilon_3 + \varepsilon_{2p-1}.$$

By virtue of Theorem 4 we obtain  $\text{ac}(P_{2p}) \geq 2p^2 - 2p + 2 - (2p - 1) + 1 + 1 = 2p^2 - 4p + 5$ .  $\square$

#### References

- [1] *G. Chartrand, D. Erwin, F. Harary, P. Zhang*: Radio labelings of graphs. *Bull. Inst. Combin. Appl.* *33* (2001), 77–85.
- [2] *G. Chartrand, D. Erwin, P. Zhang*: Radio antipodal colorings of cycles. *Congr. Numerantium* *144* (2000), 129–141.
- [3] *G. Chartrand, D. Erwin, P. Zhang*: Radio antipodal colorings of graphs. *Math. Bohem.* *127* (2002), 57–69.
- [4] *G. Chartrand, L. Nebeský, P. Zhang*: Radio  $k$ -colorings of paths. *Discuss. Math. Graph Theory* *24* (2004), 5–21.
- [5] *D. Kuo, J.-H. Yan*: On  $L(2, 1)$ -labelings of Cartesian products of paths and cycles. *Discrete Math.* *283* (2004), 137–144.
- [6] *D. Liu, X. Zhu*: Multi-level distance labelings for paths and cycles. To appear in *SIAM J. Discrete Math.*

*Author's address*: Riadh Khennoufa, Olivier Togni, LE2I, UMR 5158 CNRS, Université de Bourgogne, BP 47870, 21078 Dijon cedex, France, e-mail: [olivier.togni@u-bourgogne.fr](mailto:olivier.togni@u-bourgogne.fr).