

# Weak solution approach in fluid mechanics

Eduard Feireisl

based on joint work with J. Březina (Tokio), P. Gwiazda, A. Świerczewska-Gwiazda (Warsaw),  
Ch.Klingenberg (Wuerzburg), S.Markfelder (Wuerzburg), M. Medviďová/Lukáčová (Mainz), E.Wiedemann  
(Bonn)

Institute of Mathematics, Academy of Sciences of the Czech Republic, Prague

**Matematika v sovremennom mire, Novosibirsk August 14 - August 18 2017**

The research leading to these results has received funding from the European Research Council  
under the European Union's Seventh Framework Programme (FP7/2007-2013)/ ERC Grant Agreement 320078

# Hierarchy of solutions

## Strong solution

Strong *a priori* bounds also on derivatives

## Weak (distributional) solutions

$L^\alpha$ ,  $\alpha > 1$  *a priori* bounds, compactness in  $L^1$

## Measure-valued solutions

$L^\alpha$ ,  $\alpha > 1$  *a priori* bounds

## Measure-valued solutions with concentration measure

$L^1$  *a priori* bounds

## Dissipative measure-valued (DMV) solutions

A form of energy balance, concentration defect dominated by dissipation defect. (DMV)-strong uniqueness principle

# Barotropic Euler/ Navier Stokes system

## Field equations

$$\partial_t \varrho + \operatorname{div}_x \mathbf{m} = 0$$

$$\partial_t \mathbf{m} + \operatorname{div}_x \left( \frac{\mathbf{m} \otimes \mathbf{m}}{\varrho} \right) + \nabla_x p(\varrho) = \begin{cases} 0 \\ \operatorname{div}_x \mathbb{S} \end{cases}$$

$$\mathbf{m} = \varrho \mathbf{u}, \quad \mathbb{S} = \mu \left( \nabla_x \mathbf{u} + \nabla_x^t \mathbf{u} - \frac{2}{3} \operatorname{div}_x \mathbf{u} \mathbb{I} \right) + \eta \operatorname{div}_x \mathbf{u} \mathbb{I}$$

## Initial conditions, periodic boundary conditions

$$\varrho(0, \cdot) = \varrho_0, \quad \mathbf{m}(0, \cdot) = \mathbf{m}_0$$

$$x \in \mathcal{T}^N, \quad N = (1), 2, 3$$

## Pressure, pressure potential

$$p = p(\varrho), \quad p'(\varrho) \geq 0, \quad P(\varrho) = \varrho \int_1^{\varrho} \frac{p(z)}{z^2} dz$$

## Dissipative (weak) solutions

### Energy inequality

$$E(\tau) + \int_0^\tau \int_\Omega \mathbb{S}(\nabla_x \mathbf{u}) : \nabla_x \mathbf{u} \, dx \, dt \leq E(0)$$

$$E = \int_\Omega \frac{1}{2} \varrho |\mathbf{u}|^2 + P(\varrho) \, dx, \quad P(\varrho) = \varrho \int_1^\varrho \frac{p(z)}{z^2} \, dz$$

# Relative energy/entropy

## Lyapunov function

$$E = \int_{\Omega} \left[ \frac{1}{2} \varrho |\mathbf{u} - \mathbf{0}|^2 + P(\varrho) - P'(\bar{\varrho})(\varrho - \bar{\varrho}) - P(\bar{\varrho}) \right] dx$$

## Coercivity of the pressure potential

$$\varrho \mapsto p(\varrho) \text{ non-decreasing} \Rightarrow \varrho \mapsto P(\varrho) \text{ convex}$$

## Relative energy (relative entropy Dafermos [1979] )

$$\begin{aligned} & \mathcal{E}(\varrho, \mathbf{u} \mid r, \mathbf{U}) \\ &= \int_{\Omega} \left[ \frac{1}{2} \varrho |\mathbf{u} - \mathbf{U}|^2 + P(\varrho) - P'(r)(\varrho - r) - P(r) \right] dx \end{aligned}$$

# Dissipative solutions

## Relative energy inequality

$$\begin{aligned} & \left[ \mathcal{E}(\varrho, \mathbf{u} \mid r, \mathbf{U}) \right]_{t=0}^{t=\tau} \\ & + \int_0^\tau \int_\Omega (\mathbb{S}(\nabla_x \mathbf{u}) - \mathbb{S}(\nabla_x \mathbf{U})) : (\nabla_x \mathbf{u} - \nabla_x \mathbf{U}) \, dx \, dt \\ & \leq \int_0^\tau \mathcal{R}(\varrho, \mathbf{u} \mid r, \mathbf{U}) \, dt \end{aligned}$$

## Test functions

$r > 0$ ,  $r, \mathbf{U}$  periodic (or other relevant b.c.)

## Remainder

$$\begin{aligned} & \int_0^\tau \mathcal{R}(\varrho, \mathbf{u} \mid r, \mathbf{U}) \, dt \\ & \int_\Omega \varrho (\partial_t \mathbf{U} + \mathbf{u} \cdot \nabla_x \mathbf{U}) \cdot (\mathbf{U} - \mathbf{u}) \, dx \\ & + \int_\Omega \mathbb{S}(\nabla_x \mathbf{U}) : (\nabla_x \mathbf{U} - \nabla_x \mathbf{u}) \, dx + \int_\Omega (\rho(r) - \rho(\varrho)) \operatorname{div}_x \mathbf{U} \, dx \\ & + \int_\Omega [(r - \varrho) \partial_t P'(r) + \nabla_x P'(r) \cdot (r \mathbf{U} - \varrho \mathbf{u})] \, dx \end{aligned}$$

# Weak (distributional) solutions

## Navier–Stokes system

- P.L.Lions [1998] - global in time existence for  $p \approx a\varrho^\gamma$ ,  $\gamma \geq \frac{9}{5}$  ( $N = 3$ )
- EF, A.Novotný, H.Petzeltová [2000] - extension to  $\gamma > \frac{N}{2}$

## Euler system

- E.Chiodaroli, EF [2014, 2015] - global in time existence of *infinitely many* weak solutions for any smooth initial data
- E.Chiodaroli, EF [2014, 2015] - global in time existence of infinitely many dissipative weak solutions for special initial data
- E.Chiodaroli, C.De Lellis, O.Kreml [2016] - global in time existence of admissible entropy solutions for Lipschitz initial data



# What is a good weak solution?

## Desired properties

- A weak solution exists globally in time for “any” choice of the initial state
- A weak solution can be identified as a limit of suitable approximate problems, e.g. by adding artificial viscosity
- The set of weak solutions is closed; a limit of a family of weak solutions is another weak solution
- A weak solution can be identified as a limit of a numerical scheme
- A weak solution is the most general object that enjoys the weak–strong uniqueness property

## Weak strong uniqueness

A weak solution coincides with a strong (classical) solution as long as the latter exists

# Measure-valued solutions

## Derivatives

Partial derivatives replaced by distributional derivatives

## Oscillations

A parameterized measure (Young measure)

$\nu_{t,x} \in \mathcal{P}(F)$ ,  $t$  – time,  $x$  – spatial variable,  $F$  – phase space

$\mathbf{U} : Q \rightarrow F$ ,  $f(\mathbf{U})(t, x)$  replaced by expectations  $\langle \nu_{t,x}; f(\mathbf{U}) \rangle$

## Concentrations

Concentration measure  $\mathcal{C} \in \mathcal{M}(Q)$

# Measure valued solutions

## Equation of continuity

$$\int_0^T \int_{\mathcal{T}^N} \langle \nu_{t,x}; \varrho \rangle \partial_t \varphi + \langle \nu_{t,x}; \mathbf{m} \rangle \cdot \nabla_x \varphi \, dx dt = \int_0^T \int_{\mathcal{T}^N} \nabla_x \varphi \cdot d\mathcal{C}_1$$

for all  $\varphi \in C_c^\infty((0, T) \times \mathcal{T}^N)$

## Momentum equation

$$\begin{aligned} \int_0^T \int_{\mathcal{T}^N} \langle \nu_{t,x}; \mathbf{m} \rangle \partial_t \varphi + \left\langle \nu_{t,x}; \frac{\mathbf{m} \otimes \mathbf{m}}{\varrho} \right\rangle : \nabla_x \varphi + \langle \nu_{t,x}; \mathbf{p}(\varrho) \rangle \operatorname{div}_x \varphi \, dx dt \\ = \int_0^T \int_{\mathcal{T}^N} \nabla_x \varphi : d\mathcal{C}_2 \end{aligned}$$

for all  $\varphi \in C_c^\infty((0, T) \times \mathcal{T}^N; \mathbb{R}^N)$

# Energy dissipation

## Energy inequality

$$\frac{d}{dt} \int_{\mathcal{T}^N} \left( \frac{1}{2} \varrho |\mathbf{u}|^2 + P(\varrho) \right) dx \leq 0$$

## Measure-valued energy inequality

$$\begin{aligned} \int_{\mathcal{T}^N} \left\langle \nu_{\tau, x}; \left( \frac{1}{2} \frac{|\mathbf{m}|^2}{\varrho} + P(\varrho) \right) \right\rangle dx + \mathcal{D}(\tau) \\ \leq \int_{\mathcal{T}^N} \left\langle \nu_0; \left( \frac{1}{2} \frac{|\mathbf{m}|^2}{\varrho} + P(\varrho) \right) \right\rangle dx \end{aligned}$$

## Dissipation defect - compatibility

$$\left| \mathcal{C}_1[0, \tau] \times \mathcal{T}^N \right| + \left| \mathcal{C}_2[0, \tau] \times \mathcal{T}^N \right| \leq \xi(\tau) \mathcal{D}(\tau), \quad \xi \in L^1(0, T)$$

# Truly measure-valued solutions

## Truly measure-valued solutions for the Euler system (with E.Chiodaroli, O.Kreml, E. Wiedemann)

There is a measure-valued solution to the compressible Euler system (without viscosity) that *is not* a limit of bounded  $L^p$  weak solutions to the Euler system.

## Weak (mv) - strong uniqueness

**Theorem - EF, P.Gwiazda, A.Świerczewska-Gwiazda, E. Wiedemann [2015]**

A measure valued and a strong solution emanating from the same initial data coincide as long as the latter exists

# Relative energy (entropy)

## Relative energy functional

$$\begin{aligned} & \mathcal{E}(\varrho, \mathbf{m} \mid r, \mathbf{U})(\tau) \\ &= \int_{\mathcal{T}^N} \left\langle \nu_{\tau, x}; \frac{1}{2} \frac{|\mathbf{m} - r\mathbf{U}|^2}{\varrho} + P(\varrho) - P'(r)(\varrho - r) - P(r) \right\rangle dx \\ &= \int_{\mathcal{T}^N} \left\langle \nu_{\tau, x}; \frac{1}{2} \frac{|\mathbf{m}|^2}{\varrho} + P(\varrho) \right\rangle dx - \int_{\Omega} \langle \nu_{\tau, x}; \mathbf{m} \rangle \cdot \mathbf{U} dx \\ &\quad + \int_{\Omega} \frac{1}{2} \langle \nu_{\tau, x}; \varrho \rangle |\mathbf{U}|^2 dx \\ &\quad - \int_{\Omega} \langle \nu_{\tau, x}; \varrho \rangle P'(r) dx + \int_{\Omega} p(r) dx \end{aligned}$$

# Relative energy (entropy) inequality

## Relative energy inequality

$$\begin{aligned} & \mathcal{E}(\varrho, \mathbf{m} \mid r, \mathbf{U})(\tau) \\ & \leq \int_{\Omega} \left\langle \nu_{0,x}; \frac{1}{2} \frac{|\mathbf{m} - r\mathbf{U}_0|^2}{\varrho} + P(\varrho) - P'(r_0)(\varrho - r_0) - P(r_0) \right\rangle dx \\ & \quad + \int_0^\tau \mathcal{R}(\varrho, \mathbf{m} \mid r, \mathbf{U}) dt \end{aligned}$$



## Remainder

$$\begin{aligned} & \mathcal{R}(\varrho, \mathbf{m} \mid r, \mathbf{U}) \\ &= - \int_0^\tau \int_\Omega \langle \nu_{t,x}, \mathbf{m} \rangle \cdot \partial_t \mathbf{U} \, dx \, dt \\ & - \int_0^\tau \int_{\bar{\Omega}} \left[ \left\langle \nu_{t,x}; \frac{\mathbf{m} \otimes \mathbf{m}}{\varrho} \right\rangle : \nabla_x \mathbf{U} + \langle \nu_{t,x}; p(\varrho) \rangle \operatorname{div}_x \mathbf{U} \right] dx \, dt \\ & + \int_0^\tau \int_\Omega [\langle \nu_{t,x}; \varrho \rangle \mathbf{U} \cdot \partial_t \mathbf{U} + \langle \nu_{t,x}; \mathbf{m} \rangle \cdot \mathbf{U} \cdot \nabla_x \mathbf{U}] \, dx \, dt \\ & + \int_0^\tau \int_\Omega \left[ \left\langle \nu_{t,x}; \left(1 - \frac{\varrho}{r}\right) \right\rangle p'(r) \partial_t r - \langle \nu_{t,x}; \mathbf{m} \rangle \cdot \frac{p'(r)}{r} \nabla_x r \right] dx \, dt \\ & + \int_0^\tau \int_{\mathcal{T}^N} \frac{1}{2} \nabla_x (|\mathbf{U}|^2 - P'(r)) \, d\mathcal{C}_1 - \int_0^\tau \int_{\mathcal{T}^N} \nabla_x \mathbf{U} \, d\mathcal{C}_2 \end{aligned}$$

## Convergence of a numerical scheme

EF, M. Lukáčová–Medvidřová [2016]

Let  $\Omega \subset \mathbb{R}^3$  be a smooth bounded domain. Let

$$1 < \gamma < 2, \Delta t \approx h, 0 < \alpha < 2(\gamma - 1).$$

Suppose that the initial data are smooth and that the compressible **Navier-Stokes system** admits a smooth solution in  $[0, T]$  in the class

$$\varrho, \nabla_x \varrho, \mathbf{u}, \nabla_x \mathbf{u} \in C([0, T] \times \bar{\Omega})$$

$$\partial_t \mathbf{u} \in L^2(0, T; C(\bar{\Omega}; \mathbb{R}^3)), \varrho > 0, \mathbf{u}|_{\partial\Omega} = 0.$$

Then the numerical solutions resulting from Karlsen-Karper FV-FE scheme converge unconditionally,

$$\varrho_h \rightarrow \varrho \text{ (strongly) in } L^\gamma((0, T) \times K)$$

$$\mathbf{u}_h \rightarrow \mathbf{u} \text{ (strongly) in } L^2((0, T) \times K; \mathbb{R}^3)$$

for any compact  $K \subset \Omega$ .

# General strategy

## Basic properties of numerical scheme

Show stability, consistency, discrete energy inequality

## Measure valued solutions

Show convergence of the scheme to a

**dissipative measure – valued solution**

## Weak-strong uniqueness

Use the weak-strong uniqueness principle in the class of measure-valued solutions. Strong and measure valued solutions emanating from the same initial data coincide as long as the latter exists

# Singular limit problem

## Scaled Euler system

$$\begin{aligned}\partial_t \varrho + \operatorname{div}_x \mathbf{m} &= 0 \\ \partial_t \mathbf{m} + \operatorname{div}_x \left( \frac{\mathbf{m} \otimes \mathbf{m}}{\varrho} \right) + \frac{1}{\varepsilon^2} \nabla_x p(\varrho) &= 0\end{aligned}$$

## Incompressible (low Mach) limit - EF, Ch.Klingenberg, S.Markfelder[2017]

Convergence to the limit system

$$\operatorname{div}_x \mathbf{v} = 0, \quad \partial_t \mathbf{v} + \operatorname{div}_x (\mathbf{v} \otimes \mathbf{v}) + \nabla_x \Pi = 0$$

for well/ill prepared initial data.

# Complete Euler system

## Field equations

$$\partial_t \varrho + \operatorname{div}_x(\varrho \mathbf{u}) = 0$$

$$\partial_t(\varrho \mathbf{u}) + \operatorname{div}_x(\varrho \mathbf{u} \otimes \mathbf{u}) + \nabla_x p(\varrho, \vartheta) = 0$$

$$\begin{aligned} \partial_t \left[ \frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e(\varrho, \vartheta) \right] + \operatorname{div}_x \left( \left[ \frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e(\varrho, \vartheta) \right] \mathbf{u} \right) \\ + \operatorname{div}_x(p(\varrho, \vartheta) \mathbf{u}) = 0 \end{aligned}$$

## Entropy inequality (admissibility)

$$\partial_t(\varrho s(\varrho, \vartheta)) + \operatorname{div}_x(\varrho s(\varrho, \vartheta) \mathbf{u}) \geq 0$$

## Constitutive relations

$$p = \varrho \vartheta, \quad e = c_v \vartheta, \quad s = \log(\vartheta^{c_v}) - \log(\varrho)$$

## A priori estimates

### Energy bounds, total mass conservation

$$\int_{\mathcal{T}^N} \varrho \, dx = \int_{\mathcal{T}^N} \varrho_0 \, dx$$
$$\int_{\mathcal{T}^N} \frac{1}{2} \varrho |\mathbf{u}|^2 + \varrho e(\varrho, \vartheta) \, dx = \int_{\mathcal{T}^N} \frac{1}{2} \varrho_0 |\mathbf{u}_0|^2 + \varrho_0 e(\varrho_0, \vartheta_0) \, dx$$

### Entropy transport

$$s(\varrho, \vartheta)(\tau, x) \geq \inf s(\varrho_0, \vartheta_0)$$

### $L^1$ estimates

$$\|\varrho\|_{L^1}, \|\varrho \mathbf{u}\|_{L^1}, \|\varrho |\mathbf{u}|^2\|_{L^1}, \|\varrho \vartheta\|_{L^1}, \|\varrho s\|_{L^1}, \|\mathbf{p}\|_{L^1}, \|\varrho \mathbf{su}\|_{L^1} \text{ bounded}$$

# MV solutions, I

## Basic state variables

density  $\varrho$ , momentum  $\mathbf{m}$ , internal energy  $E = \varrho e(\varrho, \vartheta)$

$$\nu_{t,x} \in \mathcal{P}([0, \infty) \times \mathbb{R}^N \times [0, \infty))$$

## Equation of continuity

$$\int_0^T \int_{\mathcal{T}^N} [\langle \nu_{t,x}; \varrho \rangle \partial_t \varphi + \langle \nu_{t,x}; \mathbf{m} \rangle \cdot \nabla_x \varphi] \, dx dt = 0$$

for any  $\varphi \in C_c^\infty((0, T) \times \mathcal{T}^N)$

## Momentum equation

$$\begin{aligned} & \int_0^T \int_{\mathcal{T}^N} \left[ \langle \nu_{t,x}; \mathbf{m} \rangle \cdot \boldsymbol{\varphi} + \left\langle \nu_{t,x}; \frac{\mathbf{m} \otimes \mathbf{m}}{\varrho} \right\rangle : \nabla_x \boldsymbol{\varphi} \right] \, dx dt \\ & + \int_0^T \int_{\mathcal{T}^N} \langle \nu_{t,x}; \mathbf{p}(\varrho, E) \rangle \operatorname{div}_x \boldsymbol{\varphi} \, dx dt = \int_0^T \int_{\mathcal{T}^N} \nabla_x \boldsymbol{\varphi} : d\mathcal{C} \end{aligned}$$

for any  $\boldsymbol{\varphi} \in C_c^\infty((0, T) \times \mathcal{T}^N; \mathbb{R}^N)$

## MV solutions, II

### Entropy balance

$$\int_0^T \int_{\mathcal{T}^N} \left[ \langle \nu_{t,x}; \varrho Z(s) \rangle \partial_t \varphi + \langle \nu_{t,x}; Z(s) \mathbf{m} \rangle \cdot \nabla_x \varphi \right] dx dt = - \int_0^T \int_{\mathcal{T}^N} \varphi d\mathcal{D}_1$$

$\mathcal{D}_1 \geq 0$ , for any  $\varphi \in C_c^\infty((0, T) \times \mathcal{T}^N)$ ,  $\varphi \geq 0$ , and any  $Z$  concave,  $Z' \geq 0$ ,  $\sup Z < \infty$

### Total energy balance

$$\left[ \int_{\Omega} \left\langle \nu_{t,x}; \frac{1}{2} \frac{|\mathbf{m}|^2}{\varrho} + E \right\rangle dx \right]_{t=0}^{t=\tau} + \mathcal{D}_2(\tau) = 0$$

### Compatibility

$$\|C\|_{\mathcal{M}([0,\tau] \times \Omega; \mathbb{R}^3 \times \mathbb{R}^3)} \leq c \int_0^\tau [\mathcal{D}_1(t) + \mathcal{D}_2(t)] dt$$



# Relative energy

## Ballistic free energy

$$H_{\Theta}(\varrho, \vartheta) = \varrho e(\varrho, \vartheta) - \Theta \varrho s(\varrho, \vartheta),$$

## Relative energy

$$\begin{aligned} & \mathcal{E}_Z(\varrho, \vartheta, \mathbf{u} \mid r, \Theta, \mathbf{U}) \\ &= \frac{1}{2} \varrho |\mathbf{u} - \mathbf{U}|^2 + \varrho e(\varrho, \vartheta) - \Theta \varrho Z(s(\varrho, \vartheta)) - \frac{\partial H_{\Theta}(r, \Theta)}{\partial \varrho} (\varrho - r) - H_{\Theta}(r, \Theta). \end{aligned}$$

# Weak strong uniqueness

## Hypotheses

$$\vartheta Ds(\varrho, \vartheta) = De(\varrho, \vartheta) + p(\varrho, \vartheta)D\left(\frac{1}{\varrho}\right)$$

$$\frac{\partial p(\varrho, \vartheta)}{\partial \varrho} > 0, \quad \frac{\partial e(\varrho, \vartheta)}{\partial \vartheta} > 0 \text{ for all } \varrho, \vartheta > 0$$

$$|p(\varrho, \vartheta)| \leq c(1 + \varrho + \varrho|s(\varrho, \vartheta)| + \varrho e(\varrho, \vartheta))$$

## Conclusion [Březina, EF 2016]

Weak(MV)–strong uniqueness holds provided the initial density and temperature are strictly positive