

Acrylamide-induced changes in femoral bone microstructure of mice

Anna Sarocka^{1*}, Ramona Babosova¹, Veronika Kovacova¹, Radoslav Omelka², Magdalena Semla³, Edyta Kapusta³, Zofia Goc³, Grzegorz Formicki³, Monika Martiniakova¹

¹Department of Zoology and Anthropology, ²Department of Botany and Genetics, Faculty of Natural Sciences, Constantine the Philosopher University in Nitra, Slovak Republic

³Department of Animal Physiology and Toxicology, Institute of Biology, Faculty of Geography and Biology, Pedagogical University in Cracow, Poland

*Corresponding author: Anna Sarocka, Department of Zoology and Anthropology, Faculty of Natural Sciences, Constantine the Philosopher University in Nitra, Nábřežie mládeže 91, 949 74 Nitra, Slovak Republic

Short title: Acrylamide and bone microstructure

Summary

Acrylamide (AA) is one of the most common toxins in foods. Its effect on bone microstructure has not been investigated. The aim of our study was to analyse the impact of acute exposure to AA on femoral bone microstructure in mice. Adult animals were treated perorally with 2 doses of AA (E1 group, 1 mg/kg b.w.) in a 24-hour period and with 3 doses of AA (E2 group, 1 mg/kg b.w.) in a 48 hour period. Mice exposed to AA had smaller sizes of primary osteon's vascular canals. Secondary osteons were significantly smaller in mice from E2 group; however their increased number (from 38 % to 77 %) was identified in both E1 and E2 groups. In these groups, a higher number of resorption lacunae (from 100 % to 122 %) was also found. The values for bone volume, trabecular number were increased and that for trabecular separation was decreased in mice administered AA. Significantly higher value of bone surface was observed in mice from E1 group whereas trabecular thickness was increased in E2 group. The effect of AA on microstructure of compact and trabecular bone tissues is different. In our study, one dose of AA was used and acute effects of AA were investigated. Therefore, further studies are needed to study mechanisms by which AA acts on bone.

Key words

acrylamide, bone, mouse, microstructure

Acrylamide (AA, CAS Reg. No. 79-06-1), α,β -unsaturated reactive molecule, is an odorless crystalline solid (Erikson 2005; Parzefall 2008; Wei *et al.* 2014). Besides its utilization in industry, AA is a contaminant in foods prepared during high-temperature cooking (Pedreschi *et al.* 2014) like fried potatoes, bakery products, breakfast cereals and coffee. It can also be absorbed across the skin (Li *et al.* 2015).

There are many reports demonstrating various toxic effects of AA in experimental animals (Takami *et al.* 2012), including reproductive toxicity (Park *et al.* 2010), genotoxicity (Li *et al.* 2015) and neurotoxicity (Tyl *et al.* 2000). AA is a substance classified as a probable human carcinogen (Mucii and Wilson, 2008).

Although AA belongs to the most commonly consumed toxins by humans, its effect on bone microstructure is still unknown. Therefore, our study was aimed to determine femoral bone microstructure of adult mice after an acute peroral exposure to AA.

In our experiment, twelve clinically healthy 12-weeks-old Swiss mice (males) were used. The animals were obtained from the accredited experimental laboratory of the Pedagogical University in Cracow. Adult mice were randomly divided into three experimental groups of 4 animals each. In E1 group males were treated perorally with 2 doses of AA (1 mg/kg b.w.) in a 24 hour period. In E2 group mice received 3 doses of AA (1 mg/kg b.w.) during a 48 hour period. The dose of AA was chosen on the basis of the experiment conducted by Doerge *et al.* (2005) and our previous experiences (Kopańska *et al.* 2015). AA was dissolved in physiological saline. The third group without AA administration served as a control one (group C). All the applied procedures were approved by the First Local Ethic Committee on Experiments on Animals in Cracow (resolution number 175/2012).

Three hours after the last dose of AA, all mice were killed and their femora were used for microscopical analyses. Thin sections from femora (70 - 80 μm) were prepared according to the methodology of Martiniaková *et al.* (2008). The qualitative histological characteristics of the compact bone were determined according to the internationally accepted classification systems of Enlow and Brown (1956) and Ricqlés *et al.* (1991). The quantitative (morphometrical) parameters of the compact bone were assessed using the software Motic Images Plus 2.0 ML (Motic China Group Co., Ltd.). We measured area, perimeter, maximum

and minimum diameters of the vascular canals of primary osteons, Haversian canals and secondary osteons in all views (*anterior, posterior, medialis, lateralis*) of thin sections.

Morphometrical characteristics of the trabecular bone and cortical bone thickness were determined using microCT system (μ CT 50, Scanco Medical). μ CT studies were performed on the distal femora to image trabecular bone morphometry. Trabecular bone was analysed in a region of interest starting 1.2 mm from the end of the growth plate and extending 1.5 mm. Following parameters were measured: bone volume, trabecular number, trabecular thickness, trabecular separation and bone surface. Cortical bone thickness was analysed in a region of interest starting 5.2 mm from the end of the growth plate and extending 1.5 mm at femoral midshaft.

Statistical analysis was performed using SPSS 8.0 software. All data were expressed as mean \pm standard deviation. The unpaired Games-Howell's and/or Tukey's tests were used for establishing statistical significance ($P < 0.05$) among all groups.

Endosteal surfaces of femora in mice from the C group consisted of non-vascular bone tissue (in *anterior, medialis* and *lateralis* views), which contained cellular lamellae and osteocytes. In *pars posterior*, primary vascular radial bone tissue was observed. This tissue included vascular canals (branching or non-branching) radiating from the marrow cavity. In the middle parts of the compact bone (mainly in *anterior* and *lateralis* views), a few primary osteons and several secondary osteons were identified. Non-vascular bone tissue was found only in *pars medialis*. Finally, the periosteal border was composed of non-vascular bone tissue (Table 1). These results are in accordance to those of other researchers (Treuting and Dintzis 2011; Enlow and Brown 1956; Reim *et al.* 2008).

In our study, differences in compact bone microstructure of mice from the E1 and E2 groups were identified. Mice from these groups had more intact secondary osteons (from 38 % to 77

%) in the middle part of the compact bone. In addition, more resorption lacunae (from 100 % to 122 %) were found in *pars anterior* of their periosteal border (Table 1).

The changes mentioned above can be explained as an adaptive response of the compact bone to AA toxicity. The results by Raju *et al.* (2015) showed that AA increases the oxidative stress and peroxidation of the lipids. Oxidative stress is characterized by an increased level of reactive oxygen species, which enhance bone resorption (Bai *et al.* 2005) and osteoclastogenesis (Garret *et al.* 1990). Also, a biochemical link between increased lipid peroxidation and reduced bone mineral density has been described (Basu *et al.* 2001).

In total, 353 vascular canals of primary osteons, 54 Haversian canals and 54 secondary osteons were measured. The results are summarized in Table 1. All measured variables (area, perimeter, maximal and minimal diameters) of the primary osteon's vascular canals were significantly decreased ($P < 0.05$) in groups E1 and E2 when compared to the C group. Significant differences were also found between E1 and E2 groups. Haversian canals values did not differ significantly among all groups. On the contrary, the values of the secondary osteons were significantly smaller ($P < 0.05$) in mice from the E2 group. Significant differences were also demonstrated between the E1 and E2 groups.

Vasoconstriction of the vascular canals of primary osteons in mice exposed to AA could be associated with deleterious effect of AA on blood vessels. According Raju *et al.* (2015) AA decreases the high density lipoprotein (HDL). Low HDL is associated with narrowing or blockage of the arteries and vessels (Miller *et al.* 1992).

Significantly reduced sizes of the secondary osteons in mice from the E2 group could be related to decreased bone mineralization. The results by Wauquier *et al.* (2009) showed that AA increased the level of the hydrogen peroxide. The hydrogen peroxide significantly decreases the bone mineralization (Arai *et al.* 2007).

Our results also showed an insignificant effect of AA administration on cortical bone thickness (0.18 ± 0.009 mm, 0.19 ± 0.006 mm, 0.17 ± 0.022 mm) in mice from the E1, E2 and C groups, respectively.

In the trabecular bone, the values for bone volume and trabecular number were significantly increased ($P<0.05$) in mice administered AA. On the contrary, the value for trabecular separation was significantly decreased in these mice. Significantly higher value ($P<0.05$) of bone surface was observed in mice from the E1 group whereas trabecular thickness was increased in mice from the E2 group. The results are summarized in Table 2. Representative reconstructed 3D images of the trabecular bone are also illustrated in Table 2.

According to Alturfan *et al.* (2011) AA significantly increases synthesis of tissue collagen in various parenchymatous organs (brain, lung, liver, kidney, testes). The collagen is a major organic component of mineralized bone matrix. Also, the results by Raju *et al.* (2015) showed that AA increases a concentration of calcium, an important component of hydroxyapatite crystals (Koutsopoulos 2002), in the blood. Therefore, we suppose that AA could influence the trabecular bone microstructure through these mechanisms.

Our results suggest that acute peroral administration to AA significantly affects microstructure of compact and trabecular bone tissues. However, the impact of AA on microarchitecture of these tissues is different. The compact bone is more resorbed, trabecular bone is more robust. Therefore, it would be necessary to study mechanisms by which AA acts on bone. These mechanisms remain unclear. Anyway, the most evident changes were observed in mice treated with 3 doses of AA in a 48 hour period. It can be concluded that the effect of AA on bone microstructure depends on the degradation rate. Our findings provide the first information related to AA impact on bone microarchitecture in experimental animals.

There are several limitations to our study. First, only one dose of AA was used. Second, we investigated only acute effect of AA on bone microstructure. Therefore, further studies are needed to find out the complex effect of AA in the bone.

Acknowledgement

This work was supported by the Slovak Research and Development Agency under the contract No. SK-PL-2015-0032. The study was also supported by the projects VEGA 1/0653/16 and KEGA 031UKF-4/2016.

References

- ALTURFAN AA, TOZAN-BECEREN A, SEHIRLI AÖ, DEMIRALP E, SENER G, OMURTAG GZ: Resveratrol ameliorates oxidative DNA damage and protects against acrylamide-induced oxidative stress in rats. *Mol Biol Rep* **39**: 4589-4596, 2012.
- ARAI M, SHIBATA Y, PUGDEE K, ABIKO Y, OGATA Y: Effects of reactive oxygen species (ROS) on antioxidant system and osteoblastic differentiation in MC3T3- E1 cells. *UIBMB Life* **59**: 27-33, 2007.
- BAI X, LU D, LIU A, ZHANG Z, LI X, ZOU Z, ZENG W, CHENG B, LUO S: Reactive Oxygen Species Stimulates Receptor Activator of NF- κ B Ligand Expression in Osteoblast. *J Biol Chem* **280**: 17497-17506, 2005.
- BASU S, MICHAELSSON K, OLOFSSON H, JOHANSSON S, MELHUS H: Association between oxidative stress and bone mineral density. *Biochem Biophys Res Commun* **288**: 275-279, 2001.
- DOERGE DR, DA COSTA GG, MCDANIEL LP, CHURCHWELL MI, TWADDLE NC, BELAND FA: DNA adducts derived from administration of acrylamide and glycidamide to mice and rats. *Mutat Res* **580**: 131-141, 2005.

ENLOW DH, BROWN SO: Comparative histological study of fossil and recent bone tissues. Part I., *Tex J Sci* **8**: 405, 1956.

ERIKSON J: Acrylamide in food products: Identification, formation and analytical technology. Ph. D. Thesis. Stockholm: Stockholm University, Department of Environmental Chemistry. 91, 2005.

GARRET IR, BOYCE BF, OREFFO RO, BONEWALD L, POSER J, MUNDY GR: Oxygen-derived free radicals stimulate osteoclastic bone resorption in rodent bone in vitro and in vivo. *J Clin Invest* **85**: 632-639, 1990.

KOPAŃSKA M, LUKÁČ N, KAPUSTA E, FORMICKI G: Acrylamide Influence on Activity of Acetylcholinesterase, Thiol Groups, and Malondialdehyde Content in the Brain of Swiss Mice. *J Biochem Mol Toxicol* **29**: 472-478, 2015.

KOUTSOPOULOS S: Synthesis and characterization of hydroxyapatite crystals: A review study on the analytical methods. *J Biomed Mater Res* **62**: 600-612, 2002.

LI J, LI D, YANG Y, XU T, LI P, HE D: Acrylamide induces locomotor defects and degeneration of dopamine neurons in *Caenorhabditis elegans*. *J Appl Toxicol* **36**: 60-67, 2015.

MARTINIAKOVÁ M, OMEĽKA R, GROSSKOPF B, SIROTKIN AV, CHRENEK P: Sex-related variation in compact bone microstructure of the femoral diaphysis in juvenile rabbits. *Acta Vet Scand* **50**: 15, 2008.

MILLER M, SEIDLER A, KWITEROVICH PO, PEARSON TA: Long-term predictors of subsequent cardiovascular events with coronary artery disease and “desirable” levels of plasma total cholesterol. *Circulation* **86**: 1165-1170, 1992.

MUCCI LA, WILSON KM: Acrylamide intake through diet and human cancer risk. *J Agric Food Chem* **56**: 6013-6019, 2008.

PARK HR, KIM MS, KIM SJ, PARK M, KONG KH, KIM HS, KWACK SJ, KANG TS, KIM SH, KIM HS, LEE J: Acrylamide induces cell death in neural progenitor cells and impairs hippocampal neurogenesis. *Toxicol Lett* **193**: 86-93, 2010.

PARZEFALL W: Minireview on the toxicity of dietary acrylamide. *Food Chem Toxicol* **46**: 1360-1364, 2008.

PEDRESCHI F, MARIOTTI MS, GRANBY K: Current issues in dietary acrylamide: formation, mitigation and risk assessment. *J Sci Agric* **94**: 9-20, 2014.

RAJU J, ROBERTS J, TAYLOR M, PATRY D, CHOMYSHYN E, CALDWELL D, COOKE G, MEHTA R: Toxicological effects of short-term dietary acrylamide exposure in male F344 rats. *Environ Toxicol Pharmacol* **39**: 85-92, 2015.

REIM NS, BREIG B, STAHR K, EBERLE J, HOEFLICH A, WOLF E, ERBEN RG: Cortical bone loss in androgen-deficient aged male rats is mainly caused by increased endocortical bone remodeling. *J Bone Miner Res* **23**: 694-704, 2008.

RICQLÉS AJ, MEUNIER FJ, CASTANET J, FRANCILLON-VIEILLOT H: Comparative microstructure of bone. In: Hall, B. K.: Bone 3, Bone Matrix and Bone Specific Products. Boca Raton: CRC Press, 1991, pp. 1-78.

TAKAMI S, IMAI T, CHO YM, OGAWA K, HIROSE M, NISHIKAWA A: Juvenile rats do not exhibit elevated sensitivity to acrylamide toxicity after oral administration for 12 weeks. *J Appl Toxicol* **32**: 959-967, 2012.




TREUTING PM, DINTZIS SM: Comparative Anatomy and Histology: a Mouse and Human Atlas. In: Academic Press, 2011, pp. 1-480, ISBN: 9780123813626.

TYL RW, FRIEDMAN MA, LOSCO PE, FISCHER LC, JOHNSON KA, STROTHER DE, WOLF CH: Rat two-generation reproduction and dominant lethal study of acrylamide in drinking water. *Reprod Toxicol* **14**: 385-401, 2000.

WAUQUIER F, LEOTOING L, COXAM V, GUICHEUX J, WITTRANT Y: Oxidative stress in bone remodeling and disease. *Trends Mol Med* **15**: 468-477, 2009.




WEI Q, LI J, LI X, ZHANG L, SHI F: Reproductive toxicity in acrylamide-treated female mice. *Reprod toxicol* **46**: 121-128, 2014.

Table 1 Results of compact bone microstructure in mice from the E1, E2 and C groups

	C group	E1 group	E2 group			
						
	1 - non-vascular bone tissue 2 - intact secondary osteons in the middle part of <i>substantia compacta</i> 3 - primary vascular radial bone tissue	1 - non-vascular bone tissue 2 - resorption lacunae near periosteal border	1 - non-vascular bone tissue 2 - intact secondary osteons in the middle part of <i>substantia compacta</i>			
Measured structures	Group	n	Area (μm ²)	Perimeter (μm)	Max. diameter (μm)	Min. diameter (μm)
Vascular canals of primary osteons	C (1)	114	48.802±11.851	25.352±3.143	4.556±0.706	3.402±0.525
	E1 (2)	115	39.135±10.124	22.336±2.994	3.889±0.624	3.176±0.469
	E2 (3)	124	30.059±7.183	19.667±2.531	3.423±0.577	2.769±0.409
	Games-Howell test		1:2 ⁺ ; 1:3 ⁺ ; 2:3 ⁺	1:2 ⁺ ; 1:3 ⁺ ; 2:3 ⁺	1:2 ⁺ ; 1:3 ⁺ ; 2:3 ⁺	1:2 ⁺ ; 1:3 ⁺ ; 2:3 ⁺
Haversian canals	C (1)	13	32.477±6.594	20.385±2.008	3.500±0.424	2.946±0.456
	E1 (2)	18	31.967±7.534	20.094±2.282	3.400±0.424	2.967±0.455
	E2 (3)	23	27.839±4.357	18.869±1.472	3.265±0.299	2.691±0.292
	Tukey test		NS	NS	NS	NS
Secondary osteons	C (1)	13	439.153±189.217	73.323±16.554	12.723±2.750	10.515±2.611
	E1 (2)	18	370.783±115.228	68.511±10.456	11.961±1.992	9.706±1.567
	E2 (3)	23	244.396±68.085	56.013±7.928	9.874±1.611	7.787±1.260
	Games-Howell and/or Tukey test		1:3 ⁺ ; 2:3 ⁺	1:3 ⁺ ; 2:3 ⁺	1:3 ⁺ ; 2:3 ⁺	1:3 ⁺ ; 2:3 ⁺

n: number of measurements; P<0.05 (+); NS: non-significant differences

Table 2 Results of trabecular bone microstructure in mice from the E1, E2 and C groups

	C group (1)	E1 group (2)	E2 group (3)	Tukey test
				
BV/TV (%)	0.08±0.02	0.13±0.01	0.18±0.04	1:2 ⁺ ; 1:3 ⁺
Tb. N. (1/mm)	3.97±0.45	4.76±0.09	4.89±0.48	1:2 ⁺ ; 1:3 ⁺
Tb. Th. (mm)	0.04±0.0014	0.04±0.0013	0.05±0.0019	1:3 ⁺ ; 2:3 ⁺
Tb. Sp. (mm)	0.25±0.03	0.20±0.01	0.19±0.02	1:2 ⁺ ; 1:3 ⁺
Bs. (mm²)	14.75±5.29	22.63±1.68	21.93±4.69	1:2 ⁺

n: number of measurements; P<0.05 (+); NS: non-significant differences; BV/TV – bone volume; Tb. N. – trabecular number; Tb. Th. – trabecular thickness; Tb. Sp. trabecular separation; Bs. – bone surface