

**Jablonski D.: Biogeografie a druhová rozmanitost obojživelníků a plazů
Balkánského poloostrova 2. (Živy 2017, 6: 314–318)**

Seznam použité a doporučené literatury:

- Ahmadzadeh, F., Flecks, M., Rödder, D., Böhme, W., Ilgaz, Ç., Harris, D.J., Engler, J.O., Üzüm, N., Carretero, M.A., 2013. Multiple dispersal out of Anatolia: biogeography and evolution of oriental green lizards. *Biol. J. Linn. Soc.* 110: 398-408.
- Andres, C., F. Franke, C. Bleidorn, D. Bernhard & M. Schlegel. 2014. Phylogenetics analysis of the *Lacerta agilis* subspecies complex. *Syst. Biodivers.* 12: 43-54.
- Arntzen, J.W., G.E. Themudo & B. Wielstra. 2007. The phylogeny of crested newts (*Triturus cristatus* superspecies): nuclear and mitochondrial genetic characters suggest a hard polytomy, in line with the paleogeography of the centre of origin. *Contrib. Zool.* 76: 261-278.
- Augé, M. 2005. Evolution des lézards du Paléogène en Europe. *Mém. Mus. Natl. Hist. Nat.* 192: 1-369.
- Avise, J.C. 2000. *Phylogeography: The History and Formation of Species*. Harvard University Press, Cambridge. MA. 447 pp.
- Babik, W., W. Branicki, J. Crnobrnja-Isailović, D. Cogălniceanu, I. Sas, K. Olgun, N.A. Poyarkov, M. Garcia-París & J.W. Arntzen. 2005. Phylogeography of two European newt species – discordance between mtDNA and morphology. *Mol. Ecol.* 14: 2475-2491.
- Babik, W., W. Branicki, M. Šandera, S. Litvinchuk, L.J. Borkin, J.T. Irwin & J. Rafiński. 2004. Mitochondrial phylogeography of the moor frog, *Rana arvalis*. *Mol. Ecol.* 13: 1469-1480.
- Baker, R.J. & R.D. Bradley. 2006. Speciation in mammals and the genetic species concept. *J. Mammal.* 87: 643-662.
- Baran, I. & Atatür, M.K. 1998. Turkish herpetofauna (Amphibians and Reptiles). Ankara (Republic of Turkey, Ministry of the Environment). 214 pp.
- Bellati, A., S. Carranza, J. Garcia-Porta, M. Fasola & R. Sindaco. 2015. Cryptic diversity within the *Anatololacerta* species complex (Squamata: Lacertidae) in the Anatolian Peninsula: Evidence from a multi-locus approach. *Mol. Phylogenet. Evol.* 82: 219-233.
- Böhme, M. 2003. The Miocene climatic optimum: evidence from the ectothermic vertebrates of Central Europe. *Palaeograp. Palaeoclim. Palaeoecol.* 195: 389-401.
- Böhme, M.U., U. Fritz, T. Kotenko, G. Džukić, K. Ljubisavljević, N. Tzankov & T.U. Berendonk. 2007. Phylogeography and cryptic variation within the *Lacerta viridis* complex (Lacertidae, Reptilia). *Zoolog. Scripta.* 36: 119-131.

- Carranza, S., E.N. Arnold & J.M. Pleguezuelos. 2006. Phylogeny, biogeography, and evolution of two Mediterranean snakes, *Malpolon monspessulanus* and *Hemorrhois hippocrepis* (Squamata, Colubridae), using mtDNA sequences. Mol. Phylogen. Evol. 40: 532-546.
- Cogălniceanu, D., L. Rozylowicz, P. Székely, C. Samoilă, F. Stanescu, M. Tudor, D. Székely & R. Iosif. 2013. Diversity and distribution of reptiles in Romania. ZooKeys. 341: 49-76.
- Cornetti, L., M. Menegon, G. Giovine, B. Heulin & C. Vernesi. 2014. Mitochondrial and Nuclear DNA Survey of *Zootoca vivipara* across the Eastern Italian Alps: Evolutionary Relationships, Historical Demography and Conservation Implications. PLOS One: e85912.
- Costello, M.J., R.M. May & N.E. Stork 2013. Can we name Earth's species before they go extinct? Science. 339: 413-416.
- Crottini, A., F. Andreone, J. Kosuch, L.J. Borkin, S.N. Litvinchuk, C. Eggert & M. Veith. 2007. Fossorial but widespread: the phylogeography of the common spadefoot toad (*Pelobates fuscus*), and the role of the Po Valley as a major source of genetic variability. Mol. Ecol. 16: 2734-2754.
- De Lattin, G. 1967. Grundriss der Zoogeographie. Jena: Gustav Fischer Verlag.
- Dermitzakis, D.M. & D.J. Papanikolaou. 1981. Paleogeography and geodynamics of the Aegean region during the Neogene. Annales Geol. Pays Helleniques. 30: 245-289.
- Dernegi, D. (Ed.). 2010. Mediterranean basin biodiversity hotspot. Critical Ecosystem. CEPF donor council, July 27, 2010.
- Douris, V., G.C. Rodakis, S. Giokas, M. Mylonas & R. Lecanidou. 1995. Mitochondrial DNA and morphological-differentiation of *Albinaria* populations (Gastropoda, Clausiliidae). J. Mollusc. Studies. 61: 65-78.
- Dufresnes, C., J. Wassef, K. Ghali, A. Brelsford, M. Stöck, P. Lymberakis, J. Crnobrnja-Isailović & N. Perrin. 2013. Conservation phylogeography: does historical diversity contribute to regional vulnerability in European tree frogs (*Hyla arborea*)? Mol. Ecol. 22: 5669-5684.
- Dufresnes, C., A. Brelsford, J. Crnobrnja-Isailović, N. Tzankov, P. Lymberakis & N. Perrin. 2015. Timeframe of speciation inferred from secondary contact zones in the European tree frog radiation (*Hyla arborea* group). BMC Evolutionary Biology. 15: 155.
- Džukić, G. & M.L. Kalezić. 2004. The Biodiversity of Amphibians and Reptiles in the Balkan Peninsula. Pp. 167-192. In: Griffiths H.I. (ed.). Balkan Biodiversity. Kluwer Academic Publishers. 357 pp.
- Eggert, C., D. Cogălniceanu, M. Veith, G. Džukić & P. Taberlet. 2006. The declining Spadefoot toad, *Pelobates fuscus* (Pelobatidae): paleo and recent environmental changes as a major influence on current population structure and status. Conserv. Genet. 7: 185-195.

- Espregueira Themudo, G., B. Wielstra & J.W. Arntzen. 2009. Multiple nuclear and mitochondrial genes resolve the branching order of a rapid radiation of crested newts (*Triturus*, Salamandridae). *Mol. Phylogenet. Evol.* 52: 321-328.
- Estes, R. 1983. *Encyclopædia of Paleoherpetology*. Part 10A. Sauria terrestria, Amphisbaenia. Fischer Verlag, Stuttgart.
- Fattorini, S. 2002. Biogeography of the tenebrionid beetles (Coleoptera, Tenebrionidae) on the Aegean Islands (Greece). *J. Biogeogr.* 29: 49-67.
- Ferchaud, A.-L., S. Ursenbacher, M. Cheylan, L. Luiselli, D. Jelić, B. Halpern, Á. Major, T. Kotenko, N. Keyan, R. Behrooz, J. Crnobrnja-Isailović, L. Tomović, I. Ghira, Y. Ioannidis, V. Arnal & C. Montgelard. 2012. Phylogeography of the *Vipera ursinii* complex (Viperidae): mitochondrial markers reveal an east–west disjunction in the Palaearctic region. *J. Biogeogr.* 39: 1836-1847.
- Fijarczyk, A., K. Nadachowska, S. Hofman, S.N. Litvinchuk, W. Babik, M. Stuglik, G. Gollmann, L. Choleva, D. Cogălniceanu, T. Vukov, G. Džukić & J.M. Szymura. 2011. Nuclear and mitochondrial phylogeography of the European fire-bellied toads *Bombina bombina* and *Bombina variegata* supports their independent histories. *Mol. Ecol.* 20: 3381-3398.
- Fritz, U., D. Guicking, H. Kami, M. Arakelyan, M. Auer, D. Ayaz, C.A. Fernández, A.G. Bakiev, A. Celani, G. Džukić, S. Fahd, P. Havaš, U. Joger, V.F. Khabibullin, L.F. Mazanaeva, P. Široký, S. Tripepi, A.V. Vélez, G.V. Antón & M. Wink. 2007. Mitochondrial phylogeography of European pond turtles (*Emys orbicularis*, *Emys trinacris*) – an update. *Amphibia-Reptilia*. 28: 418-426.
- Fritz, U., M. Auer, A. Bertolero, M. Cheylan, T. Fattizzo, A.K. Hundsdörfer, M.M. Sampayo, J.L. Pretus, P. Široký & M. Wink. 2006. A rangewide phylogeography of Hermann's tortoise, *Testudo hermanni* (Reptilia: Testudines: Testudinidae): implications for taxonomy. *Zoolog. Scripta*. 35: 531-543.
- Fritz, U., T. Fattizzo, D. Guicking, S. Tripepi, M.G. Pennisi, P. Lenk, U. Joger & M. Wink. 2005. A new cryptic species of pond turtle from southern Italy, the hottest spot in the range of the genus *Emys* (Reptilia, Testudines, Emydidae). *Zool. Scripta*. 34: 351-371.
- Gaston, K.J. & R. David. 1994. Hotspots across Europe. *Biodiversity Letters*. 2: 108-116.
- Gauthier, J., M. Kearney, J.A. Maisano, O. Rieppel & A.D.B. Behlke. 2012. Assembling the squamate tree of life: perspectives from the phenotype and the fossil record. *Bull. Peabody Mus. Nat. Hist.* 53: 3-308.
- Gómez, A. & D.H. Lunt. 2004. Chapterer IV: Refugia within refugia: Patterns of phylogeographic concordance in the Iberian Peninsula. Pp. 1-42. In: S. Weiss & N. Ferrand (ed.): *Phylogeography in Southern European Refugia. Evolutionary Perspectives on Origins and Conservation of European Biodiversity*. Dordrecht, The Netherlands: Springer Publishers.

- Grillitsch, H. & A. Cabela. 1990. Zum systematischen Status der Blindschleichen (Squamata: Anguidae) der Peloponnes und der südlichen Ionischen Inseln (Griechenland). *Herpetozoa*. 2: 131-153.
- Guicking, D., U. Joger & M. Wink. 2009. Cryptic diversity in a Eurasian water snake (*Natrix tessellata*, Serpentes: Colubridae): evidence from mitochondrial sequence data and nuclear ISSR-PCR fingerprinting. *Org. Divers. Evol.* 9: 201-214.
- Gvoždík, V., D. Canestrelli, M. García-París, J. Moravec, G. Nascetti, E. Recuero, J. Teixeira & P. Kotlík. 2015. Speciation history and widespread introgression in the European short-call tree frogs (*Hyla arborea* sensu lato, *H. intermedia* and *H. sarda*). *Mol. Phylogenet. Evol.* 83: 143-155.
- Gvoždík, V., D. Jandzík, B. Cordos, I. Rehák & P. Kotlík. 2012. A mitochondrial DNA phylogeny of the endangered vipers of the *Vipera ursinii* complex. *Mol. Phylogenet. Evol.* 62: 1019-1024.
- Gvoždík, V., D. Jandzík, P. Lymberakis, D. Jablonski & J. Moravec. 2010. Slow Worm, *Anguis fragilis* (Reptilia: Anguidae) as a species complex: Genetic structure reveals deep divergences. *Mol. Phylogenet. Evol.* 55: 460-472.
- Gvoždík, V., N. Benkovský, A. Crottini, A. Bellati, J. Moravec, A. Romano, R. Sacchi & D. Jandzik. 2013. An ancient lineage of slow worms, genus *Anguis* (Squamata: Anguidae), survived in the Italian Peninsula. *Mol. Phylogenet. Evol.* 69: 1077-1092.
- Harris, D.J., V. Batista, P. Lymberakis & M.A. Carretero. 2004. Complex estimates of evolutionary relationships in *Tarentola mauritanica* (Reptilia: Gekkonidae) derived from mitochondrial DNA sequences. *Mol. Phylogenet. Evol.* 30: 855-859.
- Hewitt, G.M. 1996. Some genetic consequences of ice ages, and their role in divergence and speciation. *Biol. J. Linn. Soc.* 58: 247-276.
- Hewitt, G.M. 1999. Post-glacial re-colonization of European biota. *Biol. J. Linn. Soc.* 68: 87-112.
- Hewitt, G.M. 2000. The genetic legacy og the Quarternary ice ages. *Nature*. 405: 907-913.
- Hewitt, G.M. 2011. Mediterranean Peninsulas: The Evolution of Hotspots. Pp. 123-147. In: F.E. Zachos & J.C. Habel (ed.): *Biodiversity Hotspots*. Berlin Heidelberg: Springer Publishers. 563 pp.
- Hofman, S., C. Spolsky, T. Uzzell, D. Cogălniceanu, W. Babik & J.M. Szymura. 2007. Phylogeography of the fire-bellied toads *Bombina*: independent Pleistocene histories inferred from mitochondrial genomes. *Mol. Ecol.* 16: 2301-2316.
- Hsü, K., L. Montadert, D. Bernuilli, M.B. Cita, A. Erickson, B.E. Garrison, R.B. Kidd, F. Mélierés, C. Müller & R. Wright. 1977. History of the Mediterranean salinity crisis. *Nature*. 267: 399-403.
- Husemann, M., T. Schmitt, F.E. Zachos, W. Ulrich & J.C. Habel. 2014. Palaearctic biogeography revisited: evidence for the existence of a North African refugium for Western Palaearctic biota. *J. Biogeogr.* 41: 81-94.

- Ivanov, M. 2007. Herpetological assemblages from the Pliocene to the middle Pleistocene in Central Europe: palaeoecological significance. *Geodiversitas*. 29: 297-320.
- Jablonski, D., D. Jandzik & V. Gvoždík. 2012. New records and zoogeographic classification of amphibians and reptiles from Bosnia and Herzegovina. *North-West. J. Zool.* 8: 324-337.
- Jablonski, D., D. Jandzik, P. Mikulíček, G. Džukić, K. Ljubisavljević, N. Tzankov, D. Jelić, E. Thanou, J. Moravec & V. Gvoždík. 2016. Contrasting evolutionary histories of the legless lizards slow worms (*Anguis*) shaped by the topography of the Balkan Peninsula. *BMC Evolutionary Biology*. 16:99.
- Jablonski, D., Najbar, B., Grochowalska, R., Gvoždík, V. & T. Strzała. 2017. Phylogeography and postglacial colonization of Central Europe by *Anguis fragilis* and *Anguis colchica*. *Amphibia-Reptilia*. 38: 562-569.
- Jesse, R., C.D. Shubart & S. Klaus. 2010. Identification of cryptic lineage within *Potamon fluviatile* (Herbst) (Crustacea: Brachyura: Potamidae). *Invertebr. Syst.* 24: 348-356.
- Joger, U., U. Fritz, D. Guicking, S.A. Kalyabina-Hauf, Z.T. Nagy & M. Wink. 2007. Phylogeography of western Palaearctic reptiles – Spatial and temporal speciation patterns. *Zool. Anz.* 246: 293-313.
- Joger, U., U. Fritz, D. Guicking, S.A. Kalyabina-Hauf, Z.T. Nagy & M. Wink. 2010. Relict populations and endemic clades in palearctic reptiles: evolutionary history and implications for conservation. Pp. 119-143. In: J.C. Habel & T. Assmann (ed.). *Relict Species: Phylogeography and Conservation Biology*. Springer-Verlag, Berlin Heidelberg. 468 pp.
- Kasapidis, P., A. Magoulas, M. Mylonas & E. Zouros. 2005. The phylogeography of the gecko *Cyrtopodion kotschy* (Reptilia: Gekkonidae) in the Aegean archipelago. *Mol. Phylogenetic Evol.* 35: 612-623.
- Kindler, C., W. Böhme, C. Corti, V. Gvoždík, D. Jablonski, D. Jandzik, M. Metallinou, P. Široký & U. Fritz. 2013. Mitochondrial phylogeography, contact zones and taxonomy of grass snakes (*Natrix natrix*, *N. megalocephala*). *Zoolog. Scripta*. 42: 458-472.
- Korniliou, P. 2017. Polytomies, signal and noise: revisiting the mitochondrial phylogeny and phylogeography of the Eurasian blindsnake species complex (Typhlopidae, Squamata). *Zoolog. Scripta*. doi:10.1111/zsc.12243.
- Korniliou, P., C. Ilgaz, Y. Kumluatas, P. Lymberakis, J. Moravec, R. Sindaco, N. Rastegar-Pouyani, M. Afroosheh, S. Giokas, S. Fraguedakis-Tsolis & B. Chondropoulos. 2012. Neogene climatic oscillations shape the biogeography and evolutionary history of the Eurasian blindsnake. *Mol. Phylogenetic Evol.* 62: 856-873.
- Korniliou, P., E. Thanou, P. Lymberakis, R. Sindaco, C. Liuzzi & S. Giokas. 2013. Mitochondrial phylogeography, intraspecific diversity and phenotypic convergence in the four-lined snake (Reptilia, Squamata). *Zoolog. Scripta* 43: 149-160.

- Kornilios, P., P. Kyriazi, N. Poulakakis, Y. Kumlutaş, C. Ilgaz, M. Mylonas & P. Lymberakis. 2010. Phylogeography of the ocellated skink *Chalcides ocellatus* (Squamata, Scincidae), with the use of mtDNA sequences: A hitch-hiker's guide to the Mediterranean. Mol. Phylogenetic Evol. 54: 445-456.
- Král, V. 1999. Fyzická geografie Evropy. Academia, Praha.
- Kryštufek, B., E.V. Bužan, W.F. Hutchinson & B. Hanfling. 2007. Phylogeography of the rare Balkan endemic Martino's vole, *Dinaromys bogdanovi*, reveals strong differentiation within the western Balkan Peninsula. Mol. Ecol. 16: 1221-1232.
- Kyriazi, P., N. Poulakakis, A. Parmakelis, P.A. Crochet, J. Moravec, N. Rastegar-Pouyani, C.S. Tsigenopoulos, A. Magoulas, M. Mylonas & P. Lymberakis. 2008. Mitochondrial DNA reveals the geological history of the snake-eyed lizards (*Ophisops elegans* and *O. occidentalis*) (Sauria: Lacertidae). Mol. Phylogenetic Evol. 49: 795-805.
- Kyriazi, P., P. Kornilios, Z.T. Nagy, N. Poulakakis, Y. Kumlutaş, Ç. Ilgaz, A. Avcı, B. Göçmen & P. Lymberakis. 2013. Comparative phylogeography reveals distinct colonization patterns of Cretan snakes. J. Biogeogr. 40: 1143-1155.
- Litvinchuk, S.N., A. Crottini, S. Federici, P. de Pous, D. Donaire, F. Andreone, M.L. Kalezić, G. Dzukić, G.A. Lada, L.J. Borkin & J.M. Rosanov. 2013. Phylogeographic patterns of genetic diversity in the common spadefoot toad, *Pelobates fuscus* (Anura: Pelobatidae), reveals evolutionary history, postglacial range expansion and secondary contact. Org. Divers. Evol. 13: 433-451.
- Ljubisavljević, K., O. Arribas, G. Džukić & S. Carranza. 2007. Genetic and morphological differentiation of Mosor rock lizard, *Dinarolacerta mosorensis* (Kolombatović, 1886), with the description of a new species from the Prokletije Mountain Massif (Montenegro) (Squamata: Lacertidae). Zootaxa. 1613: 1-22.
- Lymberakis, P. & N. Poulakakis. 2010. Three Continents Claiming an Archipelago: The Evolution of Aegean's Herpetofaunal Diversity. Diversity 2: 233-255.
- Lymberakis, P., N. Poulakakis, A. Kaliontzopoulou, E. Valakos & M. Mylonas. 2008. Two new species of *Podarcis* (Squamata; Lacertidae) from Greece. Syst. Biodivers. 6: 307-318.
- Macey, R.J., J.A. Schulte, A. Larson, B.S. Tuniyev, N. Orlov & T.J. Papenfuss. 1999. Molecular phylogenetics, tRNA evolution, and historical biogeography in Anguid lizards and related taxonomic families. Mol. Phylogenetic Evol. 12: 250-272.
- Mačát, Z., M. Starcová, J. Červenka, D. Jablonski & M. Šandera. 2014. A molecular assessment and first record of *Tarentola mauritanica* (Squamata: Phyllodactylidae) on Corfu, Greece. Salamandra. 50: 172-176.
- Mantziou, G., N. Poulakakis, P. Lymberakis, E. Valakos & M. Mylonas. 2004. The inter- and intraspecific status of Aegean *Mauremys rivulata* (Chelonia, Bataguridae) as inferred by mitochondrial DNA sequences. Herpetol. J. 14: 34-45.

- Marzahn, E., W. Mayer, U. Joger, Ç. Ilgaz, D. Jablonski, C. Kindler, Y. Kumlutaş, A. Nistri, N. Schneeweiß, M. Vamberger, A. Žagar & U. Fritz. 2016. Phylogeography of the *Lacerta viridis* complex: mitochondrial and nuclear markers provide taxonomic insights. *Journal of Zoological Systematics and Evolutionary Research*. 54: 85-105.
- Mayer, W., H. Grillitsch & A. Cabela. 1991. Proteinelektrophoretische Untersuchungen zur Systematik der südgriechischen Blindschleiche (Squamata: Anguidae). *Herpetozoa*. 4: 157-165.
- Médail, F. & K. Diadema. 2009. Glacial refugia influence plant diversity patterns in the Mediterranean Basin. *J. Biogeogr.* 36: 1333-1345.
- Messerli, B. 1967. Die eiszeitliche und die gegenwärtige Vergletscherung im Mittelmeeraum. *Geogr. Helv.* 22: 105–228
- Mizsei, E., D. Jablonski, S.A. Roussos, M. Dimaki, Y. Ioannidis, G. Nilson & Z.T. Nagy. 2017. Nuclear markers support the mitochondrial phylogeny of *Vipera ursinii-renardi* complex (Squamata: Viperidae) and species status for the Greek meadow viper. *Zootaxa*. 4227: 75-88
- Mizsei, E., D. Jablonski, Z. Végvári, S. Lengyel & M. Szabolcs. 2017b. Distribution and diversity of reptiles in Albania: a novel database from a Mediterranean hotspot. *Amphibia-Reptilia*. 38: 157-173.
- Młynarski, M. & Z. Szyndlar. 1989. Plazy i gady – Amphibia et Reptilia. Pp. 69-88. In: Kowalski, K. (ed.): Historia i ewolucje lądowej fauny Polski. *Folia Quaternaria*. 59-60.
- Mráz, P. & M. Ronikier. 2016. Biogeography of the Carpathians: evolutionary and spatial facets of biodiversity. *Biol. J. Linn. Soc.* 119: 528-559.
- Musilová, R., V. Zavadil, S. Marková & P. Kotlík. 2010. Relics of the Europe's warm past: Phylogeography of the Aesculapian snake. *Mol. Phylogen. Evol.* 57: 1245-1252.
- Nagy, Z.T., M. Bellaagh, M. Wink, A. Paunović & Z. Korsós. 2010. Phylogeography of the Caspian whipsnake in Europe with emphasis on the westernmost populations. *Amphibia-Reptilia*. 31: 455-461.
- Nagy, Z.T., U. Joger, D. Guicking & M. Wink. 2002. Phylogeography of the European Whip Snake, *Coluber (Hierophis) viridiflavus* as inferred from nucleotide sequences of the mitochondrial cytochrome *b* gene and ISSR genomic fingerprinting. *Biota*. 3: 109-118.
- Pabijan, M., P. Zieliński, K. Dudek, M. Chloupek, K. Sotiropoulos, M. Liana & W. Babik. 2015. The dissection of a Pleistocene refugium: phylogeography of the smooth newt, *Lissotriton vulgaris*, in the Balkans. *J. Biogeogr.* 42: 671-683.
- Parmakelis, A., I. Pfenninger, L. Spanos, G. Papagiannakis, C. Louis & M. Mylonas. 2005. Inference of a radiation in *Mastus* (Gastropoda, Pulmonata, Enidae) on the island of Crete. *Evolution*. 59: 991-1005.
- Parmakelis, A., I. Stathi, L. Spanos, C. Louis & M. Mylonas. 2006a. Phylogeography of *Iurus dufoureius*. *J. Biogeogr.* 33: 251-260.

- Parmakelis, A., I. Stathi, M. Chatzaki, S. Simaiakis, L. Spanos, C. Louis & M. Mylonas. 2006b. Evolution of *Mesobuthus gibbosus* (Brullé, 1832) (Scorpiones: Buthidae) in the northeastern Mediterranean region. Mol. Ecol. 15: 2883-2894.
- Podnar, M., B.B. Mađarić, W. Mayer. 2014. Non-concordant phylogeographical patterns of three widely codistributed endemic Western Balkans lacertid lizards (Reptilia, Lacertidae) shaped by specific habitat requirements and different responses to Pleistocene climatic oscillations. J. Zoolog. Syst. Evol. Res. 52: 119-129.
- Podnar, M., W. Mayer & N. Tvrtković. 2004. Mitochondrial phylogeography of the Dalmatian wall lizard, *Podarcis melisellensis* (Lacertidae). Org. Divers. Evol. 4: 307-317.
- Podnar, M., W. Mayer & N. Tvrtković. 2005. Phylogeography of the Italian wall lizard, *Podarcis sicula*, as revealed by mitochondrial DNA sequences. Mol. Ecol. 14: 575-588.
- Popov, S.V., F. Rögl, A.Y. Rozanov, F.F. Steiniger, I.G. Shcherba & M. Kovac. 2004. Lithological-paleogeographic maps of Paratethys: 10 maps Late Eocene to Pliocene. Courier Forschungsinstitut Senckenberg 250: 1-46.
- Popov, S.V., I.G. Shcherba, L.B. Ilyina, L.A. Nevesskaya, N.P. Paramonova, S.O. Khondkarian & I. Magyar. 2006. Late Miocene to Pliocene palaeogeography of the Paratethys and its relation to the Mediterranean. Palaeogeogr. Palaeoclimatol. Palaeoecol. 238: 91-106.
- Poulakakis, N., P. Kapli, P. Lymberakis, A. Trichas, K. Vardinoyiannis, S. Sfenthourakis & M. Mylonas. 2015. A review of phylogeographic analyses of animal taxa from the Aegean and surrounding regions. Journal of Zoological Systematics and Evolutionary Research. 53: 18-32.
- Poulakakis, N., P. Lymberakis, A. Antouniou, D. Chalkia Zouros, M. Mylonas & E. Valakos. 2003. Molecular phylogeny and biogeography of the wall-lizard *Podarcis erhardii* (Squamata: Lacertidae). Mol. Phylogen. Evol. 28: 38-46.
- Poulakakis, N., P. Lymberakis, C.S. Tsigenopoulos, A. Magoulas & M. Mylonas. 2005. Phylogenetic relationships and evolutionary history of snake-eyed skink *Ablepharus kitaibelii* (Sauria: Scincidae). Mol. Phylogen. Evol. 34: 245-256.
- Poulakakis, N., P. Lymberakis, E. Valakos, E. Zouros & M. Mylonas. 2005. Phylogenetic relationships and biogeography of *Podarcis* species from the Balkan Peninsula, by bayesian and maximum likelihood analyses of mitochondrial DNA sequences. Mol. Phylogen. Evol. 37: 845-857.
- Poulakakis, N., P. Lymberakis, E. Valakos, P. Pafilis, E. Zouros & M. Mylonas. 2005. Phylogeography of Balkan wall lizard (*Podarcis tauricus*) and its relatives inferred from, E. mitochondrial DNA sequences. Mol. Ecol. 14: 2433-2443.
- Poulakakis, N., V. Pakaki, M. Mylonas & P. Lymberakis. 2008. Molecular phylogeny of the Greek legless skink *Ophiomorus punctatissimus* (Squamata: Scincidae): The impact of the Mid-Aegean trench in its phylogeography. Mol. Phylogen. Evol. 47: 396-402.

- Provan, J. & K.D. Bennett. 2008. Phylogeographic insights into cryptic glacial refugia. *Trends Ecol. Evol.* 23: 564-571.
- Psonis, N., A. Antoniou, O. Kukushkin, D. Jablonski, B. Petrov, J. Crnobrnja-Isailović, K. Soutiopoulos, I. Gherghel, P. Lymberakis & N. Poulakakis. 2017. Hidden diversity in the *Podarcis tauricus* (Sauria, Lacertidae) species subgroup in the light of multilocus phylogeny and species delimitation. *Mol. Phylogenet. Evol.* 106: 6-17
- Rato, C., M.A.L. Zuffi, C. Corti, S. Fornasiero, A. Gentilli, E. Razzetti, S. Scali, M.A. Carretero & D.J. Harris. 2009. Phylogeography of the European Whip Snake, *Hierophis viridiflavus* (Colubridae), using mtDNA and nuclear DNA sequences. *Amphibia-Reptilia* 30: 283-289.
- Rato, C., S. Carranza & D.J. Harris. 2011. When selection deceives phylogeographic interpretation: The case of the Mediterranean house gecko, *Hemidactylus turcicus* (Linnaeus, 1758). *Mol. Phylogenet. Evol.* 58: 365-373.
- Rato, C., S. Carranza, A. Perea, M.A. Carretero & D.J. Harris. 2010. Conflicting patterns of nucleotide diversity between mtDNA and nDNA in the Moorish gecko, *Tarentola mauritanica*. *Mol. Phylogenet. Evol.* 56: 962-971.
- Recuero, E., D. Buckley, M. Garcia-Paris, J.W. Arntzen, D. Cogalniceanu & I. Martínez-Solano. 2014. Evolutionary history of *Ichthyosaura alpestris* (Caudata, Salamandridae) inferred from the combined analysis of nuclear and mitochondrial markers. *Mol. Phylogenet. Evol.* 81: 207-220.
- Recuero, E., D. Canestrelli, J. Vörös, K. Szabó, N.A. Poyarkov, J.W. Arntzen, J. Crnobrnja-Isailović, A.A. Kidov, D. Cogalniceanu, F.P. Caputo, G. Nescetti & I. Martínez-Solano. 2012. Multilocus species tree analyses resolve the radiation of the widespread *Bufo bufo* species group (Anura, Bufonidae). *Mol. Phylogenet. Evol.* 62: 71-86.
- Reed, J.M., B. Kryštufek & W.J. Eastwood. 2004. The physical geography of the Balkans and nomenclature of place names. Pp. 9-22. In: Griffiths H.I. (ed.). *Balkan Biodiversity*. Kluwer Academic Publishers.
- Reyjol, Y., B. Hugueny, D. Pont, P.G. Bianco, U. Beier, N. Caiola, F. Casals, I. Cowx, A. Economou, T. Ferreira, G. Haidvogl, R. Noble, A. de Sostoa, T. Vigneron & T. Virbickas. 2007. Patterns in species richness and endemism of European freshwater fish. *Global Ecol. Biogeogr.* 16: 65-75.
- Riberon, A., C. Miaud, K. Grossenbacher & P. Taberlet. 2001. Phylogeography of the Alpine salamander, *Salamandra atra* (Salamandridae) and the influence of the Pleistocene climatic oscillations on population divergence. *Mol. Ecol.* 10: 2555-2560.
- Sagonas, K., N. Poulakakis, P. Lymberakis, A. Parmakelis, P. Pafilis & E.D. Valakos. 2014. Molecular systematics and historical biogeography of the green lizards (*Lacerta*) in Greece: Insights from mitochondrial and nuclear DNA. *Mol. Phylogenet. Evol.* 76: 144-154.

- Salvi, D., D.J. Harris, A. Kaliontzopoulou, M.A. Carretero & C. Pinho. 2013. Persistence across Pleistocene ice ages in Mediterranean and extra-Mediterranean refugia: phylogeographic insights from the common wall lizard. *BMC Evolutionary Biology* 13: 147.
- Savić, I.R. 2008. Diversification of the Balkan fauna: its origin, historical development and present status. pp. 57-79. In: Makarov, Dimitrijević (Eds.), *Advances in Arachnology and Developmental Biology. Papers dedicated to Prof. Dr. Božidar Ćurčić*. Inst.Zool., Belgrade; BAS, Sofia, Fac. Life sci., Vienna; SASA, Belgrade & UNESCO MAB Committee, Serbia. ViennaBelgrade-Sofia, Monographs, 12.
- Sfenthourakis, S. 1996. A biogeographical analysis of terrestrial isopods (Isopoda, Oniscidea) from the central Aegean islands (Greece). *J. Biogeogr.* 23: 687-698.
- Schmitt, T. & Z. Varga. 2012. Extra-Mediterranean refugia: The rule and not the exception? *Front. Zool.* 9: 22.
- Schmitt, T. 2007. Molecular biogeography of Europe: Pleistocene cycles and postglacial trends. *Front. Zool.* 4: 11.
- Sillero, N., J. Campos, A. Bonardi, C. Corti, R. Creemers, P.-A. Crochet, J. Crnobrnja-Isailović, M. Denoël, F.G. Ficetola, J. Gonçalves, S. Kuzmin, P. Lymberakis, P. de Pous, A. Rodríguez, R. Sindaco, J. Speybroek, B. Toxopeus, D.R. Vieites & M. Vences. 2014. Updated distribution and biogeography of amphibians and reptiles of Europe. *Amphibia-Reptilia*. 35: 1-31.
- Simaiakis, S.M., A. Dimopoulou, A. Mitrakos, M. Mylonas & A. Parmakelis. 2012. The evolutionary history of the Mediterranean centipede *Scolopendra cingulata* (Latreille, 1829) (Chilopoda: Scolopendridae) across the Aegean archipelago. *Biol. J. Linn. Soc.* 105: 507-521.
- Simpson, G.G. 1951. The species concept. *Evolution*. 5: 285-298.
- Sindaco, R. & V.K. Jeremčenko. 2008. The reptiles of the Western Palearctic. Edizioni Belvedere, Latina (Italy), 579 pp.
- Skourtanioti, E. P. Kapli, Ç. Ilgaz, Y. Kumlutas, A. Avci, F. Ahmadzadeh, J. Crnobrnja-Isailović, I. Gherghel, P. Lymberakis & N. Poulakakis. 2016. A reinvestigation of phylogeny and divergence times of the *Ablepharus kitaibelii* species complex (Sauria, Scincidae) based on mtDNA and nuDNA genes. *Mol. Phylogenetic Evol.* 103: 199-214.
- Sommer, R.S., A. Persson, N. Wieseke & U. Fritz. 2007. Holocene recolonization and extinction of the pond turtle, *Emys orbicularis* (L., 1758), in Europe. *Quaternary Sci. Rev.* 26: 3099-3107.
- Sommer, R.S., C. Lindqvist, A. Persson, H. Bringsøe, A.G.J. Rhodin, N. Schneeweiss, P. Široký, L. Bachmann & U. Fritz. 2009. Unexpected early extinction of the European pond turtle (*Emys orbicularis*) in Sweden and climatic impact on its Holocene range. *Mol. Ecol.* 18: 1252-1262.

- Sotiropoulos, K., K. Eleftherakos, G. Džukić, M.L. Kalezić, A. Legakis & R.M. Polymeni. 2007. Phylogeny and biogeography of the alpine newt *Mesotriton alpestris* (Salamandridae, Caudata), inferred from mtDNA sequences. Mol. Phylogenetic Evol. 45: 211-226.
- Speybroeck, J., W. Beukema, B. Bok, J. Van Der Voort & I. Velikov. 2016. Field guide to the amphibians and reptiles of Britain and Europe. London, Bloomsbury.
- Stewart, J.R., A.M. Lister, I. Barnes & L. Dalén. 2010. Refugia revisited: individualistic responses of species in space and time. Proc. R. Soc. B. 277: 661-671.
- Stöck, M., C. Dufresnes, S.N. Litvinchuk, P. Lymberakis, S. Biollay, M. Berroneau, A. Borzée, K. Ghali, M. Ogielska & N. Perrin. 2012. Cryptic diversity among Western Palearctic tree frogs: Postglacial range expansion, range limits, and secondary contacts of three European tree frog lineages (*Hyla arborea* group). Mol. Phylogenetic Evol. 65: 1-9.
- Stöck, M., S. Dubey, C. Klütsch, S.N. Litvinchuk, U. Scheidt & N. Perrin. 2008. Mitochondrial and nuclear phylogeny of circum-Mediterranean tree frogs from the *Hyla arborea* group. Mol. Phylogenetic Evol. 49: 1019-1024.
- Stojanov, A., N. Tzankov & B. Naumov. 2011. Die Amphibien und Reptilien Bulgariens. Edition Chimaira, Frankfurt am Main. 582 pp.
- Stümpel, N., M. Rajabizadeh, A. Avci, W. Wüster & U. Joger. 2016. Phylogeny and diversification of mountain vipers (*Montivipera*, Nilson et al., 2001) triggered by multiple Plio-Pleistocene refugia and high-mountain topography in the Near and Middle East. Mol. Phylogenetic Evol. 101: 336-351.
- Surget-Groba, Y., B. Heulin, C.-P. Guillaume, M. Puky, D. Semenov, V. Orlova, L. Kupriyanova, I. Ghira & B. Šmajda. 2006. Multiple origins of viviparity, or reversal from viviparity to oviparity? The European common lizard (*Zootoca vivipara*, Lacertidae) and the evolution of parity. Biol. J. Linn. Soc. 87: 1-11.
- Szabó, K. & J. Vörös. 2014. Distribution and hybridization of *Anguis fragilis* and *A. colchica* in Hungary. Amphibia-Reptilia 35: 135-140.
- Taberlet, P., L. Fumagalli, A.G. Wust-Saucy & J.F. Cosson. 1998. Comparative phylogeography and postglacial colonization routes in Europe. Mol. Ecol. 7: 453-464.
- Taskavak, E., M.J. Reimann, & W. Polder. 1999. First record of the Nile Soft-Shelled Turtle, *Trionyx triunguis*, from Kos Island, Greece, with comments on its occurrence in the eastern Mediterranean. Chelonian Conserv. Bi. 3: 510-511.
- Thanou, E., S. Giokas & P. Korniliou. 2014. Phylogeography and genetic structure of the slow worms *Anguis cephalonica* and *Anguis graeca* (Squamata: Anguidae) from the southern Balkan Peninsula. Amphibia-Reptilia. 35: 263-269.
- Tomović, L., R. Ajtić, K. Ljubisavljević, A. Urošević, D. Jović, I. Krizmanić, N. Labus, S. Đorđević, M.L. Kalezić, T. Vukov & G. Džukić. 2014. Reptiles in Serbia – distribution and diversity patterns. Bull. Nat. Hist. Mus. 7: 129-158.

- Ursenbacher, S., M. Carlsson, V. Helfer, H. Tegelström & L. Fumagalli. 2006. Phylogeography and Pleistocene refugia of the adder (*Vipera berus*) as inferred from mitochondrial DNA sequences data. *Mol. Ecol.* 15: 3425-3437.
- Ursenbacher, S., S. Schweiger, L. Tomović, J. Crnobrnja-Isailović, L. Fumagalli & W. Mayer. 2008. Molecular phylogeography of the nose-horned viper (*Vipera ammodytes*, Linnaeus (1758)): Evidence for high genetic diversity and multiple refugia in the Balkan peninsula. *Mol. Phylogen. Evol.* 46: 1116-1128.
- Valakos, E.D., P. Pafilis, K. Sotiropoulos, P. Lymberakis, P. Maragou & J. Foufopoulos. 2008. The Amphibians and Reptiles of Greece. Chimaira, Frankfurt am Main. 463 pp.
- Vamberger, M., H. Stuckas, D. Ayaz, P. Lymberakis, P. Široký & U. Fritz. 2014. Massive transoceanic gene flow in a freshwater turtle (Testudines: Geoemydidae: *Mauremys rivulata*). *Zool. Scripta.* 43: 313-322.
- Veith, M., J. Kosuch & M. Vences. 2003. Climatic oscillations triggered post-Messinian speciation of Western Palearctic brown frogs (Amphibia, Ranidae). *Mol. Phylogen. Evol.* 26: 310-327.
- Vences, M., J.S. Hauswaldt, S. Steinfartz, O. Rupp, A. Goesmann, S. Künzel, P. Orozco-terWengel, D.R. Vieites, S. Nieto-Roman, S. Hass, C. Laugsch, M. Gehara, S. Bruchmann, M. Pabijan, A.K. Ludewig, D. Rudert, C. Angelini, L.J. Borkin, P.A. Crochet, A. Crottini, A. Dubois, G.F. Ficetola, P. Galán, P. Geniez, M. Hachtel, O. Jovanovic, S.N. Litvinchuk, P. Lymberakis, A. Ohler & N.A. Smirnov. 2013. Radically different phylogeographies and patterns of genetic variation in two European brown frogs, genus *Rana*. *Mol. Phylogen. Evol.* 68: 657-670.
- Vidal, N., J. Marin, J. Sassi, F.U. Battistuzzi, S. Donnellan, A.J. Fitch, B.G. Fry, F.J. Vonk, R.C. Rodriguez de la Vega, A. Couloux & S.B. Hedges. 2012. Molecular evidence for an Asian origin of monitor lizards followed by Tertiary dispersals to Africa and Australia. *Biol. Lett.* 8: 853-855.
- Vörös, J., P. Mikulíček, Á. Major, E. Recuero & J.W. Arntzen. Phylogeographic analysis reveals northerly refugia for the riverine amphibian *Triturus dobrogicus* (Caudata: Salamandridae). *Biol. J. Linn. Soc.* 119: 974-991.
- Wielstra, B., S.N. Litvinchuk, B. Naumov, N. Tzankov & J.W. Arntzen. 2013. A revised taxonomy of crested newts in the *Triturus karelinii* group (Amphibia: Caudata: Salamandridae), with the description of a new species. *Zootaxa.* 3682: 441-453.
- Wielstra, B., W. Babik & J.W. Arntzen. 2015. The crested newt *Triturus cristatus* recolonized temperate Eurasia from an extra-Mediterranean glacial refugium. *Biol. J. Linn. Soc.* 114: 574-587.
- Wiley, E.O. 1978. Evolutionary species concept reconsidered. *Systematic Zoology.* 27: 17-26.