

## ON SOLUTIONS OF THE DIFFERENCE EQUATION

$$x_{n+1} = x_{n-3}/(-1 + x_n x_{n-1} x_{n-2} x_{n-3})$$

CENGİZ CİNAR, RAMAZAN KARATAS, İBRAHİM YALÇINKAYA, Konya

(Received March 13, 2006)

*Abstract.* We study the solutions and attractivity of the difference equation  $x_{n+1} = x_{n-3}/(-1 + x_n x_{n-1} x_{n-2} x_{n-3})$  for  $n = 0, 1, 2, \dots$  where  $x_{-3}, x_{-2}, x_{-1}$  and  $x_0$  are real numbers such that  $x_0 x_{-1} x_{-2} x_{-3} \neq 1$ .

*Keywords:* difference equation, recursive sequence, solutions, equilibrium point

*MSC 2000:* 39A11

## 1. INTRODUCTION

A lot of work has been done concerning the attractivity and solutions of the rational difference equations, for example in [1]–[9]. In [3] Cinar studied the positive solutions of the difference equation  $x_{n+1} = x_{n-1}/(1 + x_n x_{n-1})$  for  $n = 0, 1, 2, \dots$  and proved by induction the formula

$$x_n = \begin{cases} x_{-1} \frac{\prod_{i=0}^{[(n+1)/2]-1} (2x_{-1} x_0 i + 1)}{\prod_{i=0}^{[(n+1)/2]-1} ((2i+1)x_{-1} x_0 + 1)} & \text{for } n \text{ odd,} \\ x_0 \frac{\prod_{i=1}^{n/2} ((2i-1)x_{-1} x_0 + 1)}{\prod_{i=1}^{n/2} (2ix_{-1} x_0 + 1)} & \text{for } n \text{ is even.} \end{cases}$$

In [6] Stević studied the stability properties of the solutions of Cinar's equation. Also in [7] Stević investigated the solutions of the difference equation  $x_{n+1} =$

$Bx_{n-1}/B + x_n$  and gave the formulas

$$x_{2n} = x_0 \left( 1 - x_1 \sum_{j=1}^n \prod_{i=1}^{2j-1} \frac{1}{1+x_i} \right),$$

$$x_{2n+1} = x_{-1} \left( 1 - \frac{x_0}{1+x_0} \sum_{j=0}^n \prod_{i=1}^{2j} \frac{1}{1+x_i} \right).$$

Moreover, in [1] Aloqeili generalized the results from [3], [6] to the  $k$ th order case and investigated the solutions, stability character and semicycle behavior of the difference equation  $x_{n+1} = x_{n-k}/(A + x_{n-k}x_n)$  where  $x_{-k}, \dots, x_0 > 0$  and  $A > 0$ ,  $k$  being any positive integer.

Our aim in this paper is to investigate the solutions of the difference equation

$$(1.1) \quad x_{n+1} = \frac{x_{n-3}}{-1 + x_n x_{n-1} x_{n-2} x_{n-3}} \quad \text{for } n = 0, 1, 2, \dots$$

where  $x_{-3}, x_{-2}, x_{-1}$  and  $x_0$  are real numbers such that  $x_0 x_{-1} x_{-2} x_{-3} \neq 1$ .

First, we give two definitions which will be useful in our investigation of the behavior of solutions of Eq. (1.1).

**Definition 1.** Let  $I$  be an interval of real numbers and let  $f: I^4 \rightarrow I$  be a continuously differentiable function. Then for every  $x_{-i} \in I$ ,  $i = 0, 1, 2, 3$ , the difference equation  $x_{n+1} = f(x_n, x_{n-1}, x_{n-2}, x_{n-3})$ ,  $n = 0, 1, 2, \dots$ , has a unique solution  $\{x_n\}_{n=-3}^{\infty}$ .

**Definition 2.** The equilibrium point  $\bar{x}$  of the equation  $x_{n+1} = f(x_n, x_{n-1}, \dots, x_{n-k})$ ,  $n = 0, 1, 2, \dots$ , is the point that satisfies the condition  $\bar{x} = f(\bar{x}, \dots, \bar{x})$ .

## 2. MAIN RESULTS

**Theorem 1.** Assume that  $x_0 x_{-1} x_{-2} x_{-3} \neq 1$  and let  $\{x_n\}_{n=-3}^{\infty}$  be a solution of Eq. (1.1). Then for  $n = 0, 1, 2, \dots$  all solutions of Eq. (1.1) are of the form

$$(2.1) \quad x_{4n+1} = x_{-3} / (-1 + x_0 x_{-1} x_{-2} x_{-3})^{n+1},$$

$$(2.2) \quad x_{4n+2} = x_{-2} / (-1 + x_0 x_{-1} x_{-2} x_{-3})^{n+1},$$

$$(2.3) \quad x_{4n+3} = x_{-1} / (-1 + x_0 x_{-1} x_{-2} x_{-3})^{n+1},$$

$$(2.4) \quad x_{4n+4} = x_0 / (-1 + x_0 x_{-1} x_{-2} x_{-3})^{n+1}.$$

**Proof.**  $x_1, x_2, x_3$  and  $x_4$  are clear from Eq. (1.1). Also, for  $n = 1$  the result holds. Now suppose that  $n > 1$  and our assumption holds for  $(n-1)$ . We shall show

that the result holds for  $n$ . From our assumption for  $(n - 1)$  we have

$$\begin{aligned}x_{4n-3} &= x_{-3}/(-1 + x_0x_{-1}x_{-2}x_{-3})^n, \\x_{4n-2} &= x_{-2}(-1 + x_0x_{-1}x_{-2}x_{-3})^n, \\x_{4n-1} &= x_{-1}/(-1 + x_0x_{-1}x_{-2}x_{-3})^n, \\x_{4n} &= x_0(-1 + x_0x_{-1}x_{-2}x_{-3})^n.\end{aligned}$$

Then, from Eq. (1.1) and the above equality, we have

$$\begin{aligned}x_{4n+1} &= x_{4n-3}/(-1 + x_{4n}x_{4n-1}x_{4n-2}x_{4n-3}) \\&= \frac{x_{-3}/(-1 + x_0x_{-1}x_{-2}x_{-3})^n}{-1 + x_0x_{-1}x_{-2}x_{-3}} = \frac{x_{-3}}{(-1 + x_0x_{-1}x_{-2}x_{-3})^{n+1}}.\end{aligned}$$

That is,

$$x_{4n+1} = \frac{x_{-3}}{(-1 + x_0x_{-1}x_{-2}x_{-3})^{n+1}}.$$

Also,

$$\begin{aligned}x_{4n+2} &= \frac{x_{4n-2}}{-1 + x_{4n+1}x_{4n}x_{4n-1}x_{4n-2}} \\&= \frac{x_{-2}(-1 + x_0x_{-1}x_{-2}x_{-3})^n}{-1 + x_0x_{-1}x_{-2}x_{-3}/(-1 + x_0x_{-1}x_{-2}x_{-3})} \\&= x_{-2}(-1 + x_0x_{-1}x_{-2}x_{-3})^{n+1}.\end{aligned}$$

Hence, we have

$$x_{4n+2} = x_{-2}(-1 + x_0x_{-1}x_{-2}x_{-3})^{n+1}.$$

Similarly,

$$\begin{aligned}x_{4n+3} &= \frac{x_{4n-1}}{-1 + x_{4n+2}x_{4n+1}x_{4n}x_{4n-1}} = \frac{x_{-1}/(-1 + x_0x_{-1}x_{-2}x_{-3})^n}{-1 + x_0x_{-1}x_{-2}x_{-3}} \\&= \frac{x_{-1}}{(-1 + x_0x_{-1}x_{-2}x_{-3})^{n+1}}.\end{aligned}$$

Consequently, we have

$$x_{4n+3} = \frac{x_{-1}}{(-1 + x_0x_{-1}x_{-2}x_{-3})^{n+1}}.$$

Now we prove the last formula. Since

$$\begin{aligned}x_{4n+4} &= \frac{x_{4n}}{-1 + x_{4n+3}x_{4n+2}x_{4n+1}x_{4n}} \\&= \frac{x_0(-1 + x_0x_{-1}x_{-2}x_{-3})^n}{-1 + x_0x_{-1}x_{-2}x_{-3}/(-1 + x_0x_{-1}x_{-2}x_{-3})} \\&= x_0(-1 + x_0x_{-1}x_{-2}x_{-3})^{n+1},\end{aligned}$$

we have

$$x_{4n+4} = x_0 (-1 + x_0 x_{-1} x_{-2} x_{-3})^{n+1}.$$

Thus, we have proved (2.1), (2.2), (2.3) and (2.4).  $\square$

**Theorem 2.** Eq. (1.1) has three equilibrium points which are  $0$ ,  $\sqrt[4]{2}$  and  $-\sqrt[4]{2}$ .

*Proof.* For the equilibrium points of Eq. (1.1) we write

$$\bar{x} = \bar{x}/(-1 + \bar{x}\bar{x}\bar{x}).$$

Then we have

$$\bar{x}^5 - 2\bar{x} = 0.$$

Thus, the equilibrium points of Eq. (1.1) are  $0$ ,  $\sqrt[4]{2}$  and  $-\sqrt[4]{2}$ .  $\square$

**Corollary 1.** Let  $\{x_n\}$  be a solution of Eq. (1.1). Assume that  $x_{-3}, x_{-2}, x_{-1}, x_0 > 0$  and  $x_{-3}x_{-2}x_{-1}x_0 > 1$ . Then all solutions of Eq. (1.1) are positive.

*Proof.* This is clear from Eqs. (2.1), (2.2), (2.3) and (2.4).  $\square$

**Corollary 2.** Let  $\{x_n\}$  be a solution of Eq. (1.1). Assume that  $x_{-3}, x_{-2}, x_{-1}, x_0 < 0$  and  $x_{-3}x_{-2}x_{-1}x_0 > 1$ . Then all solutions of Eq. (1.1) are negative.

*Proof.* This is clear from Eqs. (2.1), (2.2), (2.3) and (2.4).  $\square$

**Corollary 3.** Let  $\{x_n\}$  be a solution of Eq. (1.1). Assume that  $x_{-3}, x_{-2}, x_{-1}, x_0 > 0$  and  $x_{-3}x_{-2}x_{-1}x_0 > 2$ . Then

$$\lim_{n \rightarrow \infty} x_{4n+1} = 0, \quad \lim_{n \rightarrow \infty} x_{4n+2} = \infty, \quad \lim_{n \rightarrow \infty} x_{4n+3} = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} x_{4n+4} = \infty.$$

*Proof.* Let  $x_{-3}, x_{-2}, x_{-1}, x_0 > 0$  and  $x_{-3}x_{-2}x_{-1}x_0 > 2$ .

Then  $x_{-3}x_{-2}x_{-1}x_0 - 1 > 1$  and Eq. (2.1), (2.2), (2.3) and (2.4) imply

$$\begin{aligned} \lim_{n \rightarrow \infty} x_{4n+1} &= \lim_{n \rightarrow \infty} \frac{x_{-3}}{(-1 + x_0 x_{-1} x_{-2} x_{-3})^{n+1}} = 0, \\ \lim_{n \rightarrow \infty} x_{4n+2} &= \lim_{n \rightarrow \infty} x_{-2} (-1 + x_0 x_{-1} x_{-2} x_{-3})^{n+1} = \infty, \\ \lim_{n \rightarrow \infty} x_{4n+3} &= \lim_{n \rightarrow \infty} \frac{x_{-1}}{(-1 + x_0 x_{-1} x_{-2} x_{-3})^{n+1}} = 0, \\ \lim_{n \rightarrow \infty} x_{4n+4} &= \lim_{n \rightarrow \infty} x_0 (-1 + x_0 x_{-1} x_{-2} x_{-3})^{n+1} = \infty. \end{aligned}$$

$\square$

**Corollary 4.** Let  $\{x_n\}$  be a solution of Eq. (1.1). Assume that  $x_{-3}, x_{-2}, x_{-1}, x_0 < 0$  and  $x_{-3}x_{-2}x_{-1}x_0 > 2$ . Then

$$\lim_{n \rightarrow \infty} x_{4n+1} = 0, \lim_{n \rightarrow \infty} x_{4n+2} = -\infty, \lim_{n \rightarrow \infty} x_{4n+3} = 0 \text{ and } \lim_{n \rightarrow \infty} x_{4n+4} = -\infty.$$

The proof is similar to that of Corollary 3. Thus it is omitted.

Now, we give the following result about the product of solutions of Eq. (1.1).

**Corollary 5.**  $\prod_{n=0}^s x_{4n+1}x_{4n+2}x_{4n+3}x_{4n+4} = (x_0x_{-1}x_{-2}x_{-3})^{s+1}$  where  $s \in \mathbb{Z}^+$ .

*Proof.* From Eqs. (2.1), (2.2), (2.3) and (2.4) we obtain

$$\begin{aligned} x_{4n+1}x_{4n+2}x_{4n+3}x_{4n+4} &= \frac{x_{-3}}{(-1 + x_0x_{-1}x_{-2}x_{-3})^{n+1}} x_{-2} (-1 + x_0x_{-1}x_{-2}x_{-3})^{n+1} \\ &\times \frac{x_{-1}}{(-1 + x_0x_{-1}x_{-2}x_{-3})^{n+1}} x_0 (-1 + x_0x_{-1}x_{-2}x_{-3})^{n+1} = x_0x_{-1}x_{-2}x_{-3} \end{aligned}$$

and the above equality yields

$$\prod_{n=0}^s x_{4n+1}x_{4n+2}x_{4n+3}x_{4n+4} = (x_0x_{-1}x_{-2}x_{-3})^{s+1}.$$

Thus, the proof is complete. □

#### References

- [1] *Aloqeili M.*: Dynamics of a  $k$ th order rational difference equation. Appl. Math. Comput. (In press.)
- [2] *Camouzis E., Ladas G., Rodrigues I. W., Northshield S.*: The rational recursive sequence  $x_{n+1} = bx_n^2/1 + x_{n-1}^2$ . Comput. Math. Appl. 28 (1994), 37–43. zbl
- [3] *Cinar C.*: On the positive solutions of the difference equation  $x_{n+1} = x_{n-1}/(1 + x_n \times x_{n-1})$ . Appl. Math. Comput. 150 (2004), 21–24. zbl
- [4] *Cinar C.*: On the positive solutions of the difference equation  $x_{n+1} = ax_{n-1}/(1 + bx_n \times x_{n-1})$ . Appl. Math. Comput. 156 (2004), 587–590. zbl
- [5] *Cinar C.*: On the difference equation  $x_{n+1} = x_{n-1}/(-1 + x_nx_{n-1})$ . Appl. Math. Comput. 158 (2004), 813–816. zbl
- [6] *Stevic S.*: More on a rational recurrence relation  $x_{n+1} = x_{n-1}/(1 + x_{n-1}x_n)$ . Appl. Math. E-Notes 4 (2004), 80–84. zbl
- [7] *Stevic S.*: On the recursive sequence  $x_{n+1} = x_{n-1}/g(x_n)$ . Taiwanese J. Math. 6 (2002), 405–414. zbl
- [8] *Stevic S.*: On the recursive sequence  $x_{n+1} = \alpha + x_{n-1}^p/x_n^p$ . J. Appl. Math. Comput. 18 (2005), 229–234. zbl
- [9] *Yang X., Su W., Chen B., Megson G., Evans D.*: On the recursive sequences  $x_{n+1} = ax_{n-1} + bx_{n-2}/(c + dx_{n-1}x_{n-2})$ . Appl. Math. Comput. 162 (2005), 1485–1497. zbl

*Authors' addresses:* Cengiz Cinar, Ramazan Karatas, Ibrahim Yalçınkaya, Selcuk University, Education Faculty, Mathematics Department, 42099, Meram Yeni Yol, Konya, Türkiye, e-mail: ccinar25@yahoo.com, rckaratas@yahoo.com, iyalcinkaya1708@yahoo.com.