



INSTITUTE OF MATHEMATICS

THE CZECH ACADEMY OF SCIENCES

**On grand and small Lebesgue
and Sobolev spaces and some
applications to PDEs**

Alberto Fiorenza

Maria Rosaria Formica

Amiran Gogatishvili

Preprint No. 95-2017

PRAHA 2017

ON GRAND AND SMALL LEBESGUE AND SOBOLEV SPACES AND SOME APPLICATIONS TO PDES

ALBERTO FIORENZA, MARIA ROSARIA FORMICA AND AMIRAN GOGATISHVILI

Abstract. This paper is essentially a survey on grand and small Lebesgue spaces, which are rearrangement-invariant Banach function spaces of interest not only from the point of view of Function Spaces theory, but also from the point of view of their applications: the corresponding Sobolev spaces are of interest, for instance, in the theory of PDEs. We discuss results of existence, uniqueness and regularity of certain Dirichlet problems, where the knowledge of these spaces plays a central role. The novelty of this paper relies in an unified treatment containing a number of equivalent quasinorms, all written making explicit the dependence of $|\Omega|$, in the discussion of the sharpness of Hölder's inequality, and in the connection of the results in PDEs with some existing literature.

This paper covers a part of subjects presented in the Conference Internationale en Analyse - Laboratoire de Mathématiques et Applications, Université de Poitiers, 29 - 30 Mars 2017, en hommage au Professeur Robert JANIN

1. Grand and small Lebesgue spaces: a short overview

1.1. The original motivation

Let $\Omega \subset \mathbb{R}^n$ be a bounded domain and $f : \Omega \rightarrow \mathbb{R}^n$, $f = (f^1, \dots, f^n)$ be a mapping of Sobolev class $W_{loc}^{1,n}(\Omega, \mathbb{R}^n)$. Let us denote by $Df(x) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ the differential and by $J(x, f) = \det Df(x)$ the Jacobian of f . After the elementary remark that by Hölder's inequality the Jacobian $J(x, f)$ is in $L_{loc}^1(\Omega)$, the first fundamental result on the integrability of the Jacobian was due to Müller ([135]):

$$f \in W^{1,n}(\Omega, \mathbb{R}^n), J(x, f) \geq 0 \text{ a.e.} \quad \Rightarrow \quad J(x, f) \in L \log L_{loc}(\Omega).$$

Mathematics subject classification (2010): 46E30, 35J65.

Keywords and phrases: grand Lebesgue spaces; small Lebesgue spaces; interpolation.

The second author has been partially supported by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM) and by Università degli Studi di Napoli Parthenope through the project "sostegno alla Ricerca individuale" (triennio 2015-2017).

The third author has been partially supported by grant no: 217282 of the Shota Rustaveli National Science Foundation (SRNSF), Operators of Fourier analysis in some classical and new function spaces and by the grant P201/13/14743S of the Grant agency of the Czech Republic and RVO: 67985840.

The authors thank professor Jean Michel Rakotoson and the University of Poitiers for the kind invitation and the warm hospitality.

As a dual result, Iwaniec and Sbordone in [96] introduced the grand Lebesgue space L^n , and proved that under the assumption $|Df| \in L^n$ the Jacobian of f is locally integrable. Such assumption, weaker than $|Df| \in L^n$, is shown to be minimal:

$$|Df| \in L^n(\Omega), J(x, f) \geq 0 \text{ a.e.} \quad \Rightarrow \quad J(x, f) \in L^1_{loc}(\Omega).$$

The space $L^n(\Omega)$ has been defined through

$$u \in L^n(\Omega) \quad \Leftrightarrow \quad \|u\|_{L^n(\Omega)} = \sup_{0 < \varepsilon < n-1} \left(\varepsilon \int_{\Omega} |u(x)|^{n-\varepsilon} dx \right)^{\frac{1}{n-\varepsilon}} < \infty$$

where the symbol \int_{Ω} denotes the integral mean: $\int_{\Omega} = \frac{1}{|\Omega|} \int_{\Omega}$.

Of course, from the point of view of function spaces, it is natural to release the exponent n from the dimension, and therefore to consider, in its place, a generic exponent p , $1 < p < \infty$.

1.2. Background

Let $\Omega \subset \mathbb{R}^n$, $n \geq 1$, be (Lebesgue) measurable, $0 < |\Omega| < \infty$, and let $f : \Omega \rightarrow \mathbb{R}$ be measurable. Let us recall the familiar notation for the standard, classical Lebesgue spaces (see e.g. the recent monograph [29] on this topic): for $1 \leq p < \infty$,

$$f \in L^p(\Omega) \quad \Leftrightarrow \quad \|f\|_{L^p(\Omega)} = \left(\int_{\Omega} |f(x)|^p dx \right)^{\frac{1}{p}} < \infty$$

while, for $p = \infty$,

$$f \in L^\infty(\Omega) \quad \Leftrightarrow \quad \|f\|_{L^\infty(\Omega)} = \operatorname{ess\,sup}_{\Omega} |f| < \infty.$$

A generalization, and at the same time an extraordinary refinement, of the theory of Lebesgue spaces is given by the theory of Orlicz spaces. It is a theory fruitful in applications because the role of the composition by a power is played by a more generic function (see e.g. [2]; for a more general notion of Orlicz space see also, for instance, the paper [97] written for the Orlicz centenary volume; see also the references therein) $\Phi : [0, \infty[\rightarrow [0, \infty[$ continuous, strictly increasing, convex, $\Phi(0) = 0$, $\Phi'(0) = 0$, $\Phi'(\infty) = \infty$. The norm of the corresponding space $L^\Phi(\Omega)$ is given by

$$f \in L^\Phi(\Omega) \quad \Leftrightarrow \quad \|f\|_{L^\Phi(\Omega)} = \inf \left\{ \lambda > 0 : \int_{\Omega} \Phi \left(\frac{|f(x)|}{\lambda} \right) dx \leq 1 \right\} < \infty$$

where, by convention, $\inf \emptyset = \infty$. The notation changes for specific choices of the function Φ : even if the norm of a function in $L^\Phi(\Omega)$ depends heavily on Φ , when $0 < |\Omega| < \infty$ the symbol $L^\Phi(\Omega)$, interpreted as set of functions, is not affected when Φ is replaced by an equivalent function at infinity (Φ_1, Φ_2 are equivalent at infinity if for some $c_1, c_2, t_0 > 0$ it is $c_1 \Phi_1(t) \leq \Phi_2 \leq c_2 \Phi_2(t)$, $t > t_0$). For instance, here there are examples of growths and symbols for the corresponding sets of functions:

$$\begin{aligned}
\Phi(t) &\approx \exp(t^\beta), & t \text{ large} &\quad \dashrightarrow \quad EXP^\beta(\Omega) & \quad (\beta > 0) \\
\Phi(t) &\approx t^p \log^{\alpha p} t, & t \text{ large} &\quad \dashrightarrow \quad L^p(\log L)^{\alpha p}(\Omega) \\
&&&&&&&&&& (p > 1, \alpha \in \mathbb{R}; p = 1, \alpha > 0) \\
\Phi(t) &\approx \frac{t^p}{\log t}, & t \text{ large} &\quad \dashrightarrow \quad \frac{L^p}{\log L}(\Omega).
\end{aligned}$$

Orlicz spaces (and therefore, in particular, Lebesgue spaces) enjoy an important property, which involves the notion of decreasing rearrangement f_* of a measurable function $f : \Omega \rightarrow \mathbb{R}$, defined as function $f_* : [0, |\Omega|] \rightarrow [0, \infty]$ through

$$f_*(t) = \inf\{\lambda \geq 0 : \mu_f(\lambda) \leq t\} \quad (\inf \emptyset = \infty),$$

where $\mu_f : [0, \infty[\rightarrow [0, |\Omega|]$ is the distribution function of $|f|$, defined by

$$\mu_f(\lambda) = |\{x \in \Omega : |f(x)| > \lambda\}|.$$

The property we highlight is to be *rearrangement-invariant*: two equimeasurable functions f, g (i.e. such that $\mu_f = \mu_g$) have the same norm: this fact is obviously a feature of all norms which can be expressed in terms of the distribution function of $|f|$. Moreover, since f and f_* have the same distribution function, it is $\|f\|_{L^p(\Omega)} = \|f_*\|_{L^p(0, |\Omega|)}$ ($1 \leq p \leq \infty$) and more generally $\|f\|_{L^\Phi(\Omega)} = \|f_*\|_{L^\Phi(0, |\Omega|)}$.

Orlicz spaces are special “Banach function spaces”, a terminology which sometimes is origin of a small confusion: the term “Banach function spaces” does not denote just Banach spaces whose elements are measurable functions (as, for instance, Sobolev spaces), but more frequently denotes Banach spaces whose elements are measurable functions satisfying some additional properties. In the theory developed in [15] a “Banach function space” is a Banach space $(X, \|\cdot\|_X)$ such that the elements of X are measurable functions $f, g : \Omega \rightarrow \mathbb{R}$ satisfying

- $|f| \leq |g|$ a.e. $\Rightarrow \|f\|_X \leq \|g\|_X$
- $|f_k| \nearrow |f|$ a.e. $\Rightarrow \|f_k\|_X \nearrow \|f\|_X$
- $\|1\|_X < \infty$
- $\int_E |f| dx \leq c(E) \|f\|_X \quad (E \subset \Omega)$.

This notion opens the way to a general theory, where many “standard” functions spaces in Analysis represent just examples. Interested readers may consult also other classical monographs, like, for instance, [124, 120, 162].

1.3. A first bundle of norms and spaces

Fix $1 < p < \infty$ and consider a measurable function $f : \Omega \rightarrow \mathbb{R}$ such that $f \in \bigcap_{0 < \varepsilon < p-1} L^{p-\varepsilon}(\Omega)$. If $f \notin L^p(\Omega)$, then

$$\left(\int_\Omega |f|^{p-\varepsilon} dx \right)^{\frac{1}{p-\varepsilon}} \uparrow \infty \quad \text{as } \varepsilon \rightarrow 0, \quad (1)$$

while, on the other hand, if $f \in L^p(\Omega)$, then by Hölder's inequality

$$\left(\int_{\Omega} |f|^{p-\varepsilon} dx \right)^{\frac{1}{p-\varepsilon}} \leq \left(\int_{\Omega} |f|^p dx \right)^{\frac{1}{p}} < \infty \quad \forall \varepsilon \in (0, p-1). \quad (2)$$

The functions f such that the blowup in (1) is controlled by a factor ε :

$$\|f\|_{L^p(\Omega)} := \sup_{0 < \varepsilon < p-1} \left(\varepsilon \int_{\Omega} |f|^{p-\varepsilon} dx \right)^{\frac{1}{p-\varepsilon}} < \infty \quad (3)$$

constitute the *grand* L^p space, denoted by $L^{p)}(\Omega)$. Of course, if there is no blowup as in (2), then (3) holds as well, so that

$$L^{p)}(\Omega) = \{f : \|f\|_{L^p(\Omega)} < \infty\} \supset L^p(\Omega).$$

The heart of the matter is the balance of the blowup of a norm (depending on a parameter, say ε) and the parameter ε itself, whose main role here is to be infinitesimal as $\varepsilon \rightarrow 0$. It is therefore natural to consider also the analogous norm, where ε is substituted by a power ε^θ , where $\theta > 0$. This has been done in [88], where the norm

$$f \in L^{p),\theta}(\Omega) \Leftrightarrow \|f\|_{L^{p),\theta}(\Omega)} = \sup_{0 < \varepsilon < p-1} \left(\varepsilon^\theta \int_{\Omega} |f|^{p-\varepsilon} dx \right)^{\frac{1}{p-\varepsilon}} < \infty \quad (4)$$

had a role in an existence and uniqueness result for the p -harmonic equation

$$\operatorname{div} |\nabla u|^{p-2} \nabla u = \operatorname{div} f.$$

The space $L^{p),\theta}(\Omega)$ (which gives back $L^{p)}(\Omega)$ when $\theta = 1$) is a rearrangement-invariant Banach function space and, by the same argument as before, it contains $L^p(\Omega)$. Moreover,

$$L^p(\Omega) \subsetneq \frac{L^p}{\log^\theta L}(\Omega) \subsetneq L^{p),\theta}(\Omega) \subsetneq \bigcap_{\alpha > 1} \frac{L^p}{\log^{\alpha\theta} L}(\Omega) \subsetneq \bigcap_{0 < \varepsilon < p-1} L^{p-\varepsilon}(\Omega). \quad (5)$$

Embeddings between grand Lebesgue spaces and Orlicz spaces have been intensively studied and fully characterized. The interested reader can see the papers [18, 87, 94, 23].

The generalization (4) can be pushed on, substituting ε^θ with a general measurable function $\delta(\varepsilon)$ which is positive a.e. ([26]). In such case, the product of the norm in $L^{p-\varepsilon}$ by $\delta(\varepsilon)$ is a measurable function, and the supremum must be changed into an essential supremum: the norm in $L^{p),\delta(\cdot)}$ is defined by

$$\|f\|_{L^{p),\delta(\cdot)}(\Omega)} = \operatorname{ess\,sup}_{0 < \varepsilon < p-1} \left(\delta(\varepsilon) \int_{\Omega} |f|^{p-\varepsilon} dx \right)^{\frac{1}{p-\varepsilon}} < \infty.$$

Of course different δ 's may generate the same space $L^{p),\delta(\cdot)}$, and in [26] it has been shown that the same class of spaces can be defined starting from the class of the increasing δ 's (hence functions defined pointwise) so that the essential supremum can be

equivalently replaced by a simple supremum. In [9] (see also [80]) also the exponent $p - \varepsilon$ has been generalized into a measurable function of ε (in the paper it is shown that this operation makes sense, and that, for instance, this way one can catch certain Orlicz spaces of exponential type for unbonded exponents; see also [12] and references therein), and recently in [8] it has been shown that again such full general definition gives back the spaces in [26] (we mention here the computation of their Boyd indices in [74] and their weighted version in [10]). Moreover, in [8] it has been shown that the same class of spaces is characterized through the norms of the type

$$\rho_\varphi(f) = \sup_{p \in [p_-, p_+]} \varphi(p) \left(\int_{\Omega} |f|^p dx \right)^{1/p},$$

where $\varphi = \varphi(\cdot)$ on $[p_-, p_+)$ is positive and decreasing (i.e. $p_- \leq p_1 < p_2 < p_+ \Rightarrow \varphi(p_1) \geq \varphi(p_2) > 0$). At last, it should be observed that such class of spaces, obtained in literature as endpoint of a sequence of generalizations, appeared maybe for the first time in the final remark in [23].

The *small Lebesgue space* $L^{(p', \theta)}(\Omega)$ (here, as usual, $p' = p/(p-1)$) is defined as the associate space of the grand Lebesgue space $L^{p, \theta}(\Omega)$, i.e. $\|\cdot\|_{L^{(p', \theta)}(\Omega)}$ is the smallest functional defined on the measurable functions such that a kind of Hölder type inequality holds:

$$\int_{\Omega} u(x)v(x)dx \leq \|u\|_{L^{(p', \theta)}(\Omega)} \|v\|_{L^{p, \theta}(\Omega)}. \quad (6)$$

As a byproduct of the general theory of Banach function spaces, $L^{(p', \theta)}(\Omega)$ is a rearrangement-invariant Banach function space. The inequality (6) gives automatically an implicit definition of the norm of $L^{(p', \theta)}(\Omega)$ (it suffices to divide by $\|u\|_{L^{(p', \theta)}(\Omega)}$ and pass to the supremum, in the left hand side, over $u \neq 0$), however, the explicit (i.e. not depending on the notion of grand Lebesgue space) characterization has been found in [57] (here we exchange p and p' for the sake of simplicity of notation):

$$\|u\|_{L^{(p, \theta)}(\Omega)} = \inf_{u = \sum_k u_k} \left\{ \sum_{k=1}^{\infty} \inf_{0 < \varepsilon < p'-1} \varepsilon^{-\frac{\theta}{p'-\varepsilon}} \left(\int_{\Omega} |u_k|^{(p'-\varepsilon)'} dx \right)^{\frac{1}{(p'-\varepsilon)'}} \right\}, \quad (7)$$

see also [67, 68, 25]. Incidentally, let us remark that in (6) equality may hold for every $v \in L^\infty(\Omega)$, as stated in [57, Lemma 2.9]: in fact, if the supremum defining $\|v\|_{L^{p, \theta}(\Omega)}$ is assumed in $\varepsilon = p-1$, then it suffices to set $u = \chi_{\{v>0\}} - \chi_{\{v<0\}}$: hence $|u| \leq 1$ and

$$\begin{aligned} \int_{\Omega} uv dx &\leq \|u\|_{L^{(p', \theta)}(\Omega)} \|v\|_{L^{p, \theta}(\Omega)} \leq \|v\|_{L^{p, \theta}(\Omega)} \inf_{0 < \varepsilon < p-1} \varepsilon^{-\frac{\theta}{p-\varepsilon}} \left(\int_{\Omega} |u|^{(p-\varepsilon)'} dx \right)^{\frac{1}{(p-\varepsilon)'}} \\ &\leq (p-1)^\theta \int_{\Omega} |v| dx \cdot \frac{1}{(p-1)^\theta} = \int_{\Omega} uv dx; \end{aligned}$$

therefore the conclusion is that the inequalities are in fact equalities (see [30] for details on this remark).

The notion of associate space has a nice behavior with respect to inclusion (note that the *inclusion* is equivalent to the *continuous embedding* in the case of Banach function spaces, see [15, Thm 1.8 p. 7]), namely, for associate spaces the inverted inclusions hold (note also that, on the contrary, this property does not hold in general for the notion of dual space). Hence ([23, 36])

$$\bigcup_{\varepsilon>0} L^{p+\varepsilon} \subsetneq \bigcup_{\beta>1} L^p(\log L)^{\frac{\beta\theta}{p'-1}} \subsetneq L^{(p,\theta)}(\Omega) \subsetneq L^p(\log L)^{\frac{\theta}{p'-1}} \subsetneq L^p(\Omega).$$

We conclude this subsection mentioning that comparisons with another class of Banach function spaces of recent interest, namely, the variable Lebesgue spaces (see e.g. the books [149, 129, 48, 39, 139, 51, 112, 113] and [98]) appear in [71, 38].

1.4. The point of view of interpolation-extrapolation theory: the first results

The structure of the norm of grand and small Lebesgue spaces is familiar to researchers in interpolation theory, from which it is possible to get the expression of a quasinorm, equivalent to that one in (4) and (7), respectively. In the case of small Lebesgue spaces, the new expression is also “nicer”, because one can get rid of decompositions and can “play” just with integrals. We are going to write the expression of such quasinorms, trying to “read” them in terms of the language of interpolation theory. We begin by recalling the classical notion of Lorentz space (see e.g. the recent monograph [139] on function spaces), starting from a classical result by Peetre in 1966 (for the sake of history, without claiming to be complete on interpolation theory, we mention the precedent papers [78, 123, 122, 21]; see also [37, 40] and references therein).

Let $1 < p < \infty$. We recall that from the equality (see e.g. [138]; [160, 5.10 p.213]; [15, Thm 6.2 p. 74])

$$\int_0^t f_*(s) ds = \inf_{f=g+h} \left\{ \|g\|_{L^1(\Omega)} + t \|h\|_{L^\infty(\Omega)} \right\} := K(f, t, L^1(\Omega), L^\infty(\Omega)),$$

with the help of the classical Hardy’s inequality, one has

$$\|f\|_{L^p(\Omega)} = \|f_*\|_{L^p(0,|\Omega|)} \approx \left\| t^{-1} \int_0^t f_* \right\|_{L^p(0,|\Omega|)} = \|t^{-1} K(f, t, L^1(\Omega), L^\infty(\Omega))\|_{L^p(0,|\Omega|)}.$$

The expression on the right hand side is a special case of the quasinorm of the Lorentz spaces, which represent a generalization of the Lebesgue spaces in a different “direction” with respect to the Orlicz spaces; for our goals we recall here few standard facts.

Fix a parameter q , $1 < q < \infty$, and set $\|f\|_{L^{p,q}(\Omega)} = \|t^{\frac{1}{p}-\frac{1}{q}} f_*(t)\|_{L^q(0,|\Omega|)}$. The parameter q , roughly speaking, gives a “tuning” of the Lebesgue space $L^p(\Omega)$: notice that, for any $1 < q < \infty$, if for instance $\Omega = (0, 1) \subset \mathbb{R}$, the “boundary” power function $t^{-\frac{1}{p}}$ remains unchanged. Proceeding as above,

$$\|f\|_{L^{p,q}(\Omega)} = \|t^{\frac{1}{p}-\frac{1}{q}} f_*(t)\|_{L^q(0,|\Omega|)} \approx \left\| t^{\frac{1}{p}-\frac{1}{q}} \cdot t^{-1} \int_0^t f_* \right\|_{L^q(0,|\Omega|)}$$

$$\begin{aligned}
&= \left\| t^{\frac{1}{p}-\frac{1}{q}} \cdot t^{-1} K(f, t, L^1(\Omega), L^\infty(\Omega)) \right\|_{L^q(0, |\Omega|)} \\
&= \left\| t^{-(1-\frac{1}{p})} \cdot t^{-\frac{1}{q}} K(f, t, L^1(\Omega), L^\infty(\Omega)) \right\|_{L^q(0, |\Omega|)} \\
&:= \|f\|_{(L^1, L^\infty)_{1-\frac{1}{p}, q}}. \tag{8}
\end{aligned}$$

Note that $\|f\|_{L^p(\Omega)} = \|f\|_{L^{p,p}(\Omega)} = \|f\|_{(L^1, L^\infty)_{1-\frac{1}{p}, p}}$. For any $1 < q < \infty$, the norm in (8) gives a space “close” to the Lebesgue space $L^p(\Omega)$. The parameter p appears in the first index, namely in the expression $1 - 1/p$ which has the advantage to be always a number between 0 and 1: when it is 0, the space is close to L^1 , the first space in the pair (L^1, L^∞) ; when it is 1, the space is close to L^∞ , the second space in the pair (L^1, L^∞) . About the parameter q , it can be shown that when q increases, the space becomes larger (with respect to inclusion).

The whole machinery can be made again considering one more parameter $\alpha \in \mathbb{R}$, which appears as power of a logarithm, put as factor of $K(f, t, L^1(\Omega), L^\infty(\Omega))$: a further refinement, again classical, is that one of the Lorentz-Zygmund spaces. For any $1 < q < \infty$, $\alpha \in \mathbb{R}$, set

$$\|f\|_{L^{p,q;\alpha}(\Omega)} = \|t^{\frac{1}{p}-\frac{1}{q}} (1 - \log t)^\alpha f_*(t)\|_{L^q(0, |\Omega|)}.$$

Proceeding again as above,

$$\begin{aligned}
\|f\|_{L^{p,q;\alpha}(\Omega)} &= \left\| t^{-(1-\frac{1}{p})} (1 - \log t)^\alpha \cdot t^{-\frac{1}{q}} K(f, t, L^1(\Omega), L^\infty(\Omega)) \right\|_{L^q(0, |\Omega|)} \\
&:= \|f\|_{(L^1, L^\infty)_{1-\frac{1}{p}, q; \alpha}}.
\end{aligned}$$

At last, we may state the characterization of grand and small Lebesgue spaces given in [61]:

$$L^{(p)} = (L^1, L^p)_{1, \infty; -\frac{1}{p}} \tag{9}$$

$$L^{(p)} = (L^p, L^\infty)_{0, 1; -\frac{1}{p}}. \tag{10}$$

From the point of view of the digression above, in (9) one can see that the first index 1 tells that the space $L^{(p)}(\Omega)$ is “close” to $L^p(\Omega)$, the second space in the pair (L^1, L^p) ; the second index ∞ tells that the space is larger (with respect to inclusion) than $L^p(\Omega)$, and, finally, the third parameter $-\frac{1}{p}$ gives the correct logarithmic tuning of the space. In an analogous way one can read the relation in (10). After the computations of $K(f, t, L^1(\Omega), L^p(\Omega))$, $K(f, t, L^p(\Omega), L^\infty(\Omega))$ which are implicit in the symbols in (9), (10), respectively, one gets the equivalent quasinorms

$$\|u\|_{L^{(p), \theta}(\Omega)} \approx \sup_{0 < t < |\Omega|} \left(1 - \log \left(\frac{t}{|\Omega|} \right) \right)^{-\frac{\theta}{p}} \left(\int_t^{|\Omega|} u_*(s)^p ds \right)^{\frac{1}{p}} \tag{11}$$

$$\|u\|_{L^{(p,\theta)}(\Omega)} \approx \int_0^{|\Omega|} \left(1 - \log\left(\frac{t}{|\Omega|}\right)\right)^{\frac{\theta}{p'}-1} \left(\int_0^t u_*(s)^p ds\right)^{\frac{1}{p}} \frac{dt}{t}.$$

A different proof of these relations, for researchers not expert in interpolation theory, appears in [47], where sharp estimates are proved by using the Euler's Gamma function. We stress that the relations above, differently from [47], are written for domains Ω having finite measure, not necessarily equal to 1. Finally, we highlight that the right hand side of (11), by a simple change of variable (see [47, Proposition 3.1]), can be written also in a different, expressive form, so that

$$\|u\|_{L^{(p,\theta)}(\Omega)} \approx \sup_{0 < \varepsilon < p-1} \left[\left(\frac{\varepsilon}{p-1}\right)^\theta \int_{|\Omega|e^{1-(p-1)/\varepsilon}}^{|\Omega|} u_*(s)^p ds \right]^{\frac{1}{p}}.$$

1.5. The development of grand and small spaces

Starting from the norms listed in subsection 1.3, in literature several authors devoted their attention either to the spaces themselves (see e.g. [105]), either to a number of generalizations and variants, applied to questions related e.g. to the boundedness of operators, to integral inequalities, to interpolation theory. Applications to certain optimization problems of the type considered in [13] will appear in a forthcoming paper.

Here we give just an idea of the large literature appeared in recent years, where generalizations and variants have been considered.

First of all, for the reader's convenience, we list some quasinorms of some spaces already mentioned above, so that it is possible to make evident the differences and the analogies between them. Assume now, for simplicity, $|\Omega| = 1$, and let $1 < p, q < \infty$, $\alpha \in \mathbb{R}$, $\theta > 0$.

We recall the quasinorms for Lorentz spaces, Zygmund spaces (which can be seen as special Orlicz spaces) and small Lebesgue spaces, respectively:

$$\begin{aligned} \|u\|_{L^{p,q}(\Omega)} &\approx \left(\int_0^1 t^{\frac{q}{p}-1} u_*(t)^q dt \right)^{\frac{1}{q}} \\ \|u\|_{L^p(\log L)^\alpha(\Omega)} &\approx \left(\int_0^1 (1 - \log t)^\alpha u_*(t)^p dt \right)^{\frac{1}{p}} \\ \|u\|_{L^{(p,\theta)}(\Omega)} &\approx \int_0^1 (1 - \log t)^{\frac{\theta}{p'}-1} \left(\int_0^t u_*(s)^p ds \right)^{\frac{1}{p}} \frac{dt}{t}. \end{aligned}$$

This last norm can be considered as special case of the so-called $GF(p, m, w)$ spaces, introduced in [70] and considered e.g. in [72, 86, 147, 58] (see these references for values and digressions on the parameters p and m , and the weight w):

$$\|u\|_{GF(p,m,w)} \approx \left[\int_0^1 w(t) \left(\int_0^t u_*(s)^p ds \right)^{\frac{m}{p}} \frac{dt}{t} \right]^{1/m}. \quad (12)$$

We mention also the following variant, appeared in [132]:

$$\|u\|_{\mathcal{H}_{x_0}^{p,m,w}(\Omega)} \approx \left[\int_0^{2\text{diam}(\Omega)} w(x_0,t)^m \left(\int_{B(x_0,t)} u(s)^p ds \right)^{\frac{m}{p}} \frac{dt}{t} \right]^{1/m};$$

moreover, the variant of $G\Gamma(p,m,w)$ with double weights appears in [59] and the $G\Gamma(p,\infty;w,v)$ spaces appeared in [83]:

$$\|u\|_{G\Gamma(p,\infty;w,v)} \approx \text{ess sup}_{t>0} w(t) \left(\int_0^t v(s) u_*(s)^p ds \right)^{\frac{1}{p}}.$$

Iteration type results, where the Lebesgue norm in the expression of the norm of grand Lebesgue spaces is replaced by the norm of grand Lebesgue spaces, are in [7] (see also [56]);

Generalizations of small Lebesgue spaces appear in two papers (see [103, 11]), where some common results have been obtained independently (see also [28]).

As to variants of grand Lebesgue spaces, we mention the bilateral grand Lebesgue spaces considered in [119], the grand Orlicz spaces in [115], [24], [52], the grand Lorentz spaces in [99], the weighted grand Lebesgue spaces in [60, 102, 101, 108], the weighted fully measurable grand Lebesgue spaces in [104], the grand Morrey spaces in [130, 164, 109, 108, 110], the grand Morrey spaces of variable exponent in [77, 136], the grand Musielak-Orlicz-Morrey spaces in [133, 125], the grand grand Morrey spaces in [143], the grand Bochner-Lebesgue spaces in [111, 100] and, finally, the grand variable exponent Lebesgue spaces in [107, 62].

Finally, we mention that abstract notions of grand spaces appear in [61, 112, 113], and a way to extend the notion of grand Lebesgue spaces when the underlying measure space has infinity measure appears in [150, 151, 163].

2. Applications to Sobolev spaces theory and to PDEs

The grand and small Lebesgue spaces generate the corresponding Sobolev analogs: for a given p , $1 < p < \infty$, the grand Sobolev space $W^{1,p}(\Omega)$ consists of all functions $u \in W^{1,1}(\Omega)$ such that $|\nabla u| \in L^p(\Omega)$; similarly, the small Sobolev space $W^{1,p}(\Omega)$ consists of all functions $u \in W^{1,1}(\Omega)$ such that $|\nabla u| \in L^p(\Omega)$. The definitions of the corresponding spaces $W_0^{1,p}(\Omega)$, $W_0^{1,p}(\Omega)$ constituted by the functions which, roughly speaking, assume value zero on the boundary, follow the same lines of the standard definitions given in the Sobolev spaces theory; the same words can be spent for grand and small Sobolev spaces of higher order. This whole class of spaces has been intensively studied and their properties found several applications in Geometric function theory and PDEs. For a review, the interested reader may consult [42] (see also [152]) and the rich bibliography therein. In the following we will confine ourselves to some results involving Sobolev embeddings and to some existence, uniqueness, regularity results for certain elliptic PDEs.

2.1. The role of grand Sobolev spaces in embedding theorems

Let us begin by recalling some classical embedding results. For $\Omega \subset \mathbb{R}^n$ open, bounded, connected, $\partial\Omega$ Lipschitz, $n \geq 2$, we have

$$W^{1,p}(\Omega) \hookrightarrow L^{\frac{np}{n-p}}(\Omega) \hookrightarrow L^{\frac{np}{n-p}}(\Omega) \quad 1 \leq p < n$$

$$W^{1,n}(\Omega) \hookrightarrow L^q(\Omega) \quad (p = n) \quad p \leq q < \infty$$

$$W^{1,p}(\Omega) \hookrightarrow \mathcal{C}^{0,1-\frac{n}{p}}(\Omega) \quad (p > n)$$

$$W^{1,n}(\Omega) \hookrightarrow EXP^{\frac{n}{n-1}}(\Omega) \quad W^{1,n}(\Omega) \not\hookrightarrow L^\infty(\Omega) \quad (p = n).$$

When derivatives are in Lorentz spaces (see [159, 46, 6, 127]), recalling that

$$L^{n,1}(\Omega) \hookrightarrow L^n(\Omega) \hookrightarrow L^{n,\infty}(\Omega),$$

we have

$$WL^{n,1}(\Omega) \hookrightarrow L^\infty(\Omega), \quad WL^{n,1}(\Omega) \hookrightarrow \mathcal{C}^0(\Omega) \quad (13)$$

and

$$WL^{n,p}(\Omega) \hookrightarrow EXP^{\frac{p}{p-1}}(\Omega), \quad WL^{n,p}(\Omega) \not\hookrightarrow L^\infty(\Omega) \quad 1 < p \leq \infty.$$

When derivatives are in Orlicz spaces $L^A(\Omega)$ (see [49, 22, 34, 32, 33]), a notion of A^* corresponding to the Sobolev conjugate exponent can be given in a sharp way, and we have, roughly speaking,

$$W^{1,A}(\Omega) \hookrightarrow L^{A^*}(\Omega) \quad A(t) \lesssim t^n (\log t)^{n-1} (\log \log t)^{n-1} \dots, \quad t \rightarrow \infty.$$

The role of grand Lebesgue spaces becomes significant if we give a look at the special case $L^A(\Omega) = \frac{L^n}{\log^\theta L}(\Omega)$: from the computation of A^* we get, in this case,

$$W_0 \frac{L^n}{\log^\theta L}(\Omega) \hookrightarrow EXP^{n/(n-1+\theta)}(\Omega) \quad \forall \theta > 0 \quad (14)$$

and now, in spite of the embedding (see (5))

$$\frac{L^n}{\log^\theta L}(\Omega) \subsetneq L^{n,\theta}(\Omega), \quad (15)$$

we have ([76, 85])

$$u \in W_0^{1,1}(\Omega), |\nabla u| \in L^{n,\theta}(\Omega) \Rightarrow u \in EXP^{n/(n-1+\theta)}(\Omega). \quad (16)$$

The imbedding (15) and the fact that on the right hand side of (14) and (16) the exponential type spaces are the same, tell that, in this critical setting, the fact to consider the gradient in grand Lebesgue spaces gives no loss of integrability.

In this framework we mention that if we write (14) in the form

$$\sup_{0 < t < 1} (1 - \log t)^{-\frac{\theta}{n}} \left(\int_t^1 |\nabla u|_*(s)^n ds \right)^{\frac{1}{n}} < \infty \Rightarrow \sup_{0 < t < 1} \frac{|u|_{**}(t)}{(1 - \log t)^{\frac{n-1+\theta}{n}}} < \infty,$$

the following generalized form holds ([58]):

$$\sup_{0 < t < 1} w(t) \left(\int_t^1 |\nabla u|_*(s)^n ds \right)^{\frac{1}{n}} < \infty \Rightarrow \sup_{0 < t < 1} \frac{w(t)|u|_{**}(t)}{(1 - \log t)^{\frac{n-1}{n}}} < \infty.$$

It should be noted, here, that the norm on the right hand side is that one of a Marcinkiewicz space (see [15]).

Let us examine now embedding theorems for small Sobolev spaces, beginning with the critical situation of derivatives in the small $L^n(\Omega)$ spaces. Recalling that

$$L^{n,\infty}(\Omega) \hookrightarrow L^n(\Omega)$$

(for results about embeddings between Lorentz and grand Lebesgue spaces see [96, 87, 3]), by associativity one gets immediately

$$L^n(\Omega) \hookrightarrow L^{n,1}(\Omega)$$

and therefore, since by (13) it is $WL^{n,1}(\Omega) \hookrightarrow \mathcal{C}^0(\Omega)$, the same holds for $W^{1,(n)}(\Omega)$ and the oscillation of these functions can be estimated as follows ([67, 68, 145]):

$$u \in W^{1,1}(\Omega), |\nabla u| \in L^n(\Omega) \Rightarrow \text{osc}_{B(x,t)} u \leq c_n |\Omega|^{1/n} \| |\nabla u| \chi_{B(x,t)} \|_{L^n(\Omega)}.$$

We conclude this subsection by discussing some compactness results and a dimension-free Sobolev estimate.

The well known classical results about compactness, for higher order Sobolev spaces, are the following: if $1 \leq p < \frac{n}{k}$, $k \in \mathbb{N}$, then (the historical references are [148] for the case $k = 1$, [114] for the case $k \geq 1$)

$$W^{k,p}(\Omega) \hookrightarrow L^q(\Omega) \quad 1 \leq q < p^* = \frac{np}{n - kp}$$

is compact, while $W^{k,p}(\Omega) \hookrightarrow L^{p^*}(\Omega)$ is not compact. With the help of the notions of grand and small Lebesgue and Sobolev spaces, the loss of compactness in the extreme case $q = p^*$ can be replaced by the following gains of compactness (see [69] for the case $k = 1$, [36] for the case $k \geq 1$): the embeddings

$$W^{k,p}(\Omega) \hookrightarrow L^{p^*}(\Omega) \quad W^{k,(p)}(\Omega) \hookrightarrow L^{p^*}(\Omega)$$

are compact. Hence, for instance, every sequence of functions equibounded in $W^{k,(p)}(\Omega)$ has a subsequence strongly convergent in $L^{p^*}(\Omega)$.

Let us spend now few words on a so-called dimension-free Sobolev estimate (for classical results on this topic, for instance the logarithmic Sobolev inequality by Gross

and its equivalence to the hypercontractive inequality by Nelson, see e.g. [1] and references therein). Let $1 < p < \infty$ be fixed, and, for any $n > p$, let B_n be a given open ball in \mathbb{R}^n of measure 1. The following version of the Sobolev inequality is well known (see e.g. [95]):

$$\|f\|_{L^{np/(n-p)}(B_n)} \leq \frac{p(n-1)}{n-p} \|\nabla f\|_{L^p(B_n)} \quad \forall f \in W_0^{1,p}(B_n), \quad \forall n > p$$

and, noticing the boundedness, with respect to the dimension, of the constant in the right hand side, we can write simply

$$\|f\|_{L^{np/(n-p)}(B_n)} \leq c_p \|\nabla f\|_{L^p(B_n)} \quad \forall f \in W_0^{1,p}(B_n), \quad \forall n > p$$

or, equivalently, taking into account of the rearrangement-invariance of the Lebesgue spaces (see subsection 1.2),

$$\|f_*\|_{L^{np/(n-p)}(0,1)} \leq c_p \|\nabla f\|_{L^p(B_n)}.$$

This inequality makes visible the gain of summability given by the Sobolev inequality, such gain being dependent of the dimension n , and being smaller and smaller as $n \rightarrow \infty$: in fact we have $\frac{np}{n-p} \searrow p$ and $\|f_*\|_{L^p(0,1)} \leq \|f_*\|_{L^{np/(n-p)}(0,1)} \forall n > p$. If, on the left hand side, one wishes to see a Lebesgue space not depending on n , the best inequality one can write is

$$\|f_*\|_{L^p(0,1)} \leq c_p \|\nabla f\|_{L^p(B_n)} \quad \forall n > p$$

which means, in fact, that there is no gain of summability (with respect to the gradient) in the scale of Lebesgue spaces. However, in the scale of rearrangement-invariant Banach function spaces, a gain of summability independent of the dimension exists: if one looks for the best (=“smallest”) space $Y_p(0, 1)$ such that

$$\|f_*\|_{Y_p(0,1)} \leq c_p \|\nabla f\|_{L^p(B_n)} \quad \forall f \in W_0^{1,p}(B_n), \quad \forall n > p,$$

one has that $Y_p(0, 1) \subset L^p(\log L)^{p/2}(0, 1)$, the exponent $p/2$ being optimal (see [126], where the authors answered positively a conjecture by Hans Triebel). The point here is that a further refinement holds: one has $Y_p(0, 1) \subset L^{(p,p')/2}(0, 1)$ and also that the inclusion $Y_p(0, 1) \subset L^p(\log L)^{p/2}(0, 1)$ is optimal in the scale of Orlicz spaces ([64], see also [128]).

3. Some applications to PDEs

In the previous section we already mentioned the paper [42] and references therein. Without claiming to be complete, we point out the presence of the notion of grand and small Lebesgue and Sobolev spaces also in several other papers, see e.g. [27, 45, 41, 53, 54, 55, 82, 89, 90, 91, 106, 134, 141, 142, 144, 147, 152, 153, 166].

In this section we will be not interested in applications to parabolic problems (however, the interested reader may consult the papers [65, 66]) and we will deal with three

particular cases of the following problem: let $\Omega \subset \mathbb{R}^n$ ($n \geq 2$) be open, bounded, connected, $\partial\Omega$ smooth, and let $a : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ be such that

$$\begin{cases} x \rightarrow a(x, \xi) & \text{measurable} \quad \forall \xi \in \mathbb{R}^n \\ \xi \rightarrow a(x, \xi) & \text{continuous} \quad \forall x \in \Omega \text{ a.e.} \end{cases}$$

$$\begin{cases} a(x, \xi)\xi \geq \alpha|\xi|^2 \\ |a(x, \xi)| \leq \beta|\xi| \end{cases} \quad \forall x \in \Omega \text{ a.e.}, \quad \forall \xi \in \mathbb{R}^n$$

$$\langle a(x, \xi) - a(x, \eta), \xi - \eta \rangle > 0, \quad \xi \neq \eta \quad (\text{strict monotonicity}).$$

We consider the Dirichlet problem:

$$\begin{cases} -\operatorname{div} a(x, \nabla u) = f & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases} \quad (17)$$

for data f lying in some Banach spaces. We recall that a function $u \in W_0^{1,1}(\Omega)$ is said to be a distributional solution if

$$\begin{cases} |a(x, \nabla u)| \in L^1(\Omega) \\ \int_{\Omega} a(x, \nabla u) \nabla \varphi \, dx = \int_{\Omega} f \varphi \, dx \quad \forall \varphi \in \mathcal{C}_0^\infty(\Omega) \end{cases}.$$

The typical result of the classical variational method (originated in [117, 157, 158] in the linear case, and extended to the nonlinear case in [118], [121, Thm 2.8 p.183], see e.g. also [75, Thm 26.14 p.205], [155, Prop. 5.1 p. 60; Thm 6.1 p.76]) is the following:

$$f \in W^{-1,2}(\Omega) = (W_0^{1,2}(\Omega))' \quad \Rightarrow \quad \exists u \text{ distributional solution in } W_0^{1,2}(\Omega).$$

In order to distinguish the numbers “2” above, let us mention that in the case of $a = a(x, \xi)$ of “growth p ” (our assumptions fit into the case $p = 2$), the above statement reads as

$$f \in W^{-1,p'}(\Omega) = (W_0^{1,p}(\Omega))' \quad \Rightarrow \quad \exists u \text{ distributional solution in } W_0^{1,p}(\Omega);$$

moreover, solutions assumed in $W_0^{1,p}(\Omega)$ are unique by the strict monotonicity.

Let us begin with the case $n = 2$ and the datum f in an Orlicz space. It is easy to find the spaces for f such that the variational method can be applied: in fact, from the Sobolev embedding

$$W_0^{1,2}(\Omega) \hookrightarrow \exp_2(\Omega) := \overline{L^\infty(\Omega)}^{EXP^2(\Omega)},$$

by duality,

$$L(\log L)^{\frac{1}{2}}(\Omega) \hookrightarrow W^{-1,2}(\Omega),$$

hence the variational method works when the datum f is in a Lebesgue or Orlicz space contained in $L(\log L)^{\frac{1}{2}}$, namely, all the spaces $L^m(\Omega)$, $m > 1$, and all the spaces $L(\log L)^\delta(\Omega)$, $\delta \geq \frac{1}{2}$. The following estimate appears in [3]:

$$\|\nabla u\|_{L^2(\log L)^{2\delta-1}(\Omega)} \leq c \|f\|_{L(\log L)^\delta(\Omega)} \quad \delta \in \left[\frac{1}{2}, 1\right]. \quad (18)$$

For completeness, we recall that in the linear case, the datum f in $L \log L(\Omega)$ was studied in [5] where sharp L^∞ estimates have been derived. Moreover, the case $f \in L(\log L)^{\frac{1}{2}}(\log \log L)^{\frac{1}{2}}(\Omega)$ has been treated in [165], the case $f \in L(\log L)^{\frac{1}{2}}(\log \log L)^{\frac{\beta}{2}}(\Omega)$ in [45], the case $f \in L(\log L)^{\frac{1}{2}}(\log \log \log L)^{\frac{\beta}{2}}(\Omega)$ in [43], the case $f \in L(\log L)^\delta(\log \log L)^{\frac{\beta}{2}}(\Omega)$ in [44], and, finally, some optimal results appear in [4].

The same procedure can be carried on in the case $n > 2$, and it may be easily seen that the variational method fails for f “free to move” in a space larger than

$$\begin{cases} L(\log L)^{\frac{1}{2}}(\Omega) & (n = 2) \\ L^{\frac{2n}{n+2}}(\Omega) & (n > 2) \end{cases}.$$

After these preliminary facts, we may therefore assert that, speaking about the spaces for the datum f , the space $L^1(\Omega)$ is never “variational”, while the space $L(\log L)(\Omega)$ is “variational” in the case $n = 2$, and it is not in the case $n > 2$. For each of these two spaces we are going to mention a result, where through the notion of grand Lebesgue space one can state an existence and uniqueness result, and a regularity result, respectively. Let us begin with the following result related to the non variational case $f \in L^1(\Omega)$:

THEOREM 1. ([73], see also [88, 131]) *If $\Omega \subset \mathbb{R}^2$ is open, bounded, connected, $\partial\Omega$ smooth, and if $a : \Omega \times \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is such that*

$$\begin{cases} x \rightarrow a(x, \xi) \text{ measurable } \forall \xi \in \mathbb{R}^2 \\ \xi \rightarrow a(x, \xi) \text{ continuous } \forall x \in \Omega \text{ a.e.} \end{cases}$$

$$\begin{cases} |a(x, \xi) - a(x, \eta)| \leq m|\xi - \eta| & \text{(Lipschitz continuity)} \\ \frac{1}{m}|\xi - \eta|^2 \leq \langle a(x, \xi) - a(x, \eta), \xi - \eta \rangle & \text{(strong monotonicity)} \\ a(x, 0) = 0 \end{cases},$$

then the Dirichlet problem

$$\begin{cases} -\operatorname{div} a(x, \nabla u) = f \in L^1(\Omega) \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

admits one and only one distributional solution $u \in W_0^{1,2}(\Omega)$.

As a consequence, from the Sobolev embedding (16), one gets $u \in EXP(\Omega)$, already known from [19, 31] in the linear case). Theorem 1 is in agreement also with [79], where existence and uniqueness of a distributional solution $u \in \bigcap_{q < 2} W_0^{1,q}(\Omega)$ has been proved in the linear case, for data f bounded Radon measure, that is, f in the dual of $\mathcal{C}(\bar{\Omega})$.

Let us consider, now, the case of the right hand side $f \in L(\log L)$ for the problem (17) in the linear case. We already stated that when $n = 2$ there is existence and uniqueness of a distributional solution; otherwise, when $n > 2$, in general, uniqueness fails if one looks for distributional solutions in spaces “much” larger than $W_0^{\frac{n}{n-1}}(\Omega)$ ([154], see e.g. also [140]). By considering appropriate notions of solutions, for several ranges for the parameter $\delta \in]0, +\infty[$, unifying results in [156, 16, 137, 81, 63, 3, 116], we can state the following

THEOREM 2. *If $\Omega \subset \mathbb{R}^n$ is open, bounded, connected, $\partial\Omega$ smooth, and if $A = (a_{ij})_{i,j=1,\dots,n}$, $a_{ij} = a_{ji}$ is sufficiently regular, then the problem*

$$\begin{cases} \operatorname{div} A(x) \nabla u = f \in L(\log L)^\delta(\Omega) \\ u = 0 \end{cases} \quad \text{on } \partial\Omega \quad \delta > 0$$

admits the existence of a solution in $W_0^{1, \frac{n}{n-1}}(\Omega)$, and the following estimate holds:

$$\|\nabla u\|_{L^{\frac{n}{n-1}}(\log L)^{\frac{n\delta}{n-1}-1}(\Omega)} \leq c \|f\|_{L(\log L)^\delta(\Omega)} \quad \delta > 0.$$

Note that (18) corresponds to the case $n = 2$, $\delta \in [1/2, 1]$, of Theorem 2. The knowledge of the small Sobolev spaces allows us to state the following improvement of the estimate in Theorem 2 (see [58]):

$$\|\nabla u\|_{L^{(n', n\delta - n + 1)}(\Omega)} \leq c \|f\|_{L(\log L)^\delta(\Omega)}, \quad \delta \in \left[\frac{n-1}{n}, +\infty \right[.$$

We conclude with the following regularity result (see [58]):

THEOREM 3. *If $\Omega \subset \mathbb{R}^n$ is open, bounded, connected, $\partial\Omega$ smooth, then there exists one and only one very weak solution u of the problem*

$$\begin{cases} -\Delta u = f \in L^1(\Omega; \delta(1 + |\log \delta|)^{\frac{n-1+\theta}{n}}) \\ u = 0 \end{cases} \quad \text{on } \partial\Omega,$$

namely, one and only one $u \in L^{(\frac{n}{n-1}, \theta)}(\Omega)$, $\theta > 0$, such that

$$-\int_{\Omega} u \Delta \varphi dx = \int_{\Omega} f \varphi dx \quad \forall \varphi \in C^2(\overline{\Omega}), \quad \varphi = 0 \text{ on } \partial\Omega.$$

Here $\delta(x) = \operatorname{dist}(x, \partial\Omega)$.

Observe that when $\theta = 1$ we have $L^{(\frac{n}{n-1}, 1)}(\Omega) \subset L^{\frac{n}{n-1}, 1}(\Omega)$, hence Theorem 3 improves the corresponding results appeared in [17, 50, 146].

4. The point of view of interpolation theory: some recent results

Using the celebrated Holmstedt's formula ([93]) and the computation of the K -functional, in the paper [59] (see also [35]) a new, recent characterization of grand and small Lebesgue spaces as interpolation spaces in the sense of Peetre, has been obtained. Namely, the following extension of formulas (9), (10) from [61], recalled in subsection 1.4, hold (here and in the remaining part of this section, for simplicity of notation, we omit to write " (Ω) " in the symbols denoting the function spaces and we assume $|\Omega| = 1$):

THEOREM 4. ([59], Theorem 3.1) *If $1 \leq p < q$, $\alpha > 0$, then*

$$L^{(q),\alpha} = (L^p, L^q)_{1,\infty; -\frac{\alpha}{q}}.$$

THEOREM 5. ([59], Proposition 4.2) *If $1 < p < q$, $\alpha > 0$, then*

$$L^{(p,\alpha)} = (L^p, L^q)_{0,1; -1+\alpha-\frac{\alpha}{p}}.$$

The main approach to get such results is based on the computation of the K -functional based on results from [14]; the following explicit equivalent expression of the K -functional between the grand Lebesgue spaces and Lebesgue spaces was found.

THEOREM 6. ([59], Theorem 3.2) *If $1 < p < q$, $\alpha > 0$, then*

$$K(f, t; L^{(p),\alpha}, L^q) \approx \sup_{0 < s < \varphi(t)} (1 - \log s)^{-\frac{\alpha}{p}} \left(\int_s^{\varphi(t)} f_*^p(x) dx \right)^{\frac{1}{p}} + t \left(\int_{\varphi(t)}^1 f_*^q(s) ds \right)^{\frac{1}{q}},$$

where φ is the inverse of the increasing function $\psi(t) = t^{\frac{1}{p}-\frac{1}{q}} (1 - \log t)^{-\frac{\alpha}{p}}$, $t \in (0, 1)$. Thus

$$t = \varphi(t)^{\frac{1}{p}-\frac{1}{q}} (1 - \log \varphi(t))^{-\frac{\alpha}{p}}.$$

As a consequence, the following new characterization of grand Lebesgue spaces holds:

THEOREM 7. ([59], Theorem 1.1) *If $1 < p < q$, $\alpha > 0$, then*

$$L^{(q),\alpha} = (L^{(p),\alpha}, L^q)_{1,\infty; -\frac{\alpha}{q}}.$$

The same paper [59] contains also the computation of the K -functional between two Grand Lebesgue spaces.

THEOREM 8. ([59], Theorem 3.3) *If $1 < p < q$, $\alpha > 0$, then*

$$K(f, t; L^{p, \alpha}, L^{q, \alpha}) \approx \sup_{0 < s < \varphi(t)} (1 - \log s)^{-\frac{\alpha}{p}} \left(\int_s^{\varphi(t)} f_*^p(x) dx \right)^{\frac{1}{p}} \\ + t \sup_{\varphi(t) < s < 1} (1 - \log s)^{-\frac{\alpha}{q}} \left(\int_s^1 f_*^q(x) dx \right)^{\frac{1}{q}},$$

where φ is the inverse of the increasing function $\psi(t) = t^{\frac{1}{p} - \frac{1}{q}} (1 - \log t)^{-\frac{\alpha}{p} + \frac{\alpha}{q}}$, $t \in (0, 1)$. Thus

$$t = \varphi(t)^{\frac{1}{p} - \frac{1}{q}} (1 - \log \varphi(t))^{-\frac{\alpha}{p} + \frac{\alpha}{q}}.$$

Again, after the computation of the K -functionals above, the following characterization of the interpolation spaces between two Grand Lebesgue spaces holds:

THEOREM 9. ([59], Theorem 1.2) *If $0 < \theta < 1$, $1 \leq r < +\infty$, $\alpha > 0$, $1 < p < q$, then*

$$\left(L^{p, \alpha}, L^{q, \alpha} \right)_{\theta, r} = L^{p\theta, r} (\log L)^{-\frac{\alpha}{p\theta}} \quad \text{where } \frac{1}{p\theta} = \frac{1 - \theta}{p} + \frac{\theta}{q}. \quad (19)$$

Readers are warned to beware of the different symbol for Lorentz-Zygmund spaces on the right hand side of (19): in this section we follow the notation commonly used in interpolation theory and adopted in [59], which is different from that one used in previous sections, which is, on the contrary, used in the theory of Orlicz spaces. According to this latter notation (introduced in subsection 1.2), the space on the right hand side of (19) should be written $L^{p\theta, r} (\log L)^{-\frac{\alpha r}{p\theta}}$.

We mention that also in [92, Theorem 1.1] the interpolation between two grand Lebesgue spaces appear, but the methods used therein are different. They employed the complex interpolation functors defined by Calderón in 1964 ([21]), obtaining that the first complex interpolation spaces between two grand Lebesgue spaces coincides with the closure of L^∞ in intermediate grand Lebesgue spaces, while the second complex interpolation spaces between two grand Lebesgue spaces coincide with the intermediate grand Lebesgue spaces.

Let us now go back to the interpolation in the sense of Peetre, analyzing the corresponding results for small Lebesgue spaces. Using the duality result on interpolation spaces, (see [20, 161]), the following characterization of the interpolation spaces of small Lebesgue spaces hold.

THEOREM 10. ([59], Theorem 3.4) *If $0 < \theta < 1$, $1 < r < +\infty$, $1 < p < q$, $\alpha > 0$, then*

$$\left(L^{(p, \alpha)}, L^{(q, \alpha)} \right)_{\theta, r} = L^{p\theta, r} (\log L)^{\frac{\alpha}{p\theta}},$$

where $\frac{1}{p\theta} = \frac{1 - \theta}{p} + \frac{\theta}{q}$, $\frac{1}{p\theta} + \frac{1}{p'\theta} = 1$.

We highlight the following consequence of Theorem 9 and Theorem 10:

THEOREM 11. ([59], Theorem 3.5) *Let $1 < a < +\infty$, $\beta \in \mathbb{R}$, $\beta \neq 0$, $1 < r < +\infty$. The Lorentz-Zygmund space $L^{a,r}(\log L)^\beta$ is an interpolation space in the sense of Peetre of two Grand Lebesgue spaces if $\beta < 0$ and of two small Lebesgue spaces if $\beta > 0$.*

The results above concern interpolation between two grand or two small Lebesgue spaces. It may be interesting to consider also the off-diagonal case: this has been obtained in the following result, where the interpolation between small Lebesgue spaces $L^{(p)}$ and grand Lebesgue spaces $L^{(q)}$ when $p < q$ is considered.

THEOREM 12. ([59], Theorem 5.1) *If $0 < \theta < 1$, $1 \leq r < +\infty$, $p < q$, then*

$$(L^{(p)}, L^{(q)})_{\theta,r} = L^{p\theta,r}(\log L)^{\alpha_\theta},$$

where $\frac{1}{p\theta} = \frac{1-\theta}{p} + \frac{\theta}{q}$, $\alpha_\theta = 1 - \theta - \frac{1}{p\theta}$.

The proof of Theorem 12 is based of the following estimate of the K -functional.

THEOREM 13. ([59], Theorem 5.2) *If $1 < p < q < +\infty$, then for all $t > 0$, $f \in L^{(p)} + L^{(q)}$*

$$\begin{aligned} K(f, t; L^{(p)}, L^{(q)}) &\approx \int_0^{\varphi(t)} (1 - \log s)^{\frac{-1}{p}} \left(\int_0^s f_*^p(\tau) d\tau \right)^{\frac{1}{p}} \frac{ds}{s} \\ &\quad + (1 - \log t)^{\frac{p-1}{p}} \left(\int_0^{\varphi(t)} f_*^p(\tau) d\tau \right)^{\frac{1}{p}} \\ &\quad + t \sup_{\varphi(t) < s < 1} (1 - \log s)^{\frac{-1}{q}} \left(\int_s^1 f_*^q(\tau) d\tau \right)^{\frac{1}{q}}, \end{aligned}$$

where φ is an invertible function from $[0, 1]$ into itself satisfying the equivalence

$$\varphi(t)^{\frac{1}{p} - \frac{1}{q}} (1 - \log \varphi(t))^{\frac{p-q+pq}{pq}} \approx t.$$

The preceding study can be extended to the case where $p = q$, where the inverse function of $\psi_1(x) = (1 - \log x)^{-1}$, say $\varphi_1(t) = e^{1-\frac{1}{t}}$, will play a fundamental role in order to express the K -functional. Note that in this case the reiteration theorem from [14] cannot be used, and the computation must be made directly.

THEOREM 14. ([59], Theorem 6.1) *If $1 < p < +\infty$, $0 < t < 1$, $f \in L^{(p)} + L^{(p)}$, then*

$$\begin{aligned} K(f, t; L^{(p)}, L^{(p)}) &\approx \sup_{0 < s < e^{1-\frac{1}{t}}} (1 - \log s)^{\frac{-1}{p}} \left(\int_s^{e^{1-\frac{1}{t}}} f_*^p(x) dx \right)^{\frac{1}{p}} \\ &\quad + t \int_{e^{1-\frac{1}{t}}}^1 (1 - \log s)^{\frac{-1}{p}} \left(\int_{e^{1-\frac{1}{t}}}^s f_*^p(x) dx \right)^{\frac{1}{p}} \frac{ds}{s}. \end{aligned}$$

From Theorem 14 we may characterize the interpolation space $(L^p, L^p)_{\theta, r}$ as follows.

THEOREM 15. ([59], Theorem 6.2) *If $1 < p < +\infty$, $0 < \theta < 1$, $1 \leq r < +\infty$, then $Z_{\theta, r} := (L^p, L^p)_{\theta, r}$ has the following equivalent norm, namely, for $f \in Z_{\theta, r}$, $\beta_\theta = \theta - \frac{1}{p} - \frac{1}{r}$*

- if $\theta < \frac{1}{p}$ then

$$\|f\|_{Z_{\theta, r}} \approx \left[\int_0^1 \left[(1 - \log t)^{\beta_\theta} t \left(\int_t^1 f_*^p \right)^{\frac{1}{p}} \frac{dt}{t} \right]^r \right]^{\frac{1}{r}};$$

- if $\theta > \frac{1}{p}$ then $Z_{\theta, r} = G\Gamma(p, r, w)$, $w(t) = t^{-1}(1 - \log t)^{\beta_\theta r}$ and

$$\|f\|_{Z_{\theta, r}} \approx \left[\int_0^1 \left[(1 - \log t)^{\beta_\theta} \left(\int_0^t f_*^p \right)^{\frac{1}{p}} \right]^r \frac{dt}{t} \right]^{\frac{1}{r}};$$

- if $\theta = \frac{1}{p}$

$$\|f\|_{Z_{\theta, r}} \approx \left[\sum_{k=0}^{+\infty} \left(\int_{2^{1-2^{k+1}}}^{2^{1-2^k}} f_*^p \right)^{\frac{r}{p}} \right]^{\frac{1}{r}}.$$

In particular

$$(L^p, L^p)_{\frac{1}{p}, p} = L^p \quad \text{and} \quad (L^p, L^p)_{\theta, p} = L^{p, p}(\log L)^{\theta - \frac{1}{p}}.$$

We conclude with the following two results, where the quasinorm of the interpolation space $(L^p, L^p)_{\theta, r}$ is given in a unified form (see next Corollary 1) and in terms of the notion of Generalized Gamma space with double weights ([59, 84]), which is a further evolution of the $G\Gamma(p, m, w)$ spaces mentioned above in (12) (see next Theorem 16).

COROLLARY 1. ([59], Corollary 6.1) *If $\theta \in (0, 1)$, $1 \leq r < +\infty$ and $f \in Z_{\theta, r}$, then*

$$\|f\|_{Z_{\theta, r}}^r \approx \sum_{k \in \mathbb{N}} 2^{kr(\theta - \frac{1}{p})} \left(\int_{t_{k+1}}^{t_k} f_*^p(y) dy \right)^{\frac{r}{p}},$$

where $t_k = 2^{1-2^k}$, $k \in \mathbb{N}$.

THEOREM 16. ([59], Theorem 6.4) *If $1 < p < +\infty$, $0 < \theta < 1$, $1 \leq r < +\infty$, then*

$$(L^p, L^p)_{\theta, r} = G\Gamma(p, r; w_1, w_2),$$

where $w_1(t) = t^{-1}(1 - \log t)^{\theta r - 1}$, $w_2(t) = (1 - \log t)^{-1}$, $t \in (0, 1)$, i.e.

$$\|f\|_{Z_{\theta, r}} \approx \left[\int_0^1 (1 - \log t)^{\theta r} \left(\int_0^t (1 - \log x)^{-1} f_*^p(x) dx \right)^{\frac{r}{p}} \frac{dt}{(1 - \log t)t} \right]^{\frac{1}{r}}.$$

REFERENCES

- [1] R. A. ADAMS AND F. H. CLARKE, *Gross's logarithmic Sobolev inequality: a simple proof*, Amer. J. Math. **101** (1979), no. 6, 1265–1269.
- [2] R. A. ADAMS AND J. J. F. FOURNIER, *Sobolev spaces*, second ed., Pure and Applied Mathematics (Amsterdam), vol. 140, Elsevier/Academic Press, Amsterdam, 2003.
- [3] A. ALBERICO, T. ALBERICO AND C. SBORDONE, *Planar quasilinear elliptic equations with right-hand side in $L(\log L)^\delta$* , Discrete Contin. Dyn. Syst. **31** (2011), no. 4, 1053–1067.
- [4] A. ALBERICO, A. CIANCHI AND C. SBORDONE, *Gradient regularity for quasilinear elliptic Dirichlet problems in the plane*, Nonlinear Anal. **145** (2016), 143–161.
- [5] A. ALBERICO AND V. FERONE, *Regularity properties of solutions of elliptic equations in \mathbf{R}^2 in limit cases*, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. **6** (1995), no. 4, 237–250 (1996).
- [6] A. ALVINO, G. TROMBETTI AND P.-L. LIONS, *On optimization problems with prescribed rearrangements*, Nonlinear Anal. **13** (1989), no. 2, 185–220.
- [7] G. ANATRIELLO, *Iterated grand and small Lebesgue spaces*, Collect. Math. **65** (2014), no. 2, 273–284.
- [8] G. ANATRIELLO, R. CHILL AND A. FIORENZA, *Identification of fully measurable grand Lebesgue spaces*, J. Funct. Spaces (2017), Art. ID 3129186, 3.
- [9] G. ANATRIELLO AND A. FIORENZA, *Fully measurable grand Lebesgue spaces*, J. Math. Anal. Appl. **422** (2015), no. 2, 783–797.
- [10] G. ANATRIELLO AND M. R. FORMICA, *Weighted fully measurable grand Lebesgue spaces and the maximal theorem*, Ric. Mat. **65** (2016), no. 1, 221–233.
- [11] G. ANATRIELLO, M. R. FORMICA AND R. GIOVA, *Fully measurable small Lebesgue spaces*, J. Math. Anal. Appl. **447** (2017), no. 1, 550–563.
- [12] S. V. ASTASHKIN, K. V. LYKOV AND M. MASTYŁO, *On extrapolation of rearrangement invariant spaces*, Nonlinear Anal. **75** (2012), no. 5, 2735–2749.
- [13] H. ATTOUCH AND R. COMINETTI, *L^p approximation of variational problems in L^1 and L^∞* , Nonlinear Anal. **36** (1999), 373–399.
- [14] J. BASTERO AND F. RUIZ, *Elementary reverse Hölder type inequalities with application to operator interpolation theory*, Proc. A.M.S. **124** (1996), no. 10, 3183–3192.
- [15] C. BENNETT AND R. SHARPLEY, *Interpolation of operators*, Pure and Applied Mathematics, vol. 129, Academic Press, Inc., Boston, MA, 1988.
- [16] L. BOCCARDO AND T. GALLOUËT, *Nonlinear elliptic equations with right-hand side measures*, Comm. Partial Differential Equations **17** (1992), no. 3–4, 641–655.
- [17] H. BREZIS, T. CAZENAVE, Y. MARTEL AND A. RAMIANDRISOA, *Blow up for $u_t - \Delta u = g(u)$ revisited*, Adv. Differential Equations **1** (1996), no. 1, 73–90.
- [18] H. BREZIS, N. FUSCO AND C. SBORDONE, *Integrability for the Jacobian of orientation preserving mappings*, J. Funct. Anal. **115** (1993), no. 2, 425–431.
- [19] H. BREZIS AND F. MERLE, *Uniform estimates and blow-up behavior for solutions of $-\Delta u = V(x)e^u$ in two dimensions*, Comm. Partial Differential Equations **16** (1991), no. 8–9, 1223–1253.
- [20] YU. A. BRUDNYI, N. KRUGLJAK, *Interpolation functors and interpolation spaces* vol 1 Translated from Russian by N. Wadhwa. Mathematical Library 47 North-Holland Publishing, Amsterdam, 1991.
- [21] C. P. CALDERÓN, *Intermediate spaces and interpolation, the complex method*, Studia Math. **24** (1964), 113–190.

- [22] C. P. CALDERÓN, *Lacunary differentiability of functions in \mathbf{R}^n* , J. Approx. Theory **40** (1984), no. 2, 148–154.
- [23] C. CAPONE AND A. FIORENZA, *On small Lebesgue spaces*, J. Funct. Spaces Appl. **3** (2005), no. 1, 73–89.
- [24] C. CAPONE, A. FIORENZA AND G. E. KARADZHOV, *Grand Orlicz spaces and global integrability of the Jacobian*, Math. Scand. **102** (2008), no. 1, 131–148.
- [25] C. CAPONE AND M. R. FORMICA, *A decomposition of the dual space of some Banach function spaces*, J. Funct. Spaces Appl. (2012), Art. ID 737534, 10.
- [26] C. CAPONE, M. R. FORMICA AND R. GIOVA, *Grand Lebesgue spaces with respect to measurable functions*, Nonlinear Anal. **85** (2013), 125–131.
- [27] M. CAROZZA, G. MOSCARIELLO AND A. PASSARELLI DI NAPOLI, *Nonlinear equations with growth coefficients in BMO*, Houston J. Math. **28** (2002), no. 4, 917–929.
- [28] R. E. CASTILLO AND H. RAFEIRO, *Inequalities with conjugate exponents in grand Lebesgue spaces*, Hacet. J. Math. Stat. **44** (2015), no. 1, 33–39.
- [29] R. E. CASTILLO AND H. RAFEIRO, *An introductory course in Lebesgue spaces*, CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, Springer, 2016.
- [30] R. ČERNÝ, *Moser-Trudinger inequality in grand Lebesgue space*, Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. **26** (2015), no. 2, 177–188.
- [31] S. CHANILLO AND Y. Y. LI, *Continuity of solutions of uniformly elliptic equations in \mathbf{R}^2* , Manuscripta Math. **77** (1992), no. 4, 415–433.
- [32] A. CIANCHI, *A sharp embedding theorem for Orlicz-Sobolev spaces*, Indiana Univ. Math. J. **45** (1996), no. 1, 39–65.
- [33] A. CIANCHI, *Optimal Orlicz-Sobolev embeddings*, Rev. Mat. Iberoamericana **20** (2004), no. 2, 427–474.
- [34] A. CIANCHI, *Continuity properties of functions from Orlicz-Sobolev spaces and embedding theorems*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **23** (1996), no. 3, 575–608.
- [35] F. COBOS AND Ó. DOMÍNGUEZ, *Approximation spaces, limiting interpolation and Besov spaces*, J. Approx. Theory **189** (2015), 43–66.
- [36] F. COBOS AND T. KÜHN, *Extrapolation results of Lions-Peetre type*, Calc. Var. Partial Differential Equations **49** (2014), no. 1-2, 847–860.
- [37] F. COBOS AND J. PEETRE, *Interpolation of compactness using Aronszajn-Gagliardo functors*, Israel J. Math. **68** (1989), 220–240.
- [38] D. CRUZ-URIBE, A. FIORENZA AND O. M. GUZMÁN, *Embeddings between grand, small, and variable Lebesgue spaces*, Math. Notes **102** (2017), no. 5, 82–91.
- [39] D. CRUZ-URIBE AND A. FIORENZA, *Variable Lebesgue Spaces: Foundations and Harmonic Analysis*, Birkhäuser/Springer, Basel, 2013.
- [40] M. CWIKEL, *Real and complex interpolation and extrapolation of compact operators*, Duke Math. J. **65** (1992), no. 2, 333–343.
- [41] L. D’ONOFRIO, C. SBORDONE AND R. SCHIATTARELLA, *Measurability properties of a homeomorphism under minimal integrability conditions on the gradient*, Boll. Unione Mat. Ital. (9) **5** (2012), no. 3, 727–730.
- [42] L. D’ONOFRIO, C. SBORDONE AND R. SCHIATTARELLA, *Grand Sobolev spaces and their applications in geometric function theory and PDEs*, J. Fixed Point Theory Appl. **13** (2013), no. 2, 309–340.
- [43] L. M. DE CAVE, L. D’ONOFRIO AND R. SCHIATTARELLA, *Regularity results for quasilinear elliptic equations in the plane*, J. Elliptic Parabol. Equ. **1** (2015), 109–121.
- [44] L. M. DE CAVE, L. D’ONOFRIO AND R. SCHIATTARELLA, *Orlicz regularity of the gradient of solutions to quasilinear elliptic equations in the plane*, Bound. Value Probl. (2016), Paper No. 103, 15.
- [45] L. M. DE CAVE AND C. SBORDONE, *Gradient regularity for solutions to quasilinear elliptic equations in the plane*, J. Math. Anal. Appl. **417** (2014), no. 2, 537–551.
- [46] R. A. DEVORE AND R. C. SHARPLEY, *On the differentiability of functions in \mathbf{R}^n* , Proc. Amer. Math. Soc. **91** (1984), no. 2, 326–328.
- [47] G. DI FRATTA AND A. FIORENZA, *A direct approach to the duality of grand and small Lebesgue spaces*, Nonlinear Anal. **70** (2009), no. 7, 2582–2592.

- [48] L. DIENING, P. HARJULEHTO, P. HÄSTÖ AND M. RŮŽIČKA, *Lebesgue and Sobolev spaces with variable exponents*, Lecture Notes in Mathematics, vol. 2017, Springer, Heidelberg, 2011.
- [49] T. K. DONALDSON AND N. S. TRUDINGER, *Orlicz-Sobolev spaces and imbedding theorems*, J. Functional Analysis **8** (1971), 52–75.
- [50] J. I. DÍAZ AND J. M. RAKOTOSON, *On the differentiability of very weak solutions with right-hand side data integrable with respect to the distance to the boundary*, J. Funct. Anal. **257** (2009), no. 3, 807–831.
- [51] D. E. EDMUNDS, J. LANG AND O. MÉNDEZ, *Differential operators on spaces of variable integrability*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2014.
- [52] F. FARRONI AND R. GIOVA, *The distance to L^∞ in the grand Orlicz spaces*, J. Funct. Spaces Appl. (2013), Art. ID 658527, 7.
- [53] F. FARRONI, L. GRECO AND G. MOSCARIELLO, *Estimates for p -Laplace type equation in a limiting case*, Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. **25** (2014), no. 4, 445–448.
- [54] F. FARRONI, L. GRECO AND G. MOSCARIELLO, *Stability for p -Laplace type equation in a borderline case*, Nonlinear Anal. **116** (2015), 100–111.
- [55] F. FARRONI AND G. MOSCARIELLO, *A quantitative estimate for mappings of bounded inner distortion*, Calc. Var. Partial Differential Equations **51** (2014), no. 3–4, 657–676.
- [56] P. FERNÁNDEZ-MARTÍNEZ AND T. SIGNES, *Limit cases of reiteration theorems*, Math. Nachr. **288** (2015), no. 1, 25–47.
- [57] A. FIORENZA, *Duality and reflexivity in grand Lebesgue spaces*, Collect. Math. **51** (2000), no. 2, 131–148.
- [58] A. FIORENZA, M. R. FORMICA AND J. M. RAKOTOSON, *Pointwise estimates for GF -functions and applications*, Differential Integral Equations **30** (2017), no. 11–12, 809–824.
- [59] A. FIORENZA, M. R. FORMICA, A. GOGATISHVILI, T. KOPALIANI AND J. M. RAKOTOSON, *Characterization of interpolation between grand, small or classical Lebesgue spaces*, Preprint arXiv:1709.05892, Nonlinear Anal., to appear.
- [60] A. FIORENZA, B. GUPTA AND P. JAIN, *The maximal theorem for weighted grand Lebesgue spaces*, Studia Math. **188** (2008), no. 2, 123–133.
- [61] A. FIORENZA AND G. E. KARADZHOV, *Grand and small Lebesgue spaces and their analogs*, Z. Anal. Anwendungen **23** (2004), no. 4, 657–681.
- [62] A. FIORENZA, V. KOKILASHVILI AND A. MESKHI, *Hardy-Littlewood maximal operator in weighted grand variable exponent Lebesgue space*, Mediterr. J. Math. **14** (2017), no. 3, Art. 118, 20.
- [63] A. FIORENZA AND M. KRBEČ, *On an optimal decomposition in Zygmund spaces*, Georgian Math. J. **9** (2002), no. 2, 271–286.
- [64] A. FIORENZA, M. KRBEČ AND H.-J. SCHMEISSER, *An improvement of dimension-free Sobolev imbeddings in $r.i.$ spaces*, J. Funct. Anal. **267** (2014), no. 1, 243–261.
- [65] A. FIORENZA, A. MERCALDO AND J. M. RAKOTOSON, *Regularity and comparison results in grand Sobolev spaces for parabolic equations with measure data*, Appl. Math. Lett. **14** (2001), no. 8, 979–981.
- [66] A. FIORENZA, A. MERCALDO AND J. M. RAKOTOSON, *Regularity and uniqueness results in grand Sobolev spaces for parabolic equations with measure data*, Discrete Contin. Dyn. Syst. **8** (2002), no. 4, 893–906.
- [67] A. FIORENZA AND J. M. RAKOTOSON, *Petits espaces de Lebesgue et quelques applications*, C. R. Math. Acad. Sci. Paris **334** (2002), no. 1, 23–26.
- [68] A. FIORENZA AND J. M. RAKOTOSON, *New properties of small Lebesgue spaces and their applications*, Math. Ann. **326** (2003), no. 3, 543–561.
- [69] A. FIORENZA AND J. M. RAKOTOSON, *Compactness, interpolation inequalities for small Lebesgue-Sobolev spaces and applications*, Calc. Var. Partial Differential Equations **25** (2006), no. 2, 187–203.
- [70] A. FIORENZA AND J. M. RAKOTOSON, *Some estimates in $GF(p, m, w)$ spaces*, J. Math. Anal. Appl. **340** (2008), no. 2, 793–805.
- [71] A. FIORENZA, J. M. RAKOTOSON AND C. SBORDONE, *Variable exponents and grand Lebesgue spaces: some optimal results*, Commun. Contemp. Math. **17** (2015), no. 6, 1550023, 14.
- [72] A. FIORENZA, J. M. RAKOTOSON AND L. ZITOUNI, *Relative rearrangement method for estimating dual norms*, Indiana Univ. Math. J. **58** (2009), no. 3, 1127–1149.

- [73] A. FIORENZA AND C. SBORDONE, *Existence and uniqueness results for solutions of nonlinear equations with right hand side in L^1* , *Studia Math.* **127** (1998), no. 3, 223–231.
- [74] M. R. FORMICA AND R. GIOVA, *Boyd indices in generalized grand Lebesgue spaces and applications*, *Mediterr. J. Math.* **12** (2015), no. 3, 987–995.
- [75] S. FUČÍK AND A. KUFNER, *Nonlinear differential equations*, *Studies in Applied Mechanics*, vol. 2, Elsevier Scientific Publishing Co., Amsterdam-New York, 1980.
- [76] N. FUSCO, P.-L. LIONS AND C. SBORDONE, *Sobolev imbedding theorems in borderline cases*, *Proc. Amer. Math. Soc.* **124** (1996), no. 2, 561–565.
- [77] T. FUTAMURA, Y. MIZUTA AND T. OHNO, *Sobolev's theorem for Riesz potentials of functions in grand Morrey spaces of variable exponent*, *Banach and function spaces IV (ISBFS 2012)*, Yokohama Publ., Yokohama, 2014, pp. 353–365.
- [78] E. GAGLIARDO, *Interpolazione di spazi di Banach e applicazioni*, *Ricerche Mat.* **9** (1960), 58–81.
- [79] T. GALLOUËT AND R. HERBIN, *Existence of a solution to a coupled elliptic system*, *Appl. Math. Lett.* **7** (1994), no. 2, 49–55.
- [80] H. GAO, C. LIU AND H. TIAN, *A generalization of exponential class and its applications*, *Abstr. Appl. Anal.* (2013), Art. ID 476309, 8.
- [81] F. GIANNETTI, *Elliptic equations with right hand side in logarithmic Orlicz spaces*, *Rend. Accad. Sci. Fis. Mat. Napoli* (4) **64** (1997), 71–81 (1998).
- [82] F. GIANNETTI, T. IWANIEC, J. ONNINEN AND A. VERDE, *Estimates of Jacobians by subdeterminants*, *J. Geom. Anal.* **12** (2002), no. 2, 223–254.
- [83] A. GOGATISHVILI, C. AYKOL AND V. S. GULIYEV, *Characterization of associate spaces of generalized weighted weak-Lorentz spaces and embeddings*, *Studia Math.* **228** (2015), no. 3, 223–233.
- [84] A. GOGATISHVILI, M. KŘEPELA, L. PICK AND F. SOUDSKÝ, *Embeddings of Lorentz-type spaces involving weighted integral means*, *J. Funct. Anal.* **273** (2017), no. 9, 2939–2980.
- [85] A. GOGATISHVILI, B. OPIC AND W. TREBELS, *Limiting reiteration for real interpolation with slowly varying functions*, *Math. Nachr.* **278** (2005), no. 1-2, 86–107.
- [86] A. GOGATISHVILI, L. PICK AND F. SOUDSKÝ, *Characterization of associate spaces of weighted Lorentz spaces with applications*, *Studia Math.* **224** (2014), no. 1, 1–23.
- [87] L. GRECO, *A remark on the equality $\det Df = \text{Det} Df$* , *Differential Integral Equations* **6** (1993), no. 5, 1089–1100.
- [88] L. GRECO, T. IWANIEC AND C. SBORDONE, *Inverting the p -harmonic operator*, *Manuscripta Math.* **92** (1997), no. 2, 249–258.
- [89] L. GRECO, T. IWANIEC, C. SBORDONE AND B. STROFFOLINI, *Degree formulas for maps with nonintegrable Jacobian*, *Topol. Methods Nonlinear Anal.* **6** (1995), no. 1, 81–95.
- [90] L. GRECO, T. IWANIEC AND U. SUBRAMANIAN, *Another approach to biting convergence of Jacobians*, *Illinois J. Math.* **47** (2003), no. 3, 815–830.
- [91] P. HAJL ASZ, T. IWANIEC, J. MALÝ AND J. ONNINEN, *Weakly differentiable mappings between manifolds*, *Mem. Amer. Math. Soc.* **192** (2008), no. 899, viii+72.
- [92] D. I. HAKIM, M. IZUKI AND Y. SAWANO, *Complex interpolation of grand Lebesgue spaces*, *Monatsh Math.* **184** (2017), 245–272.
- [93] T. HOLMSTEDT, *Interpolation of quasi-normed spaces*, *Math. Scand.* **26** (1970), 177–199.
- [94] T. IWANIEC, P. KOSKELA AND J. ONNINEN, *Mappings of finite distortion: monotonicity and continuity*, *Invent. Math.* **144** (2001), no. 3, 507–531.
- [95] T. IWANIEC AND G. MARTIN, *Geometric function theory and non-linear analysis*, *Oxford Mathematical Monographs*, The Clarendon Press, Oxford University Press, New York, 2001.
- [96] T. IWANIEC AND C. SBORDONE, *On the integrability of the Jacobian under minimal hypotheses*, *Arch. Rational Mech. Anal.* **119** (1992), no. 2, 129–143.
- [97] T. IWANIEC AND C. SBORDONE, *New and old function spaces in the theory of PDEs and nonlinear analysis*, *Orlicz centenary volume*, *Banach Center Publ.*, vol. 64, Polish Acad. Sci. Inst. Math., Warsaw, 2004, pp. 85–104.
- [98] M. IZUKI, E. NAKAI AND Y. SAWANO, *Function spaces with variable exponents—an introduction—*, *Sci. Math. Jpn.* **77** (2014), no. 2, 187–315.
- [99] P. JAIN AND S. KUMARI, *On grand Lorentz spaces and the maximal operator*, *Georgian Math. J.* **19** (2012), no. 2, 235–246.
- [100] P. JAIN AND M. SINGH, *Hilbert inequality on grand function spaces*, *Ricerche Mat.* **to appear**.

- [101] P. JAIN, M. SINGH AND A. P. SINGH, *Hardy-type integral inequalities for quasi-monotone functions*, Georgian Math. J. (2016), to appear.
- [102] P. JAIN, M. SINGH AND A. P. SINGH, *Hardy type operators on grand Lebesgue spaces for non-increasing functions*, Trans. A. Razmadze Math. Inst. **170** (2016), no. 1, 34–46.
- [103] P. JAIN, M. SINGH AND A. P. SINGH, *Duality of fully measurable grand Lebesgue space*, Trans. A. Razmadze Math. Inst. **171** (2017), no. 1, 32–47.
- [104] P. JAIN, M. SINGH AND A. P. SINGH, *Integral operators on fully measurable weighted grand Lebesgue spaces*, Indag. Math. **28** (2017), 516–526.
- [105] P. JAIN, M. SINGH, A. P. SINGH AND V. D. STEPANOV, *Sawyer’s duality principle for grand Lebesgue spaces*, to appear.
- [106] L. KLEPRLÍK, *Composition operators on W^1X are necessarily induced by quasiconformal mappings*, Cent. Eur. J. Math. **12** (2014), no. 8, 1229–1238.
- [107] V. KOKILASHVILI AND A. MESKHI, *On weighted Bernstein type inequality in grand variable exponent Lebesgue spaces*, Math. Inequal. Appl. **18** (2015), no. 3, 991–1002.
- [108] V. KOKILASHVILI, A. MESKHI AND H. RAFEIRO, *Boundedness of commutators of singular and potential operators in generalized grand Morrey spaces and some applications*, Studia Math. **217** (2013), no. 2, 159–178.
- [109] V. KOKILASHVILI, A. MESKHI AND H. RAFEIRO, *Riesz type potential operators in generalized grand Morrey spaces*, Georgian Math. J. **20** (2013), no. 1, 43–64.
- [110] V. KOKILASHVILI, A. MESKHI AND H. RAFEIRO, *Estimates for nondivergence elliptic equations with VMO coefficients in generalized grand Morrey spaces*, Complex Var. Elliptic Equ. **59** (2014), no. 8, 1169–1184.
- [111] V. KOKILASHVILI, A. MESKHI AND H. RAFEIRO, *Grand Bochner-Lebesgue space and its associate space*, J. Funct. Anal. **266** (2014), no. 4, 2125–2136.
- [112] V. KOKILASHVILI, A. MESKHI, H. RAFEIRO AND S. SAMKO, *Integral operators in non-standard function spaces. Vol. 1*, Operator Theory: Advances and Applications, vol. 248, Birkhäuser/Springer, 2016, Variable exponent Lebesgue and amalgam spaces.
- [113] V. KOKILASHVILI, A. MESKHI, H. RAFEIRO AND S. SAMKO, *Integral operators in non-standard function spaces. Vol. 2*, Operator Theory: Advances and Applications, vol. 249, Birkhäuser/Springer, 2016, Variable exponent Hölder, Morrey-Campanato and grand spaces.
- [114] W. KONDRACHOV, *Sur certaines propriétés des fonctions dans l’espace*, C. R. (Doklady) Acad. Sci. URSS (N. S.) **48** (1945), 535–538.
- [115] P. KOSKELA AND X. ZHONG, *Minimal assumptions for the integrability of the Jacobian*, Ricerche Mat. **51** (2002), no. 2, 297–311 (2003).
- [116] E. H. LAAMRI AND D. MESKINE, *Existence and regularity of solutions to a class of elliptic reaction-diffusion systems with general nonlinearities*, to appear.
- [117] P. D. LAX AND A. N. MILGRAM, *Parabolic equations*, Contributions to the theory of partial differential equations, Annals of Mathematics Studies, no. 33, Princeton University Press, Princeton, N. J., 1954, pp. 167–190.
- [118] J. LERAY AND J.-L. LIONS, *Quelques résultats de Višik sur les problèmes elliptiques nonlinéaires par les méthodes de Minty-Browder*, Bull. Soc. Math. France **93** (1965), 97–107.
- [119] E. LIFLYAND, E. OSTROVSKY AND L. SIROTA, *Structural properties of bilateral grand Lebesgue spaces*, Turkish J. Math. **34** (2010), no. 2, 207–219.
- [120] J. LINDENSTRAUSS AND L. TZAFRIRI, *Classical Banach spaces. II*, Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas], vol. 97, Springer-Verlag, Berlin-New York, 1979, Function spaces.
- [121] J.-L. LIONS, *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod; Gauthier-Villars, Paris, 1969.
- [122] J.-L. LIONS AND J. PEETRE, *Propriétés d’espaces d’interpolation*, C. R. Acad. Sci. Paris **253** (1961), 1747–1749.
- [123] J.-L. LIONS AND J. PEETRE, *Sur une classe d’espaces d’interpolation*, Inst. Hautes Études Sci. Publ. Math. (1964), no. 19, 5–68.
- [124] W. A. J. LUXEMBURG AND A. C. ZAAANEN, *Riesz spaces. Vol. I*, North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., New York, 1971, North-Holland Mathematical Library.

- [125] F.-Y. MAEDA, Y. MIZUTA, T. OHNO AND T. SHIMOMURA, *Sobolev and Trudinger type inequalities on grand Musielak-Orlicz-Morrey spaces*, Ann. Acad. Sci. Fenn. Math. **40** (2015), no. 1, 403–426.
- [126] J. MARTÍN AND M. MILMAN, *Pointwise symmetrization inequalities for Sobolev functions and applications*, Adv. Math. **225** (2010), no. 1, 121–199.
- [127] J. MARTÍN AND M. MILMAN, *Fractional Sobolev inequalities: symmetrization, isoperimetry and interpolation*, Astérisque (2014), no. 366, 1–136.
- [128] J. MARTÍN AND M. MILMAN, *Integral isoperimetric transference and dimensionless Sobolev inequalities*, Rev. Mat. Complut. **28** (2015), no. 2, 359–392.
- [129] A. MESKHI, *Measure of non-compactness for integral operators in weighted Lebesgue spaces*, Nova Science Publishers, Inc., New York, 2009.
- [130] A. MESKHI, *Maximal functions, potentials and singular integrals in grand Morrey spaces*, Complex Var. Elliptic Equ. **56** (2011), no. 10-11, 1003–1019.
- [131] A. MIRANVILLE, A. PIETRUS AND J. M. RAKOTOSON, *Equivalence of formulation and uniqueness in a T -set*, Nonlinear Anal. T.M.A. **46** (2001), 609–627.
- [132] Y. MIZUTA AND T. OHNO, *Sobolev's theorem and duality for Herz-Morrey spaces of variable exponent*, Ann. Acad. Sci. Fenn. Math. **39** (2014), no. 1, 389–416.
- [133] Y. MIZUTA AND T. OHNO, *Trudinger's exponential integrability for Riesz potentials of functions in generalized grand Morrey spaces*, J. Math. Anal. Appl. **420** (2014), no. 1, 268–278.
- [134] G. MOSCARIELLO, *Existence and uniqueness for elliptic equations with lower-order terms*, Adv. Calc. Var. **4** (2011), no. 4, 421–444.
- [135] S. MÜLLER, *A surprising higher integrability property of mappings with positive determinant*, Bull. Amer. Math. Soc. (N.S.) **21** (1989), no. 2, 245–248.
- [136] T. OHNO AND T. SHIMOMURA, *Sobolev embeddings for Riesz potentials of functions in grand Morrey spaces of variable exponents over non-doubling measure spaces*, Czechoslovak Math. J. **64(139)** (2014), no. 1, 209–228.
- [137] A. PASSARELLI DI NAPOLI AND C. SBORDONE, *Elliptic equations with right hand side in $L(\log L)^\alpha$* , Rend. Accad. Sci. Fis. Mat. Napoli (4) **62** (1995), 301–314 (1996).
- [138] J. PEETRE, *Espaces d'interpolation et théorème de Soboleff*, Ann. Inst. Fourier (Grenoble) **16** (1966), no. fasc. 1, 279–317.
- [139] L. PICK, A. KUFNER, O. JOHN AND S. FUČÍK, *Function spaces. Vol. 1*, extended ed., De Gruyter Series in Nonlinear Analysis and Applications, vol. 14, Walter de Gruyter & Co., Berlin, 2013.
- [140] A. PRIGNET, *Remarks on existence and uniqueness of solutions of elliptic problems with right-hand side measures*, Rend. Mat. Appl. (7) **15** (1995), no. 3, 321–337.
- [141] T. RADICE AND G. ZECCA, *Existence and uniqueness for nonlinear elliptic equations with unbounded coefficients*, Ric. Mat. **63** (2014), no. 2, 355–367.
- [142] T. RADICE AND G. ZECCA, *The maximum principle of Alexandrov for very weak solutions*, J. Differential Equations **256** (2014), no. 3, 1133–1150.
- [143] H. RAFEIRO, *A note on boundedness of operators in grand grand Morrey spaces*, Advances in harmonic analysis and operator theory, Oper. Theory Adv. Appl., vol. 229, Birkhäuser/Springer Basel AG, Basel, 2013, pp. 349–356.
- [144] J. M. RAKOTOSON, *Some new applications of the pointwise relations for the relative rearrangement*, Adv. Differential Equations **7** (2002), no. 5, 617–640.
- [145] J. M. RAKOTOSON, *Réarrangement relatif*, Mathématiques & Applications (Berlin) [Mathematics & Applications], vol. 64, Springer, Berlin, 2008, Un instrument d'estimations dans les problèmes aux limites. [An estimation tool for limit problems].
- [146] J. M. RAKOTOSON, *New Hardy inequalities and behaviour of linear elliptic equations*, J. Funct. Anal. **263** (2012), no. 9, 2893–2920.
- [147] J. M. RAKOTOSON, *Linear equation with data in non standard spaces*, Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. **26** (2015), no. 3, 241–262.
- [148] F. RELICH, *Ein Satz über mittlere Konvergenz*, Nachr. Akad. Wiss. Göttingen. Math.-Phys. Kl. **1930** (1930), 30–35.
- [149] M. RŮŽIČKA, *Electrorheological fluids: modeling and mathematical theory*, Lecture Notes in Mathematics, vol. 1748, Springer-Verlag, Berlin, 2000.
- [150] S. G. SAMKO AND S. M. UMARKHADZHIEV, *On Iwaniec-Sbordone spaces on sets which may have infinite measure*, Azerb. J. Math. **1** (2011), no. 1, 67–84.

- [151] S. G. SAMKO AND S. M. UMARKHADZHIEV, *On Iwaniec-Sbordone spaces on sets which may have infinite measure: addendum*, Azerb. J. Math. **1** (2011), no. 2, 2.
- [152] C. SBORDONE, *Grand Sobolev spaces and their applications to variational problems*, Matematiche (Catania) **51** (1996), no. 2, 335–347 (1997).
- [153] C. SBORDONE, *New estimates for div-curl products and very weak solutions of PDEs*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **25** (1997), no. 3-4, 739–756 (1998), Dedicated to Ennio De Giorgi.
- [154] J. SERRIN, *Pathological solutions of elliptic differential equations*, Ann. Scuola Norm. Sup. Pisa (3) **18** (1964), 385–387.
- [155] R. E. SHOWALTER, *Monotone operators in Banach space and nonlinear partial differential equations*, Mathematical Surveys and Monographs, vol. 49, American Mathematical Society, Providence, RI, 1997.
- [156] G. STAMPACCHIA, *Some limit cases of L^p -estimates for solutions of second order elliptic equations*, Comm. Pure Appl. Math. **16** (1963), 505–510.
- [157] G. STAMPACCHIA, *Formes bilinéaires coercitives sur les ensembles convexes*, C. R. Acad. Sci. Paris **258** (1964), 4413–4416.
- [158] G. STAMPACCHIA, *Le problème de Dirichlet pour les équations elliptiques du second ordre à coefficients discontinus*, Ann. Inst. Fourier (Grenoble) **15** (1965), no. fasc. 1, 189–258.
- [159] E. M. STEIN, *Editor’s note: the differentiability of functions in \mathbf{R}^n* , Ann. of Math. (2) **113** (1981), no. 2, 383–385.
- [160] E. M. STEIN AND G. WEISS, *Introduction to Fourier analysis on Euclidean spaces*, Princeton University Press, Princeton, N.J., 1971, Princeton Mathematical Series, No. 32.
- [161] L. TARTAR, *An introduction to Sobolev spaces and interpolation spaces*, 2007, Springer.
- [162] H. TRIEBEL, *Theory of function spaces*, Modern Birkhäuser Classics, Birkhäuser/Springer Basel AG, Basel, 2010, Reprint of 1983 edition, also published in 1983 by Birkhäuser Verlag.
- [163] S. M. UMARKHADZHIEV, *Generalization of the notion of grand Lebesgue space*, Russian Math. (Iz. VUZ) **58** (2014), no. 4, 35–43, Translation of Izv. Vyssh. Uchebn. Zaved. Mat. **2014**, no. 4, 42–51.
- [164] X. F. YE, *Boundedness of commutators of singular and potential operators in grand Morrey spaces*, Acta Math. Sinica (Chin. Ser.) **54** (2011), no. 2, 343–352.
- [165] G. ZECCA, *Regularity results for planar quasilinear equations*, Rend. Mat. Appl. (7) **30** (2010), no. 3-4, 329–343.
- [166] G. ZECCA, *Existence and uniqueness for nonlinear elliptic equations with lower-order terms*, Nonlinear Anal. **75** (2012), no. 2, 899–912.

Alberto Fiorenza, Dipartimento di Architettura, Università di Napoli Federico II, via Monteoliveto, 3, I-80134 Napoli, Italy
and
Istituto per le Applicazioni del Calcolo “Mauro Picone”, Consiglio Nazionale delle Ricerche, via Pietro Castellino, 111 I-80131 Napoli, Italy,
e-mail: fiorenza@unina.it

Maria Rosaria Formica, Università degli Studi di Napoli Parthenope, via Generale Parisi 13, 80132, Napoli, Italy,
e-mail: mara.formica@uniparthenope.it

Amiran Gogatishvili, Institute of Mathematics of the Czech Academy of Sciences, Žitná 25, 115 67 Prague 1, Czech Republic
e-mail: gogatish@math.cas.cz

Corresponding Author: Alberto Fiorenza