

**J. Malíček: Středoevropské pralesy a lišeňíky I. Příklady nejcennějších lokalit a ekologie lesních lišeňíků (Živa 2017, 4: 152–155)**

**Použitá a doporučená literatura:**

- Albrecht J. et al. (2003): Českobudějovicko [Českobudějovicko region] – In: Mackovčin P., Sedláček M. (eds), Chráněná území ČR, svazek VIII [Protected areas of the Czech Republic, vol. VIII], AOPK ČR, Brno.
- Bässler C., Cadotte M.W., Beudert B., Heibl C., Blaschke M., Bradtka J.H., Langbehn T., Werth S. & Müller J. (2016): Contrasting patterns of lichen functional diversity and species richness across an elevation gradient. – Ecography 39: 689–698.
- Cieśliński S. & Tobolewski Z. (1988): Porosty (Lichenes) Puszczy Białowieskiej i jej zachodniego przedpola [Lichens (Lichenes) of the Białowieża Forest and its western foreland]. – Phytocenosis 1 (N.S.), Suppl. Cartographiae Geobotanicae 1: 1–116.
- Esseen P.-A. (2006): Edge influence on the old-growth forest indicator lichen *Alectoria sarmentosa* in natural ecotones. – Journal of Vegetation Science, 17: 185–194.
- Esseen P.-A., Renhorn K.-E. & Pettersson R.B. (1996): Epiphytic lichen biomass in managed and old-growth boreal forests: effect of branch quality. – Ecological Applications, 6: 228–238.
- Hauck M., de Bruyn U. & Leuschner C. (2013): Dramatic diversity losses in epiphytic lichens in temperate broad-leaved forests during the last 150 years. – Biological Conservation, 157: 136–145.
- Hochebner T., Frank G. & Rotheneder G. (2015): Monitoring der Spechte (Picidae) im Wildnisgebiet Dürrenstein. – Silva Fera, 4: 41–69.
- Holec J., Kříž M., Pouzar Z., Šandová M. (2015): Boubínský prales virgin forest, a Central European refugium of boreal-montane and old-growth forest fungi. – Czech Mycology, 67: 157–226.
- Kuusinen M. (1996): Cyanobacterial macrolichens on *Populus tremula* as indicators of forest continuity in Finland. – Biological Conservation, 75: 43–49.
- Löbel S., Snäll T. & Rydin H. (2006): Species richness patterns and metapopulation process - evidence from epiphyte communities in boreo-nemoral forests. – Ecography, 29: 169–182.
- Ložek V. (2004): Středoevropské bezlesí v čase a prostoru, IV. Vývoj v poledové době. – Ochrana přírody, 59: 99–105.
- Mackovčin P., Jatiová M., Demek J., Slavík P. a kol. (2007): Brněnsko. – In: Mackovčin P. & Sedláček M. (eds.), Chráněná území ČR, svazek IX. AOPK ČR a EkoCentrum Brno, Praha, 932 pp.
- Nascimbene J., Brunialti G., Ravera S., Frati L. & Caniglia G. (2010a): Testing *Lobaria pulmonaria* (L.) Hoffm. as an indicator of lichen conservation importance of Italian forests. – Ecological Indicators, 10: 353–360.

- Nascimbene J., Thor G. & Nimis P.L. (2013): Effects of forest management on epiphytic lichens in temperate deciduous forests of Europe –Areview. – Forest Ecology and Management, 298: 27–38.
- Nilsson S.G., Arup U., Baranowski R. & Ekman S. (1995): Tree-Dependent Lichens and Beetles as Indicators in Conservation Forests. – Conservation Biology, 9: 1208–1215.
- Paillet Y., Bergès L., Hjältén J., Ódor P., Avon C., Bernhardt-Römermann M., Bijlsma R.J., De Bruyn L., Fuhr M., Grandin U., Kanka R., Lundin L., Luque S., Magura T., Matesanz S., Mészáros I., Sebastià M.T., Schmidt W., Standovár T., Tóthmérész B., Uotila A., Valladares F., Vellak K. & Virtanen R. (2010): Biodiversity Differences between Managed and Unmanaged Forests: Meta-Analysis of Species Richness in Europe. – Conservation Biology, 24: 101–112.
- Pokorný P. & Kuneš P. (2005): Holocene acidification process recorded in three pollen profiles from Czech sandstone and river terrace environments. – Ferrantia, 44: 107–113.
- Pokorný P. (2011): Neklidné časy. Kapitoly ze společných dějin přírody a lidí. – Dokorán, Praha.
- Radkau J. (2011): Wood: a history. – Polity Press, Cambridge.
- Svoboda D., Peksa O. & Veselá J. (2010): Epiphytic lichen diversity in central European oak forests: Assessment of the effects of natural environmental factors and human influences. – Environmental Pollution, 158: 812–819.
- UNESCO (2013): Primeval Beech Forests of the Carpathians and the Ancient Beech Forests of Germany (Slovakia/Germany/Ukraine) – In: World heritage WHC, p. 60–13/37.COM.7B.Add.
- Vrška T., Šamonil P., Unar P., Hort L., Adam D., Král K., Janík D. (2012): Development dynamics of virgin forest reserves in the Czech Republic, vol. 3, Šumava Mts. and Český les Mts. – Diana, Stožec, Boubín virgin forest, Milešice virgin forest). – Academia, Praha.
- Wirth V., Hauck M., de Bruyn U., Schiefelbein U., John V. & Otte V. (2009): Flechten aus Deutschland mit Verbreitungsschwerpunkt im Wald. – Herzogia, 22: 79–107.
- Zedda L. (2002): The epiphytic lichens on *Quercus* in Sardinia (Italy) and their value as ecological indicators. – Englera, 24: 1–457.

[www.pralesy.cz](http://www.pralesy.cz)

[www.pralesy.sk](http://www.pralesy.sk)