Reparace DNA a efekty HDAC a PRMT1 inhibitorů

Eva Bártová Institute of Biophysics Academy of Sciences of the Czech Republic Brno

DNA repair

Single-strand damage
❖Base_excision_repair (BER), which_repairs
damage to a single base caused by oxidation, Single-strand damage

*Base excision repair (BER), which repairs

damage to a single base caused by oxidation,

alkylation, hydrolysis, or deamination.

*Nucleotide, excision, repair, (NEB), which Single-strand damage

❖ Base excision repair (BER), which repairs

damage to a single base caused by oxidation,

alkylation, hydrolysis, or deamination.
◆ Nucleotide excision repair (NER), which

recognizes bully, belix d Single-strand damage

*Base excision repair (BER), which repairs

damage to a single base caused by oxidation,

alkylation, hydrolysis, or deamination.

*Nucleotide excision repair (NER), which

recognizes bulky, helix-dis

Single-strand damage

❖ Base excision repair (BER), which repairs

damage to a single base caused by oxidation,

alkylation, hydrolysis, or deamination.

❖ Nucleotide excision repair (NER), which

recognizes bulky, helix-Single-strand damage

❖ Base excision repair (BER), which repairs

damage to a single base caused by oxidation,

alkylation, hydrolysis, or deamination.

❖ Nucleotide excision repair (NER), which

recognizes bulky, helix-Single-strand damage

*Base excision repair (BER), which repairs

damage to a single base caused by oxidation,

alkylation, hydrolysis, or deamination.

*Nucleotide excision repair (NER), which

recognizes bulky, helix-dis

Single-strand damage

*Base excision repair (BER), which repairs

damage to a single base caused by oxidation,

alkylation, hydrolysis, or deamination.

*Nucleotide excision repair (NER), which

recognizes bulky, helix-dis Single-strand damage

*Base excision repair (BER), which repairs

damage to a single base caused by oxidation,

alkylation, hydrolysis, or deamination.

*Nucleotide excision repair (NER), which

recognizes bulky, helix-dis Single-strand damage

*Base excision repair (BER), which repairs

damage to a single base caused by oxidation,

alkylation, hydrolysis, or deamination.

*Nucleotide excision repair (NER), which

recognizes bulky, helix-dis nucleotides. In the excision repair (BER), which repairs

damage to a single base caused by oxidation,

alkylation, hydrolysis, or deamination.

A Nucleotide excision repair (NER), which

recognizes bulky, helix-distorting lesions damage to a single base caused by oxidation,
alkylation, hydrolysis, or deamination.

◆Nucleotide excision repair (NER), which
recognizes bulky, helix-distorting lesions such
as pyrimidine dimers and 6,4 photoproducts.

◆ A Nucleotide excision repair (NER), which
recognizes bulky, helix-distorting lesions such
as pyrimidine dimers and 6,4 photoproducts.
A Mismatch repair (MMR), which corrects
errors of DNA replication and recombination
tha as pyrimidine dimers and 6,4 photoproducts.

◆Mismatch repair (MMR), which corrects

errors of DNA replication and recombination

that result in mispaired (but undamaged)

nucleotides.

Double-strand breaks

◆non-homo

(MMEJ)

http://commons.wikimedia.org/wiki/File:DNA_Repair.jpg

Experiments: Jana Suchánková and Gabriela Šustáčková

GFP-HP1 β / mCherry-Lamin A

Experiments: Petra Sehnalová and Eva Bártová

53BP1

$HP1\beta$ and A-type lamins

 $HP1\beta$

UBF1/2 YFP-PRMT1

Šustáčková et al., JCP (2011) and experiments of Petra Sehnalová and Soňa Legartová

 $0 \quad \mu m \quad 5$

GFP-HP1ß/yH2AX/Nucleus

Šustáčková et al., JCP (2011)

European Journal of Histochemistry 2014; volume 58:2389

PRMT1 arginine methyltransferase accumulates in cytoplasmic bodies that respond to selective inhibition and DNA damage

J. Suchánková,¹ S. Legartová,¹

P. Sehnalová,¹ S. Kozubek,¹ S. Valente,² D. Labella,² A. Mai,² C. Eckerich,³ F.O. Fackelmayer,³ D.V. Sorokin,⁴ E. Bártová¹

¹Institute of Biophysics, Academy of Sciences of the Czech Republic, Brno, **Czech Republic**

²Pasteur Institute - Fondazione Cenci Bolognetti, Department of Chemistry and Drug Technology, University of Rome La Sapienza, Italy

³Institute of Molecular Biology and **Biotechnology, Department of Biomedical** Research (IMBB-FORTH), Foundation for Research and Technology Hellas, Ioannina, Greece

⁴Faculty of Informatics, Masaryk University, Brno, Czech Republic respond to DNA injury in the cell nucleus, and to treatment with various PRMT1 inhibitors.

Introduction

Chromatin structure and function is controlled by many enzymes.¹ Protein arginine methyltransferases (PRMTs) methylate histones and other regulatory and structural proteins, with particular activity in the nucleus.^{2,3} The PRMT family consists of 11 different methyltransferases (PRMT1-11) that control cellular processes such as transcription, RNA processing, nucleocytoplasmic shuttling of proteins, and DNA repair.⁴⁷ Reflecting these diverse functions, several PRMTs are located in both the cytoplasm and the nucleus, but display cell-type-specific differences in the ratio of nuclear versus cytoplasmic PRMTs' distribution.⁸ Arginine methyltransferases in the nucleus act as epigenetic factors that induce transcriptional activation or silencing depending on the affected residue in core histones, and the symmetric or asymmetric nature of the methylation.⁹ For example, PRMT1 and PRMT5 can both dimethylate arginine 3 of histone H4 (H4R3). However, PRMT5 methylates H4R3

Correspondence: Eva Bártová, Institute of Biophysics, Academy of Sciences of the Czech Republic, v.v.i., Královopolská 135, CZ-612 65, Brno, Czech Republic. Tel. +420.5.41517141 - Fax: +420.5.41240498. E-mail: bartova@ibp.cz

Keywords: Epigenetics, PRMTs, epi-drugs, arginine methylation, DNA repair.

Conflict of interests: the authors declare no potential conflict of interests.

Funding: our work was supported by the following agencies: Grant Agency of Czech Republic (Grant N. P302/10/1022, P302/12/G157, and 13-07822S). EB, FOF, and AM are members of the EU-COST Action TD0905; EB is a principal investigator and coordinator of the EU Marie Curie Project PIRS-ES-GA-2010-269156-LCS. The postdoctoral fellowship of DVS was guaranteed by the Education for Competitiveness Operational Programme (ECOP), N. CZ.1.07/2.3.00/30.0030.

Contributions: JS, cell culture cultivation, treatments, plasmid DNA isolation, cell transfection, live cell studies after UV-A irradiation, immunofluorescence studies; SL, immunofluorescence and intermediated irradiation of the cells with yrays; SK, GACR projects P302/10/1022 and P302/12/G157 coordination; CE, provided YFP-PRMT1 plasmid DNA; FOF, manuscript contribu-

YFP-PRMT1 in U2OS cells

Disassembly of PRMTsomes

Conclusions

Conclusions
1. HP1β function at DNA lesions can be
affected by HDAC inhibitors.
2. Irradiation of the cell nucleus by UVA
lasers caused disappearance of PRMT1.

1. HP1β function at DNA lesions can be

affected by HDAC inhibitors.

2. Irradiation of the cell nucleus by UVA

lasers caused disappearance of PRMT1-

positive cytoplasmic bodies

3. Disappearance of PRMT1-positive

cy 2. Irradiation of the cell nucleus by UVA

lasers caused disappearance of PRMT1-

positive cytoplasmic bodies

3. Disappearance of PRMT1-positive

cytoplasmic bodies is really fast afte γ-

irradition or the cell treatmen

Conclusions
1. HP1β function at DNA lesions can be
affected by HDAC inhibitors.
2. Irradiation of the cell nucleus by UVA 2. Irradiation of the cell nucleus by UVA
2. Irradiation of the cell nucleus by UVA
2. Irradiation of the cell nucleus by UVA
1
positive cytoplasmic bodies **Conclusions**

1. HP1β function at DNA lesions can be

affected by HDAC inhibitors.

2. Irradiation of the cell nucleus by UVA

lasers caused disappearance of PRMT1-

positive cytoplasmic bodies

3. Disappearance of PRMT1-**Conclusions**

1. HP1β function at DNA lesions can be

affected by HDAC inhibitors.

2. Irradiation of the cell nucleus by UVA

lasers caused disappearance of PRMT1-

positive cytoplasmic bodies

3. Disappearance of PRMT1-**Conclusions**

1. HP1β function at DNA lesions can be

affected by HDAC inhibitors.

2. Irradiation of the cell nucleus by UVA

lasers caused disappearance of PRMT1-

positive cytoplasmic bodies

3. Disappearance of PRMT1 Conclusions
1. HP1β function at DNA lesions can be
affected by HDAC inhibitors.
2. Irradiation of the cell nucleus by UVA
lasers caused disappearance of PRMT1-
positive cytoplasmic bodies
3. Disappearance of PRMT1-positiv Conclusions

1. HP1β function at DNA lesions can be

affected by HDAC inhibitors.

2. Irradiation of the cell nucleus by UVA

lasers caused disappearance of PRMT1-

positive cytoplasmic bodies

3. Disappearance of PRMT1-po 1. HPTP function at DNA lesions can be
affected by HDAC inhibitors.
2. Irradiation of the cell nucleus by UVA
lasers caused disappearance of PRMT1-
positive cytoplasmic bodies
3. Disappearance of PRMT1-positive
cytoplasmi Example the communitiests.

2. Irradiation of the cell nucleus by UVA

lasers caused disappearance of PRMT1-

positive cytoplasmic bodies

3. Disappearance of PRMT1-positive

cytoplasmic bodies is really fast afte y-

irra

Scanning in 2D and 3D by confocal microscope

Laser beam moves firstly along x axis and then starts with new line in y axis.

Finishing scanning of one thin optical slice in xy plane, the scanning plane is moved in z axis to other slice

 $^{\circ}$ pos=1

Deconvolution

a

a

ig. 3: Via deconvolution, artefacts can be computed out of fluorescence images, a). These artefacts

are caused by the stray light from non-focused areas above and below the focus level. These phenom-

and referred t

Subdiffraction Multicolor Imaging of the Nuclear Periphery with 3D **Structured Illumination Microscopy**

Lothar Schermelleh,^{1*} Peter M. Carlton,^{2*} Sebastian Haase,^{2,4} Lin Shao,² Lukman Winoto,² Peter Kner,² Brian Burke,³ M. Cristina Cardoso,⁴ David A. Agard,²
Mats G. L. Gustafsson,⁵ Heinrich Leonhardt,¹*† John W. Sedat²*†

Fig. 1. Subdiffraction resolution imaging with 3D-SIM. (A and **B**) Cross section through a DAPI-stained C2C12 cell nucleus acquired with conventional wide-field illumination (A) and with structured illumination (B), showing the striped interference pattern (inset). The renderings to the right illustrate the respective support of detection in frequency space. The axes k_n , k_n and k_n indicate spatial frequencies along the x , y , and z directions. The surfaces of the renderings represent the corresponding resolution limit. The depression of the frequency support ("missing cone") in z direction in (A) indicates the restriction in axial resolution of conventional wide-field microscopy. With 3D-SIM, the axial support is extended but remains within the resolution limit. (C) Five phases of the sine wave pattern are recorded at each z position, allowing the shifted components to be separated and returned to their proper location in frequency space. Three image stacks are recorded with the diffraction grating sequentially rotated into three positions 60° apart. resulting in nearly rotationally symmetric support over a larger region of frequency space. (D) The same cross section of the reconstructed 3D-SIM image shows enhanced image details compared with the original image (insets). The increase in resolution is shown in frequency space on the right, with the coverage extending two times farther from the origin. Scale bars indicate 5 um.

Stimulated Emission Depletion microscopy, or STED
microscopy, is a fluorescence microscopy technique that
uses the non-linear de-excitation of fluorescent dyes to Stimulated Emission Depletion microscopy, or STED
microscopy, is a fluorescence microscopy technique that
uses the non-linear de-excitation of fluorescent dyes to
overcome the resolution limit imposed by diffraction with Stimulated Emission Depletion microscopy, or STED
microscopy, is a fluorescence microscopy technique that
uses the non-linear de-excitation of fluorescent dyes to
overcome the resolution limit imposed by diffraction with
s Stimulated Emission Depletion microscopy, or STED
microscopy, is a fluorescence microscopy technique that
uses the non-linear de-excitation of fluorescent dyes to
overcome the resolution limit imposed by diffraction with
s Stimulated Emission Depletion microscopy, or STED
microscopy, is a fluorescence microscopy technique that
uses the non-linear de-excitation of fluorescent dyes to
overcome the resolution limit imposed by diffraction with
s Stimulated Emission Depletion microscopy, or STED
microscopy, is a fluorescence microscopy technique that
uses the non-linear de-excitation of fluorescent dyes to
overcome the resolution limit imposed by diffraction with
s

STED

High-speed live-cell imaging

Li et al., Science (2015)

Leica STED – STimulated Emission Depletion SUPERRESOLUTION (subdiffraction) in xy plane

Hell, S. W. and J. Wichmann (1994). Opt. Lett. "Breaking the diffraction resolution limit by stimulated emission"

neurobiology membrane biology membrane rafts intracellular transport

Willig Kiret al. *Natale 2006* Lin W et al. *PNAS* 2007
Kittel RJ et al. *Science* 2006 Seebach J *Cardiovas. Res.* 2007
Fitzner D et al. *EMBO* J 2006 Sieber JJ *Science* 2007

Kimura and Cook (2001)

Types of fluorochromes

sp. (Dendra2)

 -0.5

 \circ

 $[µm]$

 0.5

 $\overline{1}$

 $\overline{1}$

 $\left[\,\mu\mathrm{m}\right]$

Experiments of Veronika Foltanková and Dmitry Sorokin

Dendra2 photo-conversion

Dendra2 is an improved version of a green-to-red photoswitchable
fluorescent protein Dendra, derived from octocoral
Dendronephthya sp. (Gurskaya *et al.*, 2006). 2 is an improved version of a green-to-red photoswitchable
fluorescent protein Dendra, derived from octocoral
Dendronephthya sp. (Gurskaya *et al.*, 2006). Dendronephthya sp. (Gurskaya et al., 2006).

Normalized excitation (thin line) and emission (thick line) for non-activated (green) and activated (red) spectra.

Dendra2-H4 photoconversion

H4-Dendra2 / H4-Dendra2

FRET (Fluorescence Resonance Energy Transfer) is a technique
for measuring interactions between two proteins in vivo. In this
technique, two different fluorescent molecules (fluorophores) are FRET (Fluorescence Resonance Energy Transfer) is a technique
for measuring interactions between two proteins in vivo. In this
technique, two different fluorescent molecules (fluorophores) are
genetically fused the two prot FRET (Fluorescence Resonance Energy Transfer) is a technique
for measuring interactions between two proteins in vivo. In this
technique, two different fluorescent molecules (fluorophores) are
genetically fused the two prot FRET (Fluorescence Resonance Energy Transfer) is a technique
for measuring interactions between two proteins in vivo. In this
technique, two different fluorescent molecules (fluorophores) are
genetically fused the two prot

7/8/2010 9:30:30 AM

5/20/2010 2:01:52 PM

t=0.000sec

B

FLIM-FRET analysis

http://www.celanphy.science.ru.nl/ Bruce%20web/construction.htm

EB group, IBP, Brno

Inactivation of X chromosome in hESCs in comparison to MEFs

3D-FISH a konfokální mikroskopie

Maximální obraz Všech řezů

Galerie optických řezů

3D reconstrukce CT Weierich et al., (2003) in press

K562 cells t(9;22)

E. Bártová et al. / Leukemia Research 29 (2005) 901-913

Bártová et al., Differentiation (2008)

Lenka Stixová, Sona Legartová, Petra Sehnalová, Jana Suchánková, Dmitry V. Sorokin

