

# University of Granada

Department of Applied Mathematics

Date: June 7, 2013

I HEREBY RECOMMEND THAT THE THESIS PREPARED UNDER MY  
SUPERVISION BY

**Manuel Zamora**

ENTITLED

**Boundary value problems for differential equations with  
singularities**

BE ACCEPTED IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR  
THE DEGREE OF

**DOCTOR IN MATHEMATICS SCIENCES**

---

Thesis Advisors Dr. Robert  
Hakl and Pedro J. Torres

---

Chairman  
Dr. Irena Rachunkova

---

Thesis Reader Dr. Antonio  
Ureña

---

Thesis Reader Dr. Milan  
Tvrđy

---

Thesis Reader Dr. Alessan-  
dro Fonda

---

Thesis Reader Dr. Franco  
Leis, Daniel

# Boundary value problems for differential equations with singularities

By

Manuel Zamora

PhD Dissertation

Submitted in Partial Fulfillment of the Requirements  
for the Degree of Doctor in Mathematics  
in the Department of Applied Mathematics at  
University of Granada, 2013

Granada, Spain

Con el fin de obtener la mención internacional en el título de doctor se han cumplido los siguientes requisitos:

- La tesis está redactada en inglés.
- Uno de los miembros del tribunal propuesto es de una universidad extranjera.
- La defensa se hará en inglés con resumen y conclusiones en español.
- Dos informes de expertos extranjeros han avalado esta tesis.
- Una parte de la tesis se ha realizado en Instituto de Matemáticas de Brno (República Checa) bajo la supervisión del profesor R. Hakl.

Me gustaría agradecer a las personas que me han ayudado en este periodo, en especial a C. Bereanu, A. Fonda, R. Hakl, R. Manásevich, P. J. Torres, y A. J. Ureña.

El doctorando MANUEL ZAMORA y los directores de la tesis PEDRO J. TORRES y ROBERT HAKL garantizamos, al firmar esta tesis doctoral, que el trabajo ha sido realizado por el doctorando bajo la dirección de los directores de la tesis y hasta donde nuestro conocimiento alcanza, en la realización del trabajo, se han respetado los derechos de otros autores a ser citados, cuando se han utilizado sus resultados o publicaciones.

En Granada, 7 de junio de 2013.

Director/es de la Tesis

Fdo.:

Doctorando

Fdo.:

# Boundary value problems for differential equations with singularities

Manuel Zamora

Departamento de Matemática Aplicada  
Universidad de Granada  
Granada, Spain  
2013

## ABSTRACT

En la Tesis se estudia tres problemas clásicos en la teoría de las ecuaciones diferenciales ordinarias singulares (uno de ellos dio origen a esta consolidada disciplina). Para ello se inicia con un capítulo general que estudia ecuaciones diferenciales singulares de tipo Liénard con término de fricción también singular, esto supone un avance interesante con relación a la extensa literatura sobre estas ecuaciones donde nunca se habían estudiado singularidades en ese término. En el segundo capítulo damos un contraejemplo, probando que una clásica conjetura aparentemente intuitiva sobre la ecuación de Lazer y Solimini no es cierta. De hecho encontramos una relación en las ecuaciones con singularidades atractivas entre la regularidad de los coeficientes y el orden de la singularidad. Más precisamente existe un valor crítico  $a^* = 1/(2p - 1)$  para el cual si  $\lambda \geq a^*$  existe una única solución periódica de

$$u'' + \frac{1}{u^\lambda} = h(t),$$

donde  $h \in L^p([0, \omega]; \mathbb{R})$ , si y solo si  $\bar{h} > 0$ ; en caso contrario, bajo la condición necesaria de valor medio positivo de  $h$ , construimos ejemplos de estas ecuaciones sin soluciones periódicas. Basándonos en los resultados abstractos del capítulo 1, en el capítulo tres estudiamos bajo qué condiciones hay existencia de soluciones periódicas en todos los

casos posibles desde el punto de vista tanto físico como matemático de las ecuaciones conocidas como de Rayleigh-Plesset (hasta la fecha actual solo habían sido estudiadas mediante simulaciones numéricas). En el último capítulo estudiamos la ecuación de Brillouin dependiente de un parámetro

$$u'' + b(1 + \cos t)u - \frac{1}{u} = 0,$$

encontrando un nuevo rango fuera del habitual  $b \in (0, 1/4)$ , donde la ecuación tiene soluciones  $2\pi$  periódicas. Además justificamos que la forma de plantear el problema hasta el momento no es la correcta si deseamos probar la conjetura que postula la existencia de al menos una  $2\pi$  periódica solución cuando  $b \in (0, 1/4)$ .

---

# Table of Contents

<b>Basic Notation</b> . . . . .	<b>1</b>
<b>Introduction</b> . . . . .	<b>3</b>
<b>1 Singular second order differential equations</b> . . . . .	<b>6</b>
1.1 Repulsive singularities . . . . .	7
1.1.1 Compact operators and Schaefer's theorem . . . . .	9
1.1.2 Auxiliary results . . . . .	11
1.1.3 A priori estimates . . . . .	16
1.1.4 Main result and consequences . . . . .	29
1.2 Attractive singularities . . . . .	32
1.2.1 The method of lower and upper functions . . . . .	33
1.2.2 Auxiliary results . . . . .	34
1.2.3 Construction of lower functions . . . . .	38
1.2.4 Construction of upper functions . . . . .	45
1.2.5 Main results and consequences . . . . .	47
<b>2 The Lazer and Solimini equation</b> . . . . .	<b>54</b>
2.1 Repulsive singularity . . . . .	55
2.2 Attractive singularity . . . . .	57
2.2.1 Auxiliary results . . . . .	59
2.2.2 Main result . . . . .	62
2.2.3 Optimality and Counter-Example . . . . .	67



<b>3</b>	<b>The Rayleigh-Plesset equation</b>	<b>70</b>
3.1	The model equation	72
3.1.1	The repulsive case	72
3.1.2	The attractive case	75
3.1.3	The case $\gamma \leq 0$	79
3.2	The physical model	79
3.3	Playing with singularities	81
3.3.1	Main result and consequences	82
3.3.2	Auxiliary propositions	89
3.3.3	Proof of the main results	103
<b>4</b>	<b>The Brillouin beam equation</b>	<b>106</b>
4.1	A corollary for the Brillouin beam equation	108
4.2	A new result for the Brillouin beam equation	109
4.2.1	A non-resonance theorem for repulsive singular equations	110
4.2.2	Main result	124
	<b>Conclusion</b>	<b>127</b>

---

# Basic Notation

$\mathbb{N}$  is the set of all natural numbers;

$\mathbb{R}$  is the set of all real numbers,  $\mathbb{R}^+ = (0, \infty)$ ,  $\mathbb{R}_+ = [0, +\infty)$ ;

$C([0, \omega]; \mathbb{R})$  is the Banach space of continuous functions  $u : [0, \omega] \rightarrow \mathbb{R}$  with the norm

$$\|u\|_\infty = \max \{ |u(t)| : t \in [0, \omega] \};$$

$C([0, \omega]; \mathbb{R}_+) = \{u \in C([0, \omega]; \mathbb{R}) : u(t) \geq 0 \text{ para } t \in [0, \omega]\}$ ;

$C(D_1; D_2)$ , where  $D_1, D_2 \subseteq \mathbb{R}$ , is the set of continuous functions  $u : D_1 \rightarrow D_2$ ;

$C^1([0, \omega]; \mathbb{R})$  is the Banach space of continuous functions  $u : [0, \omega] \rightarrow \mathbb{R}$  with continuous derivative, with the norm

$$\|u\|_{C^1} = \|u\|_\infty + \|u'\|_\infty;$$

$C^1([0, \omega]; \mathbb{R}_+) = \{u \in C^1([0, \omega]; \mathbb{R}) : u(t) \geq 0 \text{ for } t \in [0, \omega]\}$ ;

$AC([0, \omega]; \mathbb{R})$  is the set of all absolutely continuous functions  $u : [0, \omega] \rightarrow \mathbb{R}$ ;

$AC^1([0, \omega]; \mathbb{R})$  is the set of all functions  $u : [0, \omega] \rightarrow \mathbb{R}$  such that  $u$  and  $u'$  are absolutely continuous;

For a number  $p \in [1, +\infty)$ ,  $L^p([0, \omega]; \mathbb{R})$  is the Banach space of Lebesgue integrable

functions  $h : [0, \omega] \rightarrow \mathbb{R}$  in the power  $p$ , endowed with the norm

$$\|h\|_p = \left( \int_0^\omega |h(t)| dt \right)^{1/p};$$

$$L([0, \omega]; \mathbb{R}) = L^1([0, \omega]; \mathbb{R});$$

For a  $h \in L([0, \omega]; \mathbb{R})$ , its mean value is defined by

$$\bar{h} = \frac{1}{\omega} \int_0^\omega h(s) ds.$$

$$L([0, \omega]; \mathbb{R}_+) = \{p \in L([0, \omega]; \mathbb{R}) : p(t) \geq 0 \text{ for almost every } t \in [0, \omega]\};$$

$f : [0, \omega] \times D_1 \rightarrow D_2$  belongs to the set of Carathéodory functions  $\text{Car}([0, \omega] \times D_1; D_2)$  if and only if  $f(\cdot, x) : [0, \omega] \rightarrow D_2$  is measurable for all  $x \in D_1$ ,  $f(t, \cdot) : D_1 \rightarrow D_2$  is continuous for almost every  $t \in [0, \omega]$ , and for each compact set  $D_0 \subseteq D_1$  it verifies

$$\sup \{ |f(\cdot, x)| : x \in D_0 \} \in L([0, \omega]; \mathbb{R}_+);$$

$$[x]_+ = \max \{x, 0\}, \quad [x]_- = \max \{-x, 0\};$$

Unless otherwise stated whenever  $u \in C([0, \omega]; \mathbb{R})$  we define the numbers

$$M = \max \{u(t) : t \in [0, \omega]\}, \quad m = \min \{u(t) : t \in [0, \omega]\}.$$

Given  $\varphi, \psi \in L([0, \omega]; \mathbb{R})$ , then

$$\begin{aligned} \Phi_+ &= \int_0^\omega [\varphi(s)]_+ ds, & \Phi_- &= \int_0^\omega [\varphi(s)]_- ds, \\ \Psi_+ &= \int_0^\omega [\psi(s)]_+ ds, & \Psi_- &= \int_0^\omega [\psi(s)]_- ds. \end{aligned}$$

---

# Introduction

Singular boundary value problems has been of substantial and rapidly growing interest for many applied mathematicians and engineers. In contrast with the regular boundary value problems, the numerical simulations of solutions to such problems usually break down near of singular points. For that we point out its importance in order to validate the numerical results.

On the other hand numerous questions in physics, chemistry, biology and economics lead to this type of problems; for example: deformations of rods, plates and shells, behaviour of plastic materials, surface waves of fluids, flows around objects in fluids or gases, shock waves in gases, movements of viscous fluids, behaviour of magnetic fields of astrophysical objects, existence and stability of periodic and quasiperiodic orbits in celestial mechanics, etc.

As a rule, there arise nonlinear differential and integral equations, variational problems and more general optimization problems. In the Thesis we use the main techniques coming from Nonlinear Analysis to obtain existence results of periodic solutions to a specific nonlinear differential equations with singularities: *Rayleigh-Plesset*, *Lazer and Solimini* and *Brillouin beam equations*, all of them being particular cases of a more general type of equations known as *Liénard equations*.

If we ignore the particular form of the problem, we can usually reduce the question to find a fixed point of a compact operator  $T : X \rightarrow X$  defined on a Banach space  $X$ . In this way, formulating particular problems abstractly in the framework has the advantages of distilling the essential and their relationships, of allowing a uniform treatment of

differing practical problems, and of enabling the use of deep and powerful mathematical methods, without which the problems could not be solved. Nevertheless it must be emphasized that the typical approach to a specific problem generally consists of two steps:

- (a) the use of precise analytical methods to obtain estimates on the solutions;
- (b) the use of general methods of functional analysis.

Based on this idea, one of the most fruitful techniques in Nonlinear Analysis – *the lower and upper functions method* – was introduced by G. Scorza Dragoni in 1931. This method was originally applied to a Dirichlet problem, but since then a large number of contributions made the theory more complete, allowing the aforementioned method to be extended to any type of boundary value problem. The construction of upper and lower functions can be regarded as a numerical approximation of solution that satisfies the equation up to an error term with a constant sign. Then the existence of a solution, together with its localization, is deduced from two of such approximations with error terms of opposite signs.

This manuscript is based on the papers [31, 30, 33, 32, 24], the main used tools are the lower and upper functions method (Chapters 1, 2, 3) and well known fixed point theorems for compact operators (Schaefer and Poincaré-Bohl fixed point theorems). As it was mentioned above, our aim is to contribute to solving three important periodic problems in a certain field of Applied Mathematics. Thus the presented work is composed by selected papers of our recent investigation making the content as concise as it is possible. We begin with a preliminary chapter (Chapter 1) devoted to the study of a very classical equation, from mathematical point of view, well-known as a Liénard type equation (see [6, 7, 11, 46, 18, 19, 16, 26, 27, 29, 40, 44, 45, 49, 48, 61]). In spite of the fact that this type of equation was investigated by many mathematicians during the last

decades, the most of their works deal with the repulsive case and/or the case when the friction-like term has no singularity. However, the physical model studied in Chapter 3 (Rayleigh-Plesset equation) justifies to consider also equations with singularities in the friction-like term. In Chapter 2, we establish a genuine relationship between the order of the singularity and the regularity of the coefficients in the classical equation proposed by Lazer and Solimini in [40]. This relation allows us to disprove an intuitive conjecture on this equation. Moreover, we expect that the same relation can be extended to a general class of the Liénard type equations, even with singular friction-like term, whenever the singularity is attractive. In the last chapter, we consider other type of singular equation with relevance in Electronics. At the beginning, we show the necessity of developing new methods to obtain something new in this field. After this, quite unexpectedly with respect to numerical and analytical results found in the literature, we establish a new range for the existence of  $2\pi$ -periodic solutions of Brillouin focusing beam equation. This is possible due to suitable non-resonance conditions acting on the rotation number of the solutions in the phase plane.

Finally, we would like to emphasize that every chapter begins with a small introduction to the problem and the actual state of art, then the used tools and the structure of the chapter are described.

For other results obtained during my PhD studies see [3, 4, 34, 57, 62, 63].

---

# CHAPTER 1

## Singular second order differential equations

In this chapter we are going to study from mathematical point of view the periodic problem for the second-order equation

$$u''(t) + f(u(t))u'(t) + g(u(t)) = h(t, u(t)) \quad \text{for a. e. } t \in [0, \omega] \quad (1.1)$$

where  $f, g \in C(\mathbb{R}^+; \mathbb{R})$  may have singularities at zero, and  $h \in \text{Car}([0, \omega] \times \mathbb{R}_+; \mathbb{R})$ ; and some consequences to the particular Liénard equation

$$u''(t) + f(u(t))u'(t) + g(u(t)) = h_0(t) \quad \text{for a. e. } t \in [0, \omega] \quad (1.2)$$

where  $f \in C(\mathbb{R}_+; \mathbb{R})$ ,  $h_0 \in L([0, \omega]; \mathbb{R})$  and  $g$  as before. In this way, we will compare our results with some classical ones.

In the related literature,  $g$  is said to present an attractive (resp. repulsive) singularity if  $\lim_{x \rightarrow 0^+} g(x) = +\infty$  (resp.  $\lim_{x \rightarrow 0^+} g(x) = -\infty$ ). By periodic solution to (1.1) we understand a function  $u : [0, \omega] \rightarrow \mathbb{R}^+$  which is absolutely continuous together with its first derivative, satisfies (1.1) almost everywhere on  $[0, \omega]$ , and verifies

$$u(0) = u(\omega), \quad u'(0) = u'(\omega). \quad (1.3)$$

Next we investigate some general criteria to guarantee existence of periodic solution to

(1.1), i.e., existence of positive solution to the boundary problem (1.1), (1.3); taking into account the type of singularity presented by (1.1).

## 1.1 Repulsive singularities

Roughly speaking, the singularity being of repulsive type, one can expect that periodic solutions exist, provided that the force is attractive at some distance from the origin. However, some care must be taken in order to avoid what seems to be a kind of *resonance* at infinity. This fact is put in evidence in the following result by Del Pino, Manásevich and Montero proved in [46] for the particular forced equation (1.1) when  $f \equiv 0$ ,  $h(t, x) = -a(t)x + h_0(t)$ , where  $a, h_0 \in L([0, \omega]; \mathbb{R})$ .

**Theorem 1.1.1** *Let the following two assumptions hold.*

1. *There are an integer  $M$  and two constants  $\alpha, \beta$  for which*

$$\left(\frac{M\pi}{\omega}\right)^2 < \alpha \leq a(t) + \lim_{x \rightarrow +\infty} \frac{g(x)}{x} \leq \beta < \left(\frac{(M+1)\pi}{\omega}\right)^2,$$

*uniformly for almost every  $t \in [0, \omega]$ .*

2. *there are positive constants  $c', c''$  and  $\nu \geq 1$ , such that*

$$\frac{c'}{x^\nu} \leq g(x) \leq \frac{c''}{x^\nu},$$

*for every  $x \in (0, \delta]$ .*

*Then, (1.1) has a periodic solution (i.e., the problem (1.1), (1.3) has at least one positive solution).*



The result in [46] is somehow related to a paper by Fabry and Habets [14], where a periodic problem without singularities is treated. Indeed, in [14] an asymmetric oscillator is considered, assuming, roughly speaking, that the nonlinearity at  $+\infty$  asymptotically lies between the asymptotes of two consecutive curves of Dancer-Fučik spectrum; cf. [23]. It is readily seen that the constants appearing in 1. correspond to these asymptotes. The analogy between these results can be explained, see, i.e. [18], by the fact that the singularity provides for the solutions a similar behavior as when a superlinear assumption on the nonlinearity at  $-\infty$  is made. However, the methods of proof followed in [46] and [14] differ considerably, despite the fact they both use topological degree theory.

Condition 2. in Theorem 1.1.1 has been improved in [59, 60], obtaining instead of it the condition

$$\lim_{x \rightarrow 0} g(x) = -\infty, \quad \int_0^1 g(x) dx = -\infty,$$

which is commonly called *strong force condition*. The previous condition has been frequently considered for the existence of periodic solutions of (1.1) (see [16, 25, 67, 65, 66, 49, 40, 26]). However, although less frequent, there are papers dealing with weak singularities, i.e., they are not fulfilling the strong force condition (see [53, 55, 54, 49]). But it will not be treated in this manuscript.

The strong force condition has as a consequence that the energy of a solution passing near the origin is arbitrarily large, allowing to have a priori estimates from below of the periodic solutions of (1.1).

In this section we want to prove a general theorem on the existence of periodic solutions to (1.1) and discuss some consequences in order to compare it with related in literature. For that we will use *Schaefer's fixed point theorem* which requires to introduce a convenient functional analysis framework for our problem.

### 1.1.1 Compact operators and Schaefer's theorem

We consider the Banach space  $X = C^1([0, \omega]; \mathbb{R}) \times \mathbb{R}$  with the norm  $\|(u, a)\| = \|u\|_{C^1} + |a|$ . The following result is known as Schaefer fixed point theorem and it is a direct consequence of the Schauder fixed point theorem (see [51], or more recent books [52], [64]).

**Theorem 1.1.2 (Schaefer[64])** *Let  $F : X \rightarrow X$  a continuous and compact operator. If there exists  $r > 0$  such that every solution of*

$$(u, a) = \lambda F(u, a) \tag{1.4}$$

for  $\lambda \in (0, 1)$  verifies

$$\|(u, a)\| \leq r, \tag{1.5}$$

then (1.4) has at least a solution for  $\lambda = 1$ .

Our aim is to apply this result to a given operator whose fixed points correspond to periodic solutions of our differential equation. In order to define such operator and prove its compactness the following definition is needed.

**Definition 1.1.1** *An operator  $H : X \rightarrow L([0, \omega]; \mathbb{R})$ , resp.  $A : X \rightarrow \mathbb{R}$  is called Carathéodory if it is continuous and for every  $r > 0$  there exists a function  $q_r \in L([0, \omega]; \mathbb{R}_+)$ , resp. a constant  $M_r \in \mathbb{R}_+$  such that*

$$|H(u, a)(t)| \leq q_r(t) \quad \text{for a. e. } t \in [0, \omega], \quad \|(u, a)\| \leq r,$$

resp.

$$|A(u, a)| \leq M_r \quad \text{for } \|(u, a)\| \leq r.$$

**Lemma 1.1.1** Let  $H : X \rightarrow L([0, \omega]; \mathbb{R})$  and  $A : X \rightarrow \mathbb{R}$  be Caratheodory operators.

Let us define the operator  $\Omega : X \rightarrow C^1([0, \omega]; \mathbb{R})$  by

$$\Omega(u, a)(t) = -\frac{1}{\omega} \left[ (\omega - t) \int_0^t sH(u, a)(s)ds + t \int_t^\omega (\omega - s)H(u, a)(s)ds \right] \quad \text{for } t \in [0, \omega].$$

Then, the operator  $F : X \rightarrow X$  given by  $F = (\Omega, A)$  is compact.

**Proof 1** It is sufficient to prove that both  $\Omega$  and  $A$  transform each bounded set of  $X$  into a precompact set. First, note that the image of each bounded set of  $X$  by  $A$  is in fact a precompact set since  $\mathbb{R}$  is a finite dimensional space and  $A$  is a Carathéodory operator.

On the other hand, the definition of  $\Omega$  involves

$$|\Omega(u, a)(t)| \leq \frac{\omega}{4} \int_0^\omega |H(u, a)(s)| ds \quad \text{for } t \in [0, \omega], \quad (1.6)$$

$$\left| \frac{d}{dt} \Omega(u, a)(t) \right| \leq \int_0^\omega |H(u, a)(s)| ds \quad \text{for } t \in [0, \omega], \quad (1.7)$$

$$\left| \frac{d^2}{dt^2} \Omega(u, a)(t) \right| \leq |H(u, a)(t)| \quad \text{for a. e. } t \in [0, \omega]. \quad (1.8)$$

Furthermore, since  $H$  is a Carathéodory operator, for every  $r > 0$  there exists a function  $q_r \in L([0, \omega]; \mathbb{R}_+)$  such that

$$|H(u, a)(t)| \leq q_r(t) \quad \text{for a. e. } t \in [0, \omega], \quad \|(u, a)\| \leq r. \quad (1.9)$$

Now let  $M \subset X$  be a bounded set. Obviously, there exists  $r > 0$  such that  $\|(u, a)\| \leq r$

for every  $(u, a) \in M$ . Then, from (1.6)–(1.9), for  $(u, a) \in M$ , we obtain

$$\begin{aligned} \|\Omega(u, a)\|_\infty &\leq \frac{\omega}{4} \|q_r\|_1, \\ \left\| \frac{d}{dt} \Omega(u, a) \right\|_\infty &\leq \|q_r\|_1, \\ \left| \frac{d^2}{dt^2} \Omega(u, a)(t) \right| &\leq q_r(t) \quad \text{for a. e. } t \in [0, \omega]. \end{aligned}$$

By Arzelà–Ascoli’s theorem, the set  $\Omega(M)$  is precompact.

The following corollary is an immediate consequence of Theorem 1.1.2 and Lemma 1.1.1.

**Corollary 1.1.1** *Let  $H : X \rightarrow L([0, \omega]; \mathbb{R})$  and  $A : X \rightarrow \mathbb{R}$  be Carathéodory operators. If there exists  $r > 0$  (not depending on  $\lambda$ ) such that every solution of the problem*

$$u''(t) = \lambda H(u, a)(t) \quad \text{for a. e. } t \in [0, \omega], \quad (1.10)$$

$$u(0) = 0, \quad u(\omega) = 0, \quad (1.11)$$

$$a = \lambda A(u, a) \quad (1.12)$$

for  $\lambda \in (0, 1)$  verifies (1.5), then (1.10)–(1.12) admits a solution for  $\lambda = 1$ .

### 1.1.2 Auxiliary results

In this subsection we will develop some preliminaries in order to prove the main theorem.

The first aim is to rewrite the problem (1.1), (1.3) as a fixed point problem.

Let us define the continuous operator  $T : X \rightarrow C^1([0, \omega]; \mathbb{R})$  by

$$T(u, a)(t) = e^a + u(t) - \min \{u(s) : s \in [0, \omega]\}.$$

For  $\lambda \in (0, 1)$  we consider the problem

$$u''(t) + \lambda f(T(u, a)(t))u'(t) + \lambda g(T(u, a)(t)) = \lambda h(t, T(u, a)(t)) + \frac{\lambda}{\omega} \left[ \int_0^\omega g(T(u, a)(s))ds - \int_0^\omega h(s, T(u, a)(s))ds \right] \quad \text{for a. e. } t \in [0, \omega], \quad (1.13)$$

$$u(0) = 0, \quad u(\omega) = 0, \quad (1.14)$$

$$a = \lambda a - \frac{\lambda}{\omega} \left[ \int_0^\omega g(T(u, a)(s))ds - \int_0^\omega h(s, T(u, a)(s))ds \right]. \quad (1.15)$$

**Remark 1.1.1** It can be easily seen that if  $(u, a) \in X$  is a solution to (1.13)–(1.15), then the function  $u$  is periodic.

**Lemma 1.1.2** *If there exists  $r > 0$  such that for each solution of (1.13)–(1.15) with  $\lambda \in (0, 1)$  is fulfilled*

$$\|(u, a)\| \leq r,$$

*then, there exists a solution of (1.1), (1.3).*

**Proof 2** We define the operators  $H : X \rightarrow L([0, \omega]; \mathbb{R})$  y  $A : X \rightarrow \mathbb{R}$  as

$$H(u, a)(t) = -f(T(u, a)(t))u'(t) - g(T(u, a)(t)) + h(t, T(u, a)(t)) + \frac{1}{\omega} \left[ \int_0^\omega g(T(u, a)(s))ds - \int_0^\omega h(s, T(u, a)(s))ds \right] \quad \text{for a. e. } t \in [0, \omega],$$

$$A(u, a) = a - \frac{1}{\omega} \left[ \int_0^\omega g(T(u, a)(s))ds - \int_0^\omega h(s, T(u, a)(s))ds \right].$$

Is clear that both  $H$  and  $A$  are Carathéodory operators. By Corollary 1.1.1, the problem (1.13)–(1.15) with  $\lambda = 1$  has got at least one solution. Furthermore, from (1.15) (with

$\lambda = 1$ ) we obtain that

$$\int_0^\omega g(T(u, a)(s))ds = \int_0^\omega h(s, T(u, a)(s))ds, \quad (1.16)$$

and, consequently, from (1.13) with  $\lambda = 1$ , (1.14) and (1.16) we prove that  $u$  is a periodic solution satisfying

$$u''(t) + f(T(u, a)(t))u'(t) + g(T(u, a)(t)) = h(t, T(u, a)(t)) \quad \text{for a. e. } t \in [0, \omega].$$

Now we define  $v$  by

$$v(t) = T(u, a)(t) \quad \text{for } t \in [0, \omega].$$

Then  $v$  is a solution of (1.1), (1.3).

The section is concluded by lemmas presenting some useful inequalities.

**Lemma 1.1.3** *Let be  $u \in AC([0, \omega]; \mathbb{R})$  such that*

$$u(0) = u(\omega). \quad (1.17)$$

*Then the inequality*

$$(M - m)^2 \leq \frac{\omega}{4} \int_0^\omega u'^2(s)ds \quad (1.18)$$

*holds.*

**Proof 3** Let us define  $\tilde{u} : [0, 2\omega] \rightarrow \mathbb{R}$  by

$$\tilde{u}(t) = \begin{cases} u(t) & \text{if } t \in [0, \omega], \\ u(t - \omega) & \text{if } t \in (\omega, 2\omega]. \end{cases} \quad (1.19)$$

Evidently, (1.17) implies that  $\tilde{u} \in AC([0, 2\omega]; \mathbb{R})$  and also there exist  $t_0 \in [0, \omega]$  and  $t_1 \in (t_0, t_0 + \omega)$  such that

$$\tilde{u}(t_0) = m, \quad \tilde{u}(t_1) = M, \quad \tilde{u}(t_0 + \omega) = m.$$

Then

$$M - m = \int_{t_0}^{t_1} \tilde{u}'(s) ds, \quad m - M = \int_{t_1}^{t_0 + \omega} \tilde{u}'(s) ds.$$

Using the Cauchy–Bunyakovskii–Schwarz inequality we prove that

$$M - m \leq \sqrt{(t_1 - t_0) \left( \int_{t_0}^{t_1} \tilde{u}'^2(s) ds \right)},$$

$$M - m \leq \sqrt{(t_0 + \omega - t_1) \left( \int_{t_1}^{t_0 + \omega} \tilde{u}'^2(s) ds \right)}.$$

Multiplying both inequalities and using that  $AB \leq \frac{1}{4}(A + B)^2$  for each  $A, B \in \mathbb{R}_+$  we can prove

$$(M - m)^2 \leq \frac{\omega}{4} \int_{t_0}^{t_0 + \omega} \tilde{u}'^2(s) ds.$$

Finally, from the last inequality, in virtue of (1.19), we obtain (1.18).

**Lemma 1.1.4** *Let  $\rho \in C(\mathbb{R}^+; \mathbb{R}^+)$  a non-decreasing function and let  $v \in AC^1([0, \omega]; \mathbb{R})$  be a positive function such that  $v(0) = v(\omega)$ ,  $v'(0) = v'(\omega)$ . Then*

$$\int_0^\omega \frac{v''(t)}{\rho(v(t))} dt \geq 0. \tag{1.20}$$

**Proof 4** There exists a sequence  $\rho_n \in C(\mathbb{R}^+; \mathbb{R}^+)$  of non-decreasing functions which

are absolutely continuous such that

$$\lim_{n \rightarrow +\infty} \|\rho_n \circ v - \rho \circ v\|_\infty = 0, \quad (1.21)$$

$$\rho_n(m_v) = \rho(m_v) \quad \text{where} \quad m_v = \min \{v(s) : s \in [0, \omega]\}.$$

After,

$$\int_0^\omega \frac{v''(t)}{\rho_n(v(t))} dt = \int_0^\omega \frac{\rho'_n(v(t))v'^2(t)}{\rho_n^2(v(t))} dt \geq 0 \quad (1.22)$$

and

$$\left| \int_0^\omega \left[ \frac{v''(t)}{\rho_n(v(t))} - \frac{v''(t)}{\rho(v(t))} \right] dt \right| \leq \frac{\|\rho_n \circ v - \rho \circ v\|_\infty}{\rho^2(m_v)} \int_0^\omega |v''(t)| dt. \quad (1.23)$$

Now from (1.21)–(1.23) we have (1.20).

**Lemma 1.1.5** *Let be  $v \in AC^1([0, \omega]; \mathbb{R})$  such that*

$$v(0) = v(\omega), \quad v'(0) = v'(\omega). \quad (1.24)$$

Then,

$$\int_0^\omega v^2(t) dt \leq \left(\frac{\omega}{\pi}\right)^2 \int_0^\omega v'^2(t) dt + 2m \int_0^\omega v(t) dt \quad (1.25)$$

where

$$m = \min \{v(t) : t \in [0, \omega]\}.$$

**Proof 5** Let  $t_m \in [0, \omega]$  be a point such that

$$v(t_m) = m, \quad (1.26)$$

and define

$$w(t) = \begin{cases} v(t) - m & \text{for } t \in [0, \omega], \\ v(t - \omega) - m & \text{for } t \in (\omega, 2\omega]. \end{cases} \quad (1.27)$$



Obviously, in accordance with (1.24) and (1.26) we have

$$w \in AC^1([0, 2\omega]; \mathbb{R}), \quad (1.28)$$

$$w(t_m) = 0, \quad w(t_m + \omega) = 0. \quad (1.29)$$

Using Wirtinger inequality, by virtue of (1.27)–(1.29), we obtain

$$\int_{t_m}^{t_m+\omega} w^2(t) dt \leq \left(\frac{\omega}{\pi}\right)^2 \int_0^\omega v'^2(t) dt. \quad (1.30)$$

On the other hand,

$$\int_{t_m}^{t_m+\omega} w^2(t) dt = \int_0^\omega (v(t) - m)^2 dt \geq \int_0^\omega v^2(t) dt - 2m \int_0^\omega v(t) dt. \quad (1.31)$$

From (1.30) and (1.31) we get (1.25).

### 1.1.3 A priori estimates

A priori estimates of possible solutions to the problem (1.13)–(1.15) with  $\lambda \in (0, 1)$  are established in this section. This will lead to a direct proof of expected main theorem.

**Lemma 1.1.6** *Let be  $h_0 \in L([0, \omega]; \mathbb{R})$  and  $\rho \in C(\mathbb{R}^+; \mathbb{R}^+)$  a non-decreasing function such that*

$$h(t, x) \leq h_0(t)\rho(x) \quad \text{for a. e. } t \in [0, \omega], \quad x \geq r, \quad (1.32)$$

*for some  $r > 0$ , and assume that*

$$g(x) \geq \bar{h}_0\rho(x) \quad \text{for } x \geq r. \quad (1.33)$$

Then, for each solution  $(u, a)$  of (1.13)–(1.15), we have

$$a \leq \ln(1 + r). \quad (1.34)$$

**Proof 6** Let us suppose that (1.34) is false. Then

$$a > \ln(1 + r) > 0, \quad (1.35)$$

$$T(u, a)(t) > 1 + r \quad \text{for } t \in [0, \omega]. \quad (1.36)$$

Using (1.35) in (1.15), we get

$$\frac{\lambda}{\omega} \left[ \int_0^\omega g(T(u, a)(s)) ds - \int_0^\omega h(s, T(u, a)(s)) ds \right] < 0. \quad (1.37)$$

From (1.13) using (1.32), (1.36) and (1.37) we obtain

$$u''(t) + \lambda f(T(u, a)(t))u'(t) + \lambda g(T(u, a)(t)) < \lambda h_0(t)\rho(T(u, a)(t)) \quad \text{for a. e. } t \in [0, \omega]. \quad (1.38)$$

Dividing by  $\rho(T(u, a)(t))$  the equation (1.38), integrating in  $[0, \omega]$ , and using (1.14), one gets

$$\int_0^\omega \frac{u''(t)}{\rho(T(u, a)(t))} dt + \lambda \int_0^\omega \frac{g(T(u, a)(t))}{\rho(T(u, a)(t))} dt < \lambda \omega \bar{h}_0.$$

In accordance with Lemma 1.1.4, Remark 1.1.1 and  $\lambda > 0$ , it gives

$$\int_0^\omega \frac{g(T(u, a)(t))}{\rho(T(u, a)(t))} dt < \omega \bar{h}_0. \quad (1.39)$$

On the other hand, by applying (1.36) and the hypothesis (1.33) we obtain

$$\omega \bar{h}_0 \leq \int_0^\omega \frac{g(T(u, a)(t))}{\rho(T(u, a)(t))} dt$$

which, however, it is a contradiction with (1.39).

**Lemma 1.1.7** *Let  $r > 0$ ,  $\eta \in \text{Car}([0, \omega] \times \mathbb{R}_+; \mathbb{R}_+)$  a non-decreasing function in the second variable such that*

$$-\eta(t, x) \leq h(t, x) \quad \text{for a. e. } t \in [0, \omega], \quad x \geq r. \quad (1.40)$$

Furthermore,

$$\limsup_{x \rightarrow 0_+} g(x) < +\infty, \quad (1.41)$$

$$g^* \stackrel{\text{def}}{=} \limsup_{x \rightarrow +\infty} \frac{[g(x)]_+}{x} < \left(\frac{\pi}{\omega}\right)^2, \quad (1.42)$$

$$\limsup_{x \rightarrow +\infty} \frac{\int_0^\omega \eta(s, x) ds}{x} < \frac{4}{\omega} \left(1 - g^* \left(\frac{\omega}{\pi}\right)^2\right). \quad (1.43)$$

Then, for each  $a_0 > 0$  there exists a constant  $K > 0$  such that any solution  $(u, a)$  of (1.13)–(1.15) with  $a \leq a_0$  verifies

$$M - m \leq K \quad (1.44)$$

where

$$M = \max \{u(s) : s \in [0, \omega]\}, \quad m = \min \{u(s) : s \in [0, \omega]\}.$$

**Proof 7** Define the truncated function

$$\tilde{\eta}(t, x) = \begin{cases} \eta(t, x) & \text{if } x \geq r, \\ \eta(t, r) & \text{if } x < r \end{cases} \quad (1.45)$$

and

$$\xi(t, x) = \tilde{\eta}(t, x) + \varphi_r(t) \quad (1.46)$$

where

$$\varphi_r(t) = \sup \{ |h(t, x)| : 0 \leq x \leq r \} \quad \text{for a. e. } t \in [0, \omega]. \quad (1.47)$$

Of course,  $\xi$  is a non-decreasing function in the second variable. Using (1.40) and (1.45)–(1.47), we obtain the inequality

$$-\xi(t, x) \leq h(t, x) \quad \text{for a. e. } t \in [0, \omega], \quad x \in \mathbb{R}_+. \quad (1.48)$$

Furthermore,

$$\limsup_{x \rightarrow +\infty} \frac{\int_0^\omega \xi(s, x) ds}{x} = \limsup_{x \rightarrow +\infty} \left( \frac{\int_0^\omega \tilde{\eta}(s, x) ds}{x} + \frac{\|\varphi_r\|_1}{x} \right) = \limsup_{x \rightarrow +\infty} \frac{\int_0^\omega \eta(s, x) ds}{x}. \quad (1.49)$$

According to (1.15) we can re-write (1.13) as

$$u''(t) + \lambda f(T(u, a)(t))u'(t) + \lambda g(T(u, a)(t)) = \lambda h(t, T(u, a)(t)) - (1 - \lambda)a. \quad (1.50)$$

Multiplying (1.50) by  $T(u, a)(t)$  and integrating on  $[0, \omega]$ , we obtain, in accordance with Remark 1.1.1,

$$\begin{aligned} - \int_0^\omega u'^2(s) ds + \lambda \int_0^\omega g(T(u, a)(s))T(u, a)(s) ds &= \lambda \int_0^\omega h(s, T(u, a)(s))T(u, a)(s) ds + \\ &\quad - (1 - \lambda)a \int_0^\omega T(u, a)(s) ds. \end{aligned}$$

Then

$$\begin{aligned} \int_0^\omega u'^2(s)ds &= \lambda \int_0^\omega g(T(u, a)(s))T(u, a)(s)ds - \lambda \int_0^\omega h(s, T(u, a)(s))T(u, a)(s)ds + \\ &\quad + (1 - \lambda)a \int_0^\omega T(u, a)(s)ds \end{aligned} \quad (1.51)$$

is fulfilled.

On the other hand, from (1.42) and (1.43) we achieve the existence of  $\varepsilon_0 > 0$  and  $r_0 > 0$  such that

$$\frac{g(x)}{x} \leq g^* + \varepsilon_0 < \left(\frac{\pi}{\omega}\right)^2 \quad \text{for } x \geq r_0 \quad (1.52)$$

and

$$\limsup_{x \rightarrow +\infty} \frac{\int_0^\omega \eta(s, x)ds}{x} < \frac{4}{\omega} \left(1 - (g^* + \varepsilon_0) \left(\frac{\omega}{\pi}\right)^2\right). \quad (1.53)$$

As follows, (1.41) implies that

$$M_g = \sup \{g(x) : x \in (0, r_0]\} < +\infty. \quad (1.54)$$

Hence, from (1.52) and (1.54) we obtain

$$g(x) \leq (g^* + \varepsilon_0)x + M_g \quad \text{for } x > 0. \quad (1.55)$$

Now, (1.55) implies

$$\int_0^\omega g(T(u, a)(s))T(u, a)(s)ds \leq (g^* + \varepsilon_0) \int_0^\omega (T(u, a)(s))^2 ds + M_g \int_0^\omega T(u, a)(s)ds. \quad (1.56)$$

Using Lema 1.1.5 in (1.56) we arrive to

$$\int_0^\omega g(T(u, a)(s))T(u, a)(s)ds \leq (g^* + \varepsilon_0) \left(\frac{\omega}{\pi}\right)^2 \int_0^\omega u'^2(s)ds + ((g^* + \varepsilon_0)2e^a + M_g) \int_0^\omega T(u, a)(s)ds. \quad (1.57)$$

If we use the inequalities (1.48), (1.57) and the hypothesis  $a \leq a_0$  in (1.51) we prove

$$\left[1 - (g^* + \varepsilon_0) \left(\frac{\omega}{\pi}\right)^2\right] \int_0^\omega u'^2(s)ds + \leq \int_0^\omega \xi(s, e^{a_0} + M - m)T(u, a)(s)ds + K_0 \int_0^\omega T(u, a)(s)ds \quad (1.58)$$

where

$$K_0 = (g^* + \varepsilon_0)2e^{a_0} + M_g + a_0.$$

Without lost generality, we can suppose that  $M \neq m$ , e. g.  $M - m > 0$ ; then, let  $\varepsilon = \frac{e^{a_0}}{M - m}$ . In addition

$$\varepsilon \rightarrow 0 \quad \text{if} \quad M - m \rightarrow +\infty, \quad (1.59)$$

and

$$T(u, a)(t) \leq (1 + \varepsilon)(M - m) \quad \text{for } t \in [0, \omega].$$

Then,

$$\left[1 - (g^* + \varepsilon_0) \left(\frac{\omega}{\pi}\right)^2\right] \int_0^\omega u'^2(s)ds \leq \left(K_0\omega + \int_0^\omega \xi(s, (1 + \varepsilon)(M - m))ds\right) (1 + \varepsilon)(M - m).$$

Using the inequality (1.18) of Lema 1.1.3, from the last inequality we get

$$\frac{4}{\omega} \left[1 - (g^* + \varepsilon_0) \left(\frac{\omega}{\pi}\right)^2\right] \leq \frac{(1 + \varepsilon)^2 (K_0\omega + \int_0^\omega \xi(s, y)ds)}{y} \quad (1.60)$$

where  $y = (1 + \varepsilon)(M - m)$ . Finally, (1.49), (1.53), (1.59) and (1.60) imply the existence of a constant  $K$  such that (1.44) is verified.

**Remark 1.1.2** Note that from the inequality (1.44), in view of (1.14), it also follows that  $\|u\|_\infty \leq K$ .

**Lemma 1.1.8** *Let us assume that*

$$\int_0^1 [f(s)]_+ ds < +\infty \quad (1.61)$$

or

$$\int_0^1 [f(s)]_- ds < +\infty. \quad (1.62)$$

Furthermore, assume that (1.41) is verified. Then, for each  $a_0 \geq 0$  and  $K > 0$  there exists a constant  $K_1 > 0$  such that every solution  $(u, a)$  of (1.13)–(1.15) with

$$\|u\|_\infty \leq K \quad \text{and} \quad a \leq a_0 \quad (1.63)$$

verifies the boundary

$$\|u'\|_\infty \leq \lambda K_1 + a_0 \omega. \quad (1.64)$$

**Proof 8** Assume that the condition (1.61) is fulfilled. Let  $(u, a)$  be a solution of (1.13)–(1.15), then  $u$  is a periodic function and in addition there exist  $t_0, t_1 \in [0, \omega]$  such that

$$u(t_0) = m, \quad u(t_1) = M \quad (1.65)$$

where

$$M = \max \{u(t) : t \in [0, \omega]\}, \quad m = \min \{u(t) : t \in [0, \omega]\}.$$

We integrate in (1.50) on the interval  $[t_0, t] \subseteq [t_0, t_0 + \omega]$  obtaining

$$\begin{aligned} \vartheta(u')(t) + \lambda \int_{t_0}^t f(\vartheta(T(u, a))(s))\vartheta(u')(s)ds + \lambda \int_{t_0}^t g(\vartheta(T(u, a))(s))ds = \\ = \lambda \int_{t_0}^t \vartheta_1(h)(s, \vartheta(T(u, a))(s))ds - (1 - \lambda)a(t - t_0) \end{aligned}$$

where  $\vartheta : C([0, \omega]; \mathbb{R}) \rightarrow C([0, 2\omega]; \mathbb{R})$  and  $\vartheta_1 : \text{Car}([0, \omega] \times \mathbb{R}_+; \mathbb{R}) \rightarrow \text{Car}([0, 2\omega] \times \mathbb{R}; \mathbb{R})$ , respectively, those are the periodic extension operators

$$\vartheta(v)(t) = \begin{cases} v(t) & \text{if } t \in [0, \omega], \\ v(t - \omega) & \text{if } t \in (\omega, 2\omega], \end{cases} \quad (1.66)$$

$$\vartheta_1(h)(t, x) = \begin{cases} h(t, x) & \text{if } t \in [0, \omega], \\ h(t - \omega, x) & \text{if } t \in (\omega, 2\omega]. \end{cases} \quad (1.67)$$

Obviously,

$$\begin{aligned} -\vartheta(u')(t) = \lambda \int_{t_0}^t f(\vartheta(T(u, a))(s))\vartheta(u')(s)ds + \lambda \int_{t_0}^t g(\vartheta(T(u, a))(s))ds \\ - \lambda \int_{t_0}^t \vartheta_1(h)(s, \vartheta(T(u, a))(s))ds + (1 - \lambda)a(t - t_0). \end{aligned} \quad (1.68)$$

Using (1.63) and (1.65) we get

$$0 < T(u, a)(t_0) \leq T(u, a)(t) \leq T(u, a)(t_1) \leq e^{a_0} + 2K \quad \text{for } t \in [0, \omega]. \quad (1.69)$$

Then, by (1.41) and the fact that  $h \in \text{Car}([0, \omega] \times \mathbb{R}_+; \mathbb{R})$ , the number  $\mu$  and the function  $\sigma$  defined by

$$\mu = \sup \{ [g(s)]_+ : s \in (0, e^{a_0} + 2K] \}, \quad \sigma(s) = \sup \{ |h(s, x)| : x \in [0, e^{a_0} + 2K] \}, \quad (1.70)$$



satisfies

$$0 \leq \mu < +\infty, \quad \sigma \in L([0, \omega]; \mathbb{R}_+). \quad (1.71)$$

Using (1.63), (1.69)–(1.71) and  $t_0 \leq t \leq t_0 + \omega$ , in the equation (1.68), we obtain

$$-\vartheta(u')(t) \leq \lambda \int_0^{e^{a_0} + 2K} [f(s)]_+ ds + \lambda \omega \mu + \lambda \|\sigma\|_1 + \omega a_0. \quad (1.72)$$

Defining  $K_1 = \int_0^{e^{a_0} + 2K} [f(s)]_+ ds + \omega \mu + \|\sigma\|_1$ , we have, from (1.72),

$$-\vartheta(u')(t) \leq \lambda K_1 + \omega a_0 \quad \text{for } t \in [t_0, t_0 + \omega]. \quad (1.73)$$

We integrate on the interval  $[t, t_1 + \omega] \subseteq [t_1, t_1 + \omega]$  the equation (1.50), obtaining

$$\begin{aligned} \vartheta(u')(t) &= \lambda \int_t^{t_1 + \omega} f(\vartheta(T(u, a))(s)) \vartheta(u')(s) ds + \lambda \int_t^{t_1 + \omega} g(\vartheta(T(u, a))(s)) ds \\ &\quad - \lambda \int_t^{t_1 + \omega} \vartheta_1(h)(s, \vartheta(T(u, a))(s)) ds + (1 - \lambda)a(t_1 + \omega - t). \end{aligned} \quad (1.74)$$

Using (1.63), (1.69)–(1.71) and  $t_1 \leq t \leq t_1 + \omega$  in the equation (1.74), we have

$$\vartheta(u')(t) \leq \lambda K_1 + \omega a_0 \quad \text{for } t \in [t_1, t_1 + \omega]. \quad (1.75)$$

From (1.73) and (1.75) we conclude that (1.64) is verified. Therefore the proof is finished for this case.

Now we suppose that (1.62) is fulfilled. By defining

$$v(t) = u(\omega - t) \quad \text{for } t \in [0, \omega] \quad (1.76)$$

we obtain that

$$v''(t) - \lambda f(T(v, a)(t))v'(t) + \lambda g(T(v, a)(t)) = \lambda \tilde{h}(t, T(v, a)(t)) - (1 - \lambda)a \quad \text{for a. e. } t \in [0, \omega],$$

where

$$\tilde{h}(t, x) = h(\omega - t, x) \quad \text{for a. e. } t \in [0, \omega], \quad x \in \mathbb{R}_+.$$

If we do an analogical reasoning, using (1.62) by (1.61), we arrive at

$$\|v'\|_\infty \leq \lambda K_1 + a_0 \omega \tag{1.77}$$

with

$$K_1 = \int_0^{e^{a_0} + 2K} [f(s)]_- ds + \omega \mu + \|\sigma\|_1.$$

Now, since (1.76) and (1.77) we obtain (1.64).

**Remark 1.1.3** If we take  $a_0 = 0$  in Lema 1.1.8, we would obtain that

$$\|u'\|_\infty \leq \lambda K_1 \tag{1.78}$$

Whenever  $(u, a)$  is a solution of (1.13)–(1.15) with  $a \leq 0$ .

**Lemma 1.1.9** *We suppose that*

$$\lim_{x \rightarrow 0^+} g(x) = -\infty, \quad \int_0^1 g(s) ds = -\infty, \tag{1.79}$$

and (1.61) or (1.62) is verified. Then, for each  $K > 0$  there exists a constant  $a_1 > 0$  such that every solution  $(u, a)$  of (1.13)–(1.15) with

$$\|u\|_\infty \leq K \quad \text{and} \quad a \leq 0 \tag{1.80}$$

admits the bound

$$-a_1 \leq a. \quad (1.81)$$

**Proof 9** We define  $\sigma$  as in (1.70) with  $a_0 = 0$ . Obviously, because (1.79) and  $h \in \text{Car}([0, \omega] \times \mathbb{R}_+; \mathbb{R})$ , we arrive to  $\sigma \in L([0, \omega]; \mathbb{R}_+)$ . Since  $(u, a)$  is solution of (1.13)–(1.15), from (1.15), by (1.70) and (1.80), we have

$$\begin{aligned} \frac{a(1-\lambda)}{\lambda} &= -\frac{1}{\omega} \left[ \int_0^\omega g(T(u, a)(s)) ds - \int_0^\omega h(s, T(u, a)(s)) ds \right] \\ &\geq -\frac{1}{\omega} \int_0^\omega g(T(u, a)(s)) ds - \frac{1}{\omega} \|\sigma\|_1, \end{aligned}$$

hence

$$-\frac{1}{\omega} \int_0^\omega g(T(u, a)(s)) ds \leq \frac{a(1-\lambda)}{\lambda} + \frac{1}{\omega} \|\sigma\|_1.$$

By (1.80) we obtain

$$-\int_0^\omega g(T(u, a)(s)) ds \leq \|\sigma\|_1. \quad (1.82)$$

On the other hand, (1.79) implies that there exists  $s_0 > 0$  such that

$$g(s) < -\frac{\|\sigma\|_1}{\omega} \leq 0 \quad \text{for } s \in (0, s_0). \quad (1.83)$$

We denote by  $t_m \in [0, \omega]$  the point where  $u(t_m) = \min \{u(t) : t \in [0, \omega]\}$ . Obviously, either

$$T(u, a)(t_m) = e^a \geq s_0, \quad (1.84)$$

or

$$T(u, a)(t_m) = e^a < s_0. \quad (1.85)$$

Clearly, if we get a bound like (1.81) in the case (1.85), that same bound would be useful also for every solution  $(u, a)$  of (1.13)–(1.15) verifying (1.84). Hence, we can suppose

that (1.85) is fulfilled without loss of generality.

If  $T(u, a)(t) < s_0$  for every  $t \in [0, \omega]$ , by using (1.82) and (1.83) we would obtain a contradiction. Then there exists points  $t_1, t_2 \in (t_m, t_m + \omega)$  such that

$$\vartheta(T(u, a))(t) < s_0 \quad \text{for } t \in [t_m, t_1), \quad \vartheta(T(u, a))(t_1) = s_0, \quad (1.86)$$

$$\vartheta(T(u, a))(t) < s_0 \quad \text{for } t \in (t_2, t_m + \omega], \quad \vartheta(T(u, a))(t_2) = s_0, \quad (1.87)$$

where  $\vartheta$  is the operator defined by (1.66). Since  $a \leq 0$ , then

$$\frac{\lambda}{\omega} \left[ \int_0^\omega g(T(u, a)(s)) ds - \int_0^\omega h(s, T(u, a)(s)) ds \right] \geq 0,$$

so

$$u''(t) + \lambda f(T(u, a)(t))u'(t) + \lambda g(T(u, a)(t)) \geq \lambda h(t, T(u, a)(t)) \quad \text{for a. e. } t \in [0, \omega].$$

Obviously,

$$\begin{aligned} [\vartheta(u')(t)]' + \lambda f(\vartheta(T(u, a))(t))\vartheta(u')(t) + \lambda g(\vartheta(T(u, a))(t)) &\geq \\ &\geq \lambda \vartheta_1(h)(t, \vartheta(T(u, a))(t)) \quad \text{for a. e. } t \in [0, 2\omega] \end{aligned} \quad (1.88)$$

where  $\vartheta$  and  $\vartheta_1$  are operators defined by (1.66) and (1.67), respectively.

first, let us assume that (1.61) is verified. Integrating on  $[t_m, t_1]$  the inequality (1.88) we obtain

$$\begin{aligned} \vartheta(u')(t_1) + \lambda \int_{t_m}^{t_1} f(\vartheta(T(u, a))(s))\vartheta(u')(s) ds + \lambda \int_{t_m}^{t_1} g(\vartheta(T(u, a))(s)) ds &\geq \\ &\geq \lambda \int_{t_m}^{t_1} \vartheta_1(h)(s, \vartheta(T(u, a))(s)) ds; \end{aligned}$$

By a variable change and using (1.85) and (1.86) we have

$$\vartheta(u')(t_1) + \lambda \int_{e^a}^{s_0} f(s)ds - \lambda \int_{t_m}^{t_1} \vartheta_1(h)(s, \vartheta(T(u, a))(s))ds \geq -\lambda \int_{t_m}^{t_1} g(\vartheta(T(u, a))(s))ds.$$

According to Lema 1.1.8, Remark 1.1.3 and the conditions (1.79) and (1.80) we obtain that there exists a constant  $K_1 > 0$  such that (1.78) is fulfilled. Using (1.78), (1.85), the inequality  $\lambda > 0$  and the fact of  $x \leq [x]_+$  for any  $x \in \mathbb{R}$  we obtain

$$- \int_{t_m}^{t_1} g(\vartheta(T(u, a))(s))ds \leq K_2 \tag{1.89}$$

where

$$K_2 = K_1 + \int_0^{s_0} [f(s)]_+ ds + \|\sigma\|_1.$$

Multiplying by  $K_1$  in the inequality (1.89), then

$$-K_1 \int_{t_m}^{t_1} g(\vartheta(T(u, a))(s))ds \leq K_2 K_1.$$

Using (1.78), (1.83) we obtain

$$- \int_{t_m}^{t_1} g(\vartheta(T(u, a))(s))\vartheta(u')(s)ds \leq K_2 K_1.$$

After a simple change of variable and using (1.85) and (1.86) we arrive to

$$- \int_{e^a}^{s_0} g(s)ds \leq K_2 K_1. \tag{1.90}$$

Using (1.79) we ensure the existence of  $a_1 > 0$  such that (1.81) is fulfilled.

As a second part, we suppose that (1.62) is true. Integrating on  $[t_2, t_m + \omega]$  the inequality (1.88) and doing similar steps as before, using (1.87) in place of (1.86), we arrive to (1.90)

with

$$K_2 = K_1 + \int_0^{s_0} [f(s)]_- ds + \|\sigma\|_1.$$

Then, the condition (1.79) implies the existence of a constant  $a_1 > 0$  such that (1.81) is fulfilled.

### 1.1.4 Main result and consequences

Now we are at the point to prove our abstract existence result to periodic solutions of (1.1) in the repulsive case and to discuss some consequences.

**Theorem 1.1.3** *Let  $\eta \in \text{Car}([0, \omega] \times \mathbb{R}_+; \mathbb{R}_+)$  non-decreasing function with respect to the second variable,  $h_0 \in L([0, \omega]; \mathbb{R})$ ,  $\rho \in C(\mathbb{R}^+; \mathbb{R}^+)$  non-decreasing and  $r > 0$  such that is fulfilled*

1.  $-\eta(t, x) \leq h(t, x) \leq h_0(t)\rho(x)$  for a. e.  $t \in [0, \omega]$ ,  $x \geq r$ ,
2.  $g(x) \geq \bar{h}_0\rho(x)$  for  $x \geq r$ ,
3.  $\lim_{x \rightarrow 0^+} g(x) = -\infty$ ,  $\int_0^1 g(x)dx = -\infty$ ,
4.  $g^* \stackrel{\text{def}}{=} \limsup_{x \rightarrow +\infty} \frac{[g(x)]_+}{x} < \left(\frac{\pi}{\omega}\right)^2$ ,
5.  $\limsup_{x \rightarrow +\infty} \frac{\int_0^\omega \eta(t, x)dt}{x} < \frac{4}{\omega} \left(1 - g^* \left(\frac{\omega}{\pi}\right)^2\right)$ ,
6.  $\int_0^1 [f(s)]_+ ds < +\infty$  or  $\int_0^1 [f(s)]_- ds < +\infty$ .

Then, there exists at least one solution of problem (1.1), (1.3).

**Proof 10** The result immediately follows from Lemma 1.1.2, Lemmas 1.1.6–1.1.9, and Remark 1.1.2.

Many classical papers consider the case where the right-hand side only depends on  $t$  and  $f$  is continuous at zero, that is,  $f \in C(\mathbb{R}_+; \mathbb{R})$  and  $\delta = 0$ . We consider this case in a separated corollary.

**Corollary 1.1.2** *Let us consider the problem (1.2), (1.3) where  $g \in C(\mathbb{R}^+; \mathbb{R})$  verifies the conditions*

1.  $\lim_{x \rightarrow 0^+} g(x) = -\infty$ ,
2.  $\int_0^1 g(x) dx = -\infty$ ,
3.  $\limsup_{x \rightarrow +\infty} \frac{g(x)}{x} < \left(\frac{\pi}{\omega}\right)^2$ ,
4. *there exists  $r > 0$  such that  $g(x) \geq \bar{h}_0$  for every  $x \geq r$ .*

*Then, there exists a positive solution of problem (1.2), (1.3).*

**Proof 11** It is enough to apply Theorem 1.1.3 with  $h(t, x) = h_0(t)$ ,  $\eta(t, x) = [h_0(t)]_-$  and  $\rho \equiv 1$ .

Let us observe that the condition 3. is in some sense optimal, since in [7] the authors have constructed an example of  $h \in C([0, \omega]; \mathbb{R})$  such that the equation

$$u'' + \left(\frac{\pi}{\omega}\right)^2 u - \frac{1}{u^3} = h(t)$$

has no periodic solution. This is because of the last equation is resonant at infinity (i.e. it does not fulfil the first hypothesis of Theorem 1.1.1 when  $a(t) = (\pi/\omega)^2$ ). Thus the phenomenon of resonance plays a crucial role to the existence of periodic solution with repulsive singularities. However, our condition 3., roughly speaking, says that (1.1) is a non-resonant equation at infinity. Although we will not consider resonant equations, in

literature there is a extensive literature about it, starting by the first paper to a non-singular equation [39] whose technique was based on original application of *Schauder fixed point theorem*. Afterwards appeared some interesting papers as [42, 37, 36, 43, 15] all of them are recommendable to read if one is interesting about the topic, in special see [42, 43] for a survey on the topic. Other recent papers to study the resonance to singular equations are [19, 7, 17].

Corollary 1.1.2 covers the classical model equation of Lazer-Solimini [40] (profoundly studied in the next chapter). It also improves the following result by Mawhin.

**Theorem 1.1.4** ([44]) *Let us assume that  $f(x) \equiv c \in \mathbb{R}$ . Fix  $0 < a < \frac{1}{2\omega^2 e^{2|c|\omega}}$  and  $b > 0$  such that*

1.  $g(x) \leq ax + b$  for  $x > 0$ ,
2.  $\lim_{x \rightarrow 0^+} g(x) = -\infty$ ,
3.  $\int_0^1 g(x) dx = -\infty$ ,
4.  $\liminf_{x \rightarrow +\infty} g(x) > \bar{h}_0$ .

*Then there exists at least one positive solution of problem (1.2), (1.3).*

Obviously, our Corollary 1.1.2 improves conditions 1. and 4. of Theorem 1.1.4. The rest of conditions are the same and they are classically assumed.

Another related result was proved by Habets and Sanchez.

**Theorem 1.1.5** ([26]) *Let there exist constants  $c > 0$  and  $0 < r_0 < 1 < r_1 < +\infty$  such that*

1.  $g(x) - h_0(t) \leq c$  for  $t \in [0, \omega]$ ,  $x > 0$ ,



2.  $g(x) < \bar{h}_0$  for all  $x < r_0$ ,

3.  $\int_0^1 g(x)dx = -\infty$ ,

4.  $g(x) > \bar{h}$  for all  $x > r_1$ .

5.  $\int_0^\omega h_0^2(s)ds < +\infty$

Then the problem (1.2), (1.3) has at least one positive solution.

Our Corollary 1.1.2 improves Theorem 1.1.5, except for the condition 1., which it is not assumed by Habets-Sanchez.

## 1.2 Attractive singularities

Intuitively speaking, if the singularity is of attractive type, one can expect to use the *lower and upper functions theory* in order to obtain existence of solutions for boundary value problems, in particular for our periodic problem. In this case, usually the expected results are "better" than when the nonlinearity presents a repulsive singularity.

These type of equations will be mainly emphasized in this work. Because it can establish an original sharp condition guaranteeing the existence of periodic solutions. More concretely, it seems that there an unusual relationship between the "order" of the singularity and the smoothness of the coeficientes in the equation. This fact will be studied in depth in the next chapters.

Such as it was mentioned we need to introduce a new tool to study existence of periodic solutions to (1.1): the lower and upper functions method. The main theorem of this part is a general existence result of periodic solutions to the above-mentioned equation based (with attractive singularity) on this method.

### 1.2.1 The method of lower and upper functions

The method of upper and lower functions is one of the most fruitful techniques in Nonlinear Analysis and the main idea can be traced back at least to Picard. The monograph [11] presents a nice and complete historical review of the subject. In our context, the definition of upper and lower functions is as follows.

**Definition 1.2.1** *A function  $\alpha \in AC^1([0, \omega]; \mathbb{R})$  is called a lower-function to the problem (1.1), (1.3) if  $\alpha(t) > 0$  for every  $t \in [0, \omega]$  and*

$$\begin{aligned}\alpha''(t) + f(\alpha(t))\alpha'(t) + g(\alpha(t)) &\geq h(t, \alpha(t)) && \text{for a. e. } t \in [0, \omega], \\ \alpha(0) = \alpha(\omega), \quad \alpha'(0) &\geq \alpha'(\omega).\end{aligned}$$

**Definition 1.2.2** *A function  $\beta \in AC^1([0, \omega]; \mathbb{R})$  is called an upper-function to the problem (1.1), (1.3) if  $\beta(t) > 0$  for every  $t \in [0, \omega]$  and*

$$\begin{aligned}\beta''(t) + f(\beta(t))\beta'(t) + g(\beta(t)) &\leq h(t, \beta(t)) && \text{for a. e. } t \in [0, \omega], \\ \beta(0) = \beta(\omega), \quad \beta'(0) &\leq \beta'(\omega).\end{aligned}$$

Next theorem is well-known in the related literature (see, e.g., [11] or more general case in [48, Theorem 8.12]).

**Proposition 1.2.1** *Let  $\alpha$  and  $\beta$  be lower and upper functions to the problem (1.1), (1.3) such that*

$$\alpha(t) \leq \beta(t) \quad \text{for } t \in [0, \omega].$$

*Then there exists a positive solution  $u$  to the problem (1.1), (1.3) such that*

$$\alpha(t) \leq u(t) \leq \beta(t) \quad \text{for } t \in [0, \omega].$$

At this point we can prove the intuitive reason why we use this method when the equation has an attractive singular non-linearity and why many times it obtained better result in this case. To see that, we propose to consider the following equation

$$u''(t) \mp ku(t) \pm \frac{1}{u^\lambda(t)} = 0 \quad \text{for } t \in [0, \omega],$$

where  $k > 0$  and  $\lambda > 0$ . Then, if the type of singularity is attractive, one can easily find constant well-ordered lower and upper functions and to apply Proposition 1.2.1. On the contrary, if the type of singularity is repulsive, it can find revers-ordered lower and upper functions, thus, in this case, we can not apply Proposition 1.2.1. Nevertheless there are theorems which ensure existence of periodic solution when we have lower and upper functions in reversed order under an additional condition which is often used to avoid resonances (see [27, 56, 1, 2]).

The objective of this section is to develop a new technique for construction of upper and lower functions to the problem (1.1), (1.3).

## 1.2.2 Auxiliary results

Given  $x_1 \in \mathbb{R}^+$  and  $x_0 \in \mathbb{R}_+$  fixed constants, let us define the operator  $T : C^1([0, \omega]; \mathbb{R}) \rightarrow C^1([0, \omega]; \mathbb{R})$  by

$$T(u)(t) = x_1 + x_0 (u(t) - \min \{u(s) : s \in [0, \omega]\}) \quad \text{for } t \in [0, \omega] \quad (1.91)$$

and consider the auxiliar problem

$$u''(t) + f(T(u)(t))u'(t) = q(t) \quad \text{for a. e. } t \in [0, \omega], \quad (1.92)$$

$$u(0) = 0, \quad u(\omega) = 0, \quad (1.93)$$

where  $f \in C(\mathbb{R}^+; \mathbb{R})$  and  $q \in L([0, \omega]; \mathbb{R})$ . By a solution to the problem (1.92), (1.93) we understand a function  $u \in AC^1([0, \omega]; \mathbb{R})$  which satisfies (1.92) almost everywhere on  $[0, \omega]$ , and verifies (1.93).

**Lemma 1.2.1** *For every possible solution  $u$  to the problem*

$$u''(t) + \lambda f(T(u)(t))u'(t) = \lambda q(t) \quad \text{for a. e. } t \in [0, \omega], \quad (1.94)$$

$$u(0) = 0, \quad u(\omega) = 0 \quad (1.95)$$

with  $\lambda \in (0, 1]$ , the estimate

$$M - m \leq \frac{\omega}{4} \max \left\{ \int_0^\omega [q(s)]_+ ds, \int_0^\omega [q(s)]_- ds \right\} \quad (1.96)$$

holds (see  $M$  and  $m$  in Basic Notation).

**Proof 12** Multiplying (1.94) by  $u$  and integrating on  $[0, \omega]$ , we get

$$-\int_0^\omega u'^2(s) ds = \lambda \int_0^\omega q(s)u(s) ds.$$

Hence,

$$\int_0^\omega u'^2(s) ds \leq \lambda \left( M \int_0^\omega [q(s)]_- ds - m \int_0^\omega [q(s)]_+ ds \right). \quad (1.97)$$

Note that (1.95) implies  $M \geq 0$ ,  $m \leq 0$ . Therefore, from (1.97) we obtain

$$\int_0^\omega u'^2(s) ds \leq \max \left\{ \int_0^\omega [q(s)]_+ ds, \int_0^\omega [q(s)]_- ds \right\} (M - m). \quad (1.98)$$

Now, (1.96) is a direct consequence of Lemma 1.1.3 and (1.98).

Next lemma is a generalized version of a lemma proved by Mawhin in [44, Lemma 6.2].

**Lemma 1.2.2** For every  $x_1 \in \mathbb{R}^+$ ,  $x_0 \in \mathbb{R}_+$  and  $q \in L([0, \omega]; \mathbb{R})$  there exists a solution  $u$  to the problem (1.92), (1.93). Furthermore,

$$u'(\omega) - u'(0) = \int_0^\omega q(s) ds \quad (1.99)$$

and (1.96) is fulfilled.

**Proof 13** Let  $u$  be a possible solution to (1.94), (1.95) with  $\lambda \in (0, 1)$ . According to Lemma 1.2.1 we have

$$\|u\|_\infty \leq \frac{\omega}{4} \|q\|_1. \quad (1.100)$$

On the other hand, it is obvious that there exists  $t_0 \in [0, \omega]$  such that

$$u'(t_0) = 0. \quad (1.101)$$

The integration of (1.94) from  $t_0$  to  $t$  with respect to (1.91), (1.101), (1.96), and the inclusion  $\lambda \in (0, 1)$ , yields

$$|u'(t)| \leq \left| \int_{t_0}^t f(T(u)(s)) u'(s) ds - \int_{t_0}^t q(s) ds \right| \leq \int_{x_1}^{x_1 + x_0 \frac{\omega}{4} \|q\|_1} |f(s)| ds + \|q\|_1 \quad \text{for } t \in [0, \omega],$$

whence we obtain

$$\|u'\|_\infty \leq \left( M_f x_0 \frac{\omega}{4} + 1 \right) \|q\|_1, \quad (1.102)$$

where

$$M_f = \max \left\{ |f(x)| : x_1 \leq x \leq x_1 + x_0 \frac{\omega}{4} \|q\|_1 \right\}.$$

Therefore, in view of (1.100) and (1.102),  $u$  satisfies  $\|u\|_{C^1} \leq r$  with

$$r = \left[ (1 + x_0 M_f) \frac{\omega}{4} + 1 \right] \|q\|_1.$$

Define  $F : C^1([0, \omega]; \mathbb{R}) \rightarrow C^1([0, \omega]; \mathbb{R})$  by

$$F(v)(t) = \frac{1}{\omega} \left[ (\omega - t) \int_0^t s(f(T(v)(s))v'(s) - q(s)) ds + t \int_t^\omega (\omega - s)(f(T(v)(s))v'(s) - q(s)) ds \right] \quad \text{for } t \in [0, \omega].$$

Then, every solution to  $F(u) = \lambda u$  with  $\lambda \in (0, 1)$  is a solution to (1.94), (1.95) and thus according to Theorem 1.1.2 the problem (1.92), (1.93) has at least one solution  $u$ . Integrating (1.92) from 0 to  $\omega$  we obtain (1.99). The estimate (1.96) immediately follows from Lemma 1.2.1.

**Remark 1.2.1** Theorem 1.1.2 is enounced to the special Banach space  $C^1([0, \omega]; \mathbb{R}) \times \mathbb{R}$  with the norm  $\|(u, a)\| = \|u\|_{C^1} + |a|$ , in this way it can be used directly to prove the results in the previous section. However it can be used in the same way to a general Banach space  $(X, \|\cdot\|)$ .

**Lemma 1.2.3** *Let  $h \in L([0, \omega]; \mathbb{R})$ . Then,*

$$\lim_{n \rightarrow +\infty} \int_0^\omega [h(s) - n]_+ ds = 0 \quad (1.103)$$

and

$$\lim_{n \rightarrow +\infty} \int_0^\omega [h(s) + n]_- ds = 0. \quad (1.104)$$

**Proof 14** Let us define

$$h_n(t) = \begin{cases} n & \text{if } h(t) > n, \\ h(t) & \text{if } h(t) \leq n, \end{cases} \quad \text{for a. e. } t \in [0, \omega], \quad n \in \mathbb{N}. \quad (1.105)$$

Then,

$$h(t) = h_n(t) + [h(t) - n]_+ \quad \text{for a. e. } t \in [0, \omega], \quad n \in \mathbb{N}. \quad (1.106)$$

Integrating (1.106) over a period,

$$\int_0^\omega h(s)ds = \int_0^\omega h_n(s)ds + \int_0^\omega [h(s) - n]_+ ds \quad \text{for } n \in \mathbb{N}. \quad (1.107)$$

On the other hand, from (1.105) and (1.106) we get

$$-[h(t)]_- \leq h_n(t) \leq h(t) \quad \text{for a. e. } t \in [0, \omega], \quad n \in \mathbb{N}$$

and

$$\lim_{n \rightarrow +\infty} h_n(t) = h(t) \quad \text{for a. e. } t \in [0, \omega].$$

Thus, according to *Lebesgue Theorem* we have

$$\lim_{n \rightarrow +\infty} \int_0^\omega h_n(s)ds = \int_0^\omega h(s)ds. \quad (1.108)$$

Now, from (1.107) and (1.108) we get (1.103). The identity (1.104) can be proved by similar arguments.

### 1.2.3 Construction of lower functions

The first result of this section gives sufficient conditions for the construction of a lower function.

**Proposition 1.2.2** *Let  $h_0 \in L([0, \omega]; \mathbb{R})$ ,  $\rho_0 \in C(\mathbb{R}^+; \mathbb{R}^+)$  be a non-decreasing function,  $0 < x_1 \leq x_2 < +\infty$ , and  $c \in \mathbb{R}$  be such that*

$$h(t, x) \leq h_0(t)\rho_0(x) \quad \text{for a. e. } t \in [0, \omega], \quad x \in [x_1, x_2], \quad (1.109)$$

$$\frac{g(x)}{\rho_0(x)} \geq c \geq \bar{h}_0 \quad \text{for } x \in [x_1, x_2], \quad (1.110)$$

and

$$\rho_0(x_2)\frac{\omega}{4}\Phi_+ \leq x_2 - x_1 \leq \rho_0(x_1)\frac{\omega}{4}\Phi_-, \quad (1.111)$$

where

$$\varphi(t) = h_0(t) - c \quad \text{for a. e. } t \in [0, \omega]. \quad (1.112)$$

Then there exists a lower function  $\alpha$  to the problem (1.1), (1.3) such that

$$x_1 \leq \alpha(t) \leq x_2 \quad \text{for } t \in [0, \omega].$$

**Proof 15** By the definition of  $\varphi$  and (1.110), we obtain  $\Phi_- \geq \Phi_+ \geq 0$ . As a first case we suppose that

$$\Phi_+ > 0.$$

Put

$$x_0 = \frac{4(x_2 - x_1)}{\omega\Phi_- \Phi_+}, \quad (1.113)$$

$$q(t) = \Phi_-[\varphi(t)]_+ - \Phi_+[\varphi(t)]_- \quad \text{for a. e. } t \in [0, \omega], \quad (1.114)$$

and let  $T : C^1([0, \omega]; \mathbb{R}) \rightarrow C^1([0, \omega]; \mathbb{R})$  be the operator defined by (1.91). Note that

$$\int_0^\omega q(s)ds = 0. \quad (1.115)$$

According to Lemma 1.2.2 there exists a solution  $u$  to (1.92), (1.93) such that (1.96) and (1.99) hold. By using (1.114) and (1.115), we obtain

$$M - m \leq \frac{\omega}{4}\Phi_+\Phi_-, \quad (1.116)$$

$$u'(0) = u'(\omega). \quad (1.117)$$



Put

$$\alpha(t) = T(u)(t) \quad \text{for } t \in [0, \omega]. \quad (1.118)$$

Then, according to (1.91)–(1.93), (1.113), (1.114) and (1.116)–(1.118) we arrive at

$$\alpha''(t) + f(\alpha(t))\alpha'(t) = x_0\Phi_-[\varphi(t)]_+ - x_0\Phi_+[\varphi(t)]_- \quad \text{for a. e. } t \in [0, \omega], \quad (1.119)$$

$$\alpha(0) = \alpha(\omega), \quad \alpha'(0) = \alpha'(\omega), \quad (1.120)$$

$$x_1 \leq \alpha(t) \leq x_2 \quad \text{for } t \in [0, \omega]. \quad (1.121)$$

Using that  $\rho_0$  is a non-decreasing function, from the inequality (1.121) we obtain

$$\rho_0(x_1) \leq \rho_0(\alpha(t)) \leq \rho_0(x_2) \quad \text{for } t \in [0, \omega]. \quad (1.122)$$

From the inequality (1.111), by virtue of (1.113), we get

$$x_0\Phi_+ \leq \rho_0(x_1), \quad \rho_0(x_2) \leq x_0\Phi_-. \quad (1.123)$$

Now (1.122) and (1.123) imply

$$x_0\Phi_+ \leq \rho_0(\alpha(t)) \leq x_0\Phi_- \quad \text{for } t \in [0, \omega]. \quad (1.124)$$

Using (1.124) in (1.119) we get

$$\alpha''(t) + f(\alpha(t))\alpha'(t) \geq \rho_0(\alpha(t))\varphi(t) \quad \text{for a. e. } t \in [0, \omega]. \quad (1.125)$$

On the other hand, we can prove, using (1.110), (1.112) and (1.121), that

$$\varphi(t) \geq h_0(t) - \frac{g(\alpha(t))}{\rho_0(\alpha(t))} \quad \text{for a. e. } t \in [0, \omega]. \quad (1.126)$$

From (1.125), on account of (1.109), (1.121) and (1.126), it follows that

$$\alpha''(t) + f(\alpha(t))\alpha'(t) + g(\alpha(t)) \geq h(t, \alpha(t)) \quad \text{for a. e. } t \in [0, \omega]. \quad (1.127)$$

Consequently, (1.120), (1.121) and (1.127) ensure us that  $\alpha$  is a lower function to the problem (1.1), (1.3).

Now, we consider the remaining case

$$\Phi_+ = 0.$$

Of course, in this case

$$\varphi(t) \leq 0 \quad \text{for a. e. } t \in [0, \omega].$$

Then, defining  $\alpha$  by

$$\alpha(t) = x_1 \quad \text{for } t \in [0, \omega] \quad (1.128)$$

we can prove easily that  $\alpha$  is a positive function which fulfils (1.120) and (1.125). Again, from (1.110), (1.112) and (1.128) we obtain (1.126) and using (1.109), (1.126) and (1.128) in (1.125) we arrive at (1.127). Finally, also in this case, (1.120), (1.127) and (1.128) imply that  $\alpha$  is a lower function to the problem (1.1), (1.3).

A simplified version of the latter proposition is presented below.

**Proposition 1.2.3** *Let  $h_0 \in L([0, \omega]; \mathbb{R})$ ,  $\rho_0 \in C(\mathbb{R}^+; \mathbb{R}^+)$  be a non-decreasing function,  $x_0 > 0$ , and  $c \in \mathbb{R}$  be such that*

$$h(t, x) \leq h_0(t)\rho_0(x) \quad \text{for a. e. } t \in [0, \omega], \quad 0 < x \leq x_0, \quad (1.129)$$

$$\frac{g(x)}{\rho_0(x)} \geq c \geq \bar{h}_0 \quad \text{for } 0 < x \leq x_0, \quad (1.130)$$

and, in addition, let there exist  $x_2 \in (0, x_0]$  such that

$$x_2 - \rho_0(x_2) \frac{\omega}{4} \Phi_+ > 0, \quad (1.131)$$

$$\rho_0(x_2) \Phi_+ \leq \rho_0 \left( x_2 - \rho_0(x_2) \frac{\omega}{4} \Phi_+ \right) \Phi_- \quad (1.132)$$

where  $\varphi(t) = h_0(t) - c$  for almost every  $t \in [0, \omega]$ . Then there exists a lower function  $\alpha$  to the problem (1.1), (1.3) with

$$0 < \alpha(t) \leq x_2 \quad \text{for } t \in [0, \omega].$$

**Proof 16** In order to apply Proposition 1.2.2, we define

$$x_1 = x_2 - \rho_0(x_2) \frac{\omega}{4} \Phi_+. \quad (1.133)$$

By (1.131),  $x_1 > 0$ . Then, it is clear that (1.129) and (1.130) imply (1.109) and (1.110).

It remains to show that (1.111) holds. Indeed, by the definition of  $x_1$  we have

$$x_2 - x_1 = \frac{\omega}{4} \rho_0(x_2) \Phi_+. \quad (1.134)$$

On the other hand, using (1.133) in (1.132) we get

$$\frac{\omega}{4} \rho_0(x_2) \Phi_+ \leq \frac{\omega}{4} \rho_0(x_1) \Phi_-. \quad (1.135)$$

Therefore, (1.134) and (1.135) imply (1.111).

The following corollaries are direct consequences of Proposition 1.2.3.

**Corollary 1.2.1** Let  $x_0 > \frac{\omega}{8} \|h_0 - \bar{h}_0\|_1$  be such that

$$g(x) \geq \bar{h}_0 \quad \text{for } 0 < x \leq x_0.$$

Then there exists a lower function  $\alpha$  to the problem (1.2), (1.3) with

$$0 < \alpha(t) \leq x_0 \quad \text{for } t \in [0, \omega]. \quad (1.136)$$

**Proof 17** The assertion immediately follows from Proposition 1.2.3 with  $h(t, x) \equiv h_0(t)$  if we put  $c = \bar{h}_0$ ,  $\rho_0 \equiv 1$ , and  $x_2 = x_0$ . Note also that in this case  $\|h_0 - \bar{h}_0\|_1 = \Phi_+ + \Phi_- = 2\Phi_+$ .

**Corollary 1.2.2** Let  $h_0 \in L([0, \omega]; \mathbb{R})$ ,  $\rho_0 \in C(\mathbb{R}^+; \mathbb{R}^+)$  be a non-decreasing function,  $x_0 > 0$ , and  $c \in \mathbb{R}$  be such that (1.129) and (1.130) are fulfilled. Let, moreover, there exist a sequence  $\{y_n\}_{n=1}^{+\infty}$  of positive numbers such that

$$\lim_{n \rightarrow +\infty} y_n = 0, \quad (1.137)$$

$$\lim_{n \rightarrow +\infty} \frac{\rho_0(y_n)}{y_n} = 0, \quad (1.138)$$

and let there exist  $\varepsilon \in (0, 1)$  and  $n_0 \in \mathbb{N}$  such that

$$\frac{\rho_0(y_n)}{\rho_0(y_n(1 - \varepsilon))} \Phi_+ \leq \Phi_- \quad \text{for } n \geq n_0 \quad (1.139)$$

where  $\varphi(t) = h_0(t) - c$  for almost every  $t \in [0, \omega]$ . Then there exists a lower function  $\alpha$  to the problem (1.1), (1.3) satisfying (1.136).

**Proof 18** According to Proposition 1.2.3, it is sufficient to prove that (1.131) and (1.132) are fulfilled for some  $x_2 \in (0, x_0]$ . According to (1.137) and (1.138), there exists  $n_1 \geq n_0$  such that

$$\begin{aligned} y_n &\leq x_0 && \text{for } n \geq n_1, \\ -\frac{\omega}{4} \Phi_+ \rho_0(y_n) &\geq -\varepsilon y_n && \text{for } n \geq n_1, \end{aligned} \quad (1.140)$$

Adding  $y_n$  to both sides of the inequality (1.140) and applying that  $\rho_0$  is a non-decreasing function, we obtain

$$\rho_0 \left( y_n - \frac{\omega}{4} \Phi_+ \rho_0(y_n) \right) \geq \rho_0(y_n(1 - \varepsilon)) \quad \text{for } n \geq n_1. \quad (1.141)$$

Now, if we put  $x_2 = y_{n_1}$  we obtain, on account of (1.139)–(1.141) that (1.131) and (1.132) are fulfilled.

**Corollary 1.2.3** *Let  $h_0 \in L([0, \omega]; \mathbb{R})$ ,  $\rho_0 \in C(\mathbb{R}^+; \mathbb{R}^+)$  be a non-decreasing function,  $x_0 > 0$ , and  $c \in \mathbb{R}$  be such that (1.129) and (1.130) are fulfilled. If  $\frac{\rho_0(x)}{x}$  is a non-increasing function and*

$$\frac{\omega}{4} \Phi_+ \Phi_- \frac{\rho_0(x_0)}{x_0} \leq \Phi_- - \Phi_+ \quad (1.142)$$

where  $\varphi(t) = h_0(t) - c$  for almost every  $t \in [0, \omega]$ , then there exists a lower function  $\alpha$  to the problem (1.1), (1.3) satisfying (1.136).

**Proof 19** According to Proposition 1.2.3, it is sufficient to prove that (1.131) and (1.132) are satisfied with  $x_2 = x_0$ . From (1.142) we easily obtain (1.131). On the other hand, since the function  $\frac{\rho_0(x)}{x}$  is non-increasing,

$$\frac{\rho_0 \left( x_0 - \rho_0(x_0) \frac{\omega}{4} \Phi_+ \right)}{x_0 - \rho_0(x_0) \frac{\omega}{4} \Phi_+} \geq \frac{\rho_0(x_0)}{x_0}.$$

Consequently,

$$\rho_0 \left( x_0 - \rho_0(x_0) \frac{\omega}{4} \Phi_+ \right) \geq \rho_0(x_0) \left( 1 - \frac{\omega}{4} \Phi_+ \frac{\rho_0(x_0)}{x_0} \right). \quad (1.143)$$

Multiplying both sides of (1.143) by  $\Phi_-$  and using the inequality (1.142) we get (1.132).

## 1.2.4 Construction of upper functions

The following assertions dealing with the existence of an upper function to the problem considered can be proved analogously to the previously results formulated, therefore, their proofs are omitted.

**Proposition 1.2.4** *Let  $h_1 \in L([0, \omega]; \mathbb{R})$ ,  $\rho_1 \in C(\mathbb{R}^+; \mathbb{R}^+)$  be a non-decreasing function,  $0 < x_1 \leq x_2 < +\infty$ , and  $c \in \mathbb{R}$  be such that*

$$h(t, x) \geq h_1(t)\rho_1(x) \quad \text{for a. e. } t \in [0, \omega], \quad x \in [x_1, x_2], \quad (1.144)$$

$$\frac{g(x)}{\rho_1(x)} \leq c \leq \bar{h}_1 \quad \text{for } x \in [x_1, x_2], \quad (1.145)$$

and

$$\rho_1(x_2)\frac{\omega}{4}\Phi_- \leq x_2 - x_1 \leq \rho_1(x_1)\frac{\omega}{4}\Phi_+ \quad (1.146)$$

where

$$\varphi(t) = h_1(t) - c \quad \text{for a. e. } t \in [0, \omega]. \quad (1.147)$$

Then there exists an upper function  $\beta$  to the problem (1.1), (1.3) such that

$$x_1 \leq \beta(t) \leq x_2 \quad \text{for } t \in [0, \omega].$$

**Proposition 1.2.5** *Let  $h_1 \in L([0, \omega]; \mathbb{R})$ ,  $\rho_1 \in C(\mathbb{R}^+; \mathbb{R}^+)$  be a non-decreasing function,  $x_0 > 0$ , and  $c \in \mathbb{R}$  be such that*

$$h(t, x) \geq h_1(t)\rho_1(x) \quad \text{for a. e. } t \in [0, \omega], \quad x \geq x_0, \quad (1.148)$$

$$\frac{g(x)}{\rho_1(x)} \leq c \leq \bar{h}_1 \quad \text{for } x \geq x_0, \quad (1.149)$$

and, in addition, let there exist  $x_1 \geq x_0$  such that

$$\rho_1 \left( x_1 + \rho_1(x_1) \frac{\omega}{4} \Phi_+ \right) \Phi_- \leq \rho_1(x_1) \Phi_+ \quad (1.150)$$

where  $\varphi(t) = h_1(t) - c$  for almost every  $t \in [0, \omega]$ . Then there exists an upper function  $\beta$  to the problem (1.1), (1.3) with

$$x_1 \leq \beta(t) \quad \text{for } t \in [0, \omega]. \quad (1.151)$$

**Corollary 1.2.4** *Let there exist  $x_0 > 0$  such that*

$$g(x) \leq \bar{h}_0 \quad \text{for } x \geq x_0.$$

*Then there exists an upper function  $\beta$  to the problem (1.2), (1.3) with*

$$\beta(t) \geq x_0 \quad \text{for } t \in [0, \omega]. \quad (1.152)$$

**Corollary 1.2.5** *Let  $h_1 \in L([0, \omega]; \mathbb{R})$ ,  $\rho_1 \in C(\mathbb{R}^+; \mathbb{R}^+)$  be a non-decreasing function,  $x_0 > 0$ , and  $c \in \mathbb{R}$  be such that (1.148) and (1.149) hold, and let there exist a sequence  $\{y_n\}_{n=1}^{+\infty}$  of positive numbers such that*

$$\lim_{n \rightarrow +\infty} y_n = +\infty, \quad (1.153)$$

$$\lim_{n \rightarrow +\infty} \frac{\rho_1(y_n)}{y_n} = 0. \quad (1.154)$$

*Furthermore, let there exist  $\varepsilon > 0$  and  $n_0 \in \mathbb{N}$  such that*

$$\frac{\rho_1(y_n(1 + \varepsilon))}{\rho_1(y_n)} \Phi_- \leq \Phi_+ \quad \text{for } n \geq n_0 \quad (1.155)$$

*where  $\varphi(t) = h_1(t) - c$  for almost every  $t \in [0, \omega]$ . Then there exists an upper function*

$\beta$  to the problem (1.1), (1.3) satisfying (1.152).

**Corollary 1.2.6** *Let  $h_1 \in L([0, \omega]; \mathbb{R})$ ,  $\rho_1 \in C(\mathbb{R}^+; \mathbb{R}^+)$  be a non-decreasing function,  $x_0 > 0$ , and  $c \in \mathbb{R}$  be such that (1.148) and (1.149) are fulfilled. If  $\frac{\rho_1(x)}{x}$  is a non-increasing function such that*

$$\frac{\omega}{4}\Phi_+\Phi_-\frac{\rho_1(x_0)}{x_0} \leq \Phi_+ - \Phi_- \quad (1.156)$$

where  $\varphi(t) = h_1(t) - c$  for almost every  $t \in [0, \omega]$ , then there exists an upper function  $\beta$  to the problem (1.1), (1.3) satisfying (1.152).

## 1.2.5 Main results and consequences

Now we can prove our abstract theorem on the existence of periodic solution to (1.1) with attractive type of singularity and to discuss some consequences.

**Theorem 1.2.1** *Let  $\rho_0, \rho_1 \in C(\mathbb{R}^+; \mathbb{R}^+)$  be non-decreasing functions,  $h_0, h_1 \in L([0, \omega]; \mathbb{R})$ , and  $0 < r_0 \leq r_1 < +\infty$  be such that*

$$h(t, x) \leq h_0(t)\rho_0(x) \quad \text{for a. e. } t \in [0, \omega], \quad 0 < x \leq r_0, \quad (1.157)$$

$$h(t, x) \geq h_1(t)\rho_1(x) \quad \text{for a. e. } t \in [0, \omega], \quad x \geq r_1, \quad (1.158)$$

and let there exist  $c_0, c_1 \in \mathbb{R}$  such that

$$\frac{g(x)}{\rho_0(x)} \geq c_0 \geq \bar{h}_0 \quad \text{for } 0 < x \leq r_0, \quad (1.159)$$

$$\frac{g(x)}{\rho_1(x)} \leq c_1 \leq \bar{h}_1 \quad \text{for } x \geq r_1. \quad (1.160)$$

Furthermore, let us suppose that  $\rho_0$  fulfils at least one of the following conditions:



a) there exists a sequence  $\{x_n\}_{n=1}^{+\infty}$  of positive numbers such that

$$\lim_{n \rightarrow +\infty} x_n = 0, \quad \lim_{n \rightarrow +\infty} \frac{\rho_0(x_n)}{x_n} = 0,$$

and there exist  $\varepsilon_0 \in (0, 1)$  and  $n_0 \in \mathbb{N}$  such that

$$\frac{\rho_0(x_n)}{\rho_0(x_n(1 - \varepsilon_0))} \Phi_+ \leq \Phi_- \quad \text{for } n \geq n_0,$$

where  $\varphi(t) = h_0(t) - c_0$  for almost every  $t \in [0, \omega]$ ;

b) the function  $\frac{\rho_0(x)}{x}$  is non-increasing and

$$\frac{\omega}{4} \Phi_+ \Phi_- - \frac{\rho_0(r_0)}{r_0} \leq \Phi_- - \Phi_+,$$

where  $\varphi(t) = h_0(t) - c_0$  for almost every  $t \in [0, \omega]$ .

Besides, let us suppose that  $\rho_1$  fulfils at least one of the following conditions:

c) there exists a sequence  $\{y_n\}_{n=1}^{+\infty}$  of positives numbers such that

$$\lim_{n \rightarrow +\infty} y_n = +\infty, \quad \lim_{n \rightarrow +\infty} \frac{\rho_1(y_n)}{y_n} = 0,$$

and there exist  $\varepsilon_1 > 0$  and  $n_1 \in \mathbb{N}$  such that

$$\frac{\rho_1(y_n(1 + \varepsilon_1))}{\rho_1(y_n)} \Psi_- \leq \Psi_+ \quad \text{for } n \geq n_1,$$

where  $\psi(t) = h_1(t) - c_1$  for almost every  $t \in [0, \omega]$ ;

d) the function  $\frac{\rho_1(x)}{x}$  is non-increasing and

$$\frac{\omega}{4} \Psi_+ \Psi_- - \frac{\rho_1(r_1)}{r_1} \leq \Psi_+ - \Psi_-,$$

where  $\psi(t) = h_1(t) - c_1$  for almost every  $t \in [0, \omega]$ .

Then there exists at least one positive solution to the problem (1.1), (1.3).

**Proof 20** According to Corollaries 1.2.2, 1.2.3, 1.2.5, and 1.2.6, the conditions of the theorem guarantee a well-ordered couple of lower and upper functions, therefore the result is a direct consequence of Proposition 1.2.1.

**Remark 1.2.2** Note that (1.159) (resp. (1.160)) implies  $\Phi_- \geq \Phi_+$  (resp.  $\Psi_+ \geq \Psi_-$ ). In addition, the conditions a) and c) are verified if, for instance,  $\Phi_- \neq \Phi_+$  and  $\Psi_- \neq \Psi_+$ ,  $\rho_i(x) = x^{\mu_i}$  ( $i = 0, 1$ ) with  $\mu_0 > 1 > \mu_1 \geq 0$ . On the other hand, conditions b) and d) are fulfilled if, for instance,  $\rho_i(x) = x$  ( $i = 0, 1$ ) and

$$\frac{\omega}{4}\Phi_+\Phi_- \leq \Phi_- - \Phi_+, \quad \frac{\omega}{4}\Psi_+\Psi_- \leq \Psi_+ - \Psi_-. \quad (1.161)$$

Next, we formulate some corollaries which can be obtained immediately from Theorem 1.2.1 and Remark 1.2.2.

**Corollary 1.2.7** Let  $h_0, h_1 \in L([0, \omega]; \mathbb{R})$ ,  $\mu_0 > 1 > \mu_1 \geq 0$  and  $0 < r_0 \leq r_1 < +\infty$  be such that

$$\begin{aligned} h(t, x) &\leq h_0(t)x^{\mu_0} && \text{for a. e. } t \in [0, \omega], \quad 0 < x \leq r_0, \\ h(t, x) &\geq h_1(t)x^{\mu_1} && \text{for a. e. } t \in [0, \omega], \quad x \geq r_1, \\ \liminf_{x \rightarrow 0^+} \frac{g(x)}{x^{\mu_0}} &> \bar{h}_0, && \limsup_{x \rightarrow +\infty} \frac{g(x)}{x^{\mu_1}} < \bar{h}_1. \end{aligned}$$

Then there exists at least one positive solution to the problem (1.1), (1.3).

**Proof 21** According to Remark 1.2.2, one can apply Theorem 1.2.1 with  $\rho_0(x) = x^{\mu_0}$ ,  $\rho_1(x) = x^{\mu_1}$ .

**Corollary 1.2.8** *Let  $h_0, h_1 \in L([0, \omega]; \mathbb{R})$  and  $0 < r_0 \leq r_1 < +\infty$  be such that*

$$\begin{aligned} h(t, x) &\leq h_0(t)x && \text{for a. e. } t \in [0, \omega], \quad 0 < x \leq r_0, \\ h(t, x) &\geq h_1(t)x && \text{for a. e. } t \in [0, \omega], \quad x \geq r_1, \\ \bar{h}_0 &< \liminf_{x \rightarrow 0_+} \frac{g(x)}{x} < +\infty, && \bar{h}_1 > \limsup_{x \rightarrow +\infty} \frac{g(x)}{x} > -\infty. \end{aligned} \quad (1.162)$$

*In addition, we suppose that*

$$\frac{\omega}{4} H_0^+ H_0^- < H_0^- - H_0^+, \quad \frac{\omega}{4} H_1^+ H_1^- < H_1^+ - H_1^-, \quad (1.163)$$

*where*

$$\begin{aligned} H_0^+ &= \int_0^\omega [h_0(t) - g_*]_+ dt, & H_0^- &= \int_0^\omega [h_0(t) - g_*]_- dt, \\ H_1^+ &= \int_0^\omega [h_1(t) - g^*]_+ dt, & H_1^- &= \int_0^\omega [h_1(t) - g^*]_- dt, \end{aligned}$$

*and*

$$g_* = \liminf_{x \rightarrow 0_+} \frac{g(x)}{x}, \quad g^* = \limsup_{x \rightarrow +\infty} \frac{g(x)}{x}.$$

*Then there exists at least one positive solution to the problem (1.1), (1.3).*

**Proof 22** From (1.162) and (1.163) we obtain that there exists  $\varepsilon > 0$  small enough such that  $\varepsilon < \min \{g_* - \bar{h}_0, \bar{h}_1 - g^*\}$  and (1.161) is verified, where

$$\varphi(t) = h_0(t) - g_* + \varepsilon \quad \text{for a. e. } t \in [0, \omega], \quad \psi(t) = h_1(t) - g^* - \varepsilon \quad \text{for a. e. } t \in [0, \omega].$$

Hence, setting  $c_0 = g_* - \varepsilon$ ,  $c_1 = g^* + \varepsilon$  and  $\rho_i(x) = x$  ( $i = 0, 1$ ), the corollary follows from Theorem 1.2.1.

**Corollary 1.2.9** *Let  $h_0, h_1 \in L([0, \omega]; \mathbb{R})$  and  $0 < r_0 \leq r_1 < +\infty$  be such that*

$$\begin{aligned} h(t, x) &\leq h_0(t)x && \text{for a. e. } t \in [0, \omega], \quad 0 < x \leq r_0, \\ h(t, x) &\geq h_1(t)x && \text{for a. e. } t \in [0, \omega], \quad x \geq r_1, \\ \lim_{x \rightarrow 0^+} \frac{g(x)}{x} &= +\infty, && \lim_{x \rightarrow +\infty} \frac{g(x)}{x} = -\infty. \end{aligned} \tag{1.164}$$

*Then there exists at least one positive solution to the problem (1.1), (1.3).*

**Proof 23** Using (1.164) and Lemma 1.2.3, we can find  $c_0 > \bar{h}_0$  and  $c_1 < \bar{h}_1$  such that (1.161) is fulfilled where  $\varphi(t) = h_0(t) - c_0$ ,  $\psi(t) = h_1(t) - c_1$ . Moreover,  $g(x) \geq c_0x$  nearby zero and  $g(x) \leq c_1x$  nearby  $+\infty$ . Hence, taking  $\rho_i(x) = x$  ( $i = 0, 1$ ), the corollary follows from Theorem 1.2.1.

In conclusion, the conditions nearby zero guarantee the existence of a positive lower function, whereas the conditions nearby infinite guarantee the existence of an upper function. Both ideas can be combined in order to get a wide variety of results.

We finish the section with two results dealing with the classical singular Liénard equation (1.2).

**Theorem 1.2.2** *Let  $\frac{\omega}{8} \|h_0 - \bar{h}_0\|_1 < r_0 \leq r_1 < +\infty$  be such that*

$$\begin{aligned} g(x) &\geq \bar{h}_0 && \text{for } 0 < x \leq r_0, \\ g(x) &\leq \bar{h}_0 && \text{for } x \geq r_1. \end{aligned}$$

*Then there exists at least one positive solution to the problem (1.2), (1.3).*

**Proof 24** It is a direct consequence of Corollaries 1.2.1 and 1.2.4.

**Theorem 1.2.3** *Let*

$$\limsup_{x \rightarrow 0^+} g(x) = +\infty$$

*and let  $r_1 > 0$  be such that*

$$g(x) \leq \bar{h}_0 \quad \text{for } x \geq r_1.$$

*If*

$$\text{ess sup } \{h_0(t) : t \in [0, \omega]\} < +\infty,$$

*then there exists at least one positive solution to the problem (1.2), (1.3).*

**Proof 25** The existence of a lower function follows from Proposition 1.2.2 with  $h(t, x) = h_0(t)$ ,  $\rho_0 \equiv 1$ ,

$$c = \text{ess sup } \{h_0(t) : t \in [0, \omega]\},$$

and  $x_1 = x_2 > 0$  sufficiently small such that  $g(x_1) \geq c$ .

The existence of an upper function follows from Proposition 1.2.5 with  $h(t, x) = h_0(t)$ ,  $h_1 \equiv h_0$ ,  $\rho_1 \equiv 1$ ,  $c = \bar{h}_0$ , and  $x_0 = x_1 = r_1$ .

Consequently, the assertion follows from Proposition 1.2.1.

It is interesting to compare our results with the existing ones in the related literature. For instance, it is easy to verify that Theorem 1.2.3 generalises in some sense the result of Lazer and Solimini [40, Theorem 2.1] for the equation

$$u'' + g(u) = h(t) \tag{1.165}$$

with attractive singularity and without friction.

On the other hand, if [48, Lemma 8.19] is applied to (1.165), the following result is obtained.

**Theorem 1.2.4** *Assume that there exist  $0 < r_0 < r_1 < +\infty$  such that*

1.  $r_0 > 2\omega\|h - \bar{h}\|_1$ ,
2.  $g(x) \geq \bar{h}$  if  $0 < x \leq r_0$ ,
3.  $g(x) \leq \bar{h}$  if  $x \geq r_1$ .

*Then the problem (1.165), (1.3) has at least one positive solution.*

Note that Theorem 1.2.2 is more general. First, it works for the equation with friction. Besides, since  $\frac{\omega}{8}\|h - \bar{h}\|_1 < 2\omega\|h - \bar{h}\|_1$ , it is evident that the assumption of Theorem 1.2.2 is better.

A related interesting result can be found in [49].

**Theorem 1.2.5** (see [49, Corollary 3.3]) *Assume that*

1.  $\limsup_{x \rightarrow +\infty} g(x) < \bar{h}_0$ ,
2. *there exists  $r > 0$  such that  $h_0(t) \leq g(r)$  for a.e.  $t \in [0, \omega]$ .*

*Then the problem (1.165), (1.3) has at least one positive solution.*

Theorem 1.2.3 shows that Theorem 1.2.5 is still valid also in the case when the term  $f(x)x'$  is incorporated to the equation, and even in the case when  $f$  has a singularity at zero.

---

## CHAPTER 2

# The Lazer and Solimini equation

In this chapter we consider a class of second order scalar differential equations with periodic forcing, zero damping, and a restoring force which become infinite at a finite displacement, which we take to be zero. As a model we have in mind the equation

$$u''(t) \pm \frac{1}{u^\lambda(t)} = h(t) \quad \text{for a. e. } t \in [0, \omega], \quad (2.1)$$

where  $h \in L([0, \omega]; \mathbb{R})$ . This simple equation can describe the dynamic of a particle with unitary mass moving on the right side of another fixed particle subjected to an external restoring force, when the positive sign is considered. However, (2.1) with negative sign can model a charged electrical particle under the influence of an electrical field generated by a fixed particle at origin with contrary electrical charge. Logically the type of external force  $h$  will determinate the dynamic of the equation. In our case, we will always assume that  $h$  is periodic (i.e. it is extended periodically on whole  $\mathbb{R}$ ).

From mathematical point of view equation (2.1) is considered as the start point to develop the actual theory of *singular equations*, however, previously, there were some authors who studied very particular singular equations, see [35, 21]. More concretely the topic of singular equation was instigated by the pioneer paper of Lazer and Solimini [40]. In spite of the passed time of the publication of the paper, there has been no contributions to this equation unless that a more general equation is considered.

In this chapter will be to present the results of Lazer and Solimini [40] with the respective

used tools. In this way we put to the reader in a suitable status to understand the main contribution of this Thesis: *to find a pioneer relation to attractive singular equations between the regularity of the external restoring force  $h$  and the order of the singularity,  $\lambda$ .*

The presented results in [40] concerns to the equation

$$u''(t) + g(u(t)) = h(t) \quad \text{for a. e. } t \in [0, \omega], \quad (2.2)$$

with  $g \in C(\mathbb{R}^+; \mathbb{R}^+)$ . This equation has essentially the same properties that (2.1) because both functions  $1/x^\lambda$  and  $g(x)$  has the some asymptotic properties. Thus the arguments to (2.1) can be easily extended to (2.2).

## 2.1 Repulsive singularity

In this part we consider a class of problems which includes (2.1) taking the minus sign, i.e.

$$u''(t) - \frac{1}{u^\lambda(t)} = h(t) \quad \text{for a. e. } t \in [0, \omega] \quad (2.3)$$

with  $h \in L([0, \omega]; \mathbb{R})$ . As always, we understand by periodic solution a positive function which is solution of the problem (2.3), (1.3).

Concerning to equation (2.2) they proved under the assumptions  $\lim_{x \rightarrow +\infty} g(x) = 0$  and

$$\lim_{x \rightarrow 0^+} g(x) = -\infty, \quad \int_0^1 g(x) dx = -\infty \quad (2.4)$$

the following theorem.

**Theorem 2.1.1** (see [40, Theorem 3.12]) *If  $g > 0$  then (2.2), (1.3) has at least one*



positive solution if and only if  $\bar{h} > 0$ .

They did a truncation to the function  $g$  in order to define it on whole  $\mathbb{R}$ . Then, using such a function, it defines a nonsingular problem which is possible to apply the existence theorem proved in [38]. Finally using the strong condition (2.4) they proved that the solution to the modified problem coincides with the solution to (2.2), (1.3). However, we can prove it by Corollary 1.1.2.

The previous result applied to equation (2.3) means:

**Corollary 2.1.1** *If  $\lambda \geq 1$  then (2.3), (1.3) has at least one positive solution if and only if  $\bar{h} < 0$ .*

Naturally it comes up the question on the "optimality" of the result. There they proved a necessary and sufficient condition under a strong condition, (2.4). The question is *does Theorem 2.1.1 remain still valid if this condition is violated?* The answer was published in the same paper proving the following result.

**Theorem 2.1.2** (see [40, Theorem 4.1]) *For every  $\omega > 0$  there exists  $M_0$  such that for any  $M > M_0$  there is, a  $T$ -periodic continuous and nonnegative function, such that (2.2), (1.3) has no solution and  $\bar{h} = M$ .*

In addition, since  $h$  is taken as a continuous function, this proves that we cannot aspire to find a relationship between the smoothness of  $h$  and the order of the singularity  $\lambda$  which allows to guarantee existence of periodic solutions to (2.3) even when the singularity is weak ( $\lambda < 1$ ). This is the reason why there are less papers dealing with weak singularities, being it a rather unexplored field.

## 2.2 Attractive singularity

Having in mind the equation (2.1) taking the sign +, i.e.

$$u''(t) + \frac{1}{u^\lambda(t)} = h(t) \quad \text{for a. e. } t \in [0, \omega] \quad (2.5)$$

with  $h \in L([0, \omega]; \mathbb{R})$ , Lazer and Solimini proved that if  $h$  is continuous and  $g$  satisfies

$$\lim_{x \rightarrow 0^+} g(x) = +\infty, \quad \lim_{x \rightarrow +\infty} g(x) = 0, \quad (2.6)$$

then there exists a positive solution to (2.1), (1.3) in the space  $C^2([0, \omega]; \mathbb{R})$  if and only if  $\bar{h} > 0$ . The way of the proof of Lazer and Solimini allows us to formulate their result as follows:

**Theorem 2.2.1** (see [40, Theorem 2.1]) *Let  $g \in C(\mathbb{R}^+; \mathbb{R}^+)$  satisfy (2.6) and let*

$$h \in L^\infty([0, \omega]; \mathbb{R}). \quad (2.7)$$

*Then there exists a positive solution  $u \in AC^1([0, \omega]; \mathbb{R})$  to (2.1), (1.3) if and only if  $\bar{h} > 0$ .*

The proof was done using a simple and elegant application of the method of lower and upper functions. Also we can obtain it as immediate consequence of our Theorem 1.2.3. Nevertheless both arguments are essentially the same, only we can do in a more general framework.

At this point, in view of the results in the previous subsection, it comes up the natural question *why did they not consider  $h \in L([0, \omega]; \mathbb{R})$ ?* There are other papers in the literature where the problem of the type (2.1), (1.3) is studied in the framework of the

Carathéodory theory, i.e., if  $h \in L([0, \omega]; \mathbb{R})$  and a positive function  $u \in AC^1([0, \omega]; \mathbb{R})$  is understood as a solution to (2.1), (1.3) (see, e.g., [26, 49, 48, 28, 31] and references therein). However, also in the works [26, 49], the boundedness of the function  $h$  is needed. In [48, 28, 31], the condition (2.7) is replaced by another condition dealing with the oscillation of the primitive of  $h$ .

On the other hand, the major part of the results dealing with the continuous input functions can be formulated also in the framework of the Carathéodory theory without any essential changes. This fact may encourage one's expectation that also Theorem 2.2.1 can be extended to the case when  $h \in L([0, \omega]; \mathbb{R})$  without any other additional conditions. Such a question was formally posted in [28, Open Problem 4.1]. Despite of all expectations, the answer is negative. The condition (2.7) is essential and, as shown we will prove (see Example 2.2.1), Theorem 2.2.1 is not valid anymore if the condition (2.7) is withdrawn unless the additional assumptions are involved—such an additional condition is, e.g., the relation (2.27) in Theorem 2.2.2 formulated below. Moreover, we prove that the condition (2.27) is optimal in a certain sense. More precisely, if we consider the equation (2.5) from our results, for every  $p \in [1, +\infty)$ , we obtain

- If  $\lambda \geq 1/(2p - 1)$  and  $h \in L^p([0, \omega]; \mathbb{R})$  then (2.5), (1.3) has a positive solution if and only if  $\bar{h} > 0$ , and such a solution is unique.
- If  $0 < \lambda < 1/(2p - 1)$  then there exists a function  $h \in L^p([0, \omega]; \mathbb{R})$  with  $\bar{h} > 0$  such that (2.5), (1.3) has no positive solution.

At this point we would like to emphasize the following: *for  $h \in L^p([0, \omega]; \mathbb{R})$ , there exists a relation between  $p$  and the order of singularity,  $\lambda$ . In other words, there exists a critical value depending on  $p$  such that if the power of the singularity  $\lambda$  is greater than or equal to this value then there exists a positive periodic solution. Moreover, if  $h \in L^\infty([0, \omega]; \mathbb{R})$ , then also  $h \in L^p([0, \omega]; \mathbb{R})$  for every  $p \in [1, +\infty)$ , and so applying*

our results for  $p$  sufficiently large we obtain that (2.5), (1.3) has a positive solution for every  $\lambda > 0$  (provided  $h \in L^\infty([0, \omega]; \mathbb{R})$ ). Thus Theorem 2.2.1 can be understood as a limit case of Theorem 2.2.2 formulated below.

Theorem 2.2.2 deals also with the uniqueness of a solution in the case when  $g$  is strictly decreasing function. This fact is worth mentioning here because in the original paper of Lazer and Solimini, the question of the uniqueness was not discussed.

## 2.2.1 Auxiliary results

The following results are important to prove our main results

**Lemma 2.2.1** *Let  $h \in L^p([0, \omega]; \mathbb{R})$  and let  $u \in AC^1([0, \omega]; \mathbb{R})$  satisfy*

$$u''(t) \leq h(t) \quad \text{for a. e. } t \in [0, \omega]. \quad (2.8)$$

Then

$$|u'(t)| \leq \left( \frac{(2p-1)}{p} \|h\|_p \right)^{p/(2p-1)} (u(t))^{(p-1)/(2p-1)} \quad \text{for } t \in [0, \omega]. \quad (2.9)$$

**Proof 26** Let  $t_0 \in \mathbb{R}$  be arbitrary but such that  $u'(t_0) \neq 0$ , and let  $\sigma = \text{sgn } u'(t_0)$ . Then there exists  $s_0 \in I(t_0, t_0 - \sigma\omega)$  such that

$$\sigma u'(t) > 0 \quad \text{for } t \in I(s_0, t_0), \quad u'(s_0) = 0.$$

Multiplying both sides of (2.8) by  $|u'(t)|^{(p-1)/p}$  and integrating over  $I(s_0, t_0)$  we get

$$\begin{aligned} \frac{p}{2p-1} |u'(t_0)|^{(2p-1)/p} &= \sigma \int_{s_0}^{t_0} u''(t) |u'(t)|^{(p-1)/p} dt \leq \sigma \int_{s_0}^{t_0} h(t) |u'(t)|^{(p-1)/p} dt \leq \\ \|h\|_p \left( \sigma \int_{s_0}^{t_0} |u'(t)| dt \right)^{(p-1)/p} &= \|h\|_p (u(t_0) - u(s_0))^{(p-1)/p} \leq \|h\|_p (u(t_0))^{(p-1)/p}. \end{aligned} \quad (2.10)$$

Thus (2.9) follows from (2.10).

**Lemma 2.2.2** *Let  $u \in AC^1([0, \omega]; \mathbb{R})$ ,  $g \in C(\mathbb{R}^+; \mathbb{R}^+)$ , and let*

$$g_*(x) \stackrel{\text{def}}{=} \inf \{g(s) : s \in (0, x]\}. \quad (2.11)$$

*Let, moreover,  $s_0, t_0 \in \mathbb{R}$  be such that  $s_0 < t_0$  and*

$$u'(s_0) = 0, \quad u'(t_0) = 0. \quad (2.12)$$

*Then*

$$\int_{s_0}^{t_0} u''(t) g_*^{p-1}(u(t)) dt \geq 0. \quad (2.13)$$

**Proof 27** Let  $g_n \in C(\mathbb{R}^+; \mathbb{R}^+)$  be a sequence of non-increasing functions which are continuous together with their derivatives and such that

$$\lim_{n \rightarrow +\infty} \|g_n^{p-1} \circ u - g_*^{p-1} \circ u\|_\infty = 0. \quad (2.14)$$

Obviously, since  $g_n$  are non-increasing, in view of (2.12) we have

$$\int_{s_0}^{t_0} u''(t) g_n^{p-1}(u(t)) dt = -(p-1) \int_{s_0}^{t_0} u'^2(t) g_n^{p-2}(u(t)) g_n'(u(t)) dt \geq 0. \quad (2.15)$$

Now, (2.13) follows from (2.14) and (2.15), because

$$\left| \int_{s_0}^{t_0} u''(t) g_n^{p-1}(u(t)) dt - \int_{s_0}^{t_0} u''(t) g_*^{p-1}(u(t)) dt \right| \leq \|g_n^{p-1} \circ u - g_*^{p-1} \circ u\|_\infty \int_{s_0}^{t_0} |u''(t)| dt.$$

**Lemma 2.2.3** *Let  $h \in L^p([0, \omega]; \mathbb{R})$ ,  $g \in C(\mathbb{R}^+; \mathbb{R}^+)$ , and let  $u \in AC^1([0, \omega]; \mathbb{R})$  be a periodic function (i.e. (1.3) holds) such that*

$$u''(t) + g(u(t)) \leq h(t) \quad \text{for a. e. } t \in [0, \omega]. \quad (2.16)$$

Let, moreover,  $s_0, t_0 \in \mathbb{R}$  be such that  $s_0 < t_0$  and (2.12) is fulfilled. Then

$$\int_{s_0}^{t_0} g(u(t))g_*^{p-1}(u(t))dt \leq \|h\|_p^p \quad (2.17)$$

where  $g_*$  is given by (2.11).

**Proof 28** Multiplying both sides of (2.16) by  $g_*^{p-1}(u(t))$  and integrating from  $s_0$  to  $t_0$ , in view of (2.12) and according to Lemma 2.2.2, we arrive at

$$\int_{s_0}^{t_0} g(u(t))g_*^{p-1}(u(t))dt \leq \int_{s_0}^{t_0} h(t)g_*^{p-1}(u(t))dt \leq \|h\|_p \left( \int_{s_0}^{t_0} g_*^p(u(t))dt \right)^{(p-1)/p},$$

whence in view of (2.11) we get (2.17).

**Lemma 2.2.4** Let  $u_i \in AC^1([0, \omega]; \mathbb{R}_+)$  ( $i = 1, 2$ ) be fulfilling (1.3) such that

$$\text{meas} \{t \in [0, \omega] : u_i(t) = 0\} = 0 \quad (i = 1, 2), \quad (2.18)$$

$$u_i''(t) + \tilde{g}(u_i(t)) = h(t) \quad \text{for a. e. } t \in [0, \omega] \quad (i = 1, 2), \quad (2.19)$$

where

$$\tilde{g}(x) = \begin{cases} g(x) & \text{for } x > 0, \\ 0 & \text{for } x = 0, \end{cases} \quad (2.20)$$

$g \in C(\mathbb{R}^+; \mathbb{R}^+)$  is a decreasing function, and  $h \in L([0, \omega]; \mathbb{R})$ . Then  $u_1 \equiv u_2$ .

**Proof 29** First assume that

$$u_1(t) \geq u_2(t) \quad \text{for } t \in [0, \omega]. \quad (2.21)$$

Put

$$z(t) = u_1(t) - u_2(t) \quad \text{for } t \in [0, \omega]. \quad (2.22)$$

Then, in view of (2.18)–(2.21) we have

$$z''(t) \geq 0 \quad \text{for a. e. } t \in [0, \omega], \quad z(0) - z(\omega) = 0 = z'(0) - z'(\omega). \quad (2.23)$$

However, (2.23) implies that  $z$  is a constant function, i.e., with respect to (2.18), (2.19), (2.20), and (2.22), we have

$$0 = u_1''(t) - u_2''(t) = -g(u_1(t)) + g(u_2(t)) \quad \text{for a. e. } t \in [0, \omega]. \quad (2.24)$$

Now from (2.24), according to the fact that  $g$  is assumed to be decreasing, it follows that  $u_1 \equiv u_2$ .

Further suppose that (2.21) is not valid. Then there exist  $t_0, t_1 \in \mathbb{R}$  such that  $t_0 < t_1$  and

$$u_1(t) > u_2(t) \quad \text{for } t \in (t_0, t_1), \quad u_1(t_0) = u_2(t_0), \quad u_1(t_1) = u_2(t_1). \quad (2.25)$$

Define  $z$  by (2.22). Then, in view of (2.18)–(2.20), and (2.25) we have

$$z''(t) \geq 0 \quad \text{for a. e. } t \in [t_0, t_1], \quad z(t_0) = 0, \quad z(t_1) = 0. \quad (2.26)$$

However, (2.26) implies  $z(t) \leq 0$  for  $t \in [t_0, t_1]$ , which, on account of (2.22) contradicts (2.25).

## 2.2.2 Main result

This subsection is devoted to prove the following theorem

**Theorem 2.2.2** *Let  $h \in L^p([0, \omega]; \mathbb{R})$  and let  $g \in C(\mathbb{R}^+; \mathbb{R}^+)$  verify (2.6). Let, more-*

over,

$$\lim_{x \rightarrow 0^+} \int_x^1 g(w(s)) g_*^{p-1}(w(s)) ds = +\infty, \quad (2.27)$$

where

$$w(s) \stackrel{\text{def}}{=} s^{(2p-1)/p} \quad \text{for } s \in \mathbb{R}^+. \quad (2.28)$$

Then the problem (2.2), (1.3) has a positive solution if and only if  $\bar{h} > 0$ . If, in addition,  $g$  is a decreasing function, then such a solution is unique.

**Proof 30** If  $u$  is a positive solution to (2.2), (1.3) then the integration of (2.2) from 0 to  $\omega$  results in

$$\int_0^\omega g(u(s)) ds = \int_0^\omega h(s) ds,$$

whence it follows that  $\bar{h} > 0$  as the function  $g$  is positive.

Now suppose that  $\bar{h} > 0$ . Together with (2.2), for every  $k \in \mathbb{N}$ , consider the auxiliary equation

$$u''(t) + g(u(t)) = h_k(t) \quad \text{for a. e. } t \in [0, \omega], \quad (2.29)$$

where

$$h_k(t) = \begin{cases} k & \text{if } h(t) > k, \\ h(t) & \text{if } h(t) \leq k, \end{cases} \quad \text{for a. e. } t \in [0, \omega], \quad k \in \mathbb{N}. \quad (2.30)$$

Obviously,

$$h_k(t) \leq h_m(t) \leq h(t) \quad \text{for a. e. } t \in [0, \omega], \quad k \leq m, \quad (2.31)$$

$$|h_k(t)| \leq |h(t)| \quad \text{for a. e. } t \in [0, \omega], \quad k \in \mathbb{N}, \quad (2.32)$$

$$\lim_{k \rightarrow +\infty} \bar{h}_k = \bar{h}. \quad (2.33)$$



According to (2.6), (2.33), and  $\bar{h} > 0$ , there exist  $x_0 > 0$  and  $k_0 \in \mathbb{N}$  such that

$$g(x) \leq \bar{h}_{k_0} \quad \text{for } x \geq x_0. \quad (2.34)$$

Let  $z$  be a solution to the Dirichlet problem

$$z''(t) = h_{k_0}(t) - \bar{h}_{k_0}, \quad z(0) = 0, \quad z(\omega) = 0 \quad (2.35)$$

and put

$$\beta(t) = \tilde{z}(t) + r \quad \text{for } t \in [0, \omega], \quad (2.36)$$

where  $\tilde{z}$  is an  $\omega$ -periodic prolongation of  $z$  to the real axis and  $r > 0$  is large enough such that

$$x_0 \leq \beta(t) \quad \text{for } t \in [0, \omega]. \quad (2.37)$$

Obviously,  $\beta \in AC^1([0, \omega]; \mathbb{R})$  and in view of (2.31) and (2.34)–(2.37),

$$\beta''(t) + g(\beta(t)) \leq h_k(t) \quad \text{for a. e. } t \in [0, \omega], \quad k \geq k_0. \quad (2.38)$$

On the other hand, on account of (2.6) and (2.30), for every  $k \geq k_0$  there exists  $x_k \in (0, x_0)$  such that

$$g(x_k) \geq h_k(t) \quad \text{for a. e. } t \in [0, \omega]. \quad (2.39)$$

If we put  $\alpha_k(t) = x_k$  for  $t \in [0, \omega]$  then, in view of (2.37) and (2.39), we have

$$\alpha_k''(t) + g(\alpha_k(t)) \geq h_k(t) \quad \text{for a. e. } t \in [0, \omega], \quad k \geq k_0, \quad (2.40)$$

$$\alpha_k(t) \leq \beta(t) \quad \text{for } t \in [0, \omega], \quad k \geq k_0. \quad (2.41)$$

Thus, for every  $k \geq k_0$ , there exists a pair of well-ordered upper and lower functions to

the problem (2.2), (1.3), which is a particular case of the problem (1.1), (1.3). According to Proposition 1.2.1, there exists a sequence of solutions  $\{u_k\}_{k=k_0}^{+\infty}$  to (2.29), (1.3) such that

$$0 < \alpha_k(t) \leq u_k(t) \leq \beta(t) \quad \text{for } t \in [0, \omega]. \quad (2.42)$$

From Lemma 2.2.1 and (2.42) it follows that

$$\|u'_k\|_\infty \leq \left( \frac{(2p-1)}{p} \|h\|_p \right)^{p/(2p-1)} \|\beta\|_\infty^{(p-1)/(2p-1)} \quad \text{for } k \geq k_0. \quad (2.43)$$

Further, we show that the set of functions  $\{u_k\}_{k=k_0}^{+\infty}$  is bounded from below. The integration of (2.29) from 0 to  $\omega$ , in view of (2.31), yields

$$\int_0^\omega g(u_k(s)) ds \leq \omega \bar{h}. \quad (2.44)$$

On the other hand, (2.6) implies the existence of  $y_0 > 0$  (which does not depend on  $k$ ) such that

$$g(x) > \bar{h} \quad \text{for } x \in (0, y_0). \quad (2.45)$$

From (2.44) and (2.45) it follows that for every  $k \geq k_0$  we have

$$\|u_k\|_\infty \geq y_0. \quad (2.46)$$

Let  $r_k \in [0, \omega]$  and  $\xi_k \in [r_k - \omega, r_k]$  be such that

$$u_k(r_k) = \max \{u_k(t) : t \in [0, \omega]\}, \quad u_k(\xi_k) = \min \{u_k(t) : t \in [0, \omega]\}. \quad (2.47)$$

Obviously,  $u'_k(\xi_k) = 0$ ,  $u'_k(r_k) = 0$ , and in view of (2.46), (2.47), and Lemmas 2.2.1 and

2.2.3 we have

$$\begin{aligned} \frac{2p-1}{p} \int_{(u_k(\xi_k))^{p/(2p-1)}}^{y_0^{p/(2p-1)}} g(w(s)) g_*^{p-1}(w(s)) ds &\leq \int_{\xi_k}^{r_k} \frac{u'_k(t) g(u_k(t)) g_*^{p-1}(u_k(t))}{(u_k(t))^{(p-1)/(2p-1)}} dt \leq \\ \left( \frac{(2p-1)}{p} \|h\|_p \right)^{p/(2p-1)} \int_{\xi_k}^{r_k} g(u_k(t)) g_*^{p-1}(u_k(t)) dt &\leq \left( \frac{(2p-1)}{p} \right)^{p/(2p-1)} \|h\|_p^{2p^2/(2p-1)}, \end{aligned}$$

where  $w$  is given by (2.28).

Thus the assumption (2.27) implies the existence of an  $\varepsilon > 0$  such that

$$\varepsilon \leq u_k(t) \quad \text{for } t \in [0, \omega], \quad k \geq k_0. \quad (2.48)$$

Finally, using (2.32), (2.42), and (2.48), from (2.29) we obtain

$$|u_k''(t)| \leq g^* + |h(t)| \quad \text{for a. e. } t \in [0, \omega], \quad k \geq k_0$$

where

$$g^* = \max \{g(x) : x \in [\varepsilon, \|\beta\|_\infty]\}.$$

Thus, the sequences  $\{u_k\}_{k=k_0}^{+\infty}$  and  $\{u'_k\}_{k=k_0}^{+\infty}$  are uniformly bounded and equicontinuous.

Therefore, according to *Arzelà–Ascoli Theorem*, we can assume without loss of generality that there exist  $u_0, v_0 \in C([0, \omega]; \mathbb{R})$  such that

$$\lim_{k \rightarrow +\infty} \|u_k - u_0\|_\infty = 0, \quad \lim_{k \rightarrow +\infty} \|u'_k - v_0\|_\infty = 0. \quad (2.49)$$

Moreover, since  $u_k$  are solutions to (2.29), (1.3), in view of (2.30), (2.48), and (2.49) we have  $u_0 \in AC^1([0, \omega]; \mathbb{R})$ ,  $u'_0 \equiv v_0$ , and  $u_0$  is a positive solution to (2.2), (1.3).

The uniqueness of a solution in the case when  $g$  is a decreasing function follows from Lemma 3.3.1.

**Remark 2.2.1** Always we can assume without lost of generality that a function  $y : [0, \omega] \rightarrow \mathbb{R}$  is defined by periodicity on whole  $\mathbb{R}$  using the introduced prolongation operators in (1.66) and (1.67). However, with the aim of keeping the exposition at a rather simple level we omite to write such an operators.

**Remark 2.2.2** Note that assumption  $h \in L^p([0, \omega]; \mathbb{R})$  in Theorem 2.2.2 can be weakened to  $h \in L([0, \omega]; \mathbb{R})$ ,  $[h]_+ \in L^p([0, \omega]; \mathbb{R})$ , where  $[h]_+$  is a non-negative part of the function  $h$ , i.e.,

$$[h]_+(t) \stackrel{\text{def}}{=} \frac{|h(t)| + h(t)}{2} \quad \text{for a. e. } t \in [0, \omega].$$

### 2.2.3 Optimality and Counter-Example

A particular case of the equation (2.2) is the equation (2.5) where  $\lambda > 0$ . For this equation, Theorems 2.2.1 and 2.2.2 yield the following assertions:

**Corollary 2.2.1** *Let*

$$h \in L^\infty([0, \omega]; \mathbb{R}), \quad \lambda > 0. \quad (2.50)$$

*Then the problem (2.5), (1.3) has a positive solution if and only if  $\bar{h} > 0$ , and such a solution is unique.*

**Corollary 2.2.2** *Let  $p \in [1, +\infty)$  and*

$$h \in L^p([0, \omega]; \mathbb{R}), \quad \lambda \geq 1/(2p - 1). \quad (2.51)$$

*Then the problem (2.5), (1.3) has a positive solution if and only if  $\bar{h} > 0$ , and such a solution is unique.*

**Remark 2.2.3** In spite of the fact that nor in Theorem 2.2.1 nor in the original result of Lazer and Solimini, the uniqueness of a positive solution to (2.2), (1.3) is discussed, Corollary 2.2.1 is valid—the uniqueness of a solution follows from Lemma 3.3.1.

Before we formulate other theorem, we introduce an example:

**Counter-example 2.2.1** Let  $p \geq 1$  and  $\lambda \in \left(0, \frac{1}{2p-1}\right)$ . Choose  $\mu \in \left(2 - \frac{1}{p\lambda}, \frac{1}{p}\right)$ ,  $\varepsilon \in \left(0, \frac{\omega}{4}\right)$ , and put

$$\varphi(t) = \begin{cases} -t^{-\mu} & \text{for } t \in (0, \varepsilon], \\ 0 & \text{for } t \in \left(\varepsilon, \frac{\omega}{2} - \varepsilon\right), \\ \left(\frac{\omega}{2} - t\right)^{-\mu} & \text{for } t \in \left[\frac{\omega}{2} - \varepsilon, \frac{\omega}{2}\right), \end{cases} \quad \varphi(t) = \varphi(\omega - t) \quad \text{for } t \in \left(\frac{\omega}{2}, \omega\right),$$

$$v(t) = \int_t^{\frac{\omega}{2}} \int_s^{\frac{\omega}{2}} \varphi(\xi) d\xi ds \quad \text{for } t \in [0, \omega).$$

If we periodically extend the functions  $\varphi$  and  $v$  to the whole real axis, we obviously obtain

$$\varphi \in L^p([0, \omega]; \mathbb{R}), \quad v \in AC^1([0, \omega]; \mathbb{R}_+), \quad v''(t) = \varphi(t) \quad \text{for a. e. } t \in [0, \omega], \quad (2.52)$$

and by a direct calculation, the following relations can be verified:

$$v(t) > 0 \quad \text{for } t \in [0, \omega/2) \cup (\omega/2, \omega], \quad v(\omega/2) = 0, \quad (2.53)$$

$$v(t) = \frac{|\omega/2 - t|^{2-\mu}}{(2-\mu)(1-\mu)} \quad \text{for } t \in \left(\frac{\omega}{2} - \varepsilon, \frac{\omega}{2} + \varepsilon\right).$$

Now it can be easily seen that

$$\frac{1}{v^\lambda} \in L^p([0, \omega]; \mathbb{R}). \quad (2.54)$$

Put

$$h(t) \stackrel{\text{def}}{=} \varphi(t) + \frac{1}{v^\lambda(t)} \quad \text{for a. e. } t \in [0, \omega]. \quad (2.55)$$

Obviously, in view of (2.52)–(2.54) we have  $h \in L^p([0, \omega]; \mathbb{R})$  and  $\bar{h} > 0$ . Consider the problem (2.5), (1.3) and suppose that there exists a positive solution  $u$  to (2.5), (1.3). According to (2.52), (2.53), (2.55), and Lemma 3.3.1, it follows that  $u(t) = v(t)$  for  $t \in [0, \omega]$ . However, that is impossible, because  $v(\omega/2) = 0$ . Thus, (2.5), (1.3) has no positive solution with  $h$  defined by (2.55).

Example 2.2.1 proves the following assertion:

**Theorem 2.2.3** *Let  $p \in [1, +\infty)$ ,  $0 < \lambda < 1/(2p - 1)$ . Then there exists  $h \in L^p([0, \omega]; \mathbb{R})$  with  $\bar{h} > 0$  such that the problem (2.5), (1.3) has no positive solution.*

According to Theorem 2.2.3, it can be seen that the condition (2.27) in Theorem 2.2.2 is essential and cannot be omitted. Moreover, Theorem 2.2.3 shows that the condition (2.51) in Corollary 2.2.2 is unimprovable.

---

## CHAPTER 3

# The Rayleigh-Plesset equation

In this section we will use our mathematical result in Chapter 1 to study the Rayleigh-Plesset equation, which models the oscillations of a spherical bubble in a liquid subjected to a periodic acoustic field. The Rayleigh-Plesset equation plays a prominent role in Dynamics of Fluids. It can be derived by taking spherical coordinates in Euler equations and assuming some physically admissible simplifications, as shown in many reviews and monographs (see for instance [9, 12, 22, 47, 58]). A variety of physical, biological and medical models rely on this equation (see bibliographies of the cited references), in connection with the physical phenomena of cavitation and sonoluminescence.

Following [22], the evolution in time of the radius  $R(t)$  of the bubble is ruled by

$$\rho \left[ R\ddot{R} + \frac{3}{2}\dot{R}^2 \right] = [P_v - P_\infty(t)] + P_{g_0} \left( \frac{R_0}{R} \right)^{3k} - \frac{2S}{R} - \frac{4\mu\dot{R}}{R}. \quad (3.1)$$

Here, at the left-hand side  $\dot{R}$  and  $\ddot{R}$  are the first and second derivatives of the bubble radius with respect to time and  $\rho$  is the density of the liquid. At the right-hand side we have four different terms. The first one is  $P_v - P_\infty(t)$ , which measures the difference between the vapour pressure  $P_v$  inside the bubble and the applied pressure, which is time-periodic. The second term is related with the non-condensability of the gas. More exactly,  $P_{g_0}$  and  $R_0$  correspond, respectively, to the gas pressure and initial radius of the bubble, while  $k$  is the polytropic coefficient, which contains information about thermic transmission behaviour of the system liquid–gas. If the behaviour is isothermal then

the coefficient  $k$  is equal to one. The most usual case considered in the cited references is when polytropic coefficient is greater than or equal to one, but possibly it is any real number. In this paper, we consider the adiabatic case (when  $k \geq 1$ ). The third term corresponds to surface tension, i.e., the energy which is needed to increase the surface of a liquid by area unit. Finally, the last term corresponds to the viscosity of liquid.

When the surface tension and viscosity effects are neglected (a physically admissible simplification for bubbles of big radius), we may obtain the classical Rayleigh equation

$$\rho \left[ R\ddot{R} + \frac{3}{2}\dot{R}^2 \right] = P_v - P_\infty(t),$$

which was proposed in 1907 by Rayleigh. Furthermore, we observe that when the applied pressure is constant, the Rayleigh equation has a first integral

$$\dot{R}^2 = \frac{2}{3} \frac{P_v - P_\infty}{\rho} \left[ 1 - \left( \frac{R_0}{R} \right)^3 \right].$$

Nevertheless, when the applied pressure  $P_\infty(t)$  is time-varying, most of the present knowledge about the dynamics of this models is based on numerical computations.

If the change of variables  $R = u^{\frac{2}{5}}$  is introduced in the Rayleigh-Plesset equation, we obtain

$$\ddot{u} = \frac{5 [P_v - P_\infty(t)]}{2\rho} u^{\frac{1}{5}} + \left( \frac{5P_{g_0} R_0^{3k}}{2\rho} \right) \frac{1}{u^{\frac{6k-1}{5}}} - \frac{5S}{u^{\frac{1}{5}}} - 4\mu \frac{\dot{u}}{u^{\frac{4}{5}}},$$

Consequently, the class of equations

$$u''(t) + \frac{cu'(t)}{u^\mu(t)} + \frac{g_1}{u^\nu(t)} - \frac{g_2}{u^\gamma(t)} = h_0(t)u^\delta(t) \quad \text{for a. e. } t \in [0, \omega] \quad (3.2)$$

with non-negative constants  $g_1, g_2, \delta, \nu > 0$  and real numbers  $c, \mu, \gamma$ , and  $h_0 \in$



$L([0, \omega]; \mathbb{R})$ , plays an important role in fluid mechanics.

The aim of this chapter is to find periodic solutions to equations (3.1) and (3.2). For that it will be convenient to consider the general equation (1.1) and to use the proved results of Chapter 1.

## 3.1 The model equation

This part is devoted to study equation (3.2). For that we will consider three cases depending on the considered type of nonlinearity: repulsive case ( $\gamma > \nu > 0$ ), attractive case ( $0 < \gamma < \nu$ ) and the case  $\gamma \leq 0$ .

### 3.1.1 The repulsive case

In this part we consider the equation (3.2) with repulsive singularity, concretely when  $\nu < \gamma$ . From Theorem 1.1.3 it finds a direct application to (3.2) in the sublinear case  $\delta < 1$ .

**Theorem 3.1.1** *Let us assume  $0 \leq \delta < 1$ ,  $\gamma > \nu$ ,  $\gamma \geq 1$ ,  $g_2 > 0$ . If  $\bar{h}_0 \leq 0$  and  $g_1 + |\bar{h}_0| > 0$ , then there exists at least one positive solution to the problem (3.2), (1.3).*

**Proof 31** It can be proved by applying Theorem 1.1.3 with  $\eta(t, x) = [h_0(t)]_- x^\delta$ ,  $\rho(x) = x^\delta$  and  $h(t, x) = h_0(t)x^\delta$ . Indeed, hypotheses 1, 3, 4, 5, and 6 of Theorem 1.1.3 are straightforward. Finally, hypothesis 2 one can easily prove by using the inequality  $g_1 + |\bar{h}_0| > 0$ .

The linear case  $\delta = 1$  is also covered by Theorem 1.1.3 as follows.

**Theorem 3.1.2** *Let us assume  $\delta = 1$ ,  $\gamma > \nu$ ,  $\gamma \geq 1$ ,  $g_2 > 0$ . If  $\bar{h}_0 \leq 0$ ,  $g_1 + |\bar{h}_0| > 0$  and*

$$\int_0^\omega [h_0(s)]_- ds < \frac{4}{\omega}, \quad (3.3)$$

*then there exists at least one positive solution to the problem (3.2), (1.3).*

**Proof 32** It can be proved by applying Theorem 1.1.3 with  $\eta(t, x) = [h_0(t)]_- x$ ,  $\rho(x) = x$ ,  $h(t, x) = h_0(t)x$  and reasoning as we did in Theorem 3.1.1.

Such as Condition 1. in Theorem 1.1.1, the above-mentioned condition (3.3) is used to avoid the resonant phenomenon at infinity. In spite of both conditions are used to the some work, they are independents and the best in some sense.

We previously mentioned the importance to take into account the strong singular condition ( $\gamma \geq 1$ ) when (3.2) is considered. However, when  $h_0$  verifies

$$h_0(t) \leq \sup \left\{ \frac{g_1}{x^{\nu+\delta}} - \frac{g_2}{x^{\gamma+\delta}} : x \in \mathbb{R}^+ \right\} \quad \text{for a. e. } t \in [0, \omega], \quad (3.4)$$

one can find a constant lower function, taking  $\alpha \equiv r_0 \in \mathbb{R}^+$  such that

$$\frac{g_1}{r_0^{\nu+\delta}} - \frac{g_2}{r_0^{\gamma+\delta}} = \sup \left\{ \frac{g_1}{x^{\nu+\delta}} - \frac{g_2}{x^{\gamma+\delta}} : x \in \mathbb{R}^+ \right\}.$$

Combining this with the obtained results to construct lower and upper functions, we can find an ordered pair of lower and upper functions under some conditions.

**Theorem 3.1.3** *Let  $0 \leq \delta < 1$ ,  $\gamma > \nu$ ,  $g_1 > 0$  and  $g_2 > 0$ . If  $\bar{h}_0 > 0$  and (3.4), then there exists at least one positive solution to the problem (3.2), (1.3).*

**Proof 33** In view of the previous discussion, we only need to prove the existence of an upper function.

To prove such a existence, we apply Corollary 1.2.5 taking  $\{y_n\}_{n=1}^{+\infty}$  an arbitrary sequence of positive numbers satisfying (1.153),  $c \in (0, \bar{h}_0)$ ,  $\rho_1(x) = x^\delta$ ,  $h_1 \equiv h_0$ ,  $x_0 > r_0$  large enough, and  $\varepsilon > 0$  small enough such that

$$(1 + \varepsilon)^\delta \Phi_- \leq \Phi_+. \quad (3.5)$$

Consequently, the assertion follows from Proposition 1.2.1.

**Theorem 3.1.4** *Let  $\delta = 1$ ,  $\gamma > \nu$ ,  $g_1 > 0$  and  $g_2 > 0$ . Assume that  $\bar{h}_0 > 0$ ,*

$$h_0(t) \leq \sup \left\{ \frac{g_1}{x^{\nu+1}} - \frac{g_2}{x^{\gamma+1}} : x \in \mathbb{R}^+ \right\} \quad \text{for a. e. } t \in [0, \omega], \quad (3.6)$$

and

$$\frac{\omega}{4} \int_0^\omega [h_0(s)]_+ ds \int_0^\omega [h_0(s)]_- ds < \int_0^\omega [h_0(s)]_+ ds - \int_0^\omega [h_0(s)]_- ds. \quad (3.7)$$

*Then there exists at least one positive solution to the problem (3.2), (1.3).*

**Proof 34** As in the proof of Theorem 3.1.3, we can check that there exists a constant  $r_0 \in \mathbb{R}^+$  such that  $\alpha(t) = r_0$  for  $t \in [0, \omega]$  is a lower function to the problem (3.2), (1.3).

On the other hand, from (3.7) it follows that there exists a sufficiently small constant  $c > 0$  such that  $c < \bar{h}_0$  and

$$\frac{\omega}{4} \Phi_+ \Phi_- \leq \Phi_+ - \Phi_-,$$

where

$$\varphi(t) = h_0(t) - c \quad \text{for a. e. } t \in [0, \omega].$$

Therefore, if we put  $\rho_1(x) = x$  and  $h_1 \equiv h_0$ , taking into account that

$$\lim_{x \rightarrow +\infty} \frac{g_1}{x^{1+\nu}} - \frac{g_2}{x^{1+\gamma}} = 0,$$

the existence of an upper function large enough follows from Corollary 1.2.6.

Consequently, the assertion follows from Proposition 1.2.1.

**Theorem 3.1.5** *Let  $\delta > 1$ ,  $\gamma > \nu$ ,  $g_1 > 0$  and  $g_2 > 0$ . If*

$$0 \leq h_0(t) \leq \sup \left\{ \frac{g_1}{x^{\nu+\delta}} - \frac{g_2}{x^{\gamma+\delta}} : x \in \mathbb{R}^+ \right\} \quad \text{for a. e. } t \in [0, \omega] \quad (3.8)$$

*and  $\bar{h}_0 > 0$ , then there exists at least one positive solution to the problem (3.2), (1.3).*

**Proof 35** Analogously to the previous proofs, there exists a constant  $r_0 \in \mathbb{R}^+$  such that the function  $\alpha(t) = r_0$  for  $t \in [0, \omega]$  is a lower function to the problem (3.2), (1.3).

On the other hand, from the first inequality of (3.8) it follows that  $h_0(t)x^\delta \geq h_0(t)$  for almost every  $t \in [0, \omega]$  and  $x \geq 1$ . Thus, the existence of an upper function to the problem (3.2), (1.3) follows from Corollary 1.2.5 by taking  $h_1 \equiv h_0$ ,  $\rho_1 \equiv 1$ , an arbitrary sequence  $\{y_n\}_{n=1}^{+\infty}$  of positive numbers such that (1.153) holds,  $c \in (0, \bar{h}_0]$ ,  $\varepsilon > 0$  arbitrary and  $x_0 > r_0$  large enough.

Consequently, the assertion follows from Proposition 1.2.1.

### 3.1.2 The attractive case

We will consider (3.2) with attractive singularity (i.e.  $\nu > \gamma$ ).

**Theorem 3.1.6** *Let  $0 \leq \delta < 1$ ,  $\gamma < \nu$  and  $g_1 > 0$ . If  $\bar{h}_0 > 0$  and*

$$\text{ess sup } \{h_0(t) : t \in [0, \omega]\} < +\infty, \quad (3.9)$$

*then there exists at least one positive solution to the problem (3.2), (1.3).*

**Proof 36** According to (3.9), we can choose  $K > 0$  such that

$$K \geq h_0(t) \quad \text{for a. e. } t \in [0, \omega].$$

As  $\lim_{x \rightarrow 0^+} \frac{g_1}{x^\nu} - \frac{g_2}{x^\gamma} = +\infty$ , there exists  $x_1 > 0$  such that

$$\frac{g_1}{x_1^{\nu+\delta}} - \frac{g_2}{x_1^{\gamma+\delta}} \geq K.$$

Obviously,  $\alpha \equiv x_1$  is a constant lower function to the problem (3.2), (1.3).

To prove the existence of an upper function we apply Corollary 1.2.5 taking  $\{y_n\}_{n=1}^{+\infty}$  an arbitrary sequence of positive numbers satisfying (1.153),  $c \in (0, \bar{h}_0)$ ,  $\rho_1(x) = x^\delta$ ,  $h_1 \equiv h_0$ ,  $x_0 > K$  large enough, and  $\varepsilon > 0$  small enough such that (3.5) holds.

Consequently, the assertion follows from Proposition 1.2.1.

**Theorem 3.1.7** *Let  $0 \leq \delta < 1$ ,  $\gamma < \nu$ ,  $g_1 > 0$  and  $g_2 > 0$ . If  $\bar{h}_0 \leq 0$ , (3.9) is fulfilled and*

$$h_0(t) \geq \inf \left\{ \frac{g_1}{x^{\nu+\delta}} - \frac{g_2}{x^{\gamma+\delta}} : x \in \mathbb{R}^+ \right\} \quad \text{for a. e. } t \in [0, \omega], \quad (3.10)$$

*then there exists at least one positive solution to the problem (3.2), (1.3).*

**Proof 37** In this case,

$$\inf \left\{ \frac{g_1}{x^{\nu+\delta}} - \frac{g_2}{x^{\gamma+\delta}} : x \in \mathbb{R}^+ \right\} > -\infty$$

and there exists  $x_0 > 0$  such that

$$\frac{g_1}{x_0^{\nu+\delta}} - \frac{g_2}{x_0^{\gamma+\delta}} = \inf \left\{ \frac{g_1}{x^{\nu+\delta}} - \frac{g_2}{x^{\gamma+\delta}} : x \in \mathbb{R}^+ \right\}.$$

According to (3.10), it can be easily verified that the function  $\beta(t) = x_0$  for  $t \in [0, \omega]$  is an upper function to the problem (3.2), (1.3).

To obtain a lower function, we proceed as in the proof of Theorem 3.1.6 choosing  $x_1$  small enough.

Consequently, the assertion follows from Proposition 1.2.1.

**Remark 3.1.1** Note that the conditions guaranteeing solvability of the problem (3.2), (1.3) in the case where  $0 \leq \delta < 1$ ,  $\gamma = \nu$ , and  $g_1 > g_2$  can be derived from Theorem 3.1.6.

However, in that case  $\gamma = \nu$  and  $g_1 < g_2$ , only the conditions sufficient for the existence of non-ordered lower and upper functions are known to the authors. Thus our analysis is incomplete and the case  $\gamma = \nu < 1$ ,  $g_1 < g_2$  remains as an open problem.

**Theorem 3.1.8** *Let  $\delta = 1$ ,  $g_1 > 0$  and  $g_2 = 0$ . If  $\bar{h}_0 > 0$  and (3.7) is fulfilled, then there exists at least one positive solution to the problem (3.2), (1.3).*

**Proof 38** Put  $\rho_0(x) = x$ . According to Lemma 1.2.3 we can choose  $c_0 > \bar{h}_0$  large enough such that the condition b) of Theorem 1.2.1 is fulfilled. Obviously, also  $r_0 > 0$  can be chosen such that (1.159) is satisfied.

On the other hand, put  $h_1 \equiv h_0$  and  $\rho_1(x) = x$ . Then, in view of (3.7), there exists a constant  $c_1 > 0$  such that  $c_1 \leq \bar{h}_1$  and the condition d) of Theorem 1.2.1 and (1.160) are fulfilled with a suitable  $r_1 > r_0$ .

Consequently, the assertion follows from Theorem 1.2.1.

**Theorem 3.1.9** *Let  $\delta = 1$ ,  $\gamma < \nu$ ,  $g_1 > 0$  and  $g_2 > 0$ . If  $\bar{h}_0 \geq 0$  and*

$$\frac{\omega}{4} \int_0^\omega [h_0(s)]_+ ds \int_0^\omega [h_0(s)]_- ds \leq \int_0^\omega [h_0(s)]_+ ds - \int_0^\omega [h_0(s)]_- ds, \quad (3.11)$$

then there exists at least one positive solution to the problem (3.2), (1.3).

**Proof 39** The proof is similar to that of Theorem 3.1.8. The only difference is that the inequality  $g_1x^{-(\nu+1)} - g_2x^{-(\gamma+1)} < 0$  for  $x$  sufficiently large allows one to choose a constant  $c_1$  equal to zero.

**Theorem 3.1.10** Let  $\delta = 1$ ,  $\gamma < \nu$ ,  $g_1 > 0$  and  $g_2 > 0$ . If  $\bar{h}_0 \leq 0$  and

$$h_0(t) \geq \inf \left\{ \frac{g_1}{x^{\nu+1}} - \frac{g_2}{x^{\gamma+1}} : x \in \mathbb{R}^+ \right\} \quad \text{for a. e. } t \in [0, \omega], \quad (3.12)$$

then there exists at least one positive solution to the problem (3.2), (1.3).

**Proof 40** As in the proof of Theorem 3.1.7, we can verify that there exists a constant  $x_1 > 0$  such that  $\beta(t) = x_1$  for  $t \in [0, \omega]$  is an upper function to the problem (3.2), (1.3).

On the other hand, put  $\rho_0(x) = x$ , and choose  $c \geq \bar{h}_0$  and  $x_0 > 0$  such that  $x_0 < x_1$  and the conditions of Corollary 1.2.3 are fulfilled. Note that the existence of  $c \geq \bar{h}_0$  large enough such that (1.142) holds follows from Lemma 1.2.3. Therefore, there exists a lower function.

Consequently, the assertion follows from Proposition 1.2.1.

**Theorem 3.1.11** Let  $1 < \delta$ ,  $\gamma < \nu$  and  $g_1 > 0$ ,  $g_2 > 0$ . If (3.10) is fulfilled, then there exists at least one positive solution to the problem (3.2), (1.3).

**Proof 41** The upper function is constructed as in the proof of Theorem 3.1.7.

On the other hand, put  $\rho_0(x) = x^\delta$ , and choose  $\{y_n\}_{n=1}^{+\infty}$  a sequence of positive numbers satisfying (1.137),  $c > \bar{h}_0$ , and  $\varepsilon \in (0, 1)$  such that

$$(1 - \varepsilon)^{-\delta} \Phi_+ \leq \Phi_-.$$

Then there exists  $x_0 > 0$  sufficiently small such that all the conditions of Corollary 1.2.2 are fulfilled. Consequently, there exists a lower function  $\alpha(t) \leq x_0$ .

Now the assertion follows from Proposition 1.2.1.

### 3.1.3 The case $\gamma \leq 0$

Finally two results dealing with the problem (3.2), (1.3) in the case when the parameter  $\gamma$  is non-positive. This case is also interesting from the physical point of view.

**Theorem 3.1.12** *Let  $0 \leq \delta < 1$ ,  $-\gamma > \delta$ ,  $g_1 > 0$  and  $g_2 > 0$ . If (3.9) is fulfilled, then there exists at least one positive solution to the problem (3.2), (1.3).*

**Proof 42** The assertion immediately follows from Theorem 1.2.1 b) and c) with  $h(t, x) = h_0(t)x^\delta$ ,  $h_1 \equiv h_0$ ,  $\rho_i(x) = x^\delta$  ( $i = 0, 1$ ),  $c_0 = \text{ess sup} \{h_0(t) : t \in [0, \omega]\}$ ,  $c_1 = \bar{h}_0 - 1$ , and  $g(x) = g_1x^{-\nu} - g_2x^{-\gamma}$ .

**Theorem 3.1.13** *Let  $0 \leq \delta < 1$ ,  $\gamma \leq 0$ ,  $|\gamma| \leq \delta$ ,  $g_1 > 0$  and  $g_2 > 0$ , and  $\bar{h}_0 > 0$ . If (3.9) is fulfilled, then there exists at least one positive solution to the problem (3.2), (1.3).*

**Proof 43** The assertion immediately follows from Theorem 1.2.1 b) and c) with  $h(t, x) = h_0(t)x^\delta$ ,  $h_1 \equiv h_0$ ,  $\rho_i(x) = x^\delta$  ( $i = 0, 1$ ),  $c_0 = \text{ess sup} \{h_0(t) : t \in [0, \omega]\}$ ,  $c_1 = 0$ , and  $g(x) = g_1x^{-\nu} - g_2x^{-\gamma}$ .

## 3.2 The physical model

A direct application of Theorem 3.1.1 gives the following result.



**Theorem 3.2.1** *Let us assume  $k \geq 1$  and  $P_v \leq \bar{P}_\infty$ . Then there exists at least one positive periodic solution to the equation (3.1).*

As far as we know, this is the first analytical proof of a well-known numerical evidence exposed in many related works, see for instance [22].

Also, Theorem 3.1.3 implies

**Theorem 3.2.2** *Let  $k > \frac{1}{3}$ ,  $P_v > \bar{P}_\infty$  and*

$$\frac{5(P_v - P_\infty(t))}{2\rho} \leq \left( \frac{6k-2}{5} \right) \left[ \frac{\left(\frac{2}{5}\right)^{\frac{2}{5}} (5S)^{\frac{6k}{5}}}{\left(\frac{6k}{5}\right)^{\frac{6k}{5}} \left(\frac{5P_{g_0} R_0^{3k}}{2\rho}\right)^{\frac{2}{5}}} \right]^{\frac{5}{6k-2}} \quad \text{for } t \in [0, \omega].$$

*Then there exists at least one positive periodic solution to the equation (3.1).*

Theorems 3.1.6 and 3.1.7, respectively, lead to

**Theorem 3.2.3** *Let  $\frac{1}{6} < k < \frac{1}{3}$ ,  $P_v > \bar{P}_\infty$ , and*

$$\text{ess inf} \{P_\infty(t) : t \in [0, \omega]\} > -\infty. \quad (3.13)$$

*Then there exists at least one positive periodic solution to the equation (3.1).*

**Theorem 3.2.4** *Let  $\frac{1}{6} < k < \frac{1}{3}$ ,  $P_v \leq \bar{P}_\infty$ , (3.13) holds, and*

$$\frac{5(P_v - P_\infty(t))}{2\rho} \geq - \left( \frac{2-6k}{5} \right) \left[ \frac{\left(\frac{6k}{5}\right)^{\frac{6k}{5}} \left(\frac{5P_{g_0} R_0^{3k}}{2\rho}\right)^{\frac{2}{5}}}{\left(\frac{2}{5}\right)^{\frac{2}{5}} (5S)^{\frac{6k}{5}}} \right]^{\frac{5}{2-6k}} \quad \text{for } t \in [0, \omega].$$

*Then there exists at least one positive periodic solution to the equation (3.1).*

Applying Theorem 3.1.6 with  $g_1 = 5S - \frac{5P_{g_0}R_0^{3k}}{2\rho}$ ,  $g_2 = 0$ , and  $\nu = 1/5$  we get

**Theorem 3.2.5** *Let  $k = \frac{1}{3}$ ,  $P_v > \bar{P}_\infty$ ,  $2\rho S > P_{g_0}R_0^{3k}$ , and let (3.13) holds. Then there exists at least one positive periodic solution to equation (3.1).*

**Remark 3.2.1** The open problem posed in Remark 3.1.1 corresponds to this last result when  $2\rho S < P_{g_0}R_0^{3k}$ .

Applying Theorems 3.1.12 and 3.1.13, we get, respectively,

**Theorem 3.2.6** *Let  $k < 0$  and let (3.13) holds. Then there exists at least one positive periodic solution to equation (3.1).*

**Theorem 3.2.7** *Let  $0 \leq k \leq \frac{1}{6}$ ,  $P_v > \bar{P}_\infty$ , and let (3.13) holds. Then there exists at least one positive periodic solution to equation (3.1).*

Let us finish by pointing out the presented results have a direct physical reading. For example, we can conclude that as a general rule a high density coefficient  $\rho$  of the liquid should benefit the presence of oscillating bubbles, an effect that seems physically plausible.

### 3.3 Playing with singularities

Taking into account that (2.1) is a particular equation of (3.2), we will use the studied theory in the previous chapter in order to motivate some questions. When  $\nu > \gamma$ ,  $g_1 > 0$ , having in mind the equation (2.5), one can expect that either  $\nu \geq 1$  or (3.9) is needed. However we should think that (3.2) has also singularity at the friction type term  $c\frac{u'}{u^\mu}$

and a new sub-linear term when  $\delta \in [0, 1)$ . Playing with these elements we can weaken the attractive singularity even when  $h_0 \in L([0, \omega]; \mathbb{R})$ .

To find a novel relation between the nonlinearities of (3.2) will be our objective, showing the following general property: *to consider singularity at friction type term helps to periodic solution exists.*

We consider the general equation (1.1) considered at the beginning of this work. By using some previous results proved in Chapter 1 (Section 1.2) and other ones news, we will obtain a general existence result to (1.1), (1.3) which we will set out in an independent section.

### 3.3.1 Main result and consequences

The next result is based on a new method of lower and upper functions which will be proven below.

**Theorem 3.3.1** *Let  $\rho_0 \in C^1(\mathbb{R}^+; \mathbb{R}^+)$  and  $\rho_1 \in C(\mathbb{R}^+; \mathbb{R}^+)$  be non-decreasing functions,  $h_0, h_1 \in L([0, \omega]; \mathbb{R})$ , and  $x_0 > 0$  be such that*

$$h_1(t)\rho_1(x) \leq h(t, x) \leq h_0(t)\rho_0(x) \quad \text{for a. e. } t \in [0, \omega], \quad x \geq x_0, \quad (3.14)$$

*and let there exist  $c_0, c_1 \in \mathbb{R}$  such that*

$$\frac{g(x)}{\rho_0(x)} \leq c_0 < \bar{h}_0 \quad \text{for } x \geq x_0, \quad (3.15)$$

$$\frac{g(x)}{\rho_1(x)} \leq c_1 \leq \bar{h}_1 \quad \text{for } x \geq x_0. \quad (3.16)$$

Let, moreover, there exist  $\lambda \in [0, 1]$  such that

$$\int_0^1 \frac{ds}{\rho_0^\lambda(s)} < +\infty, \quad (3.17)$$

$$\lim_{x \rightarrow 0^+} \frac{g(x)}{\rho_0^\lambda(x)} = +\infty, \quad (3.18)$$

and let either

$$\int_0^1 \left( \frac{[f(s)]_+}{\rho_0^\lambda(s)} + \frac{[g(s)]_+}{\rho_0^{2\lambda}(s)} \right) ds = +\infty, \quad \int_0^1 \frac{[f(s)]_-}{\rho_0^\lambda(s)} ds < +\infty \quad (3.19)$$

or

$$\int_0^1 \left( \frac{[f(s)]_-}{\rho_0^\lambda(s)} + \frac{[g(s)]_+}{\rho_0^{2\lambda}(s)} \right) ds = +\infty, \quad \int_0^1 \frac{[f(s)]_+}{\rho_0^\lambda(s)} ds < +\infty. \quad (3.20)$$

Furthermore, let us suppose that  $\rho_0$  fulfills at least one of the following conditions:

(a) there exists a sequence  $\{y_n\}_{n=1}^{+\infty}$  of positive numbers such that

$$\lim_{n \rightarrow +\infty} y_n = +\infty, \quad \lim_{n \rightarrow +\infty} \frac{\rho_0^{1-\lambda}(y_n)}{\sigma(y_n)} = 0, \quad (3.21)$$

and there exist  $\varepsilon_0 > 0$ ,  $\varepsilon_1 \in (0, \varepsilon_0]$ , and  $n_0 \in \mathbb{N}$  such that

$$\frac{\rho_0^{1-\lambda}((1 + \varepsilon_0)y_n)}{\rho_0^{1-\lambda}(y_n)} \Phi_- \leq \Phi_+ - \varepsilon_0 \quad \text{for } n \geq n_0, \quad (3.22)$$

$$(1 + \varepsilon_1)\sigma(y_n) \leq \sigma((1 + \varepsilon_0)y_n) \quad \text{for } n \geq n_0, \quad (3.23)$$

where  $\varphi(t) = h_0(t) - c_0$  for almost every  $t \in [0, \omega]$  and

$$\sigma(x) = \int_0^x \frac{ds}{\rho_0^\lambda(s)}; \quad (3.24)$$

(b) the function  $\frac{\rho_0^{1-\lambda}(x)}{\sigma(x)}$  is non-increasing and

$$\frac{\omega}{4}\Phi_+\Phi_-\frac{\rho_0^{1-\lambda}(x_0)}{\sigma(x_0)} < \Phi_+ - \Phi_-, \quad (3.25)$$

where  $\varphi(t) = h_0(t) - c_0$  for almost every  $t \in [0, \omega]$  and  $\sigma$  is given by (3.24).

Besides, let us suppose that  $\rho_1$  fulfills at least one of the following conditions:

(c) there exists a sequence  $\{z_n\}_{n=1}^{+\infty}$  of positive numbers such that

$$\lim_{n \rightarrow +\infty} z_n = +\infty, \quad \lim_{n \rightarrow +\infty} \frac{\rho_1(z_n)}{z_n} = 0,$$

and there exist  $\varepsilon_2 > 0$  and  $n_1 \in \mathbb{N}$  such that

$$\frac{\rho_1(z_n(1 + \varepsilon_2))}{\rho_1(z_n)}\Psi_- \leq \Psi_+ \quad \text{for } n \geq n_1,$$

where  $\psi(t) = h_1(t) - c_1$  for almost every  $t \in [0, \omega]$ ;

(d) the function  $\frac{\rho_1(x)}{x}$  is non-increasing and

$$\frac{\omega}{4}\Psi_+\Psi_-\frac{\rho_1(x_0)}{x_0} \leq \Psi_+ - \Psi_-,$$

where  $\psi(t) = h_1(t) - c_1$  for almost every  $t \in [0, \omega]$ .

Then there exists at least one solution to the problem (1.1), (1.3).

**Remark 3.3.1** Under the assumptions of Theorem 2.2.2, note that there exists a suitable  $\varepsilon_1$  such that (3.23) holds, e.g., if

$$\limsup_{x \rightarrow +\infty} \frac{\rho_0^\lambda((1 + \varepsilon_0)x)}{\rho_0^\lambda(x)} < 1 + \varepsilon_0.$$

Indeed, if (3.21) holds then

$$\int_0^{+\infty} \frac{ds}{\rho_0^\lambda(s)} = +\infty,$$

so that using the previous part (3.23) is proven by a classical arguments.

For the equation (3.2), from Theorem 3.3.1 we get the following assertion.

**Corollary 3.3.1** *Let  $g_1 > 0$ ,  $g_2 \geq 0$ ,  $0 \leq \delta < 1$ ,  $\nu > \gamma$  and*

$$\text{either } (\mu + \delta) \operatorname{sgn} |c| \geq 1 \quad \text{or} \quad \nu + 2\delta \geq 1.$$

*If*

$$\bar{h}_0 > - \lim_{x \rightarrow +\infty} \frac{g_2}{x^{\gamma+\delta}},$$

*then (3.2), (1.3) has at least one solution.*

**Remark 3.3.2** In the previous chapter, there is proven, among others, that the equation

$$u'' + \frac{1}{u^\nu} = h_0(t) \tag{3.26}$$

with  $h_0 \in L([0, \omega]; \mathbb{R})$  and  $\bar{h}_0 > 0$ , has a positive  $\omega$ -periodic solution if  $\nu \geq 1$ . Moreover, there is also introduced an example showing that for any  $\nu \in (0, 1)$ , there exists  $h_0 \in L([0, \omega]; \mathbb{R})$  with  $\bar{h}_0 > 0$  such that (3.26), (1.3) has no positive solution.

Corollary 3.3.1 says that if a friction-like term or sub-linear term are added to (3.26), the condition  $\nu \geq 1$  can be weakened. For example

$$u'' + \frac{u'}{u^\mu} + \frac{1}{u^\nu} = h_0(t)$$

has a positive solution satisfying (1.3) for any  $\nu > 0$  if  $\mu \geq 1$ , provided  $\bar{h}_0 > 0$ . Also the

equation

$$u'' + \frac{1}{u^\nu} = h_0(t)u^\delta$$

subjected to the boundary conditions (1.3) is solvable for any  $\nu > 0$  if  $\delta \in [1/2, 1)$ , provided  $\bar{h}_0 > 0$ .

**Corollary 3.3.2** *Let  $g_1 > 0$ ,  $g_2 \geq 0$ ,  $\nu > \gamma$ . Let, moreover, either  $g^* = -\infty$  or*

$$\bar{h}_0 > g^* > -\infty, \quad \frac{\omega}{4}\Phi_+\Phi_- < \Phi_+ - \Phi_-, \quad (3.27)$$

where  $\varphi(t) = h_0(t) - g^*$  for almost every  $t \in [0, \omega]$ , and

$$g^* = - \lim_{x \rightarrow +\infty} \frac{g_2}{x^{\gamma+1}}.$$

Then the problem (3.2), (1.3) with  $\delta = 1$  has at least one solution.

**Remark 3.3.3** According to [29] and Theorem 2.2.1, it can be easily verified that the problem

$$u'' + \frac{g_1}{u^\nu} = h_0(t)u; \quad u(0) = u(\omega), \quad u'(0) = u'(\omega) \quad (3.28)$$

with  $g_1 > 0$  and  $\nu > 0$ , has a positive solution if and only if the inclusion  $\mathcal{L}[0, -h_0] \in V^-$  holds (see notation in [29]).

Indeed, according to [29, Definition 1.1], the inclusion  $\mathcal{L}[0, -h_0] \in V^-$  implies the existence of a positive solution  $v$  to the problem

$$v'' = h_0(t)v - g_1; \quad u(0) = u(\omega), \quad u'(0) = u'(\omega).$$

Therefore there exist  $x > 0$  and  $y > 0$  such that  $x^{1+\nu} \leq v(t) \leq y^{1+\nu}$  for  $t \in [0, \omega]$ . By

setting

$$\alpha(t) \stackrel{\text{def}}{=} \frac{v(t)}{y^\nu}, \quad \beta(t) \stackrel{\text{def}}{=} \frac{v(t)}{x^\nu} \quad \text{for } t \in [0, \omega],$$

one can easily realized that  $\alpha$  and  $\beta$  are lower and upper functions to (3.28), respectively, satisfying (2.29). Now the existence of a positive solution to (3.28) follows from Theorem 2.2.1.

On the other hand, the existence of a positive solution to (3.28) implies the inclusion  $\mathcal{L}[0, -h_0] \in V^-$  (see [29, Theorem 2.1]).

However, one of the optimal effective conditions guaranteeing such an inclusion is  $h_0 \neq 0$  and

$$\frac{\omega}{4} \int_0^\omega [h_0(s)]_+ ds \int_0^\omega [h_0(s)]_- ds \leq \int_0^\omega [h_0(s)]_+ ds - \int_0^\omega [h_0(s)]_- ds.$$

(see [29, Corollary 2.5]). Therefore, the condition (3.27) is natural, in a certain sense.

When the right-hand side of the equation (1.3) does not depend on  $u$ , i.e., when  $h(t, x) \equiv h_0(t)$ , then (1.3) has the form (1.2). From Theorem 3.3.1, for the equation (1.2) we get the following assertion.

**Corollary 3.3.3** *Let there exist  $x_0 > 0$  and  $c_0 \in \mathbb{R}$  such that*

$$g(x) \leq c_0 < \bar{h}_0 \quad \text{for } x \geq x_0$$

*and let*

$$\lim_{x \rightarrow 0_+} g(x) = +\infty.$$

*Let, moreover, either*

$$\int_0^1 ([f(s)]_+ + [g(s)]_+) ds = +\infty, \quad \int_0^1 [f(s)]_- ds < +\infty$$



or

$$\int_0^1 ([f(s)]_- + [g(s)]_+) ds = +\infty, \quad \int_0^1 [f(s)]_+ ds < +\infty.$$

Then there exists at least one solution to the problem (1.2), (1.3).

In the following result, the assumptions do not depend on the friction-like term. On the other hand, a certain smallness of oscillation of the primitive to  $h_0$  is supposed. Clearly, Theorems 3.3.1 and 3.3.2 are independent.

**Theorem 3.3.2** *Let  $\rho_0 \in C^1(\mathbb{R}^+; \mathbb{R}^+)$  and  $\rho_1 \in C(\mathbb{R}^+; \mathbb{R}^+)$  be non-decreasing functions,  $h_0, h_1 \in L([0, \omega]; \mathbb{R})$ , and  $0 < x_0 \leq x_1 < +\infty$  be such that*

$$h(t, x) \leq h_0(t)\rho_0(x) \quad \text{for a. e. } t \in [0, \omega], \quad 0 < x \leq x_0, \quad (3.29)$$

$$h(t, x) \geq h_1(t)\rho_1(x) \quad \text{for a. e. } t \in [0, \omega], \quad x \geq x_1. \quad (3.30)$$

Let, moreover,

$$\frac{\omega}{8} \|h_0 - \bar{h}_0\|_1 < \int_0^{x_0} \frac{ds}{\rho_0(s)} < +\infty \quad (3.31)$$

$$\frac{g(x)}{\rho_0(x)} \geq \bar{h}_0 \quad \text{for } 0 < x \leq x_0, \quad (3.32)$$

and let there exist  $c_1 \in \mathbb{R}$  such that

$$\frac{g(x)}{\rho_1(x)} \leq c_1 \leq \bar{h}_1 \quad \text{for } x \geq x_1. \quad (3.33)$$

Besides, let us suppose that  $\rho_1$  fulfills at least one of the conditions (c) or (d) of Theorem 3.3.1. Then there exists at least one solution to the problem (1.1), (1.3).

In particular case when the equation (1.1) has the form (3.2), the following assertion immediately follows from Theorem 3.3.2.

**Corollary 3.3.4** *Let  $0 \leq \delta < 1$ , and let  $0 < x_0 \leq x_1 < +\infty$  be such that*

$$(1 - \delta) \frac{\omega}{8} \|h_0 - \bar{h}_0\|_1 < x_0^{1-\delta}.$$

*Let, moreover,*

$$\begin{aligned} \frac{g_1}{x^{\nu+\delta}} - \frac{g_2}{x^{\gamma+\delta}} &\geq \bar{h}_0 & \text{if } & 0 < x \leq x_0, \\ \frac{g_1}{x^{\nu+\delta}} - \frac{g_2}{x^{\gamma+\delta}} &\leq \bar{h}_0 & \text{if } & x \geq x_1. \end{aligned}$$

*Then the problem (3.2), (1.3) has at least one solution.*

**Remark 3.3.4** The consequence of Theorem 3.3.2 for the problem (1.2), (1.3) coincides with the obtained result Theorem 1.2.2.

### 3.3.2 Auxiliary propositions

In what follows we will show the existence of a solution to the equation

$$u''(t) + f(u(t))u'(t) + g(u(t)) = h_0(t)\rho_0(u(t)) \quad \text{for a. e. } t \in [0, \omega], \quad (3.34)$$

satisfying the boundary conditions (1.3). Here,  $\rho_0 \in C(\mathbb{R}^+; \mathbb{R}^+)$  is a non-decreasing function,  $h_0 \in L([0, \omega]; \mathbb{R})$ , and  $f, g \in C(\mathbb{R}^+; \mathbb{R})$ . Together with (3.34), for every  $k \in \mathbb{N}$ , consider the auxiliary equation

$$u''(t) + f(u(t))u'(t) + g(u(t)) = h_{0k}(t)\rho_0(u(t)) \quad \text{for a. e. } t \in [0, \omega], \quad (3.35)$$

where

$$h_{0k}(t) = \begin{cases} k & \text{if } h_0(t) > k \\ h_0(t) & \text{if } h_0(t) \leq k \end{cases} \quad \text{for a. e. } t \in [0, \omega], \quad k \in \mathbb{N}. \quad (3.36)$$

Obviously,

$$h_{0k}(t) \leq h_{0m}(t) \leq h_0(t) \quad \text{for a. e. } t \in [0, \omega], \quad k \leq m, \quad (3.37)$$

and

$$\lim_{k \rightarrow +\infty} \bar{h}_{0k} = \bar{h}_0. \quad (3.38)$$

The following three results respectively correspond to Corollaries 1.2.5, 1.2.6 and 1.2.1.

However a suitable formulation should be done into in our actual framework.

**Lemma 3.3.1** *Let  $x_0 > 0$  and  $c \in \mathbb{R}$  be such that*

$$\frac{g(x)}{\rho_0(x)} \leq c \leq \bar{h}_0 \quad \text{for } x \geq x_0. \quad (3.39)$$

*Let, moreover, there exist a sequence  $\{y_n\}_{n=1}^{+\infty}$  of positive numbers such that*

$$\lim_{n \rightarrow +\infty} y_n = +\infty, \quad \lim_{n \rightarrow +\infty} \frac{\rho_0(y_n)}{y_n} = 0, \quad (3.40)$$

*and let there exist  $\varepsilon > 0$  and  $n_0 \in \mathbb{N}$  such that*

$$\frac{\rho_0((1 + \varepsilon)y_n)}{\rho_0(y_n)} \Phi_- \leq \Phi_+ \quad \text{for } n \geq n_0,$$

*where  $\varphi(t) = h_0(t) - c$  for almost every  $t \in [0, \omega]$ . Then there exists an upper function  $\beta$  to the problem (3.34), (1.3) satisfying*

$$\beta(t) \geq x_0 \quad \text{for } t \in [0, \omega]. \quad (3.41)$$

**Lemma 3.3.2** *Let  $x_0 > 0$  and  $c \in \mathbb{R}$  be such that (3.39) holds. If  $\frac{\rho_0(x)}{x}$  is a non-increasing function such that*

$$\frac{\omega}{4}\Phi_+\Phi_-\frac{\rho_0(x_0)}{x_0} \leq \Phi_+ - \Phi_-$$

*where  $\varphi(t) = h_0(t) - c$  for almost every  $t \in [0, \omega]$ , then there exists an upper function  $\beta$  to the problem (3.34), (1.3) satisfying (3.41).*

**Lemma 3.3.3** *Let  $x_0 > \frac{\varepsilon}{8}\|h_0 - \bar{h}_0\|_1$  be such that*

$$g(x) \geq \bar{h}_0 \quad \text{for } 0 < x \leq x_0.$$

*Then there exists a lower function  $\alpha$  to the problem (1.36), (1.3) with*

$$0 < \alpha(t) \leq x_0 \quad \text{for } t \in [0, \omega]. \quad (3.42)$$

**Lemma 3.3.4** *Let  $x_0 > 0$  and  $c_0 \in \mathbb{R}$  be such that (3.15) holds. Let, moreover, there exist a sequence  $\{y_n\}_{n=1}^{+\infty}$  of positive numbers such that (3.40) is fulfilled, and let there exist  $\varepsilon > 0$  and  $n_0 \in \mathbb{N}$  such that*

$$\frac{\rho_0((1+\varepsilon)y_n)}{\rho_0(y_n)}\Phi_- \leq \Phi_+ - \varepsilon \quad \text{for } n \geq n_0, \quad (3.43)$$

*where  $\varphi(t) = h_0(t) - c_0$  for almost every  $t \in [0, \omega]$ . Then there exist  $k_0 \in \mathbb{N}$  and an upper function  $\beta$  to the problems (3.35), (1.3) for  $k \geq k_0$  satisfying (3.41).*

**Proof 44** Put

$$\varphi_k(t) = h_{0k}(t) - c_0 \quad \text{for a. e. } t \in [0, \omega], \quad (3.44)$$

$$\Phi_{k+} = \int_0^\omega [\varphi_k(s)]_+ ds, \quad \Phi_{k-} = \int_0^\omega [\varphi_k(s)]_- ds. \quad (3.45)$$

Then, obviously, in view of (3.36), we have

$$\lim_{k \rightarrow +\infty} \Phi_{k+} = \bar{\Phi}_+, \quad \lim_{k \rightarrow +\infty} \Phi_{k-} = \bar{\Phi}_- \quad (3.46)$$

and, consequently, on account of (3.15), (3.38), (3.43), and (3.46), there exists  $k_0 \in \mathbb{N}$  such that

$$\frac{g(x)}{\rho_0(x)} \leq c_0 \leq \bar{h}_{0k_0} \leq \bar{h}_0 \quad \text{for } x \geq x_0, \quad (3.47)$$

$$\frac{\rho_0((1+\varepsilon)y_n)}{\rho_0(y_n)} \Phi_{k_0-} \leq \Phi_{k_0+} \quad \text{for } n \geq n_0. \quad (3.48)$$

Therefore, according to Lemma 3.3.1, there exists an upper function  $\beta$  to (3.35), (1.3) with  $k = k_0$  satisfying (3.41). Obviously, in view of (3.37) and the non-negativity of  $\rho_0$  it follows that  $\beta$  is also an upper function to (3.35), (1.3) for  $k \geq k_0$ .

**Lemma 3.3.5** *Let  $x_0 > 0$  and  $c_0 \in \mathbb{R}$  be such that (3.15) holds. If  $\frac{\rho_0(x)}{x}$  is a non-increasing function such that*

$$\frac{\omega}{4} \Phi_+ \Phi_- - \frac{\rho_0(x_0)}{x_0} < \Phi_+ - \Phi_- \quad (3.49)$$

where  $\varphi(t) = h_0(t) - c_0$  for almost every  $t \in [0, \omega]$ , then there exist  $k_0 \in \mathbb{N}$  and an upper function  $\beta$  to the problems (3.35), (1.3) for  $k \geq k_0$  satisfying (3.41).

**Proof 45** Define  $\varphi_k$ ,  $\Phi_{k+}$ , and  $\Phi_{k-}$  by (3.44) and (3.45). Then, obviously, in view of (3.36), we have that (3.46) holds, and, consequently, on account of (3.15), (3.38), (3.46), and (3.49), there exists  $k_0 \in \mathbb{N}$  such that (3.47) is valid and

$$\frac{\omega}{4} \Phi_{k_0+} \Phi_{k_0-} - \frac{\rho_0(x_0)}{x_0} \leq \Phi_{k_0+} - \Phi_{k_0-} \quad (3.50)$$

Therefore, according to Lemma 3.3.2, there exists an upper function  $\beta$  to (3.35), (1.3)

with  $k = k_0$  satisfying (3.41). Obviously, in view of (3.37) and the non-negativity of  $\rho_0$  it follows that  $\beta$  is also an upper function to (3.35), (1.3) for  $k \geq k_0$ .

**Lemma 3.3.6** *Let*

$$\liminf_{x \rightarrow 0^+} g(x) > -\infty, \quad (3.51)$$

*and let either*

$$\int_0^1 [f(s)]_+ ds < +\infty \quad (3.52)$$

*or*

$$\int_0^1 [f(s)]_- ds < +\infty. \quad (3.53)$$

*Then, for every  $K > 0$ , there exists a constant  $K_1 > 0$  such that for any  $k \in \mathbb{N}$  and any positive solution  $u$  of (3.35), (1.3) with*

$$\|u\|_\infty \leq K \quad (3.54)$$

*we have the estimate*

$$\|u'\|_\infty \leq K_1. \quad (3.55)$$

**Proof 46** Assume that (3.53) is fulfilled. Let  $u$  be a positive solution to (3.35), (1.3) satisfying (3.54). Then there exist  $t_0, t_1 \in [0, \omega]$  such that

$$u(t_0) = \min \{u(t) : t \in [0, \omega]\}, \quad u(t_1) = \max \{u(t) : t \in [0, \omega]\}. \quad (3.56)$$

Define the operator  $\vartheta$  of  $\omega$ -periodic prolongation by

$$\vartheta(v)(t) = \begin{cases} v(t) & \text{if } t \in [0, \omega] \\ v(t - \omega) & \text{if } t \in (\omega, 2\omega] \end{cases}. \quad (3.57)$$

Then, obviously, from (3.35) and (1.3) it follows that

$$\begin{aligned} \vartheta(u)''(t) + f(\vartheta(u)(t))\vartheta(u)'(t) + g(\vartheta(u)(t)) &= \vartheta(h_{0k})(t)\rho_0(\vartheta(u)(t)) \\ &\text{for a. e. } t \in [0, 2\omega]. \end{aligned} \quad (3.58)$$

The integration of (3.58) from  $t_0$  to  $t$ , on account of (3.56), yields

$$\begin{aligned} \vartheta(u)'(t) &= - \int_{t_0}^t f(\vartheta(u)(s))\vartheta(u)'(s)ds - \int_{t_0}^t g(\vartheta(u)(s))ds \\ &\quad + \int_{t_0}^t \vartheta(h_{0k})(s)\rho_0(\vartheta(u)(s))ds \quad \text{for } t \in [t_0, t_0 + \omega]. \end{aligned} \quad (3.59)$$

From (3.54), (3.56), and (3.57) it follows that

$$0 < \vartheta(u)(t_0) \leq \vartheta(u)(t) \leq K \quad \text{for } t \in [t_0, t_0 + \omega]. \quad (3.60)$$

Put

$$\mu = \sup \{ [g(s)]_- : s \in (0, K] \}. \quad (3.61)$$

According to (3.51) we have

$$0 \leq \mu < +\infty. \quad (3.62)$$

Thus, using (3.36), (3.53), (3.54), and (3.60)–(3.62) in (3.59) we arrive at

$$\vartheta(u)'(t) \leq \int_0^K [f(s)]_- ds + \omega\mu + \|h_0\|_1\rho_0(K) \quad \text{for } t \in [t_0, t_0 + \omega]. \quad (3.63)$$

Put

$$K_1 = \int_0^K [f(s)]_- ds + \omega\mu + \|h_0\|_1\rho_0(K).$$

Then, on account of (3.57) and (3.63) we have

$$u'(t) \leq K_1 \quad \text{for } t \in [0, \omega]. \quad (3.64)$$

On the other hand, the integration of (3.58) from  $t$  to  $t_1 + \omega$ , with respect to (3.56), results in

$$\begin{aligned} \vartheta(u)'(t) = & \int_t^{t_1+\omega} f(\vartheta(u)(s))\vartheta(u)'(s)ds + \int_t^{t_1+\omega} g(\vartheta(u)(s))ds \\ & - \int_t^{t_1+\omega} \vartheta(h_{0k})(s)\rho_0(\vartheta(u)(s))ds \quad \text{for } t \in [t_1, t_1 + \omega]. \end{aligned} \quad (3.65)$$

Now, using (3.36), (3.53), (3.54), and (3.60)–(3.62) in (3.65) we obtain

$$-\vartheta(u)'(t) \leq K_1 \quad \text{for } t \in [t_1, t_1 + \omega]. \quad (3.66)$$

Therefore, in view of (3.57), from (3.66) we get

$$-u'(t) \leq K_1 \quad \text{for } t \in [0, \omega] \quad (3.67)$$

Consequently, (3.64) and (3.67) results in (3.55).

Now suppose that (3.52) is fulfilled. Put

$$v(t) = u(\omega - t) \quad \text{for } t \in [0, \omega]. \quad (3.68)$$

Then, according to (3.35) we have

$$v''(t) - f(v(t))v'(t) + g(v(t)) = \tilde{h}_{0k}(t)\rho_0(v(t)) \quad \text{for a. e. } t \in [0, \omega], \quad (3.69)$$



where

$$\tilde{h}_{0k}(t) = h_{0k}(\omega - t) \quad \text{for a. e. } t \in [0, \omega].$$

Analogously to the above-proven, using (3.52) instead of (3.53), we obtain

$$\|v'\|_\infty \leq K_1 \tag{3.70}$$

with

$$K_1 = \int_0^K [f(s)]_+ ds + \omega\mu + \|h_0\|_1 \rho_0(K).$$

Thus, (3.68) and (3.70) yields (3.55).

**Lemma 3.3.7** *Let*

$$\lim_{x \rightarrow 0_+} g(x) = +\infty \tag{3.71}$$

*and let either*

$$\int_0^1 ([f(s)]_+ + [g(s)]_+) ds = +\infty, \quad \int_0^1 [f(s)]_- ds < +\infty \tag{3.72}$$

*or*

$$\int_0^1 ([f(s)]_- + [g(s)]_+) ds = +\infty, \quad \int_0^1 [f(s)]_+ ds < +\infty \tag{3.73}$$

*Then, for every  $K > 0$  there exists a constant  $a > 0$  such that for any  $k \in \mathbb{N}$  and any positive solution  $u$  of (3.35), (1.3) satisfying (3.54) we have the estimate*

$$a \leq u(t) \quad \text{for } t \in [0, \omega]. \tag{3.74}$$

**Proof 47** Let  $u$  be a positive solution to (3.35), (1.3) satisfying (3.54). Thus, the integration of (3.35) from 0 to  $\omega$ , in view of (1.3) and (3.37), yields

$$\int_0^\omega g(u(s)) ds \leq \|h_0\|_1 \rho_0(K). \tag{3.75}$$

On the other hand, (3.71) implies the existence of  $x_0 \in \mathbb{R}^+$  such that

$$g(x) > \frac{\|h_0\|_1 \rho_0(K)}{\omega} \geq 0 \quad \text{for } x \in (0, x_0). \quad (3.76)$$

Let  $t_m \in [0, \omega]$  be such that

$$u(t_m) = \min \{u(t) : t \in [0, \omega]\}. \quad (3.77)$$

Obviously, either

$$u(t_m) \geq x_0$$

or

$$u(t_m) < x_0. \quad (3.78)$$

Obviously, it is sufficient to show the estimate (3.74) is valid just in the case when (3.78) is fulfilled. Let, therefore, (3.78) hold.

If  $u(t) < x_0$  for  $t \in [0, \omega]$ , then applying (3.76) in (3.75) we obtain a contradiction. Thus, there exist points  $t_1, t_2 \in (t_m, t_m + \omega)$  such that

$$\vartheta(u)(t) < x_0 \quad \text{for } t \in [t_m, t_1), \quad \vartheta(u)(t_1) = x_0, \quad (3.79)$$

$$\vartheta(u)(t) < x_0 \quad \text{for } t \in (t_2, t_m + \omega], \quad \vartheta(u)(t_2) = x_0, \quad (3.80)$$

where  $\vartheta$  is the operator defined by (3.57). Obviously, (3.58) holds.

Assume that (3.72) holds. Then, according to Lemma 3.3.6, there exists  $K_1 > 0$  such that (3.55) holds. The integration of (3.58) from  $t_m$  to  $t_1$ , in view of (3.37), (3.54),

(3.55), (3.72), (3.76), (3.77), and (3.79) results in

$$\begin{aligned} \vartheta(u)'(t_1) + \int_{\vartheta(u)(t_m)}^{x_0} [f(s)]_+ ds + \frac{1}{K_1} \int_{\vartheta(u)(t_m)}^{x_0} [g(s)]_+ ds \\ \leq \int_0^{x_0} [f(s)]_- ds + \|h_0\|_1 \rho_0(K). \end{aligned} \quad (3.81)$$

Note that in view of (3.79) we have  $\vartheta(u)'(t_1) \geq 0$ . Consequently, from (3.81) we obtain

$$\int_{\vartheta(u)(t_m)}^{x_0} ([f(s)]_+ + [g(s)]_+) ds \leq K_2 \quad (3.82)$$

where

$$K_2 = (K_1 + 1) \left( \int_0^{x_0} [f(s)]_- ds + \|h_0\|_1 \rho_0(K) \right).$$

Note that  $K_2$  does not depend on  $k$ . Therefore, if we apply (3.72) in (3.82), it can be easily seen, with respect to (3.77), that there exists a constant  $a > 0$  such that (3.74) holds.

If (3.73) holds, we integrate (3.58) from  $t_2$  to  $t_m + \omega$  and apply the similar steps as above, just using (3.80) instead of (3.79), finally we arrive at

$$\int_{\vartheta(u)(t_m + \omega)}^{x_0} ([f(s)]_- + [g(s)]_+) ds \leq K_2$$

with

$$K_2 = (K_1 + 1) \left( \int_0^{x_0} [f(s)]_+ ds + \|h_0\|_1 \rho_0(K) \right).$$

Therefore, also in this case there exists a constant  $a > 0$  such that (3.74) holds.

**Lemma 3.3.8** *Let  $x_0 > 0$  and  $c_0 \in \mathbb{R}$  be such that (3.15) holds. Let, moreover, (3.71) be fulfilled, and let either (3.72) or (3.73) be valid. Let, in addition, there exist a sequence  $\{y_n\}_{n=1}^{+\infty}$  of positive numbers such that (3.40) holds, and let there exist  $\varepsilon > 0$  and  $n_0 \in \mathbb{N}$*

such that (3.43) is fulfilled, where  $\varphi(t) = h_0(t) - c_0$  for almost every  $t \in [0, \omega]$ . Then there exists a positive solution  $u$  to (3.34), (1.3).

**Proof 48** According to Lemma 3.3.4, there exists  $k_0 \in \mathbb{N}$  and an upper function  $\beta$  to the problems (3.35), (1.3) for  $k \geq k_0$  satisfying (3.41). On the other hand, in view of (3.37) and (3.71) there exist  $x_k \in (0, x_0]$  for  $k \geq k_0$  such that

$$g(x_k) \geq h_{0k}(t)\rho_0(x_k) \quad \text{for a. e. } t \in [0, \omega].$$

Thus, if we put  $\alpha_k(t) = x_k$  for  $t \in [0, \omega]$ , according to Theorem 2.2.1, there exists a solution  $u_k$  to (3.35), (1.3) for  $k \geq k_0$  satisfying

$$0 < \alpha_k(t) \leq u_k(t) \leq \beta(t) \quad \text{for } t \in [0, \omega]. \quad (3.83)$$

Moreover, according to Lemmas 3.3.6 and 3.3.7, in view of (3.83), there exist constants  $K > 0$ ,  $K_1 > 0$ , and  $a > 0$ , not depending on  $k$ , such that

$$\|u_k\|_\infty \leq K, \quad \|u'_k\|_\infty \leq K_1, \quad \text{for } k \geq k_0, \quad (3.84)$$

$$a \leq u_k(t) \quad \text{for } t \in [0, \omega], \quad k \geq k_0, \quad (3.85)$$

$$|u''_k(t)| \leq f_0 K_1 + g_0 + |h_0(t)|\rho_0(K) \quad \text{for a. e. } t \in [0, \omega], \quad k \geq k_0, \quad (3.86)$$

where

$$f_0 = \max \{|f(x)| : x \in [a, K]\}, \quad g_0 = \max \{|g(x)| : x \in [a, K]\}.$$

Therefore, according to Arzelà–Ascoli Theorem, there exists  $u_0 \in C([0, \omega]; \mathbb{R})$  and  $v_0 \in C([0, \omega]; \mathbb{R})$  such that

$$\lim_{k \rightarrow +\infty} \|u_k - u_0\|_\infty = 0, \quad \lim_{k \rightarrow +\infty} \|u'_k - v_0\|_\infty = 0. \quad (3.87)$$

Moreover, since  $u_k$  are solutions to (3.35), (1.3), in view of (3.36), (3.85), and (3.87), we have  $u_0 \in AC^1([0, \omega]; \mathbb{R})$ ,  $u'_0 \equiv v_0$ , and  $u_0$  is a positive solution to (3.34), (1.3).

The following assertion can be proven analogously to Lemma 3.3.8, just Lemma 3.3.5 is used instead of Lemma 3.3.4.

**Lemma 3.3.9** *Let  $x_0 > 0$  and  $c_0 \in \mathbb{R}$  be such that (3.15) holds. Let, moreover, (3.71) be fulfilled, and let either (3.72) or (3.73) be valid. Let, in addition,  $\frac{\rho_0(x)}{x}$  be a non-increasing function and let (3.49) be fulfilled, where  $\varphi(t) = h_0(t) - c_0$  for almost every  $t \in [0, \omega]$ . Then there exists a positive solution  $u$  to (3.34), (1.3).*

**Lemma 3.3.10** *Let  $\rho_0 \in C^1(\mathbb{R}^+; \mathbb{R}^+)$  be non-decreasing,  $x_0 > 0$ , and  $c_0 \in \mathbb{R}$  be such that (3.15) holds. Let, moreover, there exist  $\lambda \in [0, 1]$  such that (3.17) and (3.18) are valid, and let either (3.19) or (3.20) be fulfilled. Let, in addition, there exist a sequence  $\{y_n\}_{n=1}^{+\infty}$  of positive numbers such that (3.21) holds and let there exist  $\varepsilon_0 > 0$ ,  $\varepsilon_1 \in (0, \varepsilon_0]$ , and  $n_0 \in \mathbb{N}$  such that (3.22) and (3.23) are fulfilled, where  $\varphi(t) = h_0(t) - c_0$  for almost every  $t \in [0, \omega]$  and  $\sigma$  is given by (3.24). Then there exists a lower function  $\alpha$  to the problem (3.34), (1.3).*

**Proof 49** Because  $\rho_0$  is a positive function, from (3.17) and (3.24) we obtain that  $\sigma$  is a positive increasing function. Therefore, there exists an inverse function  $\sigma^{-1}$  to  $\sigma$  which is also increasing. Moreover, in view of (3.17) and (3.24), it follows that

$$\lim_{x \rightarrow 0_+} \sigma(x) = 0, \quad \lim_{x \rightarrow 0_+} \sigma^{-1}(x) = 0, \quad \lim_{x \rightarrow +\infty} \sigma(x) = +\infty, \quad \lim_{x \rightarrow +\infty} \sigma^{-1}(x) = +\infty. \quad (3.88)$$

Consider the auxiliary equation

$$u''(t) + f(\sigma^{-1}(u(t)))u'(t) + \frac{g(\sigma^{-1}(u(t)))}{\rho_0^\lambda(\sigma^{-1}(u(t)))} = h_0(t)\rho_0^{1-\lambda}(\sigma^{-1}(u(t)))$$

for a. e.  $t \in [0, \omega]$ . (3.89)

Put  $z = \sigma(x)$ ,  $z_0 = \sigma(x_0)$ . Then from (3.15) we get

$$\frac{g(\sigma^{-1}(z))}{\rho_0^\lambda(\sigma^{-1}(z))} \leq c_0 < \bar{h}_0 \quad \text{for } z \geq z_0$$

(3.90)

and, in view of (3.88), from (3.18) we have

$$\lim_{z \rightarrow 0^+} \frac{g(\sigma^{-1}(z))}{\rho_0^\lambda(\sigma^{-1}(z))} = +\infty.$$

(3.91)

Furthermore, the substitution  $r = \sigma(s)$  in (3.19), resp (3.20), with respect to (3.24), yields

$$\int_0^1 \left( [f(\sigma^{-1}(r))]_+ + \frac{[g(\sigma^{-1}(r))]_+}{\rho_0^\lambda(\sigma^{-1}(r))} \right) dr = +\infty, \quad \int_0^1 [f(\sigma^{-1}(r))]_- dr < +\infty,$$

(3.92)

resp.

$$\int_0^1 \left( [f(\sigma^{-1}(r))]_- + \frac{[g(\sigma^{-1}(r))]_+}{\rho_0^\lambda(\sigma^{-1}(r))} \right) dr = +\infty, \quad \int_0^1 [f(\sigma^{-1}(r))]_+ dr < +\infty$$

(3.93)

Moreover, put  $z_n = \sigma(y_n)$  for  $n \in \mathbb{N}$ . Then from (3.21), in view of (3.88), we get

$$\lim_{n \rightarrow +\infty} z_n = +\infty, \quad \lim_{n \rightarrow +\infty} \frac{\rho_0^{1-\lambda}(\sigma^{-1}(z_n))}{z_n} = 0.$$

(3.94)

Finally, (3.23) results in

$$\sigma^{-1}((1 + \varepsilon_1)z_n) \leq (1 + \varepsilon_0)y_n \quad \text{for } n \geq n_0,$$

and so, since  $\rho_0$  is a non-decreasing function, from (3.22) we obtain

$$\frac{\rho_0^{1-\lambda}(\sigma^{-1}((1 + \varepsilon_1)z_n))}{\rho_0^{1-\lambda}(\sigma^{-1}(z_n))} \Phi_- \leq \Phi_+ - \varepsilon_1 \quad \text{for } n \geq n_0. \quad (3.95)$$

Therefore, applying Lemma 3.3.8, according to (3.90)–(3.95), there exists a positive solution  $u$  to the problem (3.89), (1.3).

Now we put  $\alpha(t) = \sigma^{-1}(u(t))$  for  $t \in [0, \omega]$ , i.e., in view of (3.24),

$$u(t) = \int_0^{\alpha(t)} \frac{ds}{\rho_0^\lambda(s)} \quad \text{for } t \in [0, \omega].$$

Obviously,  $\alpha \in AC^1([0, \omega]; \mathbb{R})$  is a positive function and

$$u'(t) = \frac{\alpha'(t)}{\rho_0^\lambda(\alpha(t))} \quad \text{for } t \in [0, \omega],$$

$$u''(t) = \frac{\alpha''(t)}{\rho_0^\lambda(\alpha(t))} - \frac{\lambda \alpha'^2(t) \rho_0'(\alpha(t))}{\rho_0^{1+\lambda}(\alpha(t))} \leq \frac{\alpha''(t)}{\rho_0^\lambda(\alpha(t))} \quad \text{for a. e. } t \in [0, \omega].$$

Thus, it can be easily seen that  $\alpha$  is a lower function to the problem (3.34), (1.3).

Analogously to the proof of Lemma 3.3.10, one can prove the following assertion applying Lemma 3.3.9 instead of Lemma 3.3.8.

**Lemma 3.3.11** *Let  $\rho_0 \in C^1(\mathbb{R}^+; \mathbb{R}^+)$  be non-decreasing,  $x_0 > 0$ , and  $c_0 \in \mathbb{R}$  be such that (3.15) holds. Let, moreover, there exist  $\lambda \in [0, 1]$  such that (3.17) and (3.18) are valid, and let either (3.19) or (3.20) be fulfilled. Let, in addition,  $\frac{\rho_0^{1-\lambda}(x)}{\sigma(x)}$  be a non-increasing function and let (3.25) be fulfilled, where  $\varphi(t) = h_0(t) - c_0$  for almost every*

$t \in [0, \omega]$  and  $\sigma$  is given by (3.24). Then there exists a lower function  $\alpha$  to the problem (3.34), (1.3).

### 3.3.3 Proof of the main results

In this section we will write the proofs of our main results using the previous proved lemmas.

**Proof 50 (Proof of Theorem 3.3.1)** According to Lemmas 3.3.1, 3.3.2, 3.3.10, and 3.3.11, the conditions of theorem guarantee a well-ordered couple of lower and upper functions, therefore the result is a direct consequence of Theorem 2.2.1.

**Proof 51 (Proof of Corollary 3.3.1)** It follows from Theorem 3.3.1 with  $h_1 \equiv h_0$ ,  $\rho_0(x) = \rho_1(x) = x^\delta$ ,  $\lambda = 1$ , and  $c_0 = c_1$  such that

$$\bar{h}_0 > c_0 > - \lim_{x \rightarrow +\infty} \frac{g_2}{x^{\gamma+\delta}}.$$

Then items *a*) and *c*) of Theorem 3.3.1 are fulfilled.

**Proof 52 (Proof of Corollary 3.3.2)** It follows from Theorem 3.3.1 with  $h_1 \equiv h_0$ ,  $\rho_0(x) = \rho_1(x) = x$ , and  $\lambda < 1$  such that  $\nu + 2\lambda \geq 1$ . Then items *b*) and *d*) of Theorem 3.3.1 are fulfilled.

**Proof 53 (Proof of Corollary 3.3.3)** It immediately follows from Theorem 3.3.1 with  $h_1 \equiv h_0$ ,  $\rho_i(x) \equiv 1$  ( $i = 0, 1$ ).

**Proof 54 (Proof of Theorem 3.3.2)** Put

$$\sigma(x) = \int_0^x \frac{ds}{\rho_0(s)} \quad \text{for } x \geq 0. \quad (3.96)$$



Because  $\rho_0$  is a positive function, from (3.31) and (3.96) we obtain that  $\sigma$  is an increasing function. Therefore, there exists an inverse function  $\sigma^{-1}$  to  $\sigma$  which is also increasing.

Consider the auxiliary equation

$$u''(t) + f(\sigma^{-1}(u(t)))u'(t) + \frac{g(\sigma^{-1}(u(t)))}{\rho_0(\sigma^{-1}(u(t)))} = h_0(t) \quad \text{for a. e. } t \in [0, \omega]. \quad (3.97)$$

Put  $z = \sigma(x)$ ,  $z_0 = \sigma(x_0)$ . Then from (3.31) and (3.32), in view of (3.96), we get

$$\begin{aligned} \frac{\omega}{8} \|h_0 - \bar{h}_0\|_1 &< z_0 \\ \frac{g(\sigma^{-1}(z))}{\rho_0(\sigma^{-1}(z))} &\geq \bar{h}_0 \quad \text{for } 0 < z \leq z_0, \end{aligned}$$

Therefore, according to Lemma 3.3.3 there exists a lower function  $w$  to the problem (3.97), (1.3) satisfying

$$0 < w(t) \leq \sigma(x_0) \quad \text{for } t \in [0, \omega]. \quad (3.98)$$

Now we put  $\alpha = \sigma^{-1}(w(t))$  for  $t \in [0, \omega]$ , i.e., in view of (3.96),

$$w(t) = \int_0^{\alpha(t)} \frac{ds}{\rho_0(s)} \quad \text{for } t \in [0, \omega].$$

Obviously, with respect to (3.98),  $\alpha \in AC^1([0, \omega]; \mathbb{R})$  is a positive function satisfying (3.42), and

$$\begin{aligned} w'(t) &= \frac{\alpha'(t)}{\rho_0(\alpha(t))} \quad \text{for } t \in [0, \omega], \\ w''(t) &= \frac{\alpha''(t)}{\rho_0(\alpha(t))} - \frac{\alpha'^2(t)\rho_0'(\alpha(t))}{\rho_0^2(\alpha(t))} \leq \frac{\alpha''(t)}{\rho_0(\alpha(t))} \quad \text{for a. e. } t \in [0, \omega]. \end{aligned}$$

Thus, on account of (3.29), (3.42), and (3.97), it can be easily seen that  $\alpha$  is a lower

function to the problem (1.1), (1.3).

The existence of an upper function  $\beta$  to (1.1), (1.3) satisfying

$$\beta(t) \geq x_1 \quad \text{for } t \in [0, \omega] \quad (3.99)$$

follows from (3.30) and Lemma 3.3.1, resp. 3.3.2.

Obviously, in view of (3.42) and (3.99), we have that (2.29) holds. Thus the theorem follows from Theorem 2.2.1.

**Proof 55 (Proof of Corollary 3.3.4)** It follows from Theorem 3.3.2 with  $h_1 \equiv h_0$ ,  $\rho_0(x) = \rho_1(x) = x^\delta$ , and  $c_1 = \bar{h}_0$ .

---

## CHAPTER 4

# The Brillouin beam equation

The last type of equation which will be studied in this work is the classical Brillouin beam equation

$$u'' + b(1 + \cos t)u = \frac{1}{u}, \quad (4.1)$$

depending on a positive parameter  $b$ . This equation is a repulsive singular perturbation of a Mathieu equation, as we will explain below. The classical problem to find  $2\pi$ -periodic solutions of (4.1) arose at the beginning in the sixties in the context of Electronics motivated by some numerical experiments realized in [5], where it was conjectured that *equation (4.1) should have a  $2\pi$ -periodic solution whenever  $b \in (0, 1/4)$ .*

By an acquired intuition in the previous chapters, one expects that under some non-resonant condition should be possible to prove the proposed conjecture. However we will explain below that this problem can be really delicate, and arising doubts on the validity of the result conjectured.

Essentially the unique research line to try to prove this result concerns to use the function  $K : [1, +\infty] \rightarrow \mathbb{R}$  defined by

$$K(\alpha) = \begin{cases} \frac{1}{\pi^2} & \alpha = 1 \\ \frac{(\alpha - 1)^{1 + \frac{1}{\alpha}}}{8\pi^{1 - \frac{1}{2\alpha}} \alpha^{1 - \frac{1}{\alpha}} (2\alpha - 1)^{\frac{1}{\alpha}}} \left( \frac{\Gamma(\frac{1}{2} - \frac{1}{2\alpha})}{\Gamma(1 - \frac{1}{2\alpha})} \right)^2 \left( \frac{\Gamma(\alpha)}{\Gamma(\frac{1}{2} + \alpha)} \right)^{\frac{1}{\alpha}} & \alpha \in (1, +\infty) \\ \frac{1}{8} & \alpha = +\infty; \end{cases}$$

with this definition, a sufficient condition in order for the Dirichlet problem ( $u(0) = 0 = u(2\pi)$ ) associated with

$$u'' + b(1 + \cos t)u = 0. \quad (4.2)$$

to have a unique solution is that

$$b < \max_{\alpha \in [0, +\infty]} K(\alpha) \approx 0.16448. \quad (4.3)$$

Thus under this non-resonance condition i.e., if  $b \in (0, 0.16448)$ , M. Zhang proved that (4.1) has at least one  $2\pi$ -periodic solution in [66]. This last result has been extended to equations where the singularity may be of weak type (see [53]).

In view of [53], one can understand that condition (4.3) implies that (4.2) has an associated Green function which has a positive sign. This makes possible to control the operator

$$T : \Omega \rightarrow C([0, 2\pi]; \mathbb{R}), \quad T[u](t) = \int_0^{2\pi} \frac{G(t, s)}{u(s)} ds,$$

where  $\Omega$  is a suitable open subset of  $C([0, 2\pi]; \mathbb{R})$  in order to apply Krasnoselskii fixed point theorem, and  $G : [0, 2\pi] \times [0, 2\pi] \rightarrow \mathbb{R}$  is the associated Green function to (4.2).

Despite of the expectations, nobody has improved the result of M. Zhang, i.e., the best range of  $b$  actually known for the  $2\pi$ -periodic solvability of (4.2) is the proven in [66], and outside from there the thing is completely unknown. Nevertheless, we can point out that the result cannot be obtained as a consequence of a general equation with strong condition singularity such as it was done in [50] (this is one of the main troubles in the related literature). Indeed, according to [67, Theorem 2.1], it was established an unanimous relation between the stability intervals for the Mathieu equation (4.3) and

the existence of periodic solutions for the Yermakov-Pinney equation

$$u'' + b(1 + \cos t)u - \frac{1}{u^3} = 0. \quad (4.4)$$

Notice that the stability intervals of the Mathieu equation

$$(\lambda_0, \lambda'_1), \quad (\lambda'_2, \lambda_1), \quad (\lambda_2, \lambda'_3), \quad \dots,$$

where  $\lambda_i$ ,  $i = 0, 1, \dots$  and  $\lambda'_i$ ,  $i = 1, 2, \dots$ , respectively, are the values of the parameter  $b$  for which equation (4.2) has, respectively, a genuine  $\pi$ -periodic solution and a genuine  $2\pi$ -periodic solution, are defined approximately by  $\lambda = 0$ ,  $\lambda'_1 \approx 1/6$ ;  $\lambda'_2 \approx 0.4$ ,  $\lambda_1 \approx 0.95, \dots$  (see [41, Theorem 2.1] and [10, Figure 1]). Thus the conjectured result in [5] cannot be extended to (4.4).

We will divide in two sections the present chapter. In the first one we use our general results proved in Chapter 1 in order to see what happen with (4.1). In the second one we show that (4.1) may have  $2\pi$ -periodic solutions also when  $b$  belongs to intervals other than  $(0, 1/4)$ . However the proposed conjecture still remained as an open problem (see [24] to know the details of the different contributions on this topic).

## 4.1 A corollary for the Brillouin beam equation

One can easily observe that (4.1) can be considered as a particular equation of (1.1), taking either  $g(x) = bx - 1/x$  and  $h(t, x) = -bx \cos t$  or  $g(x) = -1/x$  and  $h(t, x) = -b(1 + \cos t)x$ . Thus applying Theorem 1.1.3 follows the following corollaries.

**Corollary 4.1.1** *If  $b \in (0, 1/(4 + \pi))$ , then (4.1) has at least one  $2\pi$ -periodic solution.*

**Proof 56** We take  $g(x) = bx - 1/x$  and  $h(t, x) = h_0(t)\rho(x)$  where

$$h_0(t) = -b \cos t \quad \text{for } t \in [0, 2\pi], \quad \rho(x) = x.$$

Defining

$$\eta : [0, 2\pi] \times \mathbb{R}^+ \rightarrow \mathbb{R}, \quad \eta(t, x) = \begin{cases} 0 & t \in [\frac{\pi}{2}, \frac{3\pi}{2}], \\ bx \cos t & t \notin [\frac{\pi}{2}, \frac{3\pi}{2}]; \end{cases}$$

it deduces that  $h(t, x) \geq -\eta(t, x)$  on  $[0, 2\pi] \times \mathbb{R}^+$  and, therefore that

$$\frac{1}{x} \int_0^{2\pi} \eta(t, x) dt = 2b.$$

Since  $\eta$  and  $\rho$  are non-decreasing functions, the proof follows easily from Theorem 1.1.3.

On the contrary, if we take  $g(x) = -1/x$  and  $h(t, x) = -b(1 + \cos t)x$ , we can only obtain in a natural way the assertion.

**Corollary 4.1.2** *If  $b \in (0, 1/\pi^2)$ , then (4.1) has at least one  $2\pi$ -periodic solution.*

As always the parameter  $b$  should be into the interval  $(0, 1/4)$ . This is because to prove Theorem 1.1.3 we have implicitly used that  $u'' + b(1 + \cos t)u = 0$  has a positive associated Green function.

## 4.2 A new result for the Brillouin beam equation

In this section we will obtain the first range of parameter  $b$  outside of the interval  $(0, 1/4)$  thanks to suitable non-resonance assumptions which can be traced back to the work [13] by Fabry. The main abstract tool to obtain such a statement is embodied by the *Poincaré-Bohl fixed point theorem*.

### 4.2.1 A non-resonance theorem for repulsive singular equations

The proof of our Theorem is based on a non-resonance result which involves nonlinearities with “atypical” linear growing type, and could have interest by itself.

**Theorem 4.2.1** *Let us assume that there exist positive constants  $A_+$ ,  $B_+$  such that*

$$\frac{1}{2\pi} \int_0^{2\pi} \min \left\{ \frac{b(1 + \cos t)}{B_+}, 1 \right\} dt > \frac{n}{2\sqrt{B_+}}, \quad (4.5)$$

$$\frac{1}{2\pi} \int_0^{2\pi} \max \left\{ \frac{b(1 + \cos t)}{A_+}, 1 \right\} dt < \frac{n+1}{2\sqrt{A_+}}, \quad (4.6)$$

for some natural number  $n$ . Then (4.1) has at least one  $2\pi$ -periodic solution.

Before introducing the main tools to prove the theorem, a couple of remarks are in order.

**Remark 4.2.1** With the aim of keeping the exposition at a rather simple level, and taking into account that our main goal will be to study the existence of  $2\pi$ -periodic solutions of (4.1), we will always consider equation (4.1) as a starting point. However, the result can be extended, with the same approach and similar computations, to more general equations like

$$u'' + q(t)u - g(t, u) = 0, \quad (4.7)$$

where  $q$  is continuous and  $2\pi$ -periodic, and  $g : [0, 2\pi] \times (0, +\infty) \rightarrow \mathbb{R}$  has a similar behavior as  $1/x^\gamma$ , with  $\gamma \geq 1$ , being allowed to grow at most sublinearly at infinity. For instance, as in [20], one can assume that there exist  $\sigma > 0$  and a continuous function  $f : (0, \sigma] \rightarrow \mathbb{R}$  such that  $g(t, x) \leq f(x)$ , whenever  $x \in (0, \sigma]$ , and

$$\lim_{r \rightarrow 0^+} f(r) = -\infty, \quad \int_0^\sigma f(r) dr = -\infty.$$

Of course, in this case  $q(t)$  will replace  $b(1 + \cos t)$ .

**Remark 4.2.2** Conditions (4.5) and (4.6) were introduced by Fabry in [13] for the equation

$$u'' + g(t, u) = 0,$$

with

$$p(t) \leq \liminf_{|x| \rightarrow +\infty} \frac{g(t, x)}{x} \leq \limsup_{|x| \rightarrow +\infty} \frac{g(t, x)}{x} \leq q(t),$$

asking that

$$\sqrt{\lambda_j} < \sup_{\xi > 0} \frac{\frac{1}{2\pi} \int_0^{2\pi} \min\{p(t), \xi\} dt}{\sqrt{\xi}}, \quad \inf_{\xi > 0} \frac{\frac{1}{2\pi} \int_0^{2\pi} \max\{q(t), \xi\} dt}{\sqrt{\xi}} < \sqrt{\lambda_{j+1}},$$

where  $\lambda_j$  is the  $j$ -th eigenvalue of the considered  $2\pi$ -periodic problem. Such conditions are usually coupled with the sign assumption

$$\liminf_{|x| \rightarrow +\infty} \operatorname{sgn} x f(t, x) > 0$$

(see for instance [14]), which, however, in the model case  $g(t, x) = b(1 + \cos t)x + f(t, x)$ , with  $\lim_{|x| \rightarrow +\infty} f(t, x) = 0$ , is not satisfied. This is one of the main difficulties of the problem considered in this chapter.

As it is easy to see, (4.5) and (4.6) are the counterpart of such conditions for the Dirichlet spectrum (which is the natural one to consider when dealing with problems with a singularity, see [66]).

**Remark 4.2.3** As a consequence of Theorem 4.2.1, we can obtain the main results in [46, 20]. Indeed, assume that there exist positive constants  $A_+$ ,  $B_+$  such that

$$B_+ \leq q(t) \leq A_+ \quad \text{for every } t \in [0, 2\pi]. \quad (4.8)$$



Then, according to [46, 20], there exists at least one  $2\pi$ -periodic solution of (4.7) under the nonresonance assumption

$$\left(\frac{n}{2}\right)^2 < B_+ \leq A_+ < \left(\frac{n+1}{2}\right)^2,$$

where  $n \in \mathbb{N}$ . It is easy to obtain this result from Theorem 4.2.1, since from (4.8) we deduce that

$$\frac{q(t)}{A_+} \leq 1 \leq \frac{q(t)}{B_+} \quad \text{for every } t \in [0, 2\pi].$$

Under (4.8), from the point of view of resonance, the results in [46, 20] are optimal, in view of the counterexample produced in [7]. Thus, Theorem 4.2.1 seems to be optimal whenever we are able to control  $q(t)$  with an estimate like (4.8), essentially requiring, in this case, a nonresonance assumption. On the other hand, the mean conditions (4.5) and (4.6) do not ask that  $q(t)$  is controlled like in (4.8), allowing it to possibly cross some eigenvalues (cf. [8]) as in our case, being  $0 \leq q(t) \leq 2b$ .

We are now going to prove Theorem 4.2.1. As it was mentioned previously, we will have to overcome the difficulty to work with nonlinearities with atypical linear growth, since, in our concrete case, the nonlinearity grows linearly towards the function  $b(1 + \cos t)x$ , which vanishes at some times. For this reason, classic arguments in literature (like the ones in [14, 19, 20]) do not extend as they are to (4.1), because it is not possible to construct an admissible spiral which allows to control the dynamic of the solutions.

We will prove Theorem 4.2.1 by means of some preliminary lemmas. To this aim, it will be convenient to introduce the "norm" application defined as

$$\mathcal{N} : \Lambda \rightarrow \mathbb{R}, \quad \mathcal{N}(x, y) := bx^2 + y^2 - 2 \ln x,$$

being  $\Lambda$  the half plane with positive abscissa, i.e.  $\Lambda = \{(x, y) \in \mathbb{R}^2 : x > 0\}$ . Fixed a value  $c$  of the function  $\mathcal{N}(x, y)$ , we will denote the corresponding level curve by  $\gamma_c$ , i.e.,

$$\gamma_c = \{(x, y) \in \Lambda : \mathcal{N}(x, y) = c\}.$$

It is worth observing that the function  $\mathcal{N}(x, y)$  reaches its minimum in the point  $P_0 = (1/\sqrt{b}, 0)$ , where it takes the value  $1 - 2 \ln(1/\sqrt{b})$  (possibly negative for some values of the parameter  $b$ ). For values of the energy greater than  $1 - 2 \ln(1/\sqrt{b})$ , the level curves of  $\mathcal{N}$  turn around  $P_0$ , being the union of two symmetric arcs joining on the  $x$ -axis.

We will look at the solutions of (4.1) in the phase plane, taking thus into account the couple  $(u, u')$ . As already mentioned, we are interested in positive solutions, so that we will take into account the dynamics of the solutions in the right half-plane.

The first lemma ensures the global continuability of the solutions, i.e., shows that the maximal domain of every solution of (4.1) is  $\mathbb{R}_+$ .

**Lemma 4.2.1** *Let  $(u, u')$  be a solution of (4.1) (not necessarily periodic). Then*

$$\mathcal{N}(u(t), u'(t)) < +\infty \quad \text{for every } t \geq 0.$$

**Proof 57** Since

$$\lim_{x \rightarrow +\infty} \frac{bx^2}{bx^2 - 2 \ln x} = 1,$$

taking  $C > \max\{1, b\}$  there exists  $K_0 > 1$  such that

$$\frac{b}{2}(x^2 + y^2) \leq \frac{C}{2}(bx^2 + y^2 - 2 \ln x) \quad \text{for every } x \geq K_0, \quad y \in \mathbb{R}. \quad (4.9)$$

For every solution  $u(t)$  of (4.1), we define the function

$$U : I \rightarrow \mathbb{R}, \quad U(t) = \frac{\mathcal{N}(u(t), u'(t))}{2} = \frac{1}{2}(bu'^2(t) + u'^2(t) - 2 \ln u(t)),$$

where  $I$  is the maximal domain of  $u(t)$ . We are going to prove that  $I = \mathbb{R}_+$ .

Since

$$U'(t) = -bu(t)u'(t) \cos t,$$

for  $t \in I$  we have that

$$U'(t) \leq \frac{b(u^2(t) + u'^2(t))}{2},$$

from which it can be deduced that

$$U'(t) \leq CU(t) + C \ln K_0 \quad \text{for every } t \in I. \quad (4.10)$$

Indeed, if  $t \geq 0$  is such that  $u(t) \geq K_0$ , then (4.9) implies that  $U'(t) \leq CU(t)$ . On the contrary, if  $u(t) \leq K_0$ , we deduce that either  $u(t) \leq 1$ , and then  $U'(t) \leq CU(t)$ , or  $1 \leq u(t) \leq K_0$ , and thus (4.10) holds. Now, according to the *Gronwall-Bellman Lemma*, the result is proven.

As it was mentioned in the previous discussion, equations like (4.1) do not admit the existence of an admissible spiral controlling the solutions. However, the following result ensures that (4.1) has the “property of elasticity”, at least locally. Roughly speaking, this means that if there is a time when the norm of the solution is large enough, then, for every preceding time instant, the solution had to be large (in norm). Precisely, we have the following.

**Lemma 4.2.2** *Let  $\rho_0 > 0$  be sufficiently large. Then, there exists  $R_1 > \rho_0$  such that, for every solution  $(u, u')$  of (4.1) satisfying*

$$\mathcal{N}(u(t_1), u'(t_1)) \geq R_1$$

for some  $t_1 > 0$ , it holds

$$\mathcal{N}(u(t), u'(t)) \geq \rho_0 \quad \text{for every } t \in [0, t_1].$$

**Proof 58** We first observe that there exists a constant  $M > 0$  such that

$$\frac{b(x^2 + y^2)}{\mathcal{N}(x, y)} < M, \quad (4.11)$$

for every  $(x, y) \in \Lambda$ . Now, choosing  $\rho_0 > 0$  sufficiently large, there exist  $u_0^- < 1 < u_0^+$  such that

$$\gamma_{\rho_0} = \text{Graph}(F_0) \cup \text{Graph}(-F_0),$$

where  $F_0 : \mathbb{R}^+ \rightarrow \mathbb{R}$  is a function such that  $F_0(u_0^-) = F_0(u_0^+) = 0$ , having constant sign on  $(u_0^-, u_0^+)$ .

Let us fix  $L_1$  satisfying

$$2L_1 \geq \max_{x \in [u_0^-, u_0^+]} bx^2 - 2 \ln x + 2\rho_0,$$

and consider the set of the couples  $(x, y) \in \gamma_{2L_1}$ : explicitly,

$$\gamma_{2L_1} = \left\{ (x, y) \in \Lambda : y = \pm \sqrt{2L_1 - (bx^2 - 2 \ln x)} \right\}.$$

Thus, there exist  $u_1^- < u_0^- < u_0^+ < u_1^+$  such that, similarly as before,

$$\gamma_{2L_1} = \text{Graph}(F_1) \cup \text{Graph}(-F_1),$$

where  $F_1 : \mathbb{R}^+ \rightarrow \mathbb{R}$  is defined by  $F_1(x) = \sqrt{2L_1 - (bx^2 - 2 \ln x)}$  (and consequently vanishes in  $u_1^-, u_1^+$ ). On the other hand, we take  $L_2 > e^{2\pi M} L_1$ , and consider the level

curve  $\gamma_{2L_2}$ , which is explicitly given by

$$\gamma_{2L_2} = \left\{ (x, y) \in \Lambda : y = \pm \sqrt{2L_2 - (bx^2 - 2 \ln x)} \right\}.$$

Finally, we fix  $R_1 > 2L_2$ , so that

$$\gamma_{2L_2} \subset \{(x, y) \in \Lambda : \mathcal{N}(x, y) \leq R_1\}.$$

Assume that there exists  $u(t)$  solving (4.1) such that  $\mathcal{N}(u(t_1), u'(t_1)) \geq R_1$ , but there is  $t_* \in [0, t_1)$  such that  $\mathcal{N}(u(t_*), u'(t_*)) \leq \rho_0$ . By continuity, we can assume that there exist  $t_* < t^*$  such that  $(u(t_*), u'(t_*)) \in \gamma_{2L_1}$  and  $(u(t^*), u'(t^*)) \in \gamma_{2L_2}$ ; setting, as in Lemma 4.2.1,  $U(t) = \mathcal{N}(u(t), u'(t))/2$ , this explicitly means that

$$L_1 < U(t) < L_2 \quad \text{for every } t \in (t_*, t^*), \quad U(t_*) = L_1, \quad U(t^*) = L_2. \quad (4.12)$$

According to (4.11) and (4.12), from the definition of  $U(t)$  we deduce that

$$U'(t) \leq MU(t), \quad \text{for every } t \in [t_*, t^*],$$

which implies, thanks to the Gronwall-Bellman Lemma, that

$$U(t) \leq e^{2\pi M} L_1 \quad \text{for every } t \in [t_*, t^*].$$

This, however, contradicts (4.12) in view of the definition of  $L_2$ .

Now, intuitively speaking, we will prove that either the solutions of (4.1) have the global elasticity property, or their norm in the instant  $t = 2\pi$  is lower than in the initial one. This property is useful, and it is similar to the one introduced in [19].

**Lemma 4.2.3** *Let  $\rho_0 > 0$  be sufficiently large. Then, there exists  $R_2 > \rho_0$  such that, for every solution  $u(t)$  of (4.1) fulfilling*

$$\max_{t \in [0, 2\pi]} \mathcal{N}(u(t), u'(t)) \geq R_2, \quad (4.13)$$

*it is either*

$$\mathcal{N}(u(t), u'(t)) \geq \rho_0 \quad \text{for every } t \in [0, 2\pi], \quad (4.14)$$

*or*

$$\mathcal{N}(u(0), u'(0)) > \mathcal{N}(u(2\pi), u'(2\pi)). \quad (4.15)$$

**Proof 59** Let us take  $R_1$  as in the statement of Lemma 4.2.2, for the fixed  $\rho_0$ . In the same way, we apply again Lemma 4.2.2, this time with  $R_1$  playing the role of  $\rho_0$ , finding the corresponding  $R_2$  for which the statement holds.

Assume now that there exists a solution  $u(t)$  of (4.1) satisfying (4.13), for which it is

$$\mathcal{N}(u(0), u'(0)) \leq \mathcal{N}(u(2\pi), u'(2\pi)). \quad (4.16)$$

Since there exists  $t_2 \in [0, 2\pi]$  such that  $\mathcal{N}(u(t_2), u'(t_2)) \geq R_2$ , Lemma 4.2.2 implies that  $\mathcal{N}(u(0), u'(0)) \geq R_1$ , so that, in view of (4.16),  $\mathcal{N}(u(2\pi), u'(2\pi)) \geq R_1$ . Consequently, using again Lemma 4.2.2, we obtain that  $\mathcal{N}(u(t), u'(t)) \geq \rho_0$  for  $t \in [0, 2\pi]$ .

We are now able to show that an adaptation of the arguments in [14, 20] to our equation allows to prove that the global elasticity property cannot be fulfilled for solutions of (4.1) with large norm which perform an integer number of revolutions when  $t$  goes from 0 to  $2\pi$ .

**Lemma 4.2.4** *Under the hypotheses of Theorem 4.2.1, there exists  $R_2 > 0$  such that,*

if  $u(t)$  is a solution of (4.1) which satisfies

$$\max_{t \in [0, 2\pi]} \mathcal{N}(u(t), u'(t)) \geq R_2$$

and  $(u(t), u'(t))$  performs an integer number of turns around  $(1, 0)$  in the time interval  $[0, 2\pi]$ , then (4.15) holds.

**Proof 60** In view of (4.5), (4.6), there exists a positive number  $\delta$  such that

$$\begin{aligned} \frac{1}{2\pi} \int_0^{2\pi} \min \left\{ \frac{b(1 + \cos t) - \delta}{B_+}, 1 \right\} dt &> \frac{n}{2\sqrt{B_+}}, \\ \frac{1}{2\pi} \int_0^{2\pi} \max \left\{ \frac{b(1 + \cos t) + \delta}{A_+}, 1 \right\} dt &< \frac{n+1}{2\sqrt{A_+}}. \end{aligned}$$

In correspondence of  $\delta$ , we can find  $K_\delta > 0$  such that

$$\begin{aligned} [b(1 + \cos t) - \delta](x - 1)^2 - K_\delta &< \left[ b(1 + \cos t)x - \frac{1}{x} \right] (x - 1) \\ &< [b(1 + \cos t) + \delta](x - 1)^2 + K_\delta \quad \text{for every } x \in [1, +\infty), \quad t \geq 0. \end{aligned} \quad (4.17)$$

Moreover, we choose  $\rho_1$  and  $B'_+$  large, in such a way that the following relations hold:

$$\left( \frac{1}{\sqrt{B_+}} + \frac{1}{\sqrt{B'_+}} \right)^{-1} \left[ \frac{1}{2\pi} \int_0^{2\pi} \min \left\{ \frac{b(1 + \cos t) - \delta}{B_+}, 1 \right\} dt - \frac{K_\delta}{\rho_1} \right] > \frac{n}{2}, \quad (4.18)$$

$$\sqrt{A_+} \left[ \frac{1}{2\pi} \int_0^{2\pi} \max \left\{ \frac{b(1 + \cos t) + \delta}{A_+}, 1 \right\} dt + \frac{K_\delta}{\rho_1} \right] < \frac{n+1}{2}. \quad (4.19)$$

In order to perform the estimates leading to the result, we first fix  $\rho_0 > 0$  sufficiently large and apply Lemma 1.15 in order to find  $R_2 > \rho_0$  such that the statement therein holds. Then, we fix a solution  $u(t)$  of (4.1) satisfying (4.13) and such that, in the phase plane, the couple  $(u(t), u'(t))$  performs an integer number of revolutions around  $(1, 0)$  -

say  $k \in \mathbb{N}$  - during the time interval  $[0, 2\pi]$ .

Thus, assume by contradiction that (4.15) is not satisfied; then, in view of Lemma 1.15,  $u(t)$  fulfills (4.14). We are now going to estimate the time needed by  $(u(t), u'(t))$  to rotate  $k$  times around the point  $(1, 0)$ , by dividing the half-plane  $\Lambda$  in vertical strips and analyzing the behavior of the solution in each strip, following the procedure used in [20].

As a first step, we perform our estimates in the strip  $\{x > 1\}$ . Passing to modified polar coordinates around  $(1, 0)$  by writing

$$-\mu u = -\mu + \rho \sin \vartheta, \quad u' = \rho \cos \vartheta, \quad (4.20)$$

where  $\mu > 0$ , we obtain

$$-\dot{\vartheta}(t) = \mu \frac{u'^2(t) - u'(t)(u(t) - 1)}{\mu^2(u(t) - 1)^2 + u'^2(t)} \quad \text{for every } t \in [0, 2\pi]. \quad (4.21)$$

Setting

$$J_+ = \{t \in [0, 2\pi] : u(t) \geq 1\}, \quad J_- = \{t \in [0, 2\pi] : u(t) < 1\},$$

in view of the properties of the modified rotation numbers (see for instance [13]) we have that

$$2\pi \cdot \frac{k}{2} = - \int_{J_+} \dot{\vartheta}(t) dt.$$

Consequently, in view of (4.17),

$$\begin{aligned} \frac{k}{2} &\geq \frac{\mu}{2\pi} \int_{J_+} \frac{u'^2 + [b(1 + \cos t) - \delta](u - 1)^2}{\mu^2(u - 1)^2 + u'^2} dt - \frac{\mu}{2\pi} \int_{J_+} \frac{K_\delta}{\mu^2(u - 1)^2 + u'^2} dt \\ &\geq \frac{\mu}{2\pi} \int_{J_+} \frac{\min \left\{ \frac{b(1 + \cos t) - \delta}{\mu^2}, 1 \right\} (u - 1)^2 + (u'/\mu)^2}{(u - 1)^2 + (u'/\mu)^2} dt - \frac{\mu}{2\pi} \int_{J_+} \frac{K_\delta}{\mu^2(u - 1)^2 + u'^2} dt. \end{aligned}$$



Taking into account that the function

$$\Psi : [0, +\infty) \rightarrow \mathbb{R}, \quad \Psi(y) = \frac{\alpha + y}{\beta + y} \quad (4.22)$$

is non-decreasing whenever  $\alpha \leq \beta$ , choosing  $\mu = \sqrt{B_+}$ ,  $\alpha = \min \left\{ \frac{b(1+\cos t) - \delta}{\mu^2}, 1 \right\} (u-1)^2$ ,  $\beta = (u-1)^2$  and  $y = (u'/\mu)^2$  we have

$$\frac{k}{2} \geq \frac{\sqrt{B_+}}{2\pi} \int_{J_+} \min \left\{ \frac{b(1+\cos t) - \delta}{B_+}, 1 \right\} dt - \frac{\sqrt{B_+}}{2\pi} \int_{J_+} \frac{K_\delta}{B_+(u-1)^2 + u'^2} dt. \quad (4.23)$$

Without loss of generality, we can assume (up to enlarging  $\rho_0$ ) that  $R_2$  is sufficiently large, so that

$$B_+(u-1)^2 + u'^2 \geq \rho_1, \quad \text{for every } t \in J_+.$$

Therefore, (4.23) implies

$$\frac{k}{2\sqrt{B_+}} \geq \frac{1}{2\pi} \int_{J_+} \min \left\{ \frac{b(1+\cos t) - \delta}{B_+}, 1 \right\} dt - \frac{K_\delta}{\rho_1}. \quad (4.24)$$

We now pass to compute the time spent by  $(u(t), u'(t))$  to perform  $k/2$  revolutions on the “left” half phase plane, i.e. when  $u \in (0, 1]$ . Preliminarily, we fix

$$\tilde{\eta} < \frac{2\pi}{\sqrt{B'_+}}, \quad K = \left( \frac{2\pi}{\tilde{\eta}} \right)^2 \quad (4.25)$$

and observe that, since

$$\lim_{x \rightarrow 0^+} b(1+\cos t)x - \frac{1}{x} = -\infty,$$

there exists  $0 < d < 1$  such that

$$b(1+\cos t)x - \frac{1}{x} < -K \quad \text{for every } x \in (0, d]. \quad (4.26)$$

In this way it is possible to define both the sets

$$J_d^- = \{t \in J_- \mid u(t) \leq d\}, \quad J_d^+ = \{t \in J_- \mid d < u(t) < 1\}$$

and, correspondingly, the time instants  $t_1, t_2, t_3$  and  $t_4$  (as in [24, Figure 1]) such that, in the time  $t_4 - t_1$ , the couple  $(u(t), u'(t))$  performs half a turn in the “left” half phase plane ( $u \in (0, 1]$ ), and

$$u(t_1) = 1 = u(t_4), \quad u(t_2) = d = u(t_3), \quad (t_1, t_2) \cup (t_3, t_4) \subseteq J_d^+, \quad [t_2, t_3] \subseteq J_d^-.$$

$$u = 1 + \rho \cos \vartheta, \quad u' = \rho \sin \vartheta,$$

we arrive at

$$-\dot{\vartheta}(t) = \frac{u'^2(t) - u'(t)(u(t) - 1)}{(u(t) - 1)^2 + u'^2(t)}. \quad (4.27)$$

In view of (4.26), we deduce that

$$-\dot{\vartheta}(t) > K \cos^2 \vartheta(t) + \sin^2 \vartheta(t), \quad t \in [t_2, t_3],$$

so that

$$\begin{aligned} t_3 - t_2 &= \int_{\vartheta(t_3)}^{\vartheta(t_2)} \frac{ds}{K \cos^2 s + \sin^2 s} \\ &= \frac{1}{\sqrt{K}} \left[ \arctan \left( \frac{\tan \vartheta(t_2)}{K} \right) - \arctan \left( \frac{\tan \vartheta(t_3)}{K} \right) \right] \\ &\leq \frac{\pi}{2\sqrt{K}}. \end{aligned}$$

According to (4.25), it follows that  $t_3 - t_2 < \tilde{\eta}/2$ ; repeating the argument for every

revolution made by  $(u, u')$  around  $(1, 0)$  yields

$$\text{meas}(J_d^-) < \frac{k}{4} \tilde{\eta}. \quad (4.28)$$

In order to compute  $t_2 - t_1$ , we observe that, thanks to (4.27), it holds

$$-\dot{\vartheta}(t) \geq \frac{u'^2(t) - \tilde{C}|1-d|}{(1-d)^2 + u'^2(t)} \quad \text{for every } t \in [t_1, t_2],$$

where  $\tilde{C} = \max_{x \in [d, 1]} 2bx + 1/x$ . Again, we assume that  $\rho_0$  is large enough, so that  $-\dot{\vartheta}(t) > 1/2$  on  $[t_1, t_2]$ , and  $t_2 - t_1 < \tilde{\eta}/4$ . Analogously, one can prove that  $t_4 - t_3 < \tilde{\eta}/4$ , having thus that

$$\text{meas}(J_d^+) < \frac{k}{2} \frac{\tilde{\eta}}{2}. \quad (4.29)$$

Thus, in view of (4.25) and (4.28), we deduce that

$$\text{meas}(J_-) = \text{meas}(J_d^+) + \text{meas}(J_d^-) < \frac{k}{2} \tilde{\eta} < k \frac{\pi}{\sqrt{B'_+}},$$

from which

$$\frac{k}{2\sqrt{B'_+}} > \frac{1}{2\pi} \text{meas}(J_-).$$

Summing up, from (4.24) we have

$$\frac{k}{2} \left( \frac{1}{\sqrt{B_+}} + \frac{1}{\sqrt{B'_+}} \right) \geq \frac{1}{2\pi} \int_0^{2\pi} \min \left\{ \frac{b(1 + \cos t) - \delta}{B_+}, 1 \right\} dt - \frac{K_\delta}{\rho_1}. \quad (4.30)$$

On the other hand, reasoning on (4.21) with a similar argument and taking (4.17) into account, we have

$$\frac{k}{2} \leq \frac{\mu}{2\pi} \int_{J_+} \frac{\max\{\frac{b(1+\cos t)+\delta}{A_+}, 1\}(u-1)^2 + (u'/\mu)^2}{(u-1)^2 + (u'/\mu)^2} dt + \frac{\mu}{2\pi} \int_{J_+} \frac{K_\delta}{\mu^2(u-1)^2 + u'^2} dt.$$

Since the function  $\Psi$  defined in (4.22) is non-increasing whenever  $\alpha \geq \beta$ , choosing  $\mu = \sqrt{A_+}$  and taking  $\alpha = \max \left\{ \frac{b(1+\cos t)+\delta}{A_+}, 1 \right\} (u-1)^2$  and  $\beta = (u-1)^2$ , we obtain

$$\frac{k}{2} \leq \frac{\sqrt{A_+}}{2\pi} \int_{J_+} \max \left\{ \frac{b(1+\cos t)+\delta}{A_+}, 1 \right\} dt + \frac{\sqrt{A_+}}{2\pi} \int_{J_+} \frac{K_\delta}{A_+(u-1)^2 + u'^2} dt.$$

Again, we can assume  $\rho_0$  (and thus  $R_2$ ) so large that

$$\sqrt{A_+}(u(t)-1)^2 + \dot{u}(t)^2 \geq \rho_1, \quad t \in J_+.$$

Hence,

$$\frac{k}{2\sqrt{A_+}} \leq \frac{1}{2\pi} \int_{J_+} \max \left\{ \frac{b(1+\cos t)+\delta}{A_+}, 1 \right\} dt + \frac{K_\delta}{\rho_1}. \quad (4.31)$$

We are now able to conclude the proof. Assume first that  $u(t)-1$  has at most  $2n$  zeros. Then  $k \leq n$ , but this contradicts (4.18) and (4.30). On the contrary, if  $u(t)-1$  has at least  $2n+2$  zeros, since  $k \in \mathbb{N}$  it has to be  $k \geq n+1$ . However, this contradicts (4.19) and (4.31). The proof is completed.

**Remark 4.2.4** In [8], the relationships between conditions (4.5) and (4.6) and the rotation number of “large” solutions of a first order planar system were highlighted. This perfectly agrees with what we have seen in the proof which has just been performed; indeed, conditions (4.5) and (4.6) force the solutions of the Cauchy problems associated with (4.1) not to perform an integer number of turns around  $(1, 0)$  in the time interval  $[0, 2\pi]$ . Thus, they turn to be hypotheses on the number of rotations made by the solutions of equation (4.1) in the phase plane.

Using the previous results, a basic application of the Poincaré-Bohl Theorem allows to prove Theorem 4.2.1.

**Proof 61 (Proof of Theorem 4.2.1)** Let us take  $R_2$  sufficiently large satisfying Lemma 4.2.4

and set  $B = \{(x, y) \in \Lambda : \mathcal{N}(x, y) \leq R_2\}$ . In view of Lemma 4.2.1, the Poincaré map

$$P : B \rightarrow \mathbb{R}^2, \quad P(x_0, y_0) = (u(2\pi), u'(2\pi)),$$

where  $(u(t), u'(t))$  is the unique solution of the problem

$$u'' + b(1 + \cos t)u - \frac{1}{u} = 0, \quad u(0) = x_0 > 0, \quad u'(0) = y_0,$$

is well defined. Moreover, if  $(x_0, y_0)$  is a fixed point of  $P$ , then it is  $(u(0), u'(0)) = (u(2\pi), u'(2\pi))$ , i.e.,  $u(t)$  is a periodic solution of (4.1). Therefore, to get the conclusion it is sufficient to prove that  $P$  has a fixed point. However, if we denote by  $\tau_1$  the unitary right translation in the plane  $(u, u')$ , the map  $\Phi := \tau_{-1} \circ P \circ \tau_1 : \tau_{-1}(B) \rightarrow \mathbb{R}^2$  satisfies all the hypotheses of the Poincaré-Bohl fixed point theorem, since  $0 \in \tau_{-1}(B)$  and  $\Phi(z) \neq \lambda z$  for every  $\lambda > 1$ , in view of Lemma 4.2.4. Consequently,  $P$  has a fixed point and the statement is proved.

## 4.2.2 Main result

In order to prove our Theorem it will be convenient, for any  $n \in \mathbb{N}$ , to define the absolutely continuous functions  $F_n, G_n : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}$  by

$$F_n(b, x) = \frac{1}{2\pi} \int_0^{2\pi} \min \left\{ \frac{b(1 + \cos t)}{\sqrt{x}}, \sqrt{x} \right\} dt - \frac{n}{2},$$

$$G_n(b, x) = \frac{1}{2\pi} \int_0^{2\pi} \max \left\{ \frac{b(1 + \cos t)}{\sqrt{x}}, \sqrt{x} \right\} dt - \frac{n+1}{2}.$$

Both functions are non-decreasing with respect to the variable  $b$ . Moreover, if there exists  $n \in \mathbb{N}$  such that  $\inf_{x>0} G_n(b, x) < 0$  and  $\sup_{x>0} F_n(b, x) > 0$ , then Theorem 4.2.1 implies that (4.1) has at least one periodic solution. Therefore, we have the following

proposition.

**Proposition 4.2.1** *Assume that there exists  $n \in \mathbb{N}$  such that*

$$b \in \left( \inf \left\{ b > 0 : \sup_{x>0} F_n(b, x) > 0 \right\}, \sup \left\{ b > 0 : \inf_{x>0} G_n(b, x) < 0 \right\} \right). \quad (4.32)$$

*Then, (4.1) has at least one  $2\pi$ -periodic solution.*

Let us first observe that, in view of the continuity and the monotonicity of the functions  $F_n, G_n$  in the variable  $b$ , there exist  $b_0^n$  and  $b_1^n$  such that

$$\left\{ b > 0 : \sup_{x>0} F_n(b, x) > 0 \right\} = (b_0^n, +\infty),$$

and

$$\left\{ b > 0 : \inf_{x>0} G_n(b, x) < 0 \right\} = (0, b_1^n).$$

The point is to prove that these two intervals contain common points, i.e.,  $b_0^n < b_1^n$ . We will show this in the case when  $n = 0$  and  $n = 1$ , and the estimates performed in this last case will allow to achieve the new result consisting in the following Theorem.

**Theorem 4.2.2** *If  $b \in [0.4705, 0.59165]$ , then (4.1) has at least one  $2\pi$ -periodic solution.*

In particular, a gross estimation of the interval in (4.32) would lead to prove existence for

$$b \in \left( \frac{n^2}{2}, \frac{(n+1)^2}{4} \left( \frac{\pi}{1+\pi} \right)^2 \right). \quad (4.33)$$

Indeed, setting  $B_+ = 2b$ , since  $b > n^2/2$  we have

$$F_n(b, B_+) = \frac{1}{2\pi} \sqrt{b} \int_0^{2\pi} \min \left\{ \frac{1 + \cos t}{\sqrt{2}}, \sqrt{2} \right\} dt - \frac{n}{2} = \sqrt{\frac{b}{2}} - \frac{n}{2} > 0.$$

On the other hand, we choose

$$A_+ = \frac{4b^2}{(n+1)^2} \left( \frac{\pi+1}{\pi} \right)^2,$$

so that, since  $b < \frac{1}{4}(n+1)^2(\pi/(1+\pi))^2$ ,

$$\begin{aligned} G_n(b, A_+) &= \frac{1}{2\pi} \int_0^{2\pi} \max \left\{ \frac{(n+1)\pi(1+\cos t)}{2(\pi+1)}, \frac{2b}{n+1} \frac{\pi+1}{\pi} \right\} dt - \frac{n+1}{2} \\ &< \frac{1}{2\pi} \frac{(n+1)\pi}{2(\pi+1)} \int_0^{2\pi} \max\{1+\cos t, 1\} dt - \frac{n+1}{2} = 0. \end{aligned}$$

Now, in order for the interval in (4.33) to be nonempty, we need

$$\frac{n^2}{2} < \frac{(n+1)^2}{4} \left( \frac{\pi}{1+\pi} \right)^2,$$

which approximately requires  $n < 1.1$ . Since  $n \in \mathbb{N}$ , we can take either  $n = 0$  or  $n = 1$ , so that the  $2\pi$ -periodic solvability of (4.1) is guaranteed whenever

$$b \in \left( 0, \frac{1}{4} \left( \frac{\pi}{1+\pi} \right)^2 \right) \cup \left( \frac{1}{2}, \left( \frac{\pi}{1+\pi} \right)^2 \right).$$

However, taking into account that  $F_1, G_1$  are non-decreasing, we can use a numerical approach to estimate the interval in (4.32) and try to compute approximately, by means of a numerical software, its endpoints, obtaining

$$\sup_{x>0} F_1(0.4705, x) > 0 \quad (\text{but } \sup_{x>0} F_1(0.47, x) < 0)$$

and

$$\inf_{x>0} G_1(0.59165, x) < 0 \quad (\text{but } \inf_{x>0} G_1(0.591, x) > 0),$$

whence the statement of Theorem 4.2.2.

---

# Conclusion

Usando técnicas básicas del análisis no lineal (teoría de sub y super soluciones y teoremas de punto fijo), los principales logros que hemos obtenido en la Tesis giran en torno a la existencia de soluciones periódicas y son:

- Encontrar criterios de existencia en ecuaciones con singularidades en el término de fricción, aplicando los resultados para estudiar modelos de cavitación determinados por las ecuaciones de Rayleigh-Plesset.
- Dar un contraejemplo para probar, a pesar de lo que uno puede esperar (ver [28, Open Problem 4.1]), que los resultados de Lazer y Solimini obtenidos en el caso repulsivo no pueden extenderse al caso atractivo.
- Establecer una relación unívoca entre el orden de la singularidad y la regularidad de los coeficientes en las ecuaciones de Lazer y Solimini de tipo atractivo, garantizando existencia de soluciones.
- Determinar relaciones entre las singularidades y las no-linealidades presentadas en ecuaciones con singularidades atractivas de tipo Liénard, obteniendo aplicaciones para las ecuaciones de Rayleigh-Plesset.
- Obtener resultados de existencia de soluciones  $2\pi$ -periódicas para la ecuación de Brillouin Beam fuera del rango usual donde se conjetura la existencia mediante simulaciones numéricas.



By the use of basic tools of Nonlinear Analysis (lower and upper solutions theory and fixed point theorems), the main results obtained in this Thesis revolve around of the existence of periodic solutions and they are:

- To find existence criteria to singular differential equations with singular friction like term, applying the results to the study of cavitation models which are determined by the known Rayleigh-Plesset equations.
- To give a counter-example in order to prove that, despite of all expectations (see [28, Open Problem 4.1]), one cannot extend in a standard way the results of Lazer and Solimini obtained to the repulsive case.
- To establish an unanimous relation between the order of singularity and the smoothness of the coefficients of Lazer and Solimini equations with attractive singularity, guaranteeing existence of solutions.
- To determinate relations between the singularities and the nonlinearity presented in singular equation of Liénard type, obtaining some applications to the Rayleigh-Plesset equations.
- To obtain results of existence of  $2\pi$ -periodic solutions to the Brillouin Beam equation outside of the usual parameter range where it is conjectured the existence of solutions by mean of numerical simulations.

---

# Bibliography

- [1] A. Adje, Sur et sous- Solutions généralisées te problèmes aux limites du second ordre, **Bull. Soc. Math. Bel. Sér B** 42 (1990), 347–368. [34](#)
- [2] J. Appell, The superposition operator in function spaces- A survey, **Exposition Math.** 6 (1988), 209–270. [34](#)
- [3] C. Bereanu, D. Gheorghe, M. Zamora, Periodic solutions for singular perturbations of the singular  $\phi$ -Laplacian operator, **Communications in Contemporary Mathematics** (2013) 1250063 doi:10.1142/s0219199712500630. [5](#)
- [4] C. Bereanu, D. Gheorghe, M. Zamora, Non-resonant boundary value problems with singular  $\phi$ -Laplacian operators, **Nonlinear Differ. Eqn. Appl.** 20 (2013), 1365–1377. [5](#)
- [5] V. Bevc, J. L. Palmer, C. Süsskind, On the design of the transition region of axisymmetric, magnetically focused beam valves, **J. British Inst. Radio Eng** 18 (1958), 696–708. [106](#), [108](#)
- [6] D. Bonheure, C. De Coster, Forced singular oscillators and the method of upper and lower solutions, **Topol. Methods Nonlinear Anal.** 22 (2003), 297–317. [4](#)
- [7] D. Bonheure, C. Fabry, D. Smets, Periodic solutions of forced isochronous oscillators at resonance. **Discrete Contin. Dyn. Syst.** 8 (2002), 907–930. [4](#), [30](#), [31](#), [112](#)

- [8] A. Boscaggin, M. Garrione, Resonance and rotation numbers for planar Hamiltonian systems: multiplicity results via the Poincaré-Birkhoff theorem, **Nonlinear Anal.** 74 (2011), 4166–4185. [112](#), [123](#)
- [9] M. P. Brenner, S. Hilgenfeldt, D. Lohse, Single bubble sonoluminescence. **Reviews of Modern Physics** 74, No. 2 (2002), 425–484. [70](#)
- [10] H. Broer, M. Levi, Geometrical aspects of stability theory for Hill’s equations, **Arch. Rational Mech. Anal.** 131 (1995), 225–240. [108](#)
- [11] C. De Coster, P. Habets, The lower and upper solutions method for boundary value problems, in Handbook of Differential Equations, ODE, Elsevier, 2004, pp. 69–160. [4](#), [33](#)
- [12] L. A. Crum, Sonochemistry and sonoluminescence. Springer, 1999. [70](#)
- [13] C. Fabry, Periodic solutions of the equation  $x'' + f(t, x) = 0$ , **Séminaire de Mathématique** 117 (1987), Louvain-la-Neuve. [109](#), [111](#), [119](#)
- [14] C. Fabry, P. Habets, Periodic solutions of second order differential equations with superlinear asymmetric nonlinearities, **Arch. Math.** 60 (1993), 266–276. [8](#), [111](#), [112](#), [117](#)
- [15] A. Fonda, Topological degree and generalized asymmetric oscillators, **Topological Methods in Nonlinear Anal.** 28 (2006), 171–188. [31](#)
- [16] A. Fonda, Periodic solutions of scalar second order differential equations with a singularity, **Mém. Cl. Sci. Acad. Roy. Belgique** (4) 8 (1993), 1–39. [4](#), [8](#)
- [17] A. Fonda, M. Garrione, A Landesman-Lazer type condition for asymptotically linear second order equations with a singularity, **Proc. Royal Soc. Edinburgh Sect. A** 142 (2012), 1263–1277. [31](#)

- [18] A. Fonda, R. Manásevich, F. Zanolin, Subharmonic solutions for some second-order differential equations with singularities, **SIAM J. Math. Anal.** 24 (1993), 1294–1311. [4](#), [8](#)
- [19] A. Fonda, A. Sfecci, A general method for the existence of periodic solutions of differential systems in the plane, **J. Differential Equations** 252 (2012), 1369–1391. [4](#), [31](#), [112](#), [116](#)
- [20] A. Fonda, R. Toader, Radially symmetric systems with a singularity and asymptotically linear growth, **Nonlinear Analysis** 74 (2011), 2485–2496. [110](#), [111](#), [112](#), [117](#), [119](#)
- [21] N. Forbat, A. Huaux, Détermination approchée et stabilité locale de la solution périodique d’une équation différentielle non linéaire, **Mém. Publ. Soc. Sci. Art. Lett. Hainaut** 76 (1962), 3–13. [54](#)
- [22] J. P. Franc, The Rayleigh-Plesset equation: A simple and powerful tool to understand various aspects of cavitation. In *Fluid Dynamics of Cavitation and Cavitating Turbopumps*, Springer Vienna, 2008. [70](#), [80](#)
- [23] S. Fučík, *Solvability of Nonlinear Equations and Boundary Value Problems*, Reider Boston, 1980. [8](#)
- [24] M. Garrione, M. Zamora, Periodic solutions of the Brillouin electron beam focusing equation (preprint). [4](#), [108](#), [121](#)
- [25] W.B. Gordon, Conservative dynamical systems involving strong forces, **Tans. Amer. Math. Soc.** 204 (1975), 113–135. [8](#)
- [26] P. Habets, L. Sanchez, Periodic solutions of some Liénard equations with singularities, **Proc. Amer. Math. Soc.** 109 (1990), 1135–1144. [4](#), [8](#), [31](#), [58](#)

- [27] R. Hakl, P. J. Torres, A combined variational-topological approach for dispersion-managed solitons in optical fiber, **Z. Angew. Math. Phys.** 62 (2011) 245–266. [4](#), [34](#)
- [28] R. Hakl, P. J. Torres, On periodic solutions of second-order differential equations with attractive-repulsive singularities, **J. Differential Equations** 248 (2010), 111-126. [58](#), [127](#), [128](#)
- [29] R. Hakl, P.J. Torres, Maximum and antimaximum principles for a second order differential operator with variable coefficients of indefinite sign, **Applied Mathematics and Computation** 217 (2011) 7599–7611. [4](#), [86](#), [87](#)
- [30] R. Hakl, P. J. Torres, M. Zamora, Periodic solutions of singular second order differential equations: the repulsive case, **Topological methods in nonlinear analysis** 39 (2012), 199-220. [4](#)
- [31] R. Hakl, P. J. Torres, M. Zamora, Periodic solutions of singular second order differential equations: upper and lower functions, **Nonlinear. Anal.** 74 (2011), 7078-7093. [4](#), [58](#)
- [32] R. Hakl, M. Zamora, Periodic solutions to the Liénard type equations with phase attractive singularities, **Boundary Value Problems** 2013:47 doi:10.1186/1687-2770-2013-47. [4](#)
- [33] R. Hakl, M. Zamora, On the open problems connected to the results of Lazer and Solimini, **Proc. Roy. Soc. Edinb., Sect. A. Math.** to appear. [4](#)
- [34] R. Hakl, M. Zamora Existence of a solution to a Dirichlet problem associated to second order equations with singularities: method of lower and upper functions, **Georgian Math. J.** to appear. [5](#)

- [35] A. Huaux, Sur l' existence d'une solution périodique de l' équation différentielle non linéaire  $x'' + 0,2x' + x/(1 - x) = (0,5) \cos \omega t$ , **Bull. Class. Sci. Acad. R. Belgique** 48 (1962), 494–504. [54](#)
- [36] A. M. Krasnosel'skii, J. Mawhin, Periodic solutions of equations with oscillating nonlinearities, **Math. Computer Modelling** 32 (2000) 1445–1455. [31](#)
- [37] E. M. Landesman, A. C. Lazer, Nonlinear perturbations of linear elliptic boundary value problems at resonance, **J. Math. Mech.** 19 (1970) 609–623. [31](#)
- [38] A. C. Lazer, On Schauder's fixed point theorem and forced second order nonlinear oscillations, **J. Math. Anal. Appl.** 21 (1968) 421–625. [56](#)
- [39] A. C. Lazer, D. E. Leach, Bounded perturbations of forced harmonic oscillators at resonance, **Ann. Mat. Pura Appl.** (4) 82, 49–68. [31](#)
- [40] A. C. Lazer, S. Solimini, On periodic solutions of nonlinear differential equations with singularities, **Proc. Amer. Math. Soc.** 99 (1987), 109–114. [4](#), [5](#), [8](#), [31](#), [52](#), [54](#), [55](#), [56](#), [57](#)
- [41] W. Magnus, S. Winkler, Hill's equation, corrected reprint of 1966 edition, Dover, New York, 1979. [108](#)
- [42] J. Mawhin, Landesman-Lazer conditions for boundary-value problems: a nonlinear version of resonance, **Boletín de la sociedad española de matemática aplicada** 16 (2000) 45–65. [31](#)
- [43] J. Mawhin, Resonance and nonlinearity: a survey, **Ukrainian Mathematical J.** 59 (2007), No.2, 197–214. [31](#)
- [44] J. Mawhin, Topological degree and boundary value problems for nonlinear differential equations. In Topological Methods for Ordinary Differential Equations (Mon-

- tecatini Terme, 1991), M. Furi and P. Zecca, Eds., Lecture Notes in Mathematics 1537, 74–142, Springer, Berlin, Germany, 1993. [4](#), [31](#), [35](#)
- [45] P. Omari, W. Ye, Necessary and sufficient conditions for the existence of periodic solutions of second-order ordinary differential equations with singular nonlinearities, **Differential Integral Equations** 8 (1995), 1843–1858. [4](#)
- [46] M. del Pino, R. Manásevich, A. Montero,  $T$ -periodic solutions for some second order differential equations with singularities, **Proc. Royal Soc. Edinburgh Sect. A** 120 (1992) 231–243. [4](#), [7](#), [8](#), [111](#), [112](#)
- [47] A. Prosperetti, Bubble dynamics: a review and some recent results. In Mechanics and physics of bubbles in liquids, L. van Wijngaarden ed., Kluwer, 1982. [70](#)
- [48] I. Rachunková, S. Staněk, M. Tvrđý, Solvability of Nonlinear Singular Problems for Ordinary Differential Equations, Contemporary Mathematics and Its Applications 5. New York, NY: Hindawi Publishing Corporation, 2008, 268 pp. [4](#), [33](#), [53](#), [58](#)
- [49] I. Rachunková, M. Tvrđý, I. Vrkoč, Existence of nonnegative and nonpositive solutions for second–order periodic boundary–value problems, **J. Differential Equations** 176 (2001), No. 2, 445–469. [4](#), [8](#), [53](#), [58](#)
- [50] J. Ren, Z. Cheng, S. Siegmund, Positive periodic solution for Brillouin electron beam focusing systems, **Discrete Cont. Dynamical Syst. B** 16 (2011), 385–392. [107](#)
- [51] H. Schaefer, Über die Methode der a priori Schranken, **Math. Ann.** 129 (1955), 415–416. [9](#)
- [52] D. R. Smart, Fixed point theorems, Cambridge Tracts in Mathematics, 66. London: Cambridge University Press (1974), 104 pp. [9](#)

- [53] P.J. Torres, Existence of one-signed periodic solutions of some second order differential equations via a Krasnoselskii fixed point theorem, **J. Differential Equations** 190 (2003), 643–662. 8, 107
- [54] P.J. Torres, Existence and stability of periodic solutions for second-order semilinear differential equations with a singular nonlinearity, **Proc. Royal Soc. Edinburgh Sect. A** 137A (2007) 195–201. 8
- [55] P.J. Torres, Weak singularities may help periodic solutions to exist, **J. Differential Equations** 232 (2007) 277-284. 8
- [56] P.J. Torres, Existence and stability of periodic solutions of a duffing equation by using a new maximum principle, **Mediterr. J. Math.** 1 (2004) 479-486. 34
- [57] P. J. Torres, A. J. Ureña, M. Zamora, Periodic and quasi-periodic motions of a relativistic particle under a central force field, **Bulletin of the London Mathematical Society** (2013) 45(1): 140-152 doi: 10.1112/blms/bds076 5
- [58] R. F. Young, Cavitation. Imperial College Press, 1999. 70
- [59] Z. Wang, Periodic solutions of the second-order differential equations with singularity, **Nonlinear Anal.** 58 (2004) 319–331. 8
- [60] Z. Wang, Existence and multiplicity of periodic solutions for the Duffing equation with singularity, **Proc. Roy. Soc. Edinburgh** 137A (2007), 1–21. 8
- [61] P. Yan, M. Zhang, Higher order nonresonance for differential equations with singularities, **Math. Methods Appl. Sci.** 26 (2003), 1067–1074. 4
- [62] M. Zamora, New periodic and quasi-periodic motions of a relativistic particle under a planar central force field with applications to scalar boundary periodic problems,



**Electronic J. Qualitative Theory of Differential Equations** No. 31 (2013), 1-16; <http://www.math.u-szeged.hu/ejqtde/> 5

- [63] M. Zamora, On a periodically forced Liénard differential equation with singular  $\phi$ -Laplacian, **Bulletin Mathématique de la Société des Sciences Mathématiques de Roumanie**. to appear 5
- [64] E. Zeidler, Nonlinear functional analysis and its applications. Volume I: Fixed-point theorems. Springer-Verlag, 1985. 9
- [65] M. Zhang, Periodic solutions of Liénard equations with singular forces of repulsive type, **J. Math. Anal. Appl** 203 (1996), 254–269. 8
- [66] M. Zhang, A relationship between the periodic and the Dirichlet BVPs of singular differential equations, **Proc. Roy. Soc. Edinburgh Sect. A** 128 (1998), 1099–1114. 8, 107, 111
- [67] M. Zhang, Periodic solutions of equations of Emarkov-Pinney type, **Adv. Non-linear Studies** 6 (2006), 57–67. 8, 107